

(Department of Public Health, Health Management and Policy, Nara Medical University School of Medicine) for the data analyses, and the Tokyo Metropolitan Government for providing data on the cedar pollen levels.

References

- Awaya A, Murayama K. 2012. Positive correlation between Japanese cedar pollen numbers and the development of Kawasaki disease. *Open Allergy J.* 5:1–10.
- Bell DS, Kahn Jr CE. 1996. Health status assessment via the World Wide Web. *Proc AMIA Annu Fall Symp.* 338–342.
- Berger U, Karatzas K, Jaeger S, Voukantsis D, Sofiev M, Brandt O, Zuberbier T, Bergmann KC. 2013. Personalized pollen-related symptom-forecast information services for allergic rhinitis patients in Europe. *Allergy.* 68:963–965.
- Cook C, Heath F, Thompson RL. 2000. A meta-analysis of response rates in web- or internet-based surveys. *Educational Psychological Meas.* 60:821–836.
- Crystal-Peters J, Crown WH, Goetzel RZ, Schutt DC. 2000. The cost of productivity losses associated with allergic rhinitis. *Am J Manag Care.* 6:373–378.
- D'Amato G, Cecchi L, Bonini S, Nunes C, Annesi-Maesano I, Behrendt H, Liccardi G, Popv T, van Cauwenberge P. 2007. Allergenic pollen and pollen allergy in Europe. *Allergy.* 62:976–990.
- D'Amato G, Liccardi G, D'Amato M, Holgate S. 2005. Environmental risk factors and allergic bronchial asthma. *Clin Exp Allergy.* 35:1113–1124.
- Dejima K, Saito Y, Shoji H. 1992. A relationship between the annual day of onset of Japanese cedar pollinosis and pollen dispersion. *Arerugi.* 41:1405–1412. Japanese.
- Ekman A, Dickman PW, Klint A, Weiderpass E, Litton JE. 2006. Feasibility of using web-based questionnaires in large population-based epidemiological studies. *Eur J Epidemiol.* 21:103–111.
- Futamura M, Ohya Y, Akashi M, Adachi Y, Odajima H, Akiyama K, Akasawa A. 2011. Age-related prevalence of allergic diseases in Tokyo schoolchildren. *Allergol Int.* 60:509–515.
- Gotoh M, Suzuki H, Okubo K. 2011. Delay of onset of symptoms of Japanese cedar pollinosis by treatment with a leukotriene receptor antagonist. *Allergol Int.* 60:483–489.
- Kaneko Y, Motohashi Y, Nakamura H, Endo T, Eboshida A. 2005. Increasing prevalence of Japanese cedar pollinosis: a meta-regression analysis. *Int Arch Allergy Immunol.* 136:365–371.
- Kawashima S, Takahashi Y. 1999. An improved simulation of mesoscale dispersion of airborne cedar pollen using a flowering-time map. *Grana.* 38:316–324.
- Kawashima S, Takahashi Y, Sahashi N. 1998. Forecast of the first day of Japanese cedar pollen release using variable air temperature patterns. *Arerugi.* 47:649–657. Japanese.
- Long AA. 2007. Findings from a 1000-patient internet-based survey assessing the impact of morning symptoms on individuals with allergic rhinitis. *Clin Ther.* 29:342–351.
- Medek DE, Kljakovic M, Fox I, Pretty DG, Prebble M. 2012. Hay fever in a changing climate: linking an internet-based diary with environmental data. *Ecohealth.* 9:440–447.
- Okamoto Y, Horiguchi S, Yamamoto H, Yonekura S, Hanazawa T. 2009. Present situation of cedar pollinosis in Japan and its immune responses. *Allergol Int.* 58:155–162.
- Okuda M. 2003. Epidemiology of Japanese cedar pollinosis throughout Japan. *Ann Allergy Asthma Immunol.* 91:288–296.
- Okubo K, Gotoh M, Shimada K, Ritsu M, Okuda M, Crawford B. 2005. Fexofenadine improves the quality of life and work productivity in Japanese patients with seasonal allergic rhinitis during the peak cedar pollinosis season. *Int Arch Allergy Immunol.* 136:148–154.
- Sakashita M, Hirota T, Harada M, Nakamichi R, Tsunoda T, Osawa Y, Kojima A, Okamoto M, Suzuki D, Kubo S, et al. 2010. Prevalence of allergic rhinitis and sensitization to common aeroallergens in a Japanese population. *Int Arch Allergy Immunol.* 151:255–261.
- Sano T, Akahane M, Sugiura H, Ohkusa Y, Okabe N, Imamura T. 2013. Internet survey of the influence of environmental factors on human health: environmental epidemiologic investigation using the web-based daily questionnaire for health. *Int J Environ Health Res.* 23:247–257.
- Sasaki K, Okamoto Y, Yonekura S, Okawa T, Horiguchi S, Chazono H, Hisamitsu M, Sakurai D, Hanazawa T, Okubo K. 2009. Cedar and cypress pollinosis and allergic rhinitis: quality of life effects of early intervention with leukotriene receptor antagonists. *Int Arch Allergy Immunol.* 149:350–358.

- Schleyer TK, Forrest JL. 2000. Methods for the design and administration of web-based surveys. *J Am Med Inform Assoc.* 7:416–425.
- Sharp TJ, Seeto C. 2010. The psychosocial impact of self-reported morning allergy symptoms: findings from an Australian internet-based survey. *J Allergy (Cairo).* 2010:710926.
- Sugiura H, Ohkusa Y, Akahane M, Sano T, Okabe N, Imamura T. 2011. Development of a web-based survey for monitoring daily health and its application in an epidemiological survey. *J Med Internet Res.* 13:e66.
- Sugiura H, Ohkusa Y, Akahane M, Sugahara T, Okabe N, Imamura T. 2010. Construction of syndromic surveillance using a web-based daily questionnaire for health and its application at the G8 Hokkaido Toyako Summit meeting. *Epidemiol Infect.* 138:1493–1502.
- Takasaki K, Enatsu K, Kumagami H, Takahashi H. 2009. Relationship between airborne pollen count and treatment outcome in Japanese cedar pollinosis patients. *Eur Arch Otorhinolaryngol.* 266:673–676.
- Tilston NL, Eames KT, Paolotti D, Ealden T, Edmunds WJ. 2010. Internet-based surveillance of Influenza-like-illness in the UK during the 2009 H1N1 influenza pandemic. *BMC Publ Health.* 10:650.

