

| Gene name     | accession number | position of complementary DNA      |
|---------------|------------------|------------------------------------|
| <i>Oasis</i>  | NM_011957.2      | nucleotides 66 to 1184             |
| <i>Bbf2h7</i> | NM_178661.3      | nucleotides 340 to 1470            |
| <i>Atf6</i>   | NM_001081304.1   | nucleotides 71 to 1165             |
| <i>Luman</i>  | NM_013497.3      | nucleotides 180 to 893             |
| <i>Crebh</i>  | NM_145365.3      | nucleotides 422 to 1372            |
| <i>Creb4</i>  | NM_030080.2      | nucleotides 210 to 1022            |
| <i>Xbp1</i>   | NM_013842.2      | nucleotides 29 to 506, 533 to 1170 |
| <i>Atf4</i>   | NM_009716.2      | nucleotides 595 to 1644            |

**Supplementary Table S1.** National Center for Biotechnology Information's accession numbers and the positions of complementary DNAs inserted into the plasmids.

| Gene name                       | Sense primer (5' - 3')    | Antisense primer (5' - 3')  |
|---------------------------------|---------------------------|-----------------------------|
| <i>Gfap</i>                     | GCAGATGAAGCCACCCTGGCTC    | CCAGATCGCAGGTCAAGGCCTGCAG   |
| <i>Nestin</i>                   | CTACCAGGAGCGCGTGGC        | TCCACAGCCAGCTGGAAGCTT       |
| <i>Bip</i>                      | GTTTGCTGAGGAAGACAAAAAGCTC | CACTTCCATAGAGTTTGCTGATAATTG |
| <i>P58<sup>PK</sup></i>         | GAGGTTTGTGTTTGGGATGCAG    | GCTCTTCAGCTGACTCAATCAG      |
| <i>Grp94</i>                    | GCCGAGAAGGCTCAAGGACAG     | CACCCGTGTCTGTGACATGCAG      |
| <i>Erdj4</i>                    | CCCCAGTGTCAAAGTGTACCAG    | AGCGTTTCCAATTTCCATAAATT     |
| <i>Edem</i>                     | AAGCCCTCTGGAAGTTGCG       | AACCCAATGGCCTGTCTGG         |
| <i>Pdi</i>                      | CAAGATCAAGCCCCACCTGAT     | AGTTCGCCCAACCAGTACTT        |
| <i>Orp150</i>                   | GATGTCTGTAGACCTGGGCAGTG   | GAAGTGCAGCTGCGGACTGATCTG    |
| <i>Chop</i>                     | GTCCAGCTGGGAGCTGGAAG      | CTGACTGGAATCTGGAGAG         |
| <i>Xbp1</i>                     | ACACGCTTGGGAATGGACAC      | CCATGGGAAGATGTTCTGGG        |
| <i><math>\beta</math>-actin</i> | TCCTCCCTGGAGAAGAGCTA      | TCCTGCTTGCTGATCCACAT        |

**Supplementary Table S2.** The primer sets used for real-time PCR and RT-PCR.

## Treatment of a Mouse Model of Spinal Cord Injury by Transplantation of Human Induced Pluripotent Stem Cell-Derived Long-Term Self-Renewing Neuroepithelial-Like Stem Cells

YUSUKE FUJIMOTO,<sup>a,b</sup> MASAHIKO ABEMATSU,<sup>b</sup> ANNA FALK,<sup>c</sup> KEITA TSUJIMURA,<sup>a</sup> TSUKASA SANOSAKA,<sup>a</sup> BERRY JULIANDI,<sup>a,d</sup> KATSUNORI SEMI,<sup>a</sup> MASAKAZU NAMIHIRA,<sup>a</sup> SETSURO KOMIYA,<sup>b</sup> AUSTIN SMITH,<sup>c</sup> KINICHI NAKASHIMA<sup>a</sup>

<sup>a</sup>Laboratory of Molecular Neuroscience, Graduate School of Biological Sciences, Nara Institute of Science and Technology, Ikoma, Japan; <sup>b</sup>Department of Orthopaedic Surgery, Graduate School of Medical and Dental Sciences, Kagoshima University, Kagoshima, Japan; <sup>c</sup>Department of Biochemistry, Wellcome Trust Centre for Stem Cell Research, Stem Cell Institute, University of Cambridge, Cambridge, United Kingdom; <sup>d</sup>Department of Biology, Bogor Agricultural University (IPB), Bogor, Indonesia

**Key Words.** Spinal cord injury • Induced pluripotent stem cells • Neural stem cells • Transplantation • Regenerative medicine

### ABSTRACT

Because of their ability to self-renew, to differentiate into multiple lineages, and to migrate toward a damaged site, neural stem cells (NSCs), which can be derived from various sources such as fetal tissues and embryonic stem cells, are currently considered to be promising components of cell replacement strategies aimed at treating injuries of the central nervous system, including the spinal cord. Despite their efficiency in promoting functional recovery, these NSCs are not homogeneous and possess variable characteristics depending on their derivation protocols. The advent of induced pluripotent stem (iPS) cells has provided new prospects for regenerative medicine. We used a recently developed robust and stable protocol for the generation of long-term, self-renewing, neuroepithelial-like stem cells from human iPS cells (hiPS-lt-NES cells),

which can provide a homogeneous and well-defined population of NSCs for standardized analysis. Here, we show that transplanted hiPS-lt-NES cells differentiate into neural lineages in the mouse model of spinal cord injury (SCI) and promote functional recovery of hind limb motor function. Furthermore, using two different neuronal tracers and ablation of the transplanted cells, we revealed that transplanted hiPS-lt-NES cell-derived neurons, together with the surviving endogenous neurons, contributed to restored motor function. Both types of neurons reconstructed the corticospinal tract by forming synaptic connections and integrating neuronal circuits. Our findings indicate that hiPS-lt-NES transplantation represents a promising avenue for effective cell-based treatment of SCI. *STEM CELLS* 2012;30:1163–1173

Disclosure of potential conflicts of interest is found at the end of this article.

### INTRODUCTION

Our current ability to reconstruct the damaged central nervous system, including spinal cord injury (SCI), is limited. SCI is one of the commonest causes of loss of movement and sensation below the level of the injured spinal cord. While partial spontaneous functional recovery has been observed in the case of moderate SCI, there is no adequate treatment to repair the injured spinal cord itself, and most patients have no therapeutic options except for general management and rehabilitation [1].

Transplantation of neural stem cells (NSCs) into the injured spinal cord has been shown to be an effective treat-

ment [2] because of their competence to differentiate into neurons and oligodendrocytes and to secrete neurotrophic factors. Transplantation into injured spinal cords of several different types of human NS/progenitor cells, derived from human fetal tissue [3, 4] or from human embryonic stem cells (hESCs) [5, 6], promotes functional recovery in animal models. Nonetheless, the mechanism underlying such functional improvement remains to be fully elucidated.

The establishment of induced pluripotent stem cells (iPSCs) offers new prospects for regenerative therapies [7, 8]. Human iPSCs (hiPSCs) can be generated from cells in adult tissue, making it possible to create iPSCs from SCI patients themselves. From the viewpoint of ethics and host immune rejection, NSCs derived from human iPSCs (hiPS-NSCs)

Author contributions: Y.F.: conception and design, collection and assembly of data, data analysis and interpretation, and manuscript writing; A.F. and A.S.: provision of study materials; M.A., K.T., T.S., K.S., and M.N.: collection and assembly of data, data analysis and interpretation; B.J.: manuscript writing; S.K.: administrative support; K.N.: conception and design, administrative support, manuscript writing, and final approval of manuscript.

Correspondence: Kinichi Nakashima, Ph.D., Laboratory of Molecular Neuroscience, Graduate School of Biological Sciences, Nara Institute of Science and Technology, 8916-5 Takayama, Ikoma 631-0192, Japan. Telephone: 81-743-72-5471; Fax: 81-743-72-5479; e-mail: kin@bs.naist.jp. Received October 18, 2011; accepted for publication February 11, 2012; first published online in *STEM CELLS EXPRESS* March 14, 2012. © AlphaMed Press 1066-5099/2012/\$30.00/0 doi: 10.1002/stem.1083



appear to be an ideal resource for transplantation therapy, and the efficacy of hiPS-NSC transplantation in SCI treatment has just begun to be investigated using the mouse model of SCI [9].

We used a recently developed protocol for the generation of long-term self-renewing neuroepithelial-like stem (lt-NES) cells, which satisfy the criteria to be defined as NSCs, from several different lines of hESCs and hiPSCs [10, 11]. hiPSC-derived lt-NES (hiPS-lt-NES) cells exhibit consistent characteristics such as continuous expandability, stable neuronal and glial differentiation ability, and the capacity to generate functional mature neurons in monolayer culture. Whereas neurosphere cultures display heterogeneous character and are sensitive to variation in methodological procedures [12], monolayer cultures offer a more homogeneous and robust cell generation [13, 14].

We report here that transplanted hiPS-lt-NES cells, derived from our robust and stable monolayer cultures, have a therapeutic potential comparable to that of NSCs from human fetal spinal cord (hsp-NSCs) for SCI in the nonobese diabetic-severe combined immunodeficient (NOD-SCID) mouse model. We further show that hiPS-lt-NES cell transplantation promotes recovery of hind limb motor function through the reconstruction of the corticospinal tract (CST), and restores disrupted neuronal circuitry in a relay manner as we have previously demonstrated in a study of mouse NSC transplantation [15]. Our results suggest that hiPS-lt-NES cells represent a promising cell source for transplantation into the injured spinal cord.

## MATERIALS AND METHODS

### Cell Culture

hsp-NSCs and hiPS-lt-NES cells were established and maintained as described previously [10, 11, 16]. The hsp-NS cell line CB660sp and hiPS-lt-NES cell line AF22 were used in the present study. hsp-NSCs were plated onto 10  $\mu$ g/ml laminin (Sigma, St. Louis, MO, <http://www.sigmaldrich.com>)-coated plates in maintenance medium, consisting of Euromed-N medium (Euroclone, Milano, Italy, <http://www.euroclonegroup.it>), 2 mM L-glutamine, 0.1 mg/ml penicillin/streptomycin (Sigma), N2 supplement (1:100), 20 ml/l B27 (all from Invitrogen, Carlsbad, CA, <http://www.invitrogen.com>), 10 ng/ml fibroblast growth factor (FGF, R&D Systems, Minneapolis, MN, <http://www.rndsystems.com>)-2, and 10 ng/ml epidermal growth factor (EGF, R&D Systems). Cells were passaged at a ratio of 1:2 every second to third day using Accutase (Sigma). hiPS-lt-NES cells were plated onto 0.1 mg/ml poly-L-ornithine and 10  $\mu$ g/ml laminin (O/L, Sigma)-coated plates in maintenance medium, consisting of Dulbecco's modified Eagle's medium/F12 (Invitrogen), 2 mM L-glutamine, 1.6 mg/ml glucose, 0.1 mg/ml penicillin/streptomycin, N2 supplement (1:100), 1  $\mu$ l/ml B27, 10 ng/ml FGF2 and 10 ng/ml EGF. Cells were passaged at a ratio of 1:3 every second to third day using trypsin. To induce in vitro differentiation, hiPS-lt-NES cells were plated onto an O/L-coated 35-mm dish at a density of  $5 \times 10^5$  cells per dish in culture medium without both EGF and FGF, containing 1% fetal bovine serum (FBS), and cultured for 4 weeks. Half of the medium was changed every 2 days and laminin (1:500) was added to the medium once a week to prevent detachment of the cells.

### Lentivirus Production and Infection of hsp-NSCs and hiPS-lt-NES Cells

Lentivirus production and infection of cells were performed as described previously [17, 18]. hsp-NSCs and hiPS-lt-NES cells were infected with lentiviruses harboring the luciferase and green

fluorescent protein (GFP) genes connected by an internal ribosomal entry site (IRES), Efp (elongation factor promoter)-luciferase-IRES-GFP (Fig. 1B). hsp-NSCs and hiPS-lt-NES cells that had undergone more than 20 passages were infected. GFP-positive cells were collected by fluorescence activated cell sorting (FACS)-Aria II CellSorter (B.D. Biosciences, San Jose, CA, <http://www.bdbiosciences.com>) and used for transplantation.

### SCI Model and Cell Transplantation

All aspects of animal care and treatment were carried out according to the guidelines of the experimental animal care committee of Nara Institute of Science and Technology. We used female NOD-SCID mice (8-10 weeks old, weighing 18-20 g, Charles River, Osaka, Japan, <http://www.crj.co.jp>). Anesthetized (ketamine 50 mg/kg, xylazine 5 mg/kg, and sodium pentobarbital 20 mg/kg) mice received laminectomies and partial laminectomies at the ninth and 10th thoracic spinal vertebrae, respectively. The dorsal surface of the dura mater was exposed and SCI was induced using an SCI device (70 kdyn; Infinite Horizon impactor, Precision Systems & Instrumentation, Lexington, KY, <http://www.presysin.com>) as previously described [15]. The muscle and skin were closed in layers. All mice subcutaneously received gentamicin (8 mg/kg) daily. The mice underwent manual bladder evacuation once a day. Seven days after injury, mice were anesthetized and transplanted with hiPS-lt-NES cells or hsp-NSCs using a glass micropipette attached to a stereotaxic injector (Narishige, Tokyo, Japan, <http://www.narishige.co.jp>). The tip of the micropipette was inserted into the injury epicenter in the injured spinal cord, and 2  $\mu$ l of culture medium lacking growth factors, with or without NSCs ( $5 \times 10^5 \mu$ l<sup>-1</sup>), was injected at a rate of 1  $\mu$ l/min.

### Behavioral Testing and Electrophysiological Recordings


We evaluated the motor function of the hind limbs for up to 10 weeks after injury. Two individuals, blinded to the treatment of the mice, examined motor function using the Basso Mouse Scale (BMS) locomotor rating scale [19]. Hind limb movements of the mice were captured using a high-definition digital camcorder. We edited these movies and exported movie files using editing software.

To examine signal conduction in motor pathways after SCI, motor-evoked potentials (MEPs) at 12 weeks after SCI were measured. Mice were anesthetized with intraperitoneally injected ketamine (100 mg/kg) and their heads were fixed in a stereotaxic frame. The skulls were tightly fixed to a stereotaxic apparatus (Narishige). Scalping and two small craniotomies were performed with a drill over the motor cortex area (Nakanishi, Tochigi, Japan; <http://www.nsk-nakanishi.co.jp>). Silver ball electrodes were placed epidurally via the holes, into which mineral oil was applied. The motor cortex was stimulated with 0.2-millisecond square wave pulses at a constant current of 10 mA using an electrical stimulator (SEM-3301, Nihon Kohden, Tokyo, Japan; <http://www.nihonkohden.co.jp>). Recording needle electrodes were inserted into the hamstring. A subcutaneous ground electrode was placed in the tail. Electrophysiological recordings were made (MED64, Alpha MED Scientific, Osaka, Japan; <http://www.amedsci.com>) and band-pass filtered at 1-10 kHz. Amplitudes from onset to peak of the negative deflection were measured. Latency of MEP was measured as the time interval between the end of stimulation and the onset of the first wave. Indicated values are the average of five experiments from each mouse.

### Antibodies

The following antibodies were used: rabbit anti-Sox2 (1:1,000, Chemicon-Millipore, Billerica, MA, <http://www.millipore.com>), mouse anti-Nestin (1:250, Chemicon), rabbit anti-brain lipid-binding protein (BLBP, 1:500, Abcam, Cambridge, U.K., <http://www.abcam.com>), mouse anti- $\beta$ -tubulin isotype III (Tuj1; 1:500,





**Figure 1.** Characterization of hiPS-It-NES cells. (A): Expansion culture. hiPS-It-NES cells can be expanded continuously in the presence of both epidermal growth factor and fibroblast growth factor. hiPS-It-NES cells were uniformly immunopositive for Sox2 (green) and Nestin (red). Hoechst (blue) shows nuclear staining. (B): The pCSII-EFp-luciferase-IRES2-GFP construct. (C): Flow cytometric analysis of GFP-positive cells in hiPS-It-NES cells. Uninfected cells (left) and cells infected with lentiviruses expressing luciferase and GFP (middle) were subjected to flow cytometric analysis. GFP-positive cells in the infected cell population were sorted on the basis of GFP fluorescence and reanalyzed for GFP expression (right). (D): After 4 weeks in differentiation conditions, infected hiPS-It-NES cells differentiated into Tuj1-positive neurons (red) and GFAP-positive astrocytes (blue). (E): Quantification of neural marker-positive cells after 4 weeks' differentiation. Lentiviral infection did not influence the differentiation tendency. Data are means  $\pm$  SD ( $n = 3$ ). Scale bars = 100  $\mu$ m. Abbreviations: EFp, elongation factor promoter; GFP, green fluorescent protein; GFAP, glial fibrillary acidic protein; hiPS-It-NES, human induced pluripotent stem cell-derived long-term self-renewing neuroepithelial-like stem; and IRES2, internal ribosomal entry site 2.

Sigma), rabbit anti- $\beta$ -tubulin isotype III (Tuj1; 1:1,000, Covance, Madison, WI, <http://www.covance.com>), rabbit anti-GFP (1:1,000, Molecular Probes, Carlsbad, CA, <http://probes.invitrogen.com>), chick anti-GFP (1:500, Aves Labs, Tigard, OR, <http://www.aveslab.com>), rabbit anti-glial fibrillary acidic protein (GFAP, 1:2,000, DAKO, Carpinteria, CA, <http://www.dako.com>), mouse anti-human-specific GFAP (hGFAP, 1:1,000, STEM123, StemCells Science, Newark, CA, <http://www.stemcellscience.com>), mouse anti-MAP2ab (1:500, Sigma), chick anti-myelin basic protein (MBP) (1:200, Aves Labs), mouse anti-adenomatous polyposis coli CC-1 (APC, 1:200, Calbiochem-Merck, Darmstadt, Germany, <http://www.merckgroup.com>), mouse anti-synaptophysin (1:200, Chemicon), mouse anti-Bassoon (Bsn, 1:400, Stressgen-Enzo Life Sciences, Plymouth Meeting, PA, <http://www.enzo-life.com>), mouse anti-human-specific synaptophysin (hSyn, 1:200, Chemicon), and mouse anti-NeuN (1:500, Millipore).

fesciences.com), mouse anti-human-specific synaptophysin (hSyn, 1:200, Chemicon), and mouse anti-NeuN (1:500, Millipore).

### Immunocytochemistry

Immunocytochemistry experiments were performed as described previously [20]. Cells were washed with phosphate buffered saline (PBS) and fixed with 4% paraformaldehyde (PFA) in PBS for 10 minutes, and then washed with PBS and incubated in blocking solution (PBS containing 10% FBS and 0.1% Triton X-100). Subsequently, cells were incubated overnight at 4°C with the primary antibodies described above. After three washes in PBS, cells were incubated for 1 hour with the following secondary antibodies: FITC-conjugated donkey anti-chick/rabbit, Cy3-



conjugated donkey anti-chick/mouse/rabbit, Cy5-conjugated donkey anti-mouse/rabbit (1:500, Jackson ImmunoResearch, West Grove, PA, <http://www.jacksonimmuno.com>). After three washes with PBS, nuclei were stained with Hoechst (bisbenzimidazole H33258 fluorochrome trihydrochloride, Calbiochem-Merck). Samples were washed with PBS and mounted on glass slides with Immu-Mount (Thermo Scientific, Waltham, MA, <http://www.thermoscientific.com>). The cells were examined using a fluorescence microscope (Axiovert 200M, Zeiss, Jena, Germany, <http://www.zeiss.com>) equipped with the appropriate epifluorescence filters.

### Immunohistochemistry

Animals were anesthetized and perfused with PBS followed by 4% PFA in 0.1 M PBS, pH 7.4. The spinal cords were dissected and postfixed overnight in the same fixative at 4°C. For cryosectioning, fixed tissues were cryoprotected in 10% sucrose in PBS overnight at 4°C, then in 20% sucrose in PBS overnight at 4°C, and embedded in optimal cutting temperature (OCT) compound (Tissue Tek, Sakura Finetek, Tokyo, Japan, <http://www.sakura-finetek.com>). Cryostat sections (20  $\mu$ m) were cut and affixed to matsunami adhesive slide (MAS)-coated glass slides (Matsunami Glass, Osaka, Japan, <http://www.matsunami-glass.co.jp>). The sections were permeabilized in PBS-T (PBS containing 0.1% Triton X-100) for 10 minutes and blocked with 10% FBS in PBS-T for 1 hour, and then incubated overnight at 4°C with the primary antibodies described above. After three washes with PBS, they were incubated in a mixture of the secondary antibodies described above for 1 hour. After a final rinse with PBS, nuclei were stained using Hoechst. Sections were mounted and examined under a fluorescence microscope (Axiovert 200M, Zeiss) and a scanning laser confocal imaging system (LSM 710, Zeiss).

### In Vivo Imaging of Transplanted Cells

In vivo imaging experiments were performed as described previously [18]. A Xenogen-IVIS 100 cooled CCD optical macroscopic imaging system (SC BioScience, Tokyo, Japan, [www.scbio.co.jp](http://www.scbio.co.jp)) was used for bioluminescence imaging. Mice were given an intraperitoneal injection of D-luciferin (150 mg/kg) and serial images were acquired from 20 minutes after administration until the maximum intensity was obtained with the field-of-view set at 10 cm. All images were analyzed using Igor (WaveMetrics, Portland, OR, <http://www.wavemetrics.com>) and Xenogen Living Image software, and optical signal intensity was expressed as photon flux in units of photons/s per cm<sup>2</sup>/steradian. To quantify the measured light, we defined regions of interest (ROIs) over the cell-implanted area and examined all values within the same ROI. The obtained photon count intensity was expressed as a percentage of the initial value.

### Anterograde Labeling of the CST

Twelve weeks after injury, descending CST fibers were labeled with biotinylated dextran amine (BDA) (MW 10,000, 10% in saline, 2  $\mu$ l per cortex; Molecular Probes) [21, 22] by injection into the left and right motor cortices [23]. The skulls of anesthetized mice were tightly fixed to a stereotaxic apparatus (Narishige). Scalping and craniotomy over the motor cortex area were carried out using a micromotor system (Nakanishi). The injection site was 2.1 mm posterior to the bregma, 2 mm lateral to the bregma, and 0.7 mm in depth [23]. For pressure injections with a 20- $\mu$ m outer diameter glass capillary attached to a microsyringe (Narishige), we used 0.1- $\mu$ l steps per minute until the desired volume (1  $\mu$ l per injection site) was injected. Two weeks later, the animals were perfused and their spinal cords were fixed as described above. Sections (30  $\mu$ m) were cut and used for immunohistochemistry. To visualize the BDA, a tyramide signal amplification fluorescence system (Perkin Elmer, Waltham, MA, <http://www.perkinelmer.com>) was used.

### Visualization of Multisynaptic Neural Pathways

To visualize selective and functional transsynaptic neural pathways, wheat germ agglutinin (WGA)-expressing recombinant adenoviruses were used [24, 25]. Twelve weeks after injury, 1  $\mu$ l of saline containing WGA-expressing virus at a titer of  $1 \times 10^{11}$  pfu/ml was injected into the bilateral motor cortices (0.5  $\mu$ l per injection site). The injection site was 2.1 mm posterior to the bregma, 2 mm lateral to the bregma, and 0.7 mm in depth. Two weeks later, animals were perfused and the spinal cords were fixed as described above. Sections (15  $\mu$ m) were cut and used for immunohistochemistry. Rabbit anti-WGA polyclonal antibody (3  $\mu$ g/ml, Sigma) was preabsorbed with 1% acetone powder of mouse brains in blocking solution overnight at 4°C.

### Ablation of Transplant-Derived Cells

Cell ablation experiments were performed as described previously [26, 27]. Diphtheria toxin (DT) was purified from conditioned medium of the PW8 strain of *Corynebacterium diphtheriae* by diethylaminoethyl Sepharose column chromatography and diluted to an appropriate concentration with saline. Seven weeks after injury, DT solution (50  $\mu$ g/kg per day  $\times$  2 days) was administered by intraperitoneal injection into seven hiPS-NES cell-transplanted mice.

### Statistical Analysis

We performed statistical analysis with an unpaired two-tailed Student's *t* test for single comparisons. For BMS and BDA fiber analysis, we used repeated-measures analysis of variance (ANOVA) with Tukey-Kramer multiple comparison test at each point (Prism, GraphPad, LA Jolla, CA, <http://www.graphpad.com>). *p* < .05 was considered significant.

## RESULTS

### Characterization of hiPS-Lt-NES Cells In Vitro

We have previously shown that hiPS-NES cells, which are reliably derived from different hiPS cell lines, exhibit consistent characteristics such as continuous expandability, stable neuronal and glial differentiation, and the capacity to generate functional mature human neurons [11]. hiPS-lt-NES cells can be expanded in the presence of FGF2 and EGF in monolayer culture, and express the NSC markers Sox2 and Nestin (Fig. 1A) and the radial glial marker BLBP (Supporting Information Fig. 1), but not the ES/iPS cell markers Oct3/4 or Nanog (not shown).

To visualize transplanted cells by both luminescence and fluorescence, hiPS-lt-NES cells were infected with lentiviruses engineered to express luciferase and GFP (Fig. 1B). Almost all hiPS-lt-NES cells were infected (Fig. 1C, middle), and we isolated only strongly GFP-expressing cells for use in subsequent in vitro and transplantation experiments (Fig. 1C, right).

We then examined whether lentiviral infection affected the differentiation potential of hiPS-lt-NES cells. Uninfected and infected hiPS-lt-NES cells were induced to differentiate by removal of growth factors and cultured in the presence of 1% FBS for 4 weeks. Both hiPS-lt-NES cell population differentiated into large and small numbers of Tuj1-positive neurons and GFAP-positive astrocytes, respectively, as observed in our previous study [11] (Fig. 1D). Quantitative analysis revealed no significant differences in the proportions of differentiated cells between uninfected and infected hiPS-lt-NES cells (Fig. 1E). Thus, we concluded that lentiviral infection did not influence the differentiation potential of hiPS-NES cells.



### Transplantation of hiPS-Lt-NES Cells into the Injured Spinal Cord Improves Functional Recovery of Hind Limbs

In the present study, we used NOD-SCID mice, which are a suitable model for xenograft research because they are constitutively immunodeficient; their pathology and innate immune response after SCI are also similar to those of other mouse strains [4, 9, 28].


Previous studies have already shown that transplantation of human NSCs from fetal spinal cord or brain tissue improves locomotor functional recovery of SCI models [3, 4]. Nevertheless, only recently has the efficacy of hiPS-NSC transplantation into SCI begun to be investigated [9]. To determine whether our hiPS-Lt-NES cells have a therapeutic capability, we first sought to compare the effects of these cells with those of human fetal spinal cord-derived NSCs (hsp-NSCs). Seventy-kilodyne contusive SCI was applied at the ninth thoracic vertebral level of mouse spinal cords, and 1 week later we injected medium alone (SCI control), or medium containing hsp-NSCs or hiPS-Lt-NES cells, into the epicenter. We then monitored the animals' hind limb motor function using the BMS [19] for at least 8 weeks. All mice showed complete paralysis at 1-day after SCI (BMS scores were 0, Supporting Information Video 1). At 8 weeks after SCI, control mice could move their hind limbs but could not support their own weight (BMS scores were approximately 2, Supporting Information Video 2). In contrast, mice that received hsp-NSCs could touch the ground with their paws and/or support their body weight using their hind limbs (BMS scores were 3-4). The hiPS-Lt-NES cell-transplanted group also showed functional recovery (BMS scores were 3-4, Supporting Information Video 3) compared to the control group, and there was no significant difference in BMS scores between the hsp-NSC- and hiPS-Lt-NES cell-transplanted groups (Fig. 2A). These results indicate that hiPS-Lt-NES cells have a comparable therapeutic effect to hsp-NSCs on SCI.

Next, to evaluate the recovery of descending pathways from the motor cortex to the hind limb motor neurons, we monitored MEPs at 12 weeks after SCI. We stimulated motor cortices electrically and recorded MEP amplitudes from the hamstring muscles. The MEP amplitudes in the hiPS-Lt-NES cell-transplanted group were significantly higher than those in the SCI control group (Fig. 2B, 2C). Furthermore, the latency of MEP response in the hiPS-Lt-NES cell-transplanted group was significantly shorter than that in the SCI control group (Fig. 2D). These results suggest that transplantation of hiPS-Lt-NES cells into the injured spinal cord stimulates the functional recovery of hind limbs.

### Transplanted hiPS-Lt-NES Cells Survive and Differentiate in the Injured Spinal Cord of NOD-SCID Mice

We have established hiPS-Lt-NES cells that express luciferase and GFP, enabling us to trace the survival of the transplanted cells with a bioluminescence imaging system which detects photon signals from living cells after administration of luciferin (a substrate of luciferase) to the mice [18]. The photon signals decreased gradually after transplantation (Fig. 3A) and approximately 20% of transplanted hiPS-Lt-NES cells survived in the injured spinal cord at 5 weeks after SCI (4 weeks after transplantation), whereafter the photon signals remained stable (Fig. 3B).

Using immunohistochemistry, we found that GFP-positive transplanted cells survived and migrated to both rostral and caudal areas around the lesion site in the injured spinal cord (Fig. 3C). High-magnification images showed that trans-




**Figure 2.** Transplantation of hiPS-Lt-NES cells into the injured spinal cord improves functional recovery of hind limbs. (A): Time course of functional recovery of hind limbs after SCI. Data are means  $\pm$  SEM (SCI control,  $n = 8$ ; hsp-NSCs,  $n = 8$ ; hiPS-Lt-NES cells,  $n = 9$ ). Mean BMS values of hsp-NSC- and hiPS-Lt-NES cell-transplanted groups were compared with those of the SCI control group. \*,  $p < .05$ . There was no significant difference between values in the hsp-NSC group and the hiPS-Lt-NES cell group. (B): Representative MEP waves of intact, SCI control, and hiPS-Lt-NES cell-treated mice at 12 weeks after injury. The motor cortices were stimulated and MEP amplitudes were recorded from hamstring muscles. Amplitudes from onset to peak of the negative deflection were measured. (C): Relative values of the mean MEP amplitudes. Values are expressed as percentages of those in intact mice. Mean relative MEP amplitude in the hiPS-Lt-NES cell group was significantly higher than that in the SCI control group. \*,  $p < .05$ . Data are means  $\pm$  SD ( $n = 3$ ). (D): Relative values of the mean MEP latency. Mean relative MEP latency in the hiPS-Lt-NES cell group was significantly shorter than that in the SCI control group. \*,  $p < .05$ . Data are means  $\pm$  SD ( $n = 3$ ). Abbreviations: BMS, Basso Mouse Scale; hiPS-Lt-NES, human induced pluripotent stem cell-derived long-term self-renewing neuroepithelial-like stem; MEP, motor-evoked potential; hsp-NSC, human fetal spinal cord-derived neural stem cell; and SCI, spinal cord injury.

planted hiPS-Lt-NES cells extended processes into both gray and white matter (Fig. 3C-1, 3C-2).

We then examined the differentiation of the transplanted hiPS-Lt-NES cells in the injured spinal cord and found a similar differentiation tendency to that observed in vitro (Fig. 1D, 1E), with many GFP-positive transplant-derived cells having become Tuj1-positive neurons (75%) at 8 weeks after SCI (Fig. 3D-3F). Twenty percent of GFP-positive transplanted cells differentiated into GFAP-positive astrocytes, while their





**Figure 3.** Transplanted human induced pluripotent stem cell-derived long-term self-renewing neuroepithelial-like stem (hiPS-lt-NES) cells survive and differentiate in the injured spinal cord of nonobese diabetic-severe combined immunodeficient mice. (A): Survival of transplanted cells was checked every week using a bioluminescence imaging system; left, 2 hours after transplantation; middle, 4 weeks after SCI; and right, 8 weeks after SCI. (B): Time course of transplanted hiPS-lt-NES cell survival in SCI model mice. Optical signal intensity was measured using the bioluminescence imaging system. Quantification of the photon intensity revealed that approximately 20% of the transplanted cells survived 5 weeks after SCI; thereafter, the photon signals remained stable. Data are means  $\pm$  SEM ( $n = 6$ ). (C): Sagittal sections of SCI model mice treated with hiPS-lt-NES cells at 8 weeks after transplantation. Sections were stained with anti-GFP antibody (green). The epicenter of the SCI is indicated (\*). Higher-magnification images of the white dotted boxes (C-1 and C-2) show GFP-positive transplant-derived cells extending their processes into gray and white matter. (D): Immunostaining images, 8 weeks after injury, of hiPS-lt-NES cells grafted into spinal cord. Spinal cord sections were stained with anti-GFP (green), hGFAP (magenta) and Tuj1 (red) antibodies and with Hoechst (blue). (E): Confocal images, 8 weeks after injury, of hiPS-lt-NES cells transplanted into spinal cord, revealing transplanted cells which were double-positive for GFP and markers of neural lineages. (F): Quantitative analyses of Tuj1-positive neurons, hGFAP-positive astrocytes, and MBP-positive oligodendrocytes as in E. Data are means  $\pm$  SD ( $n = 3$ ). Scale bars = 500  $\mu$ m in C, 100  $\mu$ m in C-1, C-2, D, and E. Abbreviations: GFP, green fluorescent protein; hGFAP, anti-human-specific glial fibrillary acidic protein; MBP, and SCI, spinal cord injury.

differentiation into MBP- or APC-positive oligodendrocytes was less than 1% (Fig. 3E, 3F, Supporting Information Fig. 2). These data indicate that transplanted hiPS-lt-NES cells differentiate preferentially into neurons in the injured spinal cord.

#### hiPS-Lt-NES Cell Transplantation Does Not Promote CST Axon Re-Extension After SCI

Since transplanted NSCs have been reported to play a supportive role in the re-extension of injured axons [29], we examined the effect of the transplanted hiPS-lt-NES cells on CST first-order axonal regeneration. We injected BDA into the bilateral motor cortices and labeled CST first-order neuron axons by anterograde tracing [21, 22]. Because BDA is not transported from first- to second-order neurons across the synapse, only the axons of first-order neurons in the CST are visualized by this method.


In the region caudal to the injury site, >50% of the labeled fibers observed at 5 mm anterior to the lesion site were

detected in the intact mice, whereas almost no BDA-traced CST fibers could be seen in the SCI control or hiPS-lt-NES cell-transplanted mice (Fig. 4A). Quantitative analysis revealed that there was no significant difference in the number of BDA-labeled fibers between the SCI control and hiPS-lt-NES cell-treated groups at any position up to 5 mm on either side of the lesion site (Fig. 4B). Although we cannot exclude possibilities such as the re-extension of descending raphespinal axons, which are also important for the motor functional recovery of hind limbs [30, 31], these results suggest that the regeneration of CST axons, if it occurs, cannot be a major contributing factor in the recovery induced by hiPS-lt-NES cell transplantation in our SCI model.

#### Local Neurons Reconstruct Disrupted CST Neuronal Circuits in a Relay Manner

Recent studies have revealed that local endogenous and transplant-derived neurons can form new neuronal circuits and make synaptic connections after SCI [32–34]. To determine





**Figure 4.** hiPS-Lt-NES cell transplantation does not promote corticospinal tract (CST) axon re-extension after SCI. **(A):** Representative pictures of BDA-labeled CST fibers (red) at 5 and 2 mm rostral and 2 mm caudal to the lesion site at 12 weeks after SCI. Hoechst (blue) shows nuclear staining. Scale bar = 50  $\mu$ m. **(B):** Quantification of the labeled CST fibers in the spinal cords of untreated, SCI control, and hiPS-Lt-NES cell-transplanted mice. The x-axis indicates specific locations along the rostro-caudal axis of the spinal cord, and the y-axis indicates the ratio of the mean number of BDA-labeled fibers at the indicated site to that at 5 mm rostral to the lesion site (thoracic vertebra [Th] 9). Intact mice were compared with SCI control mice. \*,  $p < .05$ ; \*\*,  $p < .01$ . There was no significant difference in the number of BDA-labeled fibers between hiPS-Lt-NES cell-treated (blue line) and SCI control groups (red line). Data are means  $\pm$  SEM ( $n = 5$ ). Abbreviations: BDA, biotinylated dextran amine; hiPS-Lt-NES, human induced pluripotent stem cell-derived long-term self-renewing neuroepithelial-like stem; and SCI, spinal cord injury.

whether disrupted CSTs were reconstructed by forming a local neuronal relay, we injected WGA-expressing adenoviruses into the motor cortex of the hind limb area at 12 weeks after SCI. WGA, a plant lectin, has been widely used as a tracer of neuronal pathways [24, 25] because it is transported in axons and dendrites, and across synapses, to second- and even third-order neurons.

After injection, we could detect WGA-immunoreactive intracellular granule-like structures in MAP2ab-positive neurons (Fig. 5A). Furthermore, we found that hiPS-Lt-NES cell-treated mice had more WGA/Map2ab double-positive cells than SCI control mice in the caudal region below the injured site (Fig. 5A, 5B). Taking these data and the results of the BDA experiment into consideration, it is suggested that WGA

was transferred to the caudal area through the lesion site via new synaptic connections, and that hiPS-Lt-NES cell transplantation promoted the CST reconstruction without CST axonal re-extension. In support of this proposition, we could observe GFP-positive transplant-derived neurons adjacent to synaptophysin-positive patches (Fig. 5C-1 and 5C-1'). Furthermore, immunohistochemistry using species-specific antibodies for presynaptic markers revealed that transplant-derived neurons made synapses with endogenous neurons, suggesting that they reconstructed disrupted CST neuronal circuits in a relay fashion (Fig. 5C-2, 5C-2', 5C-3, and 5C-3').

### hiPS-Lt-NES Cell Transplantation Promotes the Survival of Endogenous Neurons

Massive endogenous cell death is observed in the injured spinal cord, probably due to environmental changes such as increased levels of inflammatory cytokines [35, 36]. Previous studies have suggested that neurotrophic factors secreted from transplanted cells could alleviate cell death and thereby contribute to improved locomotor function after SCI [37]. We therefore wanted to examine whether transplanted hiPS-Lt-NES cells influence the survival of endogenous neurons. We performed immunohistochemistry and counted the number of NeuN-positive/GFP-negative endogenous neurons around the lesion site (Fig. 6A) and found that the number of endogenous neurons in the hiPS-Lt-NES cell-treated mice was significantly higher than that in the SCI control (Fig. 6B). These findings indicate that transplantation of hiPS-Lt-NES cells alters the environment of the injured spinal cord and promotes the survival of endogenous neurons.


### Transplanted Cells Contribute Directly to Functional Recovery of Hind Limb Movement in SCI Model Mice

Ablation of specific cells is a useful method to evaluate their functional contribution in animal models. Mouse and rat cells are less sensitive to DT than human cells, which express human heparin-binding epidermal growth factor-like growth factor (hHB-EGF) as a DT receptor [26, 27]. We administered DT to hiPS-Lt-NES cell-transplanted mice at 7 weeks after SCI. Using an in vivo imaging system, we were able to trace the survival of transplanted cells while evaluating hind limb function. Almost all the transplanted hiPS-Lt-NES cells were specifically ablated following DT administration (Fig. 7A). After ablation of the transplanted cells, the BMS score of hiPS-Lt-NES cell-treated mice dropped to a level similar to that of SCI control mice, although the reduction was not large enough to make the score differ significantly from that attained by hiPS-Lt-NES cell-treated mice without DT administration (Fig. 7B). This may be because transplanted hiPS-Lt-NES cells promoted the survival of endogenous neurons (Fig. 6) that participated in the reconstruction of disrupted neuronal circuitry. Nevertheless, these results suggest that transplanted hiPS-Lt-NES cells play direct and important roles in the functional recovery of the injured spinal cord.


## DISCUSSION

The advent of cell reprogramming and the establishment of iPSCs have provided new prospects for stem cell transplantation therapy [7, 8], and a large number of different types of iPSCs have already been generated by various methods [38–40]. Transplantation of NSCs into the injured spinal cord has been shown to be effective, and with recent advances in stem cell biology, particularly in human ES/iPSCs, we anticipate





**Figure 5.** Local neurons reconstruct disrupted corticospinal tract neuronal circuits in a relay manner. (A): Representative pictures of WGA-labeled neuronal cell bodies located in the caudal area (Th11 to lumbar vertebra [L] 1) at 12 weeks after spinal cord injury (SCI; left, intact mice; middle, SCI control mice; and right, hiPS-It-NES cell-treated mice). Spinal cord sections were stained with anti-WGA (red) and anti-MAP2ab (neuronal marker, blue) antibodies. WGA immunoreactivity was observed as intracellular granule-like structures. (B): WGA-positive cells/MAP2ab-positive neurons in the caudal area were quantified. Values are expressed as percentages of those in intact mice. The percentage of WGA-positive cells in hiPS-It-NES cell-treated mice was significantly higher than that in SCI control mice.  $* p < .05$ . Data are means  $\pm$  SD ( $n = 3$ ). (C): Spinal cord sections were stained with anti-GFP and Tuj1 antibodies, Hoechst (blue), and Syn, Bsn, or hSyn antibodies. (C-1) Representative confocal image showing that GFP (green) and Tuj1 (red) double-positive transplant-derived neurons were adjacent to synaptophysin-positive patches (white). (C-1') High-magnification view of boxed area in (C-1). (C-2) Host and graft synapses were distinguished using a monoclonal antibody for the presynaptic marker Bsn that selectively recognizes rat and mouse, but not human, epitopes. Confocal image showing that GFP (green) and Tuj1 (red) double-positive transplant-derived neurons were in contact with Bsn-positive host synapses (white). (C-2') High-magnification view of boxed area in (C-2). (C-3) Confocal images showing that GFP (green)-negative and Tuj1 (red)-positive endogenous neurons were in contact with hSyn-positive synapses (white). (C-3') High-magnification view of boxed area in (C-3). Scale bars = 10  $\mu$ m. Abbreviations: Bsn, anti-Bassoon; GFP, green fluorescent protein; hiPS-It-NES, human induced pluripotent stem cell-derived long-term self-renewing neuroepithelial-like stem; hSyn, anti-human-specific synaptophysin; Syn, anti-synaptophysin; and WGA, wheat germ agglutinin.



**Figure 6.** Transplanted hiPS-It-NES cells promote the survival of endogenous neurons. (A): Representative sagittal sections of SCI control and hiPS-It-NES cell-transplanted mice at 12 weeks after SCI. Sections were stained with anti-GFAP (white), anti-NeuN (mature neuronal marker, red) and anti-GFP (green) antibodies. Two 500- $\mu$ m regions, at the rostral and caudal edge of the lesioned site (double-headed arrows), were selected for the assessment. The epicenter of the SCI is indicated (\*). Scale bars = 500  $\mu$ m. (B): Quantification of NeuN-positive/GFP-negative endogenous neurons. Values are expressed as percentages of those in intact mice. The number of NeuN-positive/GFP-negative endogenous neurons in hiPS-It-NES-treated mice was significantly higher than that in SCI control mice.  $* p < .05$ . Data are means  $\pm$  SD ( $n = 3$ ). Abbreviations: GFAP, glial fibrillary acidic protein; GFP, green fluorescent protein; hiPS-It-NES, human induced pluripotent stem cell-derived long-term self-renewing neuroepithelial-like stem; and SCI, spinal cord injury.