


Contents lists available at ScienceDirect

Biochemical and Biophysical Research Communications

journal homepage: www.elsevier.com/locate/ybbrc

Genistein promotes DNA demethylation of the steroidogenic factor 1 (SF-1) promoter in endometrial stromal cells

Hiroshi Matsukura^a, Ken-ichi Aisaki^b, Katsuhide Igarashi^b, Yuko Matsushima^b, Jun Kanno^b, Masaaki Muramatsu^a, Katsuko Sudo^{a,c}, Noriko Sato^{a,*}

^a Department of Molecular Epidemiology, Medical Research Institute, Tokyo Medical and Dental University, 2-3-10 Kanda-surugadai, Chiyoda-ku, Tokyo 101-0062, Japan

^b Division of Cellular and Molecular Toxicology, National Institute of Health Sciences, 1-18-1 Kamiyoga, Setagaya-ku, Tokyo 158-8501, Japan

^c Animal Research Center, Tokyo Medical University, 6-1-1 Shinjuku, Shinjuku-ku, Tokyo 160-8402, Japan

ARTICLE INFO

Article history:

Received 21 July 2011

Available online 29 July 2011

Keywords:

Genistein

DNA methylation

Ovariectomized mice

Primary culture

Steroidogenic factor 1

High-resolution melting analysis

ABSTRACT

It has recently been demonstrated that genistein (GEN), a phytoestrogen in soy products, is an epigenetic modulator in various types of cells; but its effect on endometrium has not yet been determined. We investigated the effects of GEN on mouse uterine cells, *in vivo* and *in vitro*. Oral administration of GEN for 1 week induced mild proliferation of the endometrium in ovariectomized (OVX) mice, which was accompanied by the induction of steroidogenic factor 1 (SF-1) gene expression. GEN administration induced demethylation of multiple CpG sites in the SF-1 promoter; these sites are extensively methylated and thus silenced in normal endometrium. The GEN-mediated promoter demethylation occurred predominantly on the luminal side, as opposed to myometrium side, indicating that the epigenetic change was mainly shown in regenerated cells. Primary cultures of endometrial stromal cell colonies were screened for GEN-mediated alterations of DNA methylation by a high-resolution melting (HRM) method. One out of 20 colony-forming cell clones showed GEN-induced demethylation of SF-1. This clone exhibited a high proliferation capacity with continuous colony formation activity through multiple serial clonings. We propose that only a portion of endometrial cells are capable of receiving epigenetic modulation by GEN.

© 2011 Elsevier Inc. All rights reserved.

1. Introduction

Genistein (GEN), a major phytoestrogen in dietary soy, is a substantial component of the typical Asian and Western vegetarian diets, as well as recently developed infant soy milk formulas. There are several well known potential health benefits of GEN intake [1,2], one of which is an apparent decreased risk of breast and prostate cancers, based on human observational studies [1,3]. But GEN also paradoxically stimulates growth of breast cancer cells in culture [2] and uterine enlargement in rodents [4]. These effects may be mediated through estrogen receptor interactions and/or modulation of endogenous estrogen metabolism [5,6]. Since GEN can bind to estrogen receptors (ERs) α and β , with a stronger affinity to ER β [5], it is categorized as a phyto-selective estrogen receptor modulator (SERM) [6,7]. The variations in GEN's agonistic or antagonistic effects may be affected by variations in endogenous estrogen levels. Previous studies have not determined whether the pleiotropic effects of GEN involve distinct epigenetic alteration.

Recently, GEN was shown to alter DNA methylation in various types of cells, including ES cells [8], but most studies have been performed using cancer cell lines [9–11]. There have been few reports of the effects of GEN on DNA methylation in intact cells or *in vivo* [12]. In the present study, we utilized a uterotrophic assay in ovariectomized (OVX) mice, as a model system to analyze epigenetic regulation by GEN.

In a previous study, high-dose GEN administration to OVX rats resulted in increased uterine weight and changed endometrial cell gene expression [6]. However, no epigenetic alterations were demonstrated under this condition. We selected the steroidogenic factor 1 (SF-1; official symbol: Nr5a1) gene as a target for the methylation analysis. SF-1 is an orphan nuclear receptor and transcription factor for key enzymes involved in steroidogenesis, such as StAR, Cyp11a1 (p450scc), Cyp17a1 (p450c17), and Cyp19a1 (aromatase) [13]. The SF-1 gene is not expressed in normal endometrium; however, SF-1 expression is reactivated in the disease state of human ectopic endometriosis, in which the SF-1 promoter is abnormally demethylated by an unknown mechanism [14]. The subsequent enhancement of steroidogenic genes and resultant local steroidogenesis are proposed to be important etiologies [15]. Therefore, we hypothesized that in mouse endometrial cells,

* Corresponding author. Fax: +81 3 5280 8058.

E-mail addresses: hmatsukura.epi@mri.tmd.ac.jp (Hiroshi Matsukura), nsato.epi@tmd.ac.jp (N. Sato).

SF-1 might be subjected to epigenetic modulation by some external stimuli. Here we show that the SF-1 promoter was demethylated *in vivo* and *in vitro* by GEN treatment. This is the first demonstration of a phytoestrogen altering the epigenetic state of adult endometrium.

2. Materials and methods

2.1. Ethics statement

All procedures described here were performed according to protocols approved by the Animal Care Committee of the National Institute of Health Sciences, and Tokyo Medical and Dental University (No. 0110306A).

2.2. Oral administration of genistein to ovariectomized mice

C57BL/6JmsSlc female mice (SLC) were used in this study. All mice were fed a phytoestrogen-free diet (Oriental Yeast) and were ovariectomized (OVX) 2 weeks prior to the genistein (GEN) treatment. OVX mice were divided into three different treatment groups, each consisting of 3–5 independent replicates, which orally received low-dose GEN (60 mg/kg/day), high-dose GEN (200 mg/kg/day), or vehicle (0.5% CMC-Na (Maruishi Pharmaceutical); 5 ml/kg/day) for 1 week. At the end of treatment (9 weeks of age), all mice were euthanized by exsanguination under ether anesthesia.

2.3. Uterotrophic assay and gene expression study after oral administration of genistein

Whole uteri were harvested, blotted, and weighted. Each uterus was divided into two horns, immediately placed into 2 ml plastic tubes of RNAlater solution (Ambion), and stored at 4 °C. From each sample, one horn was processed for mRNA expression analyses; RNAlater was replaced with 1.0 ml of RLT buffer (Qiagen), and the horn was homogenized by addition of a 5 mm diameter Zirconium bead (Funakoshi) and shaking with a MixerMill 300 (Qiagen) at 20 Hz for 5 min (only the outermost row of the shaker box was used). Further sample preparation and analysis were performed as previously described [16]. mRNA expressions were analyzed using Affymetrix Murine Genome 430 2.0 GeneChips, and calculated as copy number per cell by the Perccellome method [16]. The second uterine horn of each sample was subjected to genomic DNA isolation.

2.4. Isolation of colony-forming cells derived from intact uteri

Five 8- to 9-week-old C57BL/6JmsSlc female mice (SLC) were euthanized by cervical dislocation and whole uteri were harvested. Uterine horns were collected in Dulbecco's modified Eagle's medium/Hams F-12 (DMEM/F-12; Nacalai Tesque) containing 0.05 mg/ml gentamicin (Sigma–Aldrich). Each horn was dissected longitudinally and the endometrial tissue was divided into two portions: the luminal side and the myometrium side. A single cell suspension of endometrial cells was obtained using enzymatic digestion and mechanical means adapted from Chan et al. [17]. The tissue samples were minced and dissociated in 500 µl DMEM/F-12 containing 0.12 mg/ml (0.56 Wunsch U/ml) Blendzyme 2 and 40 µg/ml deoxyribonuclease type I (both from Roche Applied Science) in a shaking incubator (~90 rpm) at 37 °C. At 15 min intervals, the digests were pipetted to promote separation and cell dissociation was monitored microscopically. After 45 min, debris was filtered out using a 40-µm sieve (BD Biosciences). The single-cell suspensions were collected in DMEM/F-12 containing

10% FBS, 0.05 mg/ml gentamicin and stored on ice. Then the sieves were backwashed, and myometrial and glandular debris were further digested to single cells for 45 min as described above. All cell suspensions were filtered as described above, and combined. To remove erythrocytes, the cells were resuspended in 500 µl of HLB solution (Immuno Biological Laboratories) and incubated for 3 min. After washing twice with PBS, viable cell numbers were counted with trypan blue (Sigma–Aldrich). Cells were seeded on gelatin (0.1%, Sigma–Aldrich)-coated dishes at various densities of $0.1\text{--}3 \times 10^5$ cell/60-mm dish. After 14 days, non-overlapping clones were distinguished. Primary cell clones were expanded in DMEM/F-12 containing 5% FBS (SAFC Biosciences) and 0.05–0.1 mg/ml gentamicin, on gelatin-coated dishes.

2.5. Serial cloning of colony-forming cell clones

Self renewal was assessed by serial cloning of individual clones as described by Gargett et al. [18]. Cells were seeded on gelatin-coated 100-mm dishes at 10 cells/cm² (600 cells/100-mm dish). Culture medium was changed every 4 days and secondary clones formed distinct colonies by 14 days after plating. Secondary clones were similarly recloned to generate tertiary clones, and were also expanded in the same manner as the primary culture.

2.6. *In vitro* genistein exposure to colony-forming cells

From 70 isolated cell clones, we selected 20 clones from colonies that were composed of fibroblastic-shaped, homogenous cells with an average doubling time of less than 100 h. The selected clonal cells, whose passage number was less than 10, were subjected to *in vitro* GEN exposure. Cells were seeded on gelatin-coated 60-mm dishes, treated with or without 10 µM of GEN (dissolved in dimethylsulfoxide (DMSO)) in DMEM/F-12 containing 5% FBS and 0.05 mg/ml gentamicin for 7 days. The final DMSO concentration was 0.02%. The culture medium was changed every 2 days.

2.7. Genomic DNA preparation and bisulfite sequencing

Genomic DNA was isolated using a QIAamp DNA Mini Kit (QIAGEN) and 180 ng–1 µg was subjected to sodium bisulfite modification with a EpiTect Bisulfite Kit (QIAGEN) according to manufacturer's protocols. Bisulfite sequencing primers are shown in Supplementary Table 1. PCR products were cloned into the pT7 blue T vector (Novagen) and transformed into *Escherichia coli*. Plasmid DNA from positive colonies was purified and sequenced at the Tokyo Medical and Dental University Genome Laboratory (Tokyo, Japan). Sequence and statistical analyses were performed with the QUantification tool for Methylation Analysis; http://quma.cd-b.riken.jp/top/quma_main_j.html [19]. The statistical significance of the difference between two bisulfite sequence groups at each CpG site was evaluated with Fisher's exact test.

2.8. Screening of DNA methylation status by high-resolution melting assay

All assays were performed on the LightCycler 480 using the LightCycler 480 High Resolution Melting Master kit, according to the manufacturer's instructions. Primers, designed using LightCycler Probe Design Software 2.0 (All, Roche Applied Science) are shown in Supplementary Table 1. All data were analyzed using LightCycler Gene Scanning Software.

2.9. Statistical analysis

Data are shown as means ± SD. Unpaired *t*-tests were used to compare the significance between two groups. Statistical analysis

was performed using Dr. SPSS 2 for Windows. Results were considered statistically significant at a P value of <0.05 .

3. Results

3.1. Effects of genistein in uteri of ovariectomized (OVX) mice

OVX mice were fed with either vehicle (control) or low (60 mg/kg) or high (200 mg/kg) doses of GEN for 7 days and blotted uterus weights were determined (Fig. 1A). Compared to the control,

low-dose GEN treatment did not significantly increase the uterus weight; high-dose treatment induced a slight but significant uterus enlargement (1.4-fold of control; $P < 0.005$). We then determined the mRNA expression levels of SF-1 (Fig. 1B) and steroidogenic genes (Fig. 1C–F) by the Percellome method. The mRNA levels of these genes were very low in the endometria from control and low-dose GEN-treatment groups, but were significantly increased (still less than one copy per cell on average) in the high-dose treatment group ($P < 0.05$), indicating that high-doses of GEN induced expression of these genes. Next, we determined the methylation status of SF-1 in




Fig. 1. Genistein induced endometrial regeneration and SF-1 mRNA expression in uterine tissue of OVX mice. (A) Blotted uterine weights were recorded. Control and GEN-treated groups comprised five and four mice, respectively. ctr; control, G 60; GEN 60 mg/kg/d, G200; GEN 200 mg/kg/d. (B–F) mRNA expressions of steroidogenic genes were determined using GeneChip analysis and were calculated by the Percellome method. Y axis indicates mRNA expression as copy number per cell. (B) SF-1 was determined by 1418315_at, (C) Cyp11a1 by 1439947_at, (D) StAR by 1418729_at, (E) Cyp17a1 by 1417017_at, and (F) Cyp19a1 by 1449920_at. *Statistically significant at $P < 0.05$. **Statistically significant at $P < 0.005$.

the total uterus tissues of the three groups of mice by bisulfite sequencing (Fig. 2A–C). There were 15 CpG sites spanning –272 to +199 of the promoter and the 5'-UTR (exon 1) region of SF-1. The percentages of total methylated CpG sites in this region, in control, and low and high-dose GEN, were 78.3%, 73.9%, and 54.4%, respectively, indicating that GEN dose-dependently induced demethylation in this region. Most CpG sites in the 5'-UTR (exon1) were demethylated by high-dose GEN. In particular, the methylation levels of 5 CpG sites between +45 and +89 were significantly lower in high-dose GEN than in control (Fisher's exact test, $P < 0.01$). We further split the endometrium to separate the luminal side (LU) from the basilar myometrial side (MY), and both specimens were separately subjected to bisulfite sequencing. This procedure was applied to the samples from the GEN-treated groups but not to control samples due to uterus atrophy. In low-dose GEN treated mice, the mean methylation levels of LU and MY were 84.8% and 65.0%, respectively (not shown). In high-dose GEN treated mice, the mean methylation levels of LU and MY were 42.1% and 66.7%, respectively (Fig. 2D). Thus, the demethylation induced by high-dose GEN occurred predominantly in the LU, rather than in the MY.

3.2. Effect of genistein on primary endometrial cell culture

In order to study the SF-1 promoter methylation at the cellular level, we employed an endometrial cell primary culture. Intact

murine endometrium were divided into LU and MY portions, and cells were separately isolated. Primary cell clones were established by colony-formation, following the plating of a serially-titrated cell suspension (see Section 2). Efficient isolated colony formation was achieved with cells seeded at a density of 4,500–15,300 cells/cm². For LU and MY, the average frequencies of colony appearance were 7.5 per 10⁵ cells and 15 per 10⁵ cells, respectively. The growth curves of representative clones derived from LU and MY are shown in Fig. 3. Cell clones with highest and lowest proliferative activities were obtained from LU and MY, separately. More highly proliferative cells were obtained from MY than from LU (Fig. 3A and C). We selected 20 highly proliferative clones for further study (see Section 2). Two rapid growing clones obtained from MY showed self-renewal activity when secondarily seeded at a very low cell density (10 cells/cm²) (not shown). To screen for primary cultured cells that responded to GEN, we set up a high-resolution melting (HRM) assay that identified region-specific methylation levels. The region analyzed by HRM assay exclusively contained the 7 CpG sites between +19 and +89 bp that were most differentially demethylated following oral administration of GEN (Fig. 2). Each clone was treated with or without GEN for 1 week; cells treated with a similar concentration of DMSO served as control. Among the 20 clones that we screened, only one GEN-treated clone (No. 16) exhibited a significant shift in the melting curve compared to control cells (Fig. 4A). This clone had the highest proliferation




Fig. 2. Genistein induced demethylation of the SF-1 gene promoter in endometrial tissues of OVX mice. The schematic diagram indicates CpG locations on the SF-1 promoter region, spanning 5'-flanking to exon 1. The CpG position relative to the first base of exon 1 (+1) is shown. The bisulfite sequencing fragment contains 15 CpG sites. The black inlay represents the mean methylation levels of each CpG, and the left panel contains the number of sequenced clones and the mean methylation level of all CpGs. Summarized methylation results are shown at the top. (A–C) Difference in methylation levels between (A) vehicle (control), (B) low-dose GEN (60 mg/kg/d), and (C) high-dose GEN (200 mg/kg/d) exposed uteri. (D) Difference in methylation levels between luminal and myometrial sides of a uterus exposed to high-dose GEN.




Fig. 3. Proliferation properties of isolated clones: one clone showed highly proliferative activity. Representative growth curves of clones harvested from intact murine endometrium are shown. Formula indicates the slope of fitted growth curves. Clones were divided into two groups: high and low proliferative (see Section 2). A total of 20 highly proliferative clones were analyzed by HRM after *in vitro* GEN treatment. (A, B) Asterisks indicate representative growth curves of cell clones isolated from luminal side. (A) Clones with higher proliferative activities, derived from luminal side. Six highly proliferative clones were obtained in total; 2 representatives are shown. (B) Clones with lower proliferative activities, derived from luminal side. (C, D) Each circle indicates representative growth curves of cell clones isolated from myometrial side. (C) Clones with higher proliferative activities, derived from the myometrial side. In total, 14 highly proliferative clones were obtained; 10 representatives are shown. The arrow indicates the clone with the most rapid growth. (D) Clones with lower proliferative activities, derived from myometrial side.

activity (Fig. 3C, arrow). GEN treatment of the other clones did not result in significant changes to the melting curve patterns (not shown). We further confirmed the methylation status of clone No. 16 by bisulfite sequencing. The percentages of CpG methylation in the SF-1-272 to +199 promoter regions for untreated and GEN-treated cells were 85.0% and 65.8%, respectively (Fig. 4B).

4. Discussion

Growing evidence suggests that the manner in which nutrients can either help maintain health, or conversely, promote disease development may be mediated by epigenetic regulation [12,20]. However, relatively little is known about tissue-specific sensitivity or how much plasticity exists in regards to the effect that a given environmental factor can exert on a certain epigenetic target

[20,21]. GEN, a non-nutrient dietary component of soy products, exhibits mixed estrogen agonist and antagonist properties, and multiple functions both *in vivo* and *in vitro* [7,22]. Several animal studies have demonstrated that GEN acts as an epigenetic modulator [20]. We focused on the effects of GEN on endometrium, because endometrium is not only hormone responsive, but also a highly proliferative organ. Epigenetic alterations of proliferative tissue or cells may then be expanded through tissue proliferation. We used OVX rodents, which are a widely used model for studying estrogen withdrawal and replacement [23], as well as for the assessment of endocrine-disrupting chemicals in the environment [4]. In our experiment, GEN induced proliferation of the endometrium and increased uterine weight (Fig. 1A) to extents similar to those previously reported in OVX rats [4]. Our findings also suggested that GEN treatment induced marked demethylation of




Fig. 4. Genistein induced demethylation of the SF-1 gene in the cell clone that showed the highest proliferative activity. (A) The GEN-mediated demethylation of CpGs (at the positions of +19, 23, 45, 48, 55, 59, and 89) in the SF-1 gene *in vitro* in one out of 20 isolated clones (Fig. 3A and C). HRM analysis enabled to clear separation of the PCR product mixtures with different CpG compositions. The methylation standards were prepared as 100% (red), 95.4% (orange), 90% (yellow), 77% (green), 54% (blue), and 9% (violet) methylated template in demethylated background. Grey and pink indicate the control and GEN-treated clone sample, respectively. (B) SF-1 DNA methylation status, obtained from bisulfate sequencing, of the clone that showed demethylation in HRM assay. Percentage under box (HRM region) indicates percent methylation of HRM region. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this paper.)

CpG sites in the SF-1 promoter (Fig. 2), and substantial increase in SF-1 mRNA level (Fig. 1B). Expressions of genes downstream of transcription factor SF-1 were also significantly enhanced (Fig. 1C–F). However, it should be noted that the induced mRNA levels were still very low (Fig. 1B–F), less than one copy per cell, as determined by the Percellome method [16]. Our results are consistent with those of a previous study reporting that a physiological concentration of GEN increased Cyp19a1 enzymatic activity in endometrial cells derived from a normal uterus, whereas GEN did not affect Cyp19a1 activity in a cell-free assay [24]. It is unknown whether GEN stimulated Cyp19a1 activity by epigenetic modulation.

In the present study, we also identified primary cultured endometrial cells that were competent for epigenetic regulation by GEN, which were present at a very low frequency. In our *in vitro* study, GEN treatment did not enhance, but rather inhibited the proliferation of colony-derived cells. Taken together, these findings indicate that a minor population of endometrium cells can respond to GEN and that, in these cells, GEN induces demethylation and

activation of SF-1, followed by the induction of the SF-1 steroidogenic cascade. This might lead to local steroidogenesis and enhanced endometrium proliferation *in vivo* in GEN-treated OVX mice. Since demethylation of the SF-1 promoter was observed at the whole tissue level (Fig. 2), and the induced expression of steroidogenic genes occurred only to a subcellular amount (Fig. 1B–F), the demethylation event in each cell may not be sufficient for the SF-1 induction.

The demethylation in the endometrium that occurred after 1 week of treatment with high-dose GEN was more prominent in the LU than in the MY. Following GEN treatment, the methylation level in the MY was similar to that in the untreated endometrium as a whole. After 7 days of GEN exposure, the endometrial cells in the LU of OVX mice were composed of regenerated cells moving from MY to LU; in light of this, our results indicate that the initial epigenetic alteration might be expanded through proliferation of regenerating cells. This is also consistent with our observation that more endometrial cells derived from MY showed higher colony-formation activity and rapid proliferation than did those derived

from LU (Fig. 3). We thus speculate that there are GEN-sensitive cells in the MY, which might contain endometrial stem cells [25].

Recently, human endometrial stem cells were identified; they reside in endometrial stromal tissue and possess fibroblastic-shape and self-renewal ability, thus forming large, densely packed, homogenous colonies [17,18]. Some candidates for murine endometrial progenitor cells have been suggested to reside in the luminal epithelial or area of adjacent to the myometrium [25]. The rapidly growing endometrial cell that we obtained had a fibroblastic-shape, formed large, relatively homogenous, densely packed colonies, and showed self-renewal activity when seeded at a very low cell density (data not shown). Although there are differences between the species, we speculate that our rapid growing cell clones may correspond to the human endometrial stromal progenitor cells [17].

Aberrant SF-1 expression with lack of promoter CpG methylation has been reported in ectopic endometriosis [14], and thus endometriosis is now considered an epigenetic disease [26]. Endometriosis is classically defined as the growth of endometrial tissue at extrauterine sites; it has been suggested that each endometriotic lesion originates from a single epigenetically deregulated endometrial progenitor cell [27]. Further studies are required to identify cells which are competent for epigenetic changes following GEN exposure, and to elucidate the relationships between these cells, GEN, and endometriosis.

In conclusion, we demonstrated that GEN demethylates the promoter region of the SF-1 gene. This is the first demonstration of phytoestrogen participation in epigenetic alterations in adult endometrial tissue. These findings are important from standpoints of nutrition, public health, and disease prevention. Further study is warranted to characterize the nature of the cells that respond to GEN in the endometrium.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.bbrc.2011.07.104.

References

- [1] H. Adlercreutz, Phyto-oestrogens and cancer, *Lancet Oncol.* 3 (2002) 364–373.
- [2] M. Messina, W. McCaskill-Stevens, J.W. Lampe, Addressing the soy and breast cancer relationship: review, commentary, and workshop proceedings, *J. Natl. Cancer Inst.* 98 (2006) 1275–1284.
- [3] A.H. Wu, R.G. Ziegler, A.M. Nomura, D.W. West, L.N. Kolonel, P.L. Horn-Ross, R.N. Hoover, M.C. Pike, Soy intake and risk of breast cancer in Asians and Asian Americans, *Am. J. Clin. Nutr.* 68 (1998) 1437S–1443S.
- [4] J. Kanno, L. Onyon, S. Peddada, J. Ashby, E. Jacob, W. Owens, The OECD program to validate the rat uterotrophic bioassay. Phase 2: dose-response studies, *Environ. Health Perspect.* 111 (2003) 1530–1549.
- [5] K. Pettersson, J.A. Gustafsson, Role of estrogen receptor beta in estrogen action, *Annu. Rev. Physiol.* 63 (2001) 165–192.
- [6] P. Diel, T. Hertrampf, J. Seibel, U. Laudenschow, S. Kolba, G. Vollmer, Combinatorial effects of the phytoestrogen genistein and of estradiol in uterus and liver of female Wistar rats, *J. Steroid Biochem. Mol. Biol.* 102 (2006) 60–70.
- [7] V. Beck, U. Rohr, A. Jungbauer, Phytoestrogens derived from red clover: an alternative to estrogen replacement therapy?, *J. Steroid Biochem. Mol. Biol.* 94 (2005) 499–518.
- [8] N. Sato, N. Yamakawa, M. Masuda, K. Sudo, I. Hatada, M. Muramatsu, Genome-wide DNA methylation analysis reveals phytoestrogen modification of promoter methylation patterns during embryonic stem cell differentiation, *PLoS One* 6 (2011) e19278.
- [9] S.M. Meeran, A. Ahmed, T.O. Tollefsbol, Epigenetic targets of bioactive dietary components for cancer prevention and therapy, *Clin. Epigenetics* 1 (2010) 101–116.
- [10] S. Majid, A.A. Dar, V. Shahryari, H. Hirata, A. Ahmad, S. Saini, Y. Tanaka, A.V. Dahiya, R. Dahiya, Genistein reverses hypermethylation and induces active histone modifications in tumor suppressor gene B-Cell translocation gene 3 in prostate cancer, *Cancer* 116 (2010) 66–76.
- [11] A.K. Jha, M. Nikbakht, G. Parashar, A. Shrivastava, N. Capalash, J. Kaur, Reversal of hypermethylation and reactivation of the RARbeta2 gene by natural compounds in cervical cancer cell lines, *Folia. Biol. (Praha)* 56 (2010) 195–200.
- [12] Y. Li, T.O. Tollefsbol, Impact on DNA methylation in cancer prevention and therapy by bioactive dietary components, *Curr. Med. Chem.* 17 (2010) 2141–2151.
- [13] K. Morohashi, S. Honda, Y. Inomata, H. Handa, T. Omura, A common trans-acting factor, Ad4-binding protein, to the promoters of steroidogenic P-450s, *J. Biol. Chem.* 267 (1992) 17913–17919.
- [14] Q. Xue, Z. Lin, P. Yin, M.P. Milad, Y.H. Cheng, E. Confino, S. Reierstad, S.E. Bulun, Transcriptional activation of steroidogenic factor-1 by hypomethylation of the 5' CpG island in endometriosis, *J. Clin. Endocrinol. Metab.* 92 (2007) 3261–3267.
- [15] S.E. Bulun, H. Utsunomiya, Z. Lin, P. Yin, Y.H. Cheng, M.E. Pavone, H. Tokunaga, E. Trukhacheva, E. Attar, B. Gurates, M.P. Milad, E. Confino, E. Su, S. Reierstad, Q. Xue, Steroidogenic factor-1 and endometriosis, *Mol. Cell. Endocrinol.* 300 (2009) 104–108.
- [16] J. Kanno, K. Aisaki, K. Igarashi, N. Nakatsu, A. Ono, Y. Kodama, T. Nagao, "Per cell" normalization method for mRNA measurement by quantitative PCR and microarrays, *BMC Genomics* 7 (2006) 64.
- [17] R.W. Chan, K.E. Schwab, C.E. Gargett, Clonogenicity of human endometrial epithelial and stromal cells, *Biol. Reprod.* 70 (2004) 1738–1750.
- [18] C.E. Gargett, K.E. Schwab, R.M. Zillwood, H.P. Nguyen, D. Wu, Isolation and culture of epithelial progenitors and mesenchymal stem cells from human endometrium, *Biol. Reprod.* 80 (2009) 1136–1145.
- [19] Y. Kumaki, M. Oda, M. Okano, QUMA: quantification tool for methylation analysis, *Nucleic Acids Res.* 36 (2008) W170–W175.
- [20] J.A. McKay, J.C. Mathers, Diet induced epigenetic changes and their implications for health, *Acta Physiol. (Oxf)* (2011) 103–118.
- [21] B.C. Christensen, E.A. Houseman, C.J. Marsit, S. Zheng, M.R. Wrensch, J.L. Wiemels, H.H. Nelson, M.R. Karagas, J.F. Padbury, R. Bueno, D.J. Sugarbaker, R.F. Yeh, J.K. Wiencke, K.T. Kelsey, Aging and environmental exposures alter tissue-specific DNA methylation dependent upon CpG island context, *PLoS Genet.* 5 (2009) e1000602.
- [22] C.B. Klein, A.A. King, Genistein genotoxicity: critical considerations of in vitro exposure dose, *Toxicol. Appl. Pharmacol.* 224 (2007) 1–11.
- [23] G. Rimoldi, J. Christoffel, D. Seidlova-Wuttke, H. Jarry, W. Wuttke, Effects of chronic genistein treatment in mammary gland, uterus, and vagina, *Environ. Health Perspect.* 115 (Suppl. 1) (2007) 62–68.
- [24] K.M. Edmunds, A.C. Holloway, D.J. Crankshaw, S.K. Agarwal, W.G. Foster, The effects of dietary phytoestrogens on aromatase activity in human endometrial stromal cells, *Reprod. Nutr. Dev.* 45 (2005) 709–720.
- [25] C.E. Gargett, H. Masuda, Adult stem cells in the endometrium, *Mol. Hum. Reprod.* 16 (2010) 818–834.
- [26] S.W. Guo, Epigenetics of endometriosis, *Mol. Hum. Reprod.* 15 (2009) 587–607.
- [27] Y. Wu, Z. Basir, A. Kajdacsy-Balla, E. Strawn, V. Macias, K. Montgomery, S.W. Guo, Resolution of clonal origins for endometriotic lesions using laser capture microdissection and the human androgen receptor (HUMARA) assay, *Fertil. Steril.* 79 (Suppl. 1) (2003) 710–717.

PKA-dependent regulation of the histone lysine demethylase complex PHF2–ARID5B

Atsushi Baba¹, Fumiaki Ohtake^{1,2}, Yosuke Okuno^{1,2}, Kenichi Yokota¹, Maiko Okada^{1,2}, Yuuki Imai^{1,3}, Min Ni³, Clifford A. Meyer⁴, Katsuhide Igarashi⁵, Jun Kanno⁵, Myles Brown³ and Shigeaki Kato^{1,2,6}

Reversible histone methylation and demethylation are highly regulated processes that are crucial for chromatin reorganization and regulation of gene transcription in response to extracellular conditions. However, the mechanisms that regulate histone-modifying enzymes are largely unknown. Here, we characterized a protein kinase A (PKA)-dependent histone lysine demethylase complex, PHF2–ARID5B. PHF2, a *jmjC* demethylase, is enzymatically inactive by itself, but becomes an active H3K9Me2 demethylase through PKA-mediated phosphorylation. We found that phosphorylated PHF2 then associates with ARID5B, a DNA-binding protein, and induce demethylation of methylated ARID5B. This modification leads to targeting of the PHF2–ARID5B complex to its target promoters, where it removes the repressive H3K9Me2 mark. These findings suggest that the PHF2–ARID5B complex is a signal-sensing modulator of histone methylation and gene transcription, in which phosphorylation of PHF2 enables subsequent formation of a competent and specific histone demethylase complex.

Post-translational modifications of histone amino-terminal tails, including reversible acetylation, methylation, phosphorylation and ubiquitylation, modulate both chromatin structure and gene regulation^{1,2}. Reversible histone methylation defines the state of chromatin; repressive histone methylation marks such as H3K9Me2 inhibit gene transcription from nearby promoters, and removal of these marks by histone lysine demethylases permit transcription. Thus, enzymes such as *jmjC* (Jumonji C)-domain-containing histone lysine demethylases seem to play pivotal roles in epigenetic control^{3–10}. Though histone marks are altered in response to extracellular signals, the underlying mechanism by which this is achieved is largely unknown.

RESULTS

Identification of the PHF2–ARID5B histone H3K9Me2 demethylase complex

During biochemical identification of transcriptional co-regulators for nuclear receptors^{11,12}, we purified a number of direct interactants for nuclear receptors. Among them, a co-activator complex for FXR (farnesoid X receptor) from HepG2 cells was selected for further analysis, because its abundance in the purified fractions was significantly increased when the cells were pretreated with forskolin (FSK), a protein kinase A (PKA) activator (Supplementary Fig. S1a). Matrix-assisted laser desorption/ionization–time of flight mass

spectrometry analysis of the purified complex^{11,12} identified three subunits: plant homeodomain (PHD) finger two (PHF2; ref. 13), AT-rich interactive domain 5B (ARID5B) isoform α and ARID5B isoform β (ref. 14; Fig. 1a and Supplementary Fig. S1b–e). The complex was still detectable after further column purification (Supplementary Fig. S1f). ARID5B contains the ARID/Bright domain, which is seen in subunits of chromatin-remodelling complexes and is considered to be a sequence-specific/nonspecific DNA-attachment domain¹⁴. PHF2 (ref. 13) contains PHD and *jmjC* domains^{4,6}. Unlike several other *jmjC* histone demethylases^{3,5,8,9}, PHF2 lacks the ARID domain. PHF2 and ARID5B genes are expressed in a broad range of tissues, including liver (Supplementary Fig. S2c), and at protein levels as well in various cell lines (Supplementary Fig. S2a,b,d). PHF2 physically interacted with FXR and HNF4 α (hepatocyte nuclear factor 4 α ; Supplementary Fig. S3a,b,d), but not with ER α (oestrogen receptor alpha) or VDR (vitamin D receptor; Supplementary Fig. S3c,d). PHF2 directly interacted with ARID5B *in vitro* (Supplementary Fig. S3e–g).

The PHF2–ARID5B complex was purified from HepG2 cells stably expressing Flag–PHF2. As *jmjC*-domain-containing proteins catalyse lysine demethylation^{3–10}, we investigated the histone lysine demethylation activity of the purified PHF2 complex^{7,10}. The PHF2 complex exhibited lysine demethylase activity *in vitro* as assessed by a formaldehyde release (Fig. 1b). Then we examined

¹Institute of Molecular and Cellular Biosciences, University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-0032, Japan. ²ERATO, Japan Science and Technology Agency, 4-1-8 Honcho, Kawaguchishi, Saitama 332-0012, Japan. ³Department of Medical Oncology, Dana-Farber Cancer Institute and Harvard Medical School, Boston, Massachusetts 02115, USA. ⁴Department of Biostatistics and Computational Biology, Dana-Farber Cancer Institute and Harvard School of Public Health, Boston, Massachusetts 02115, USA. ⁵Division of Cellular and Molecular Toxicology, National Institute of Health Sciences, 1-18-1 Kamiyoga, Setagaya-ku, Tokyo 158-8501, Japan.

⁶Correspondence should be addressed to S.K. (e-mail: uskato@mail.ecc.u-tokyo.ac.jp)




Figure 1 Identification of the PHF2–ARID5B histone H3K9Me2 demethylase complex. (a) A schematic representation of domains of PHF2 and ARID5B. (b–d) Histone H3K9Me2 demethylase activity of the purified PHF2 complex *in vitro*. (b) Native histones were incubated with PHF2 complex, and formaldehyde release was analysed by fluorescence detection. Shown is the average \pm s.d. ($n = 3$) of the relative activity. (c) The PHF2 complex was incubated with purified native histones (left) or mononucleosomes (right) for demethylation reaction, and each modification was determined by western blotting as indicated. (d) Intensity of histone marks in the demethylation assay of native histones by PHF2 (in c and Supplementary Fig. S8a) was quantified, and averages \pm s.d. of three independent experiments are shown. The asterisk shows $P < 0.05$ in Student's *t*-test. (e) Recombinant PHF2

the site-specificity of lysine demethylation using native histones or purified mononucleosomes as substrates. *In vitro*, the PHF2 complex demethylated dimethylated Lys 9 on histone H3 (H3K9Me2), but neither mono- nor trimethylated H3K9 (H3K9Me1, Me3) (Fig. 1c,d). Methylated H3K4, K27, K36 and K79 and H4K20 seemed unlikely PHF2 substrates (Fig. 1c,d). Thus it is likely that PHF2 substrate recognition is more specific than the closely related PHF subfamily members such as PHF8 and KDM7/KIAA1718 (refs 15–17). This demethylase activity was not detected in recombinant PHF2 protein (Fig. 1e). PHF2 did not demethylate a methylated peptide (amino acids 1–21 of H3) containing K9Me2 (Supplementary Fig. S4a). Such substrate specificity is known for Lid2 (ref. 18) and yeast Lsd1 (ref. 19), and possibly due to requirement of other regions of H3. Immunostaining analysis of 293F cells confirmed that PHF2 almost entirely demethylated H3K9Me2 in the presence of FSK, but had only marginal effects on H3K27Me2 and the other tested methylated

protein is catalytically inactive. Recombinant PHF2 protein (lower panel) was subjected to *in vitro* demethylation assay (upper panels). (f) 293F cells were transfected with Flag–PHF2 expression vectors for 24 h and treated with FSK for 6 h, then cells were fixed and immunostained with the indicated antibodies. Note the decreased level of H3K9Me2 in the anti-Flag-positive cells (arrowheads). Scale bars, 40 μ m. (g) 293F cells were transfected with either the Flag–PHF2 Δ JmjC or the HA–PHF2^{H249A} mutant, and immunostaining was carried out as in f. Scale bars, 40 μ m. (h) H3K9Me2 demethylation activity of PHF2 *in vivo*. 293F cells transfected with the indicated expression vectors or short interfering RNAs (siRNAs), and were treated with FSK for 6 h. The soluble chromatin fraction was subjected to western blotting as indicated. Uncropped images of blots are shown in Supplementary Fig. S9.

histones (Fig. 1f and Supplementary Fig. S4b). The conserved histidine residue within the PHF2 jmjC domain was required for enzymatic activity (Fig. 1g). In 293F cells, overexpression of PHF2 in the presence of FSK decreased H3K9Me2, whereas knockdown of PHF2 did not lead to global increase of this mark (Fig. 1h), raising the possibility that other H3K9 demethylase(s) may compensate for PHF2 in these cells.

PHF2 is phosphorylated by PKA at a specific serine residue

In a co-immunoprecipitation assay, association of PHF2 and ARID5B was inducible on FSK treatment of immortalized hepatocytes²⁰ both in the absence (Fig. 2a) and presence (Supplementary Fig. S4c) of a proteasomal inhibitor, MG132. Consistently, abundance of these proteins was not clearly affected by FSK treatment (Supplementary Fig. S4d). Therefore, we reasoned that phosphorylation of the factor(s) by PKA was a trigger to induce association. Endogenous PHF2, but not ARID5B, was phosphorylated by FSK treatment of the cells (Fig. 2b,c).




Figure 2 PKA-dependent complex assembly of PHF2-ARID5B. (a) Assembly of the endogenous PHF2-ARID5B complex. Hepatocytes were treated with the indicated compounds for 2 h and were subjected to immunoprecipitation (IP) and immunoblotting (IB) with the indicated antibodies. Specificity of antibodies was confirmed in Supplementary Fig. S2a,b. (b,c) Phosphorylation of endogenous PHF2 by PKA. Hepatocytes (b) or 293F cells (c) were treated with vehicle, FSK, or FSK + H89 as indicated for 2 h, then lysates were immunoprecipitated using either anti-phospho-serine/threonine or anti-PHF2 antibodies. The eluates were subjected to western blotting as indicated. (d-f) 293F

FSK-dependent PHF2 phosphorylation was negated by the PKA inhibitor H89, but not by a MAP kinase inhibitor (MAPKi; Fig. 2d). Cyclohexamide (CHX) did not attenuate PHF2 phosphorylation, indicating that this event does not mediate *de novo* protein synthesis (Fig. 2e). Knockdown of PKA $\alpha/\beta/\gamma$, but not of extracellular signal-regulated kinases ERK1/2, abrogated PHF2 phosphorylation (Fig. 2d,f and Supplementary Fig. S5a). PKA phosphorylated recombinant PHF2 in *in vitro* phosphorylation assays (Fig. 2g). Using recombinant PHF2 deletion mutants, the carboxy-terminal region of PHF2 was mapped to be phosphorylated by PKA (Supplementary Fig. S5b,d), and bears four conserved, PKA consensus serine residues (Supplementary Fig. S5c). Replacement of all of four serine residues by alanines (4SA: Ser 757/Ser 899/Ser 954/Ser 1056) fully abrogated PKA phosphorylation of PHF2 (Fig. 2h). Further mapping revealed that Ser 1056 seemed to be a major phosphorylation site (Fig. 2h).

PKA-dependent demethylase activity of PHF2-ARID5B complex

We then asked if the demethylase activity of PHF2 for H3K9Me2 was also PKA dependent with proteins purified from 293F cells as well as recombinant proteins. Purified PHF2 protein, phosphorylated by PKA *in vitro*, exhibited H3K9Me2 demethylase activity (Fig. 3a-c). Conversely, PHF2 phosphorylated by FSK treatment in cells exhibited

cells transfected with Flag-PHF2 and indicated siRNAs were treated with the indicated inhibitors, and immunoprecipitated with anti-Flag antibodies. Immunoprecipitates were subjected to immunoblotting with an anti-phospho-PKA substrate antibody. (d, lower panel) Averages \pm s.d. The other data sets are shown in Supplementary Fig. S8b,c. The asterisks show $P < 0.05$ in Student's *t*-test. (g) *In vitro* kinase assay with recombinant PHF2 and PKA proteins. (h) PHF2 is phosphorylated by PKA at Ser 757/899/954/1056. In the 4SA mutants, Ser 757/899/954/1056 were replaced with alanines. Uncropped images of blots are shown in Supplementary Fig. S9.

the demethylase activity, but *in vitro* dephosphorylation by BAP (bacterial alkaline phosphatase) negated the demethylase activity (Fig. 3b,c). ARID5B was dispensable for the enzymatic activity of PHF2 *in vitro* (Fig. 3b,c), because ARID5B was absent from the purified PHF2 protein fraction used (Supplementary Fig. S5f). Similarly, recombinant PHF2 was activated by PKA phosphorylation (Fig. 3d-f). PHF2 phosphorylation mutants were not activated by PKA (Fig. 3g). Consistent with the *in vitro* analyses, H3K9Me2 demethylation by overexpression of PHF2-ARID5B in cells was significantly enhanced by FSK treatment, but 4SA was not activated by FSK (Fig. 3h). Subnuclear localization patterns of the PHF2^{4SA} mutant did not look identical to that of wild-type PHF2 (Fig. 3h), raising the possibility that PHF2 phosphorylation is indispensable for chromatin association. Together, these findings suggest that PKA-mediated phosphorylation induces the demethylase activity of PHF2, unlike the other characterized jmjC demethylases exhibiting constitutive enzymatic activities both *in vivo* and *in vitro* as recombinant proteins⁴⁻¹⁰.

ARID5B directs PKA-dependent promoter targeting of PHF2

We examined how PKA-mediated activation of the PHF2-ARID5B complex activated transcription at endogenous promoters. In immortalized hepatocytes, glucagon-PKA signalling regulates glucose