

- for IL-21-induced secretion of IgE from human naive B cells. *Blood*. 112:1784–1793. <http://dx.doi.org/10.1182/blood-2008-02-142745>
- Avery, D.T., E.K. Deenick, C.S. Ma, S. Suryani, N. Simpson, G.Y. Chew, T.D. Chan, U. Palendira, J. Bustamante, S. Boisson-Dupuis, et al. 2010. B cell-intrinsic signaling through IL-21 receptor and STAT3 is required for establishing long-lived antibody responses in humans. *J. Exp. Med.* 207:155–171. <http://dx.doi.org/10.1084/jem.20091706>
- Banchereau, J., F. Brière, Y.J. Liu, and F. Rousset. 1994. Molecular control of B lymphocyte growth and differentiation. *Stem Cells*. 12:278–288. <http://dx.doi.org/10.1002/stem.5530120304>
- Bernasconi, N.L., E. Traggiai, and A. Lanzavecchia. 2002. Maintenance of serological memory by polyclonal activation of human memory B cells. *Science*. 298:2199–2202. <http://dx.doi.org/10.1126/science.1076071>
- Bernasconi, N.L., N. Onai, and A. Lanzavecchia. 2003. A role for Toll-like receptors in acquired immunity: up-regulation of TLR9 by BCR triggering in naive B cells and constitutive expression in memory B cells. *Blood*. 101:4500–4504. <http://dx.doi.org/10.1182/blood-2002-11-3569>
- Boisson-Dupuis, S., X.F. Kong, S. Okada, S. Cypowyj, A. Puel, L. Abel, and J.L. Casanova. 2012. Inborn errors of human STAT1: allelic heterogeneity governs the diversity of immunological and infectious phenotypes. *Curr. Opin. Immunol.* 24:364–378. <http://dx.doi.org/10.1016/j.coi.2012.04.011>
- Bryant, V.L., C.S. Ma, D.T. Avery, Y. Li, K.L. Good, L.M. Corcoran, R. de Waal Malefyt, and S.G. Tangye. 2007. Cytokine-mediated regulation of human B cell differentiation into Ig-secreting cells: predominant role of IL-21 produced by CXCR5+ T follicular helper cells. *J. Immunol.* 179:8180–8190.
- Casanova, J.L., S.M. Holland, and L.D. Notarangelo. 2012. Inborn errors of human JAKs and STATs. *Immunity*. 36:515–528. <http://dx.doi.org/10.1016/j.immuni.2012.03.016>
- Chapgier, A., S. Boisson-Dupuis, E. Jouanguy, G. Vogt, J. Feinberg, A. Prochnicka-Chaloufour, A. Casrouge, K. Yang, C. Soudais, C. Fieschi, et al. 2006. Novel STAT1 alleles in otherwise healthy patients with mycobacterial disease. *PLoS Genet.* 2:e131. <http://dx.doi.org/10.1371/journal.pgen.0020131>
- Chapgier, A., X.F. Kong, S. Boisson-Dupuis, E. Jouanguy, D. Averbuch, J. Feinberg, S.Y. Zhang, J. Bustamante, G. Vogt, J. Lejeune, et al. 2009. A partial form of recessive STAT1 deficiency in humans. *J. Clin. Invest.* 119:1502–1514. <http://dx.doi.org/10.1172/JCI37083>
- Chevalier, N., D. Jarrossay, E. Ho, D.T. Avery, C.S. Ma, D. Yu, F. Sallusto, S.G. Tangye, and C.R. Mackay. 2011. CXCR5 expressing human central memory CD4 T cells and their relevance for humoral immune responses. *J. Immunol.* 186:5556–5568. <http://dx.doi.org/10.4049/jimmunol.1002828>
- Craxton, A., D. Magaletti, E.J. Ryan, and E.A. Clark. 2003. Macrophage- and dendritic cell-dependent regulation of human B-cell proliferation requires the TNF family ligand BAFF. *Blood*. 101:4464–4471. <http://dx.doi.org/10.1182/blood-2002-10-3123>
- Darce, J.R., B.K. Arendt, X. Wu, and D.F. Jelinek. 2007. Regulated expression of BAFF-binding receptors during human B cell differentiation. *J. Immunol.* 179:7276–7286.
- Davey, A.M., and S.K. Pierce. 2012. Intrinsic differences in the initiation of B cell receptor signaling favor responses of human IgG(+) memory B cells over IgM(+) naive B cells. *J. Immunol.* 188:3332–3341. <http://dx.doi.org/10.4049/jimmunol.1102322>
- Deenick, E.K., J. Hasbold, and P.D. Hodgkin. 1999. Switching to IgG3, IgG2b, and IgA is division linked and independent, revealing a stochastic framework for describing differentiation. *J. Immunol.* 163:4707–4714.
- Diehl, S.A., H. Schmidlin, M. Nagasawa, S.D. van Haren, M.J. Kwakkenbos, E. Yasuda, T. Beaumont, F.A. Scheeren, and H. Spits. 2008. STAT3-mediated up-regulation of BLIMP1 is coordinated with BCL6 down-regulation to control human plasma cell differentiation. *J. Immunol.* 180:4805–4815.
- Dupuis, S., C. Dargemont, C. Fieschi, N. Thomassin, S. Rosenzweig, J. Harris, S.M. Holland, R.D. Schreiber, and J.L. Casanova. 2001. Impairment of mycobacterial but not viral immunity by a germline human STAT1 mutation. *Science*. 293:300–303. <http://dx.doi.org/10.1126/science.1061154>
- Ellyard, J.I., D.T. Avery, T.G. Phan, N.J. Hare, P.D. Hodgkin, and S.G. Tangye. 2004. Antigen-selected, immunoglobulin-secreting cells persist in human spleen and bone marrow. *Blood*. 103:3805–3812. <http://dx.doi.org/10.1182/blood-2003-09-3109>
- Engels, N., L.M. König, C. Heemann, J. Lutz, T. Tsubata, S. Griep, V. Schrader, and J. Wienands. 2009. Recruitment of the cytoplasmic adaptor Grb2 to surface IgG and IgE provides antigen receptor-intrinsic costimulation to class-switched B cells. *Nat. Immunol.* 10:1018–1025. <http://dx.doi.org/10.1038/ni.1764>
- Erttinger, R., G.P. Sims, A.M. Fairhurst, R. Robbins, Y.S. da Silva, R. Spolski, W.J. Leonard, and P.E. Lipsky. 2005. IL-21 induces differentiation of human naive and memory B cells into antibody-secreting plasma cells. *J. Immunol.* 175:7867–7879.
- Geha, R.S., H.H. Jabara, and S.R. Brodeur. 2003. The regulation of immunoglobulin E class-switch recombination. *Nat. Rev. Immunol.* 3:721–732. <http://dx.doi.org/10.1038/nri1181>
- Good, K.L., and S.G. Tangye. 2007. Decreased expression of Kruppel-like factors in memory B cells induces the rapid response typical of secondary antibody responses. *Proc. Natl. Acad. Sci. USA*. 104:13420–13425. <http://dx.doi.org/10.1073/pnas.0703872104>
- Good, K.L., V.L. Bryant, and S.G. Tangye. 2006. Kinetics of human B cell behavior and amplification of proliferative responses following stimulation with IL-21. *J. Immunol.* 177:5236–5247.
- Good, K.L., D.T. Avery, and S.G. Tangye. 2009. Resting human memory B cells are intrinsically programmed for enhanced survival and responsiveness to diverse stimuli compared to naive B cells. *J. Immunol.* 182:890–901.
- Goodnow, C.C., C.G. Vinuesa, K.L. Randall, F. Mackay, and R. Brink. 2010. Control systems and decision making for antibody production. *Nat. Immunol.* 11:681–688. <http://dx.doi.org/10.1038/ni.1900>
- Griffin, D.O., N.E. Holodick, and T.L. Rothstein. 2011. Human B1 cells in umbilical cord and adult peripheral blood express the novel phenotype CD20+ CD27+ CD43+ CD70-. *J. Exp. Med.* 208:67–80. <http://dx.doi.org/10.1084/jem.20101499>
- Hirata, O., S. Okada, M. Tsumura, R. Kagawa, M. Miki, H. Kawaguchi, K. Nakamura, S. Boisson-Dupuis, J.L. Casanova, Y. Takihara, and M. Kobayashi. 2013. Heterozygosity for the Y701C STAT1 mutation in a multiplex kindred with multifocal osteomyelitis. *Haematologica*. 98:1641–1649. <http://dx.doi.org/10.3324/haematol.2013.083741>
- Hodgkin, P.D., J.H. Lee, and A.B. Lyons. 1996. B cell differentiation and isotype switching is related to division cycle number. *J. Exp. Med.* 184:277–281. <http://dx.doi.org/10.1084/jem.184.1.277>
- Holland, S.M., F.R. DeLeo, H.Z. Elloumi, A.P. Hsu, G. Uzel, N. Brodsky, A.F. Freeman, A. Demidowich, J. Davis, M.L. Turner, et al. 2007. STAT3 mutations in the hyper-IgE syndrome. *N. Engl. J. Med.* 357:1608–1619. <http://dx.doi.org/10.1056/NEJMoa073687>
- Horikawa, K., S.W. Martin, S.L. Pogue, K. Silver, K. Peng, K. Takatsu, and C.C. Goodnow. 2007. Enhancement and suppression of signaling by the conserved tail of IgG memory-type B cell antigen receptors. *J. Exp. Med.* 204:759–769. <http://dx.doi.org/10.1084/jem.20061923>
- Ives, M.L., C.S. Ma, U. Palendira, A. Chan, J. Bustamante, S. Boisson-Dupuis, P.D. Arkwright, D. Engelhard, D. Averbuch, K. Magdorf, et al. 2013. Signal transducer and activator of transcription 3 (STAT3) mutations underlying autosomal dominant hyper-IgE syndrome impair human CD8(+) T-cell memory formation and function. *J. Allergy Clin. Immunol.* 132:400–411. <http://dx.doi.org/10.1016/j.jaci.2013.05.029>
- Jabara, H.H., D.R. McDonald, E. Janssen, M.J. Massaad, N. Ramesh, A. Borzutzky, I. Rauter, H. Benson, L. Schneider, S. Baxi, et al. 2012. DOCK8 functions as an adaptor that links TLR-MyD88 signaling to B cell activation. *Nat. Immunol.* 13:612–620. <http://dx.doi.org/10.1038/ni.2305>
- Kotlarz, D., R. Beier, D. Murugan, J. Diestelhorst, O. Jensen, K. Boztug, D. Pfeifer, H. Kreipe, E.D. Pfister, U. Baumann, et al. 2012. Loss of interleukin-10 signaling and infantile inflammatory bowel disease: implications for diagnosis and therapy. *Gastroenterology*. 143:347–355. <http://dx.doi.org/10.1053/j.gastro.2012.04.045>
- Kotlarz, D., N. Ziętara, G. Uzel, T. Weidemann, C.J. Braun, J. Diestelhorst, P.M. Krawitz, P.N. Robinson, J. Hecht, J. Puchalka, et al. 2013. Loss-of-function mutations in the IL-21 receptor gene cause a primary immunodeficiency

- syndrome. *J. Exp. Med.* 210:433–443. <http://dx.doi.org/10.1084/jem.20111229>
- Leung, D.Y., D.M. Ambrosino, R.D. Arbeit, J.L. Newton, and R.S. Geha. 1988. Impaired antibody responses in the hyperimmunoglobulin E syndrome. *J. Allergy Clin. Immunol.* 81:1082–1087. [http://dx.doi.org/10.1016/0091-6749\(88\)90873-1](http://dx.doi.org/10.1016/0091-6749(88)90873-1)
- Litinskiy, M.B., B. Nardelli, D.M. Hilbert, B. He, A. Schaffer, P. Casali, and A. Cerutti. 2002. DCs induce CD40-independent immunoglobulin class switching through BLYS and APRIL. *Nat. Immunol.* 3:822–829. <http://dx.doi.org/10.1038/ni829>
- Liu, Y.J., C. Barthélémy, O. de Bouteiller, C. Arpin, I. Durand, and J. Banchereau. 1995. Memory B cells from human tonsils colonize mucosal epithelium and directly present antigen to T cells by rapid up-regulation of B7-1 and B7-2. *Immunity*. 2:239–248. [http://dx.doi.org/10.1016/1074-7613\(95\)90048-9](http://dx.doi.org/10.1016/1074-7613(95)90048-9)
- Ma, C.S., N.J. Hare, K.E. Nichols, L. Dupré, G. Andolfi, M.G. Roncarolo, S. Adelstein, P.D. Hodgkin, and S.G. Tangye. 2005. Impaired humoral immunity in X-linked lymphoproliferative disease is associated with defective IL-10 production by CD4+ T cells. *J. Clin. Invest.* 115:1049–1059.
- Ma, C.S., S. Pittaluga, D.T. Avery, N.J. Hare, I. Maric, A.D. Klion, K.E. Nichols, and S.G. Tangye. 2006. Selective generation of functional somatically mutated IgM+CD27+, but not Ig isotype-switched, memory B cells in X-linked lymphoproliferative disease. *J. Clin. Invest.* 116:322–333. <http://dx.doi.org/10.1172/JCI25720>
- Ma, C.S., G.Y. Chew, N. Simpson, A. Priyadarshi, M. Wong, B. Grimbacher, D.A. Fulcher, S.G. Tangye, and M.C. Cook. 2008. Deficiency of Th17 cells in hyper IgE syndrome due to mutations in STAT3. *J. Exp. Med.* 205:1551–1557. <http://dx.doi.org/10.1084/jem.20080218>
- Ma, C.S., D.T. Avery, A. Chan, M. Batten, J. Bustamante, S. Boisson-Dupuis, P.D. Arkwright, A.Y. Kreins, D. Averbuch, D. Engelhard, et al. 2012. Functional STAT3 deficiency compromises the generation of human T follicular helper cells. *Blood*. 119:3997–4008. <http://dx.doi.org/10.1182/blood-2011-11-392985>
- Macallan, D.C., D.L. Wallace, Y. Zhang, H. Ghattas, B. Asquith, C. de Lara, A. Worth, G. Panayiotakopoulos, G.E. Griffin, D.F. Tough, and P.C. Beverley. 2005. B-cell kinetics in humans: rapid turnover of peripheral blood memory cells. *Blood*. 105:3633–3640. <http://dx.doi.org/10.1182/blood-2004-09-3740>
- Martin, S.W., and C.C. Goodnow. 2002. Burst-enhancing role of the IgG membrane tail as a molecular determinant of memory. *Nat. Immunol.* 3:182–188. <http://dx.doi.org/10.1038/ni752>
- Mazerolles, F., C. Picard, S. Kracker, A. Fischer, and A. Durandy. 2013. Blood CD4+CD45RO+CXCR5+ T cells are decreased but partially functional in signal transducer and activator of transcription 3 deficiency. *J. Allergy Clin. Immunol.* 131:1146–1156. e1–e5. <http://dx.doi.org/10.1016/j.jaci.2012.12.1519>
- Minegishi, Y., M. Saito, S. Tsuchiya, I. Tsuge, H. Takada, T. Hara, N. Kawamura, T. Ariga, S. Pasic, O. Stojkovic, et al. 2007. Dominant-negative mutations in the DNA-binding domain of STAT3 cause hyper-IgE syndrome. *Nature*. 448:1058–1062. <http://dx.doi.org/10.1038/nature06096>
- Nilsson, A., A. de Milito, F. Mowafi, G. Winberg, O. Björk, E.Z. Wolpert, and F. Chioldi. 2005. Expression of CD27–CD70 on early B cell progenitors in the bone marrow: implication for diagnosis and therapy of childhood ALL. *Exp. Hematol.* 33:1500–1507. <http://dx.doi.org/10.1016/j.exphem.2005.10.005>
- Notarangelo, L.D., G. Lanzi, S. Peron, and A. Durandy. 2006. Defects of class-switch recombination. *J. Allergy Clin. Immunol.* 117:855–864. <http://dx.doi.org/10.1016/j.jaci.2006.01.043>
- Nutt, S.L., N. Taubenheim, J. Hasbold, L.M. Corcoran, and P.D. Hodgkin. 2011. The genetic network controlling plasma cell differentiation. *Semin. Immunol.* 23:341–349. <http://dx.doi.org/10.1016/j.smim.2011.08.010>
- Pène, J., J.F. Gauchat, S. Lécart, E. Drouet, P. Guglielmi, V. Boulay, A. Delwail, D. Foster, J.C. Lecron, and H. Yssel. 2004. Cutting edge: IL-21 is a switch factor for the production of IgG1 and IgG3 by human B cells. *J. Immunol.* 172:5154–5157.
- Poudrier, J., and T. Owens. 1994. Co-stimulation by anti-immunoglobulin is required for B cell activation by CD40low T cells. *Eur. J. Immunol.* 24:2993–2999. <http://dx.doi.org/10.1002/eji.1830242121>
- Rajewsky, K. 1996. Clonal selection and learning in the antibody system. *Nature*. 381:751–758. <http://dx.doi.org/10.1038/381751a0>
- Recher, M., L.J. Berglund, D.T. Avery, M.J. Cowan, A.R. Gennery, J. Smart, J. Peake, M. Wong, S.Y. Pai, S. Baxi, et al. 2011. IL-21 is the primary common γ chain-binding cytokine required for human B-cell differentiation in vivo. *Blood*. 118:6824–6835. <http://dx.doi.org/10.1182/blood-2011-06-362533>
- Sampaio, E.P., H.I. Bax, A.P. Hsu, E. Kristosturyan, J. Pechacek, P. Chandrasekaran, M.L. Paulson, D.L. Dias, C. Spalding, G. Uzel, et al. 2012. A novel STAT1 mutation associated with disseminated mycobacterial disease. *J. Clin. Immunol.* 32:681–689. <http://dx.doi.org/10.1007/s10875-012-9659-2>
- Sheerin, K.A., and R.H. Buckley. 1991. Antibody responses to protein, polysaccharide, and phi X174 antigens in the hyperimmunoglobulinemia E (hyper-IgE) syndrome. *J. Allergy Clin. Immunol.* 87:803–811. [http://dx.doi.org/10.1016/0091-6749\(91\)90126-9](http://dx.doi.org/10.1016/0091-6749(91)90126-9)
- Tangye, S.G. 2011. Staying alive: regulation of plasma cell survival. *Trends Immunol.* 32:595–602. <http://dx.doi.org/10.1016/j.it.2011.09.001>
- Tangye, S.G., and D.M. Tarlinton. 2009. Memory B cells: effectors of long-lived immune responses. *Eur. J. Immunol.* 39:2065–2075. <http://dx.doi.org/10.1002/eji.200939531>
- Tangye, S.G., Y.J. Liu, G. Aversa, J.H. Phillips, and J.E. de Vries. 1998. Identification of functional human splenic memory B cells by expression of CD148 and CD27. *J. Exp. Med.* 188:1691–1703. <http://dx.doi.org/10.1084/jem.188.9.1691>
- Tangye, S.G., D.T. Avery, E.K. Deenick, and P.D. Hodgkin. 2003a. Intrinsic differences in the proliferation of naive and memory human B cells as a mechanism for enhanced secondary immune responses. *J. Immunol.* 170:686–694.
- Tangye, S.G., D.T. Avery, and P.D. Hodgkin. 2003b. A division-linked mechanism for the rapid generation of Ig-secreting cells from human memory B cells. *J. Immunol.* 170:261–269.
- Tangye, S.G., E.K. Deenick, U. Palendira, and C.S. Ma. 2012. T cell–B cell interactions in primary immunodeficiencies. *Ann. N. Y. Acad. Sci.* 1250:1–13. <http://dx.doi.org/10.1111/j.1749-6632.2011.06361.x>
- Warnatz, K., L. Bossaller, U. Salzer, A. Skrabl-Baumgartner, W. Schwinger, M. van der Burg, J.J. van Dongen, M. Orlowska-Volk, R. Knoth, A. Durandy, et al. 2006. Human ICOS deficiency abrogates the germinal center reaction and provides a monogenic model for common variable immunodeficiency. *Blood*. 107:3045–3052. <http://dx.doi.org/10.1182/blood-2005-07-2955>
- Yefenof, E., V.M. Sanders, J.W. Uhr, and E.S. Vitetta. 1986. In vitro activation of murine antigen-specific memory B cells by a T-dependent antigen. *J. Immunol.* 137:85–90.
- Zeng, R., R. Spolski, E. Casas, W. Zhu, D.E. Levy, and W.J. Leonard. 2007. The molecular basis of IL-21-mediated proliferation. *Blood*. 109:4135–4142. <http://dx.doi.org/10.1182/blood-2006-10-054973>

