

Yozawa R, Inamasu E, Kojo M, Morodomi Y, Shiraishi Y, Takenaka T, Yamaguchi M, Shimokawa M, Seto T, Ichinose Y	and treatment in patients with malignant pleural mesothelioma pretreated with platinum plus pemetrexed chemotherapy.			3-10619-10145	
Toyokawa G, Takenoyama M, Ichinose Y	Do mutations of the enhancer of zeste homolog 2 gene exist in small-cell lung cancer?	J Thorac Oncol	8	e103	2013
Hishida,T., Yoshida,J.,Ohe,Y., Aokage,K., Ishii,G., Nagai,K.	Surgical Outcomes after Initial Surgery for Clinical Single-station N2 Non-small-cell Lung Cancer.	Jpn J Clin Oncol	44 (1)	85-92	2014
Ichinokawa,H., Ishii,G., Nagai,K., Kawase,A., Yoshida,J., Nishimura,M., Hishida,T., Ogasawara,N., Tsuchihara,K., Ochiai,A.	Distinct clinicopathologic characteristics of lung mucinous adenocarcinoma with KRAS mutation.	Hum Pathol	44 (12)	2636-42	2013
Kinoshita,T., Yoshida,J., Ishii,G., Aokage,K., Hishida,T., Nagai,K.	The differences of biological behavior based on the clinicopathological data between resectable large-cell neuroendocrine carcinoma and small-cell lung carcinoma.	Clin Lung Cancer	14 (5)	535-40	2013
Kaseda,K., Ishii,G., Aokage,K., Takahashi,A., Kuwata,T., Hishida,T., Yoshida,J., Kohno,M., Nagai,K., Ochiai,A.	Identification of intravascular tumor microenvironment features predicting the recurrence of pathological stage I lung adenocarcinoma.	Cancer Sci	104 (9)	1262-9	2013
Aokage,K., Yoshida,J., Ishii,G., Matsumura,Y., Haruki,T., Hishida,T., Nagai,K.	Identification of early T1b lung adenocarcinoma based on thin-section computed tomography findings.	J Thorac Oncol	8 (10)	1289-94	2013
Kawase,A., Yoshida,J., Miyaoka,E., Asamura,H., Fujii,Y., Nakanishi,Y., Eguchi,K., Mori,M., Sawabata,N., Okumura,M., Yokoi,K., Japanese Joint Committee of Lung Cancer Registry.	Visceral pleural invasion classification in non-small-cell lung cancer in the 7th edition of the tumor, node, metastasis classification for lung cancer: validation analysis based on a large-scale nationwide database.	J Thorac Oncol	8 (5)	606-11	2013
Kinoshita,T., Ishii,G., Hiraoka,N., Hirayama,S., Yamauchi,C., Aokage,K., Hishida,T., Yoshida,J., Nagai,K., Ochiai,A.	Forkhead box P3 regulatory T cells coexisting with cancer associated fibroblasts are correlated with a poor outcome in lung adenocarcinoma.	Cancer Sci	104 (4)	409-15	2013
Ono,S., Ishii,G., Nagai,K., Takuwa,T., Yoshida,J., Nishimura,M., Hishida,T., Aokage,K., Fujii,S., Ikeda,N., Ochiai,A.	Podoplanin-positive cancer-associated fibroblasts could have prognostic value independent of cancer cell phenotype in stage I lung squamous cell carcinoma: usefulness of combi	Chest	143 (4)	963-70	2013

	ning analysis of both cancer cell phenotype and cancer-associated fibroblast phenotype.				
Hishida,T., Yoshida, J., Maeda,R., Ishii,G., Aokage,K., Nishimura,M., Nagai,K.	Prognostic impact of intratumoural microvascular invasion and microlymphatic permeation on node-negative non-small-cell lung cancer: which indicator is the stronger prognostic factor?	Eur J Cardiothorac Surg	43 (4)	772-7	2013
Kumanogoh A and Kikutani H.	Immunological functions of the neuropilins and plexins as receptors for semaphorin.	Nature Reviews Immunology	13 Pages:	802	2013
Nojima S, Toyofuku T, Kamao H, Ishigami C, Kaneko J, Okuno T, Takamatsu H, Ito D, Kang S, Kimura T, Yoshida Y, Morimoto K, Maeda Y, Ogata A, Ikawa M, Morii E, Aozasa K, Takagi J, Takahashi M, Kumanogoh A.	A point mutation in Semaphorin 4A associates with defective endosomal sorting and causes retinal degeneration.	Nature communications.	4	1406	2013
Jin Y, Tachibana I, Takeda Y, He P, Kang S, Suzuki M, Kuhara H, Tetsumoto S, Tsujino K, Minami T, Iwasaki T, Nakanishi K, Kohmo S, Hirata H, Takahashi R, Inoue K, Nagatomo I, Kida H, Kijima T, Ito M, Saya H, Kumanogoh A.	Statins Decrease Lung Inflammation in Mice by Upregulating Tetraspanin CD9 in Macrophages.	PloS one	8(9)	e73706	2013
Minami T, Kijima T, Kohmo S, Arase H, Otani Y, Nagatomo I, Takahashi R, Miyake K, Higashiguchi M, Morimura O, Ihara S, Tsujino K, Hirata H, Inoue K, Takeda Y, Kida H, Tachibana I, Kumanogoh A.	Overcoming chemoresistance of small-cell lung cancer through stepwise HER2-targeted antibody-dependent cell-mediated cytotoxicity and VEGF-targeted antiangiogenesis.	Scientific reports.	3	2669	2013
Nishino K, Imamura F, Morita S, Mori M, Komuta K, Kijima T, Namba Y, Kumagai T, Yamamoto S, Tachibana I, Nakazawa Y, Uchida J, Minami S, Takahashi R, Yano Y, Okuyama T, Kumanogoh A.	A retrospective analysis of 335 Japanese lung cancer patients who responded to initial gefitinib treatment.	Lung cancer.	82	299	2013
He P, Kuhara H, Ta	Calretinin mediates apop	FEBS open	3	225.	2013.

chibana I, Jin Y, Takeda Y, Tetsumoto S, Minami T, Kohmo S, Hirata H, Takahashi R, Inoue K, Nagatomo I, Kida H, Kijima T, Naka T, Morii E, Kawase I, Kumano A.	tosis in small cell lung cancer cells expressing tetraspanin CD9.	bio.			
Coenen A, Honda O, van der Jagt EJ, Tomiyama N.	Computer-assisted solid lung nodule 3D volumetry on CT: influence of scan mode and iterative reconstruction: a CT phantom study	Jpn J Radio l.	31	677-84	2013
Kono T, Hata T, Morita S, Munemoto Y, Matsui T, Kojima H, Takemoto H, Fukunaga M, Nagata N, Shimada M, Sakamoto J, Mishima H	Goshajinkigan oxaliplatin neurotoxicity evaluation (GONE): a phase 2, multicenter, randomized, double-blind, placebo-controlled trial of goshajinkigan to prevent oxaliplatin-induced neuropathy	Cancer Chemother Pharmacol	72(6)	1283-90	2013
Oba K, Paoletti X, Alberts S, Bang YJ, Benedetti J, Bleiberg H, Catalano P, Lordick F, Michiels S, Morita S, Ohashi Y, Pignon JP, Rougier P, Sasako M, Sakamoto J, Sargent D, Shitara K, Cutsem EV, Buyse M, Burzykowski T; on behalf of the GASTRIC group	Disease-Free Survival as a Surrogate for Overall Survival in Adjuvant Trials of Gastric Cancer: A Meta-Analysis	J Natl Cancer Inst	105(21)	1600-7	2013
Paoletti X, Oba K, Bang YJ, Bleiberg H, Boku N, Bouchet O, Catalano P, Fuse N, Michiels S, Moehler M, Morita S, Ohashi Y, Ohtsu A, Roth A, Rougier P, Sakamoto J, Sargent D, Sasako M, Shitara K, Thuss-Patience P, Van Cutsem E, Burzykowski T, Buyse M; on behalf of the GASTRIC group	Progression-Free Survival as a Surrogate for Overall Survival in Advanced/Recurrent Gastric Cancer Trials: A Meta-Analysis	J Natl Cancer Inst	105(21)	1667-70	2013
Hazama S, Mishima H, Tsunedomi R, Okuyama Y, Kato T, Takahashi KI, Nozawa H, Ando H, Kobayashi M, Takemoto H, Nagata N, Kanekiyo S, Inoue Y, Hamamoto Y, Fujita Y, Hino da Y, Okayama N, Oba K, Sakamoto J, Oka M	UGT1A1*6, 1A7*3, and 1A9*22 genotypes predict severe neutropenia in FOLFIRI-treated metastatic colorectal cancer in two prospective studies in Japan	Cancer Sci	104(12)	1662-9	2013

Kitagawa Y, Takeuchi H, Takagi Y, Natsugoe S, Terashima M, Murakami N, Fujimura T, Tsujimoto H, Hayashi H, Yoshimizu N, Takagane A, Mohri Y, Nobeshima K, Uenosono Y, Kinami S, Sakamoto J, Morita S, Aikou T, Miwa K, Kitajima M	Sentinel node mapping for gastric cancer: a prospective multicenter trial in Japan	J Clin Oncol	31(29)	3704-10	2013
Khamlub S, Harun-Or-Rashid M, Sarker MAB, Hirosawa T, Outavong P, Sakamoto J	Job satisfaction of health-care workers at health centers in Vientiane capital and Bolikhamsai province, Lao PDR	Nagoya J Med Sci	75(3-4)	233-41	2013
Srisawang P, Harun-Or-Rashid M, Hirosawa T, Sakamoto J	Knowledge, attitudes and barriers of physicians, policy makers/regulators regarding use of opioids for cancer pain management in Thailand	Nagoya J Med Sci	75(3-4)	201-12	2013
Serker MAB, Rahman M, Harun-Or-Rashid M, Hossain S, Kasuya H, Sakamoto J, Hamajima N	Association of smoked and smokeless tobacco use with migraine: a hospital-based case-control study in Dhaka, Bangladesh	Tob Induc Dis	11(1)	15	2013
Tezuka T, Hamada C, Ishida H, Ooshiro M, Matsuoka H, Kawasaki S, Mishima H, Maeda K, Sakamoto J, Koda K	Phase II clinical study of modified FOLFOX7 (intermittent oxaliplatin administration) plus bevacizumab in patients with unresectable metastatic colorectal cancer—CRAFT study	Invest New Drugs	31(5)	1321-9	2013
Yoshikawa T, Tanabe K, Nishikawa K, Ito Y, Matsui T, Kimura Y, Hirabayashi N, Mikata S, Iwahashi M, Fukushima R, Takiguchi N, Miyashiro I, Morita S, Miyashita Y, Tsuburaya A, Sakamoto J	Induction of a pathological complete response by four courses of neoadjuvant chemotherapy for gastric cancer: early results of the randomized phase II COMPASS trial	Ann Surg Oncol	21(1)	213-9	2013
Tsuburaya A, Nagata N, Cho H, Hirabayashi N, Kobayashi M, Kojima H, Munakata Y, Fukushima R, Kameda Y, Shimoda T, Oba K, Sakamoto J	Phase II trial of paclitaxel and cisplatin as neoadjuvant chemotherapy for locally advanced gastric cancer	Cancer Chemother Pharmacol	71(5)	1309-14	2013
Okada K, Kawai M, Uesaka K, Kodera Y, Nagano H, Murakami Y, Morita S, Sakamoto J, Yamaue H; JAPAN-PD Investig	Effect of Daikenchuto (TJ-100) on postoperative bowel motility and on prevention of paralytic ileus after pancreaticoduodenectomy: a multicenter, ran	Jpn J Clin Oncol	43(4)	436-8	2013

ators	domized, placebo-controlled phase II trial (the JAPAN-PD study)				
Hamrah MS, Harun-Or-Rashid M, Hirose T, Sakamoto J, Hashemi H, Emami MH, Shariati M, Fotouhi A.	Smoking and associated factors among the population aged 40-64 in Shahrood, Iran.	Asian Pacific J Cancer Prev	14(3)	1919-1923	2013
Shitara K, Morita S, Fujitani K, Kadowaki S, Takiguchi N, Hirabayashi N, Takahashi M, Takagi M, Tokunaga Y, Fukushima R, Munakata Y, Nishikawa K, Takagane A, Tanaka T, Sekishita Y, Kang Y, Sakamoto J, Tsuburaya A.	Combination Chemotherapy with S-1 plus Cisplatin for Gastric Cancer that Recurs after Adjuvant Chemotherapy with S-1: Multi-institutional Retrospective Analysis.	Gastric Cancer	15	245-251	2012
Yamada A, Ishikawa T, Ota I, Kimura M, Shimizu D, Tanabe M, Chishima T, Sasaki T, Ichikawa Y, Morita S, Yoshimura K, Takabe K, Endo I.	High expressions of ATP-binding cassette transporter ABCC11 in the breast tumors are associated with aggressive subtypes and worse disease-free survival.	Breast Cancer Research and Treatment	137	773-782	2013.

書籍

著者氏名	論文タイトル名	書籍全体の編集者名	書 籍 名	出 版 社 名	出版地	出版年	ページ
<u>Ikeda N</u> , Lam S	Optical Coherence Tomography	Diaz-Jimenez J P, Rodriguez A N	Interventions in Pulmonary Medicine	Springer	アメリカ	2013	231-236
多田弘人	最新肺癌学－基礎と臨床の最新研究動向－肺癌の治療戦略 外科治療 Ⅲ期肺癌の外科治療	日本臨床社 日本臨床社	日本臨床	日本臨床社	日本	2013	445-449

#### IV. 研究成果の刊行物・別刷

# Wilms Tumor Gene (WT1) Peptide-based Cancer Vaccine Combined With Gemcitabine for Patients With Advanced Pancreatic Cancer

Sumiyuki Nishida,\*† Shigeo Koido,‡ Yutaka Takeda,§ Sadamu Homma,|| Hideo Komita,‡  
Akitaka Takahara,‡ Satoshi Morita,¶ Toshinori Ito,§§ Soyoko Morimoto,† Kazuma Hara,\*\*  
Akihiro Tsuboi,\* Yoshihiro Oka,†† Satoru Yanagisawa,‡‡ Yoichi Toyama,‡‡  
Masahiro Ikegami,§§ Toru Kitagawa,§ Hidetoshi Eguchi,§ Hiroshi Wada,§ Hiroaki Nagano,§  
Jun Nakata,† Yoshiki Nakae,† Naoki Hosen,\*\* Yusuke Oji,|| Toshio Tanaka,††¶¶  
Ichiro Kawase,† Atsushi Kumanogoh,††† Junichi Sakamoto,### Yuichiro Doki,§ Masaki Mori,§  
Toshifumi Ohkusa,‡ Hisao Tajiri,‡ and Haruo Sugiyama\*\*

**Summary:** Wilms tumor gene (*WT1*) protein is an attractive target for cancer immunotherapy. We aimed to investigate the feasibility of a combination therapy consisting of gemcitabine and WT1 peptide-based vaccine for patients with advanced pancreatic cancer and to make initial assessments of its clinical efficacy and immunologic response. Thirty-two HLA-A\*24:02<sup>+</sup> patients with advanced pancreatic cancer were enrolled. Patients received HLA-A\*24:02-restricted, modified 9-mer WT1 peptide (3 mg/body) emulsified with Montanide ISA51 adjuvant (WT1 vaccine) intradermally biweekly and gemcitabine (1000 mg/m<sup>2</sup>) on days 1, 8, and 15 of a 28-day cycle. This combination therapy was well tolerated. The frequencies of grade 3–4 adverse events for this combination therapy were similar to those for gemcitabine alone. Objective response rate was 20.0% (6/30 evaluable patients). Median survival time and 1-year survival rate were 8.1 months and 29%, respectively. The association between longer survival and positive delayed-type hypersensitivity to WT1 peptide was statistically significant, and longer survivors featured a higher frequency of memory-phenotype WT1-specific cytotoxic T lymphocytes both before and after treatment. WT1 vaccine in combination with gemcitabine was well tolerated for patients with advanced pancreatic cancer. Delayed-type hypersensitivity-positivity to WT1 peptide and a higher frequency of memory-phenotype WT1-specific cytotoxic T lymphocytes could be useful prognostic markers for

survival in the combination therapy with gemcitabine and WT1 vaccine. Further clinical investigation is warranted to determine the effectiveness of this combination therapy.

**Key Words:** Wilms tumor gene (WT1), WT1 peptide vaccine, cancer immunotherapy, pancreatic cancer, gemcitabine

(*J Immunother* 2014;37:105–114)

Pancreatic cancer remains a malignancy with high mortality.<sup>1</sup> Gemcitabine has been the standard first-line treatment for patients with advanced pancreatic cancer, but featured a median overall survival time (MST) of about 6 months and a 1-year overall survival (OS) rate of ≤20%.<sup>2</sup> Although many trials of gemcitabine-based combination therapies with cytotoxic or biological agents have been attempted, these therapies, with the exception of erlotinib,<sup>3</sup> have not achieved any survival results superior to those attained with gemcitabine alone.<sup>1</sup> Prognosis of patients with pancreatic cancer thus remains extremely poor, so that novel treatments are urgently needed to improve survival.

Among promising therapeutic strategies, active cancer immunotherapies, such as peptide-based cancer vaccines against tumor-associated antigens (TAAs), which elicit TAA-specific cytotoxic T lymphocytes (CTLs) that eventually eradicate cancer cells, have been and are being developed.<sup>4</sup> However, because their clinical efficacy has been limited,<sup>5,6</sup> several approaches have been tried to improve their efficacy. One approach is the use of combination therapies with certain chemotherapeutic agents, including gemcitabine, which can stimulate the immune system.<sup>7–9</sup> An additional benefit is that chemotherapy makes the tumor cells susceptible to CTL response,<sup>10,11</sup> whereas cancer immunotherapy can sensitize the tumor cells to subsequent chemotherapeutic agents. For this reason, cancer vaccine in combination with certain chemotherapeutic agents can be expected to exert synergistic effects.

The Wilms tumor gene (*WT1*) is highly expressed in various kinds of malignancies and has been found to perform oncogenic rather than tumor-suppressor functions in tumorigenesis.<sup>12,13</sup> Moreover, both cellular and humoral immune responses against the WT1 protein are naturally elicited in cancer patients, indicating that the *WT1* gene product is actually immunogenic.<sup>14–18</sup> In view of these

Received for publication June 3, 2013; accepted December 16, 2013.  
From the Departments of \*Cancer Immunotherapy; §Surgery; †Respiratory Medicine, Allergy and Rheumatic Diseases; \*\*Functional Diagnostic Science; ||Cancer Stem Cell Biology; #Complementary and Alternative Medicine; ¶¶Clinical Application of Biologics, Graduate School of Medicine; ††Department of Immunopathology, WPI Immunology Frontier Research Center, Osaka University, Osaka; Departments of ‡Internal Medicine, Division of Gastroenterology and Hepatology; ||Oncology, Institute of DNA Medicine; Departments of ‡‡Surgery; §§Pathology, The Jikei University School of Medicine, Tokyo; ¶¶Department of Biostatistics and Epidemiology, Yokohama City University Medical Center, Kanagawa; and ###Department of Young Leaders' Program in Medical Administration, Nagoya University Graduate School of Medicine, Showa-ku, Nagoya, Japan.

S.N. and S.K. contributed equally.

Trial registration ID: UMIN000001187.

Reprints: Haruo Sugiyama, Department of Functional Diagnostic Science, Graduate School of Medicine, Osaka University, 1-7 Yamada-Oka, Suita City, Osaka 565-0871, Japan (e-mail: sugiyama@sahs.med.osaka-u.ac.jp).

Supplemental Digital Content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's Website, [www.immunotherapy-journal.com](http://www.immunotherapy-journal.com).

Copyright © 2014 by Lippincott Williams & Wilkins



findings, we and others have been performing clinical studies of the efficacy of WT1 peptide–based immunotherapies for patients, including children, with various kinds of malignancies.<sup>13,19–26</sup>

This report describes a phase 1 clinical study of a WT1 peptide–based cancer vaccine combined with gemcitabine for patients with advanced pancreatic cancer. The main objective of this study was to investigate the feasibility of this combination therapy and to make initial assessments of its clinical efficacy and the immunologic response to WT1 peptide.

## MATERIALS AND METHODS

### Patient Characteristics

Patients with pathologically or cytologically confirmed, measurable, locally advanced, or metastatic pancreatic adenocarcinoma or with recurrent disease were recruited for this noncomparative, open-label, phase 1 study at 2 centers: Osaka University Hospital and Jikei University Kashiwa Hospital, in Japan. Another major eligibility criterion was HLA-A\*24:02 positivity. We chose this phenotype because about 60% of Japanese population had this phenotype. Other eligibility criteria included age of 20 years and older, 75 years and younger, Karnofsky performance status 60%–100%, no previous history of treatment for locally advanced or metastatic disease, a minimum 6-month interval from completion of any previous treatment for recurrent disease, a life expectancy of  $\geq 3$  months, and adequate organ functions. This study was approved by the ethical review boards of the 2 centers and performed in accordance with the Helsinki Declaration. All patients provided written informed consent.

### WT1-Peptide–based Cancer Vaccine (WT1 Vaccine)

A HLA-A\*24:02-restricted, modified 9-mer WT1 peptide (mp235; CYTWNQMNL; Peptide Institute Inc., Osaka, Japan) was generated according to the Good Manufacturing Practice Guidelines. In our previous report about the first clinical use of WT1 peptide,<sup>19</sup> the dose-

escalation of WT1 peptide from 0.3 to 3.0 mg was designed to decide the recommended dose in combination with the incomplete Freund's adjuvant (Montanide ISA51; Seppic, Paris, France), and 3 mg of WT1 peptide in combination with Montanide ISA51 was decided to be well tolerated. In our present study, we chose WT1 vaccine composed of 3 mg of WT1 peptide and Montanide ISA51 adjuvant. WT1 vaccine was prepared, according to our previous report.<sup>19</sup> WT1 peptide of 3 mg was dissolved in a small volume of dimethyl sulfoxide (DMSO; Sigma, St Louis, MO). The solution was then diluted to 400  $\mu$ L with 5% glucose and finally emulsified with an equal weight of Montanide ISA51 adjuvant.

### Treatment

Gemcitabine was intravenously administered at a dose of 1000 mg/m<sup>2</sup> on days 1, 8, and 15 of a 28-day cycle. WT1 vaccine was intradermally administered at 6 different sites (bilateral upper arms, lower abdomen, and femoral regions) on days 1 and 15 of a 28-day cycle. The initial treatment protocol was planned as 2 courses. Patients without early progressive disease upon the completion of protocol treatment could receive additional treatment until the occurrence of disease progression, unacceptable adverse events, or withdrawal of consent.

### Study Assessment

Toxicity was graded using the National Cancer Institute's Common Toxicity Criteria of Adverse Events (CTCAE version 3.0). Dose-limiting toxicity (DLT) was defined as the following adverse events, during the first 2 courses, which were possibly, probably, or definitely related to treatment: grade 4 hematological toxicity lasting  $> 7$  days, grade 3 or worse neutropenia accompanied by high fever ( $\geq 38^{\circ}\text{C}$ ) or infection (febril neutropenia), and any nonhematological toxicity of grade 3 or worse in other organ systems, including vaccine-injection sites. Biliary tract infection secondary to biliary obstruction was not considered to be a DLT unless it occurred in conjunction with grade  $\geq 3$  neutropenia. Computed tomography was performed every 4 weeks during the protocol treatment and




FIGURE 1. Study profile.



every 6–8 weeks during the additional treatment until disease progression, and tumor response was assessed by the investigators according to the Response Evaluation Criteria in Solid Tumors criteria. Stable disease (SD) was defined as a disease that was stable for  $\geq 8$  weeks after the beginning of treatment. The concentration of the tumor marker carbohydrate antigen 19-9 (CA19-9) was measured at baseline and each course.

WT1-specific Immunologic Assessment

As WT1-specific immunologic assessment, delayed-type hypersensitivity (DTH) to WT1 peptide and the WT1 peptide/HLA-A\*24:02 tetramer assay was examined. DTH was examined on day 1 of each course during the protocol treatment and optionally at suitable time during the additional treatment. All DTH tests were performed and measured by the investigators. Briefly, 30  $\mu$ g of WT1 peptide in saline and saline alone were intradermally injected in the forearm, and the maximum diameter of erythema and other skin reaction, including induration, were measured after 48 hours. DTH-positivity was defined as erythema  $\geq 2$  mm in diameter, which size was the minimum size measurable with a ruler at the clinical practice.

Peripheral blood (PB) mononuclear cells for WT1 peptide/HLA-A\*24:02 tetramer assay were collected on day 1 of each course during the protocol treatment and appropriately during the additional treatment, and cryopreserved until use. The following tetramer and monoclonal antibodies were used: PE-conjugated WT1<sub>235</sub> tetramer [HLA-A\*24:02-restricted natural 9-mer WT1 peptide (CMT WNQMNL)] (MBL, Nagoya, Japan), anti-CD4-FITC, anti-CD16-FITC, anti-CD45RA-APC (BioLegend, San Diego, CA), anti-CD19-FITC, anti-CCR7-PE-Cy7 (BD Pharmingen, San Diego, CA), anti-CD3-PerCP, anti-CD8-APC-Cy7, anti-CD14-FITC (BD Biosciences, San Jose, CA), and anti-CD56-FITC (eBioscience, San Diego, CA). Lineage antigen (CD4, CD14, CD16, CD19, and CD56)-negative, CD3<sup>+</sup>, CD8<sup>+</sup>, and WT1<sub>235</sub> tetramer<sup>+</sup> lymphocytes were defined as WT1 tetramer<sup>+</sup> CD3<sup>+</sup> CD8<sup>+</sup> T lymphocytes (WT1-CTLs). Data acquisition were performed on a FACS Aria instrument (BD Biosciences), and data analysis were performed with FACS Diva software (BD Biosciences).

Statistical Analysis

The safety profile constituted the primary end point. A treatment schedule was considered to be acceptable if the probability of developing DLT was estimated to be  $< 20\%$ . If the estimated probability of DLT occurrence was 10%, the upper limit of the 90% (one-sided) confidence interval (CI) of DLT probability was  $< 20\%$ , based on the projected sample size of 20 patients. For a more accurate determination of the associations with clinical efficacy and immunologic parameters, in total 32 patients were enrolled (8 patients were further enrolled after the completion of safety assessment with the initial 24 patients as shown in Fig. 1). The secondary end points included objective response, CA19-9 response, defined as a decrease in CA19-9 concentration of at least 50% in the patients with  $\geq 100$  U/mL of CA19-9 at baseline, progression-free survival defined as time from date of beginning of the treatment to date of disease progression as confirmed by the investigators or death without progression, OS, immunologic responses to WT1 peptide, and correlations between clinical benefit response (CBR)<sup>2</sup> and quality of life (QOL) assessed using by the Functional Assessment of Cancer Therapy-General

(FACT-G) measurement system.<sup>27</sup> The nonparametric, Wilcoxon signed-rank test or Mann-Whitney *U* test was used to calculate *P* values for change in immune cells because the data were skewed. We judged *P* values of  $< 0.01$  to be significant.  $\chi^2$  test was used to calculate *P* values for associations between DTH and clinical efficacy. The statistical analyses were performed with SAS for Windows version 9.2 (SAS Institute Inc., Cary, NC). Correlations between CBR and the physical and functional scores based on replies to the FACT-G QOL questionnaire were analyzed with a linear mixed-effects model, for which SAS for Windows release 9.1 (SAS Institute Inc.) was used.

RESULTS

Patient Characteristics

A total of 63 patients with advanced pancreatic cancer, whose median age was 63.0 years old, were screened and checked a phenotype in HLA-A locus. Twenty-two patients failed to enroll in this trial because of lack of HLA-A\*24:02 phenotype. A total of 32 HLA-A\*24:02<sup>+</sup> patients with advanced pancreatic cancer were finally enrolled in this trial between 2008 and 2010. Of 32 patients, 28 had inoperable advanced pancreatic cancer (6 locally advanced and 22 metastatic diseases), and the remaining 4 had recurrent

TABLE 1. Patients Characteristics at Baseline

Characteristics	N (%)
Age (y)	
Median	60.0
Range	41–75
Sex	
Male	17 (53.1)
Female	15 (46.9)
Karnofsky performance status (%)	
$\sim 70$	7 (21.9)
80	10 (31.3)
90	12 (37.5)
100	3 (9.4)
Disease extent	
Inoperable advanced disease	28 (87.5)
Locally advanced	6 (18.8)
Metastatic	22 (68.8)
Recurrent disease	4 (12.5)
Local relapse	1 (3.1)
Distant metastasis	3 (9.4)
Primary tumor site	
Head	15 (46.9)
Body/tail	17 (53.1)
Metastatic sites	
Liver	17 (53.1)
Distant lymph node	16 (50.0)
Lung	7 (21.9)
Peritoneum	6 (18.8)
Others*	4 (12.5)
CA19-9 concentration at baseline (U/mL)	
Median	248
Range (U/mL)	$< 5$ –75,050
$\leq 5$	3 (9.4)†
6–99	10 (31.3)
100–999	7 (21.9)
1000–9999	5 (15.6)
$\geq 10,000$	7 (21.9)

\*Other metastatic sites included bone, ovary, or adrenal gland.  
†All patients had the Lewis blood group-negative phenotype.  
CA19-9 indicates carbohydrate antigen 19-9.

disease. Table 1 summarizes the patient baseline characteristics. Three patients did not complete the first 2 courses of treatment: 2 patients showed rapid disease progression, and 1 refused to continue the treatment. It was determined by the supervising Data Safety and Monitoring Board that the elimination of these cases was unlikely to be or was not related to the protocol treatment. Of the initial 24 patients, 21 could thus be used for assessment of DLT, 29 of all 32 patients for assessment of adverse events (Fig. 1).

## Safety

Administration of WT1 vaccine in combination with gemcitabine was well tolerated. All adverse events are listed in Table 2. The initial assessment of safety for 21 patients found that a grade 4 central nervous system cerebrovascular ischemia considered to be a DLT had occurred in 1 patient. The most commonly reported adverse event was skin toxicity related to WT1 vaccine. All patients developed grade 1 or 2 skin reactions with swelling, redness, erythema, and induration with or without involvement of small vesicles at the local vaccine-injection sites. Hematological abnormalities were similar to those observed with the administration of gemcitabine alone, and none of the patients developed DLTs associated with hematological abnormalities or febrile neutropenia. Eight grade 3 non-hematological adverse events (1 instance of hyponatremia and 7 hepatobiliary/pancreas infections) were detected and attributed to complications associated with disease progression or biliary obstruction. Other major non-hematological adverse events included grade 1 or 2 skin rash, anorexia, nausea, and fever, all of which were previously reported as major adverse events associated with

gemcitabine. Hepatic transaminase elevation was principally related to disease progression and/or hepatobiliary infection. Except for local skin reactions, none of the patients experienced adverse events considered to be related to WT1 vaccination.

## Clinical Response and Survival Analysis

The clinical efficacy results for all 32 patients are summarized in Table 3. Two patients were excluded from some of these analyses. One patient, who had followed a satisfactory and interesting treatment course and finally undergone a surgical resection (Supplementary Figure 1, Supplemental Digital Content 1, <http://links.lww.com/JIT/A317> and Table 3), was excluded from the evaluations of response and survival because the diagnosis of pancreatic cancer could not be pathologically confirmed due to the lack of viable tumor cells in the resected specimens. The other patient was excluded from the evaluation of response because of withdrawal of consent before the first evaluation. Thus, of the total of 32 patients, 30 could be used to evaluate response to treatment and 31 to assess survival. Six of 30 patients (20.0%) reached partial response (PR), and 16 of them (53.3%) showed SD at least for  $\geq 8$  weeks (Table 3). Median progression-free survival was 4.2 months (95% CI, 3.6–4.6) (Fig. 2A) and MST was 8.1 months (95% CI, 6.3–10.0) (Fig. 2B). Six-month and 1-year OS rates were 71.0% (95% CI, 54.9–87.1) and 29.0% (95% CI, 12.9–45.1), respectively (Fig. 2B).

Ten of 19 patients with  $\geq 100$  U/mL of CA19-9 at baseline (52.6%) showed a decrease in CA19-9 serum concentration of at least 50% (Table 3).

**TABLE 2.** Adverse Events Reported in 29 Patients who Completed the First 2 Courses of Treatment

	Grades				N (%)		
	1	2	3	4	Any Grade (N = 29)	Grade 3 or 4 (N = 29)	DLT (N = 21)
Hematological abnormalities							
Neutropenia	3	6	13	0	22 (75.9)	13 (44.8)	0 (0.0)
Leukocytopenia	4	12	8	0	24 (82.8)	8 (27.6)	0 (0.0)
Lymphopenia	3	12	8	0	23 (79.3)	8 (27.6)	0 (0.0)
Anemia	6	15	2	0	23 (79.3)	2 (6.9)	0 (0.0)
Thrombocytopenia	15	6	1	0	22 (75.9)	1 (3.4)	0 (0.0)
Nonhematological events							
CNS ischemia	0	0	1	0	1 (3.4)	1 (3.4)	1 (4.8)
Hepatobiliary tract infection with normal ANC	0	1	7	0	8 (27.6)	7 (24.1)	0 (0.0)
Hyponatremia	3	0	1	0	4 (13.8)	1 (3.4)	0 (0.0)
Hypoalbuminemia	9	4	0	0	13 (44.8)	0 (0.0)	0 (0.0)
Alanine aminotransferase	9	4	0	0	13 (44.8)	0 (0.0)	0 (0.0)
Aspartate aminotransferase	10	1	0	0	11 (37.9)	0 (0.0)	0 (0.0)
Bilirubin	2	4	0	0	6 (20.7)	0 (0.0)	0 (0.0)
Hyperkalemia	3	0	0	0	3 (10.3)	0 (0.0)	0 (0.0)
Hemorrhage in urinary tracts	2	1	0	0	3 (10.3)	0 (0.0)	0 (0.0)
Proteinuria	2	0	0	0	2 (6.9)	0 (0.0)	0 (0.0)
Hypokalemia	1	0	0	0	1 (3.4)	0 (0.0)	0 (0.0)
Anorexia	9	0	0	0	9 (31.0)	0 (0.0)	0 (0.0)
Rash*	5	3	0	0	8 (27.6)	0 (0.0)	0 (0.0)
Fever	6	1	0	0	7 (24.1)	0 (0.0)	0 (0.0)
Nausea	7	0	0	0	7 (24.1)	0 (0.0)	0 (0.0)
Diarrhea	2	1	0	0	3 (10.3)	0 (0.0)	0 (0.0)

Adverse events were graded using the National Cancer Institute Common Toxicity Criteria of Adverse Events (CTCAE version 3.0).

\*Exclude skin reaction at WT1 vaccine-injection sites.

ANC indicates absolute neutrophil count; CNS, central nervous system; DLT, dose-limiting toxicity.

Correlations between CBR and either physical or functional scores assessed with the FACT-G QOL questionnaire were analyzed. For assessment of CBR, 16 of the initial 24 patients (66.7%) could be used. Nine (56.3%) of these patients (3 with PR, 5 with SD, and 1 with progressive disease) were classified as CBR responders (data not shown). CBR responders showed improvement in physical and functional scores during the first 2 courses, whereas both scores for CBR nonresponders tended to become worse (Supplementary Figure 2, Supplemental Digital Content 2, <http://links.lww.com/JIT/A318>).

WT1-specific Immune Response

Exploratory analyses of the immune response consisted of assessment of DTH to WT1 peptide and WT1 tetramer + CD3 + CD8 + T lymphocytes (WT1-CTLs) in PB of all 32 patients. All patients were DTH-negative at baseline, but 31 were at least once assessed as DTH after WT1 vaccination and 18 patients (58.1%) showed DTH-positivity, all of which conversion was detected during the protocol treatment. All of the DTH-positive patients showed at least ≥ 4 mm diameter of erythema, which was a length that was easy enough to measure. Next, for

evaluation of associations between survival and DTH, the patients were classified into 4 groups according to survival time: Superior (>12 mo), good (8–12 mo), moderate (4–8 mo), and poor (≤4 mo) responders. These categories were based on the following findings: (i) MST for best supportive care only is no more than 3–4 months<sup>1</sup>; (ii) MST of our patients was 8.1 months; and (iii) survival time of > 12 months generally indicates that the treatment has been beneficial. DTH-positivity of superior and good responders was 68.7% (11/16), whereas that of poor responders was 0% (0/7). The association between DTH-positivity and longer survival time was statistically significant

TABLE 3. Summary of Clinical Efficacy Results

	All Patients	DTH Positive	DTH Negative
Best overall response [N (%)]			
Complete response	0 (0.0)	0 (0.0)	0 (0.0)
Partial response	6 (20.0)	3 (17.6)	3 (23.1)
Stable disease*	16 (53.3)	12 (70.6)	4 (30.8)
Progressive disease	8 (26.7)	2 (11.8)	6 (46.2)
Excluded Not evaluable	1† 1	1	0
CA19-9 response (≥100 U/mL at baseline)			
Positive‡ [N (%)]	10 (52.6)	7 (63.6)	3 (42.9)
PFS	N = 31	N = 17	N = 13
Range (d)	21–1504 +	55–1504 +	21–373
Median PFS (mo)	4.2 (1.1–7.4)	5.4 (2.6–8.2)	2.9 (–1.6 to 7.1)
3-mo PFS (%)	67 (50–84)	82 (64–100)	46 (9–73)
OS	N = 31	N = 17	N = 13
Range (d)	30–1504 +	154–1504 +	30–443
Median OS (mo)	8.1 (6.3–10.0)	10.9 (1.2–20.7)	3.9 (–3.0 to 10.7)
6-mo OS (%)	71 (55–87)	88 (73–104)	46 (19–73)
12-mo OS (%)	29 (13–45)	47 (18–65)	7.7 (–6.8 to 22)

(): 95% CI.  
\*Stable disease conformation is determined at least for ≥ 8 weeks.  
†This patient was reached partial response after 3 courses of treatment, and finally underwent the surgical resection. This patient was excluded the analysis of clinical response, PFS, and OS.  
‡“Positive” CA19-9 response is defined as a ≥ 50% decrease in CA19-9 concentration after treatment.  
CA19-9 indicates carbohydrate antigen 19-9; CI, confidence interval; DTH, delayed-type hypersensitivity; OS, overall survival; PFS, progression-free survival.




FIGURE 2. Kaplan-Meier survival curves. A, Progression-free survival (N=31). B, Overall survival (N=31). C, Overall survival in DTH-positive (gray line) or DTH-negative patients (black line). DTH indicates delayed-type hypersensitivity.

( $\chi^2 = 15.908$ ,  $P = 0.0012$ ) (Table 4). Therefore, survival was retrospectively reanalyzed in terms of DTH-positivity or DTH-negativity. MST was 3.9 and 10.9 months for DTH-negative (N = 13) and DTH-positive (N = 17) patients, respectively, with a statistically significant difference ( $P = 0.0030$ ) (Fig. 2C and Table 3).

The number of WT1-CTLs and the percentages of naive (CD45RA<sup>+</sup> CCR7<sup>+</sup>), memory (CD45RA<sup>+</sup> CCR7<sup>+</sup> and CD45RA<sup>+</sup> CCR7<sup>+</sup>), and effector (CD45RA<sup>+</sup> CCR7<sup>+</sup>) phenotypes in WT1-CTLs did not show any significant changes during the protocol treatment by the analysis using all patients (Supplementary Table 1, Supplemental Digital Content 3, <http://links.lww.com/JIT/A319> and Supplementary Table 2, Supplemental Digital Content 4, <http://links.lww.com/JIT/A320>). Next, these immunologic parameters were compared between patients showing DTH-positivity and DTH-negativity. The difference in the number of WT1-CTLs was not statistically significant (Supplementary Table 1, Supplemental Digital Content 3, <http://links.lww.com/JIT/A319>). Phenotype analysis of WT1-CTLs showed that the percentage of naive-phenotype was higher in DTH-positive than in DTH-negative patients at baseline (Fig. 3A). After treatment, DTH-positive patients showed a significantly higher percentage of memory-phenotype and consequently a lower percentage of effector-phenotype WT1-CTLs than did their DTH-negative counterparts (Fig. 3A and Supplementary Table 2, Supplemental Digital Content 4, <http://links.lww.com/JIT/A320>). Furthermore, the percentage of memory-phenotype WT1-CTLs for the superior responders seemed to be relatively higher than that of effector-phenotype WT1-CTLs (Fig. 3B), whereas this tendency was quite the opposite for the poor responders (Fig. 3B and Supplementary Table 3, Supplemental Digital Content 5, <http://links.lww.com/JIT/A321>).

Case Report

A 44-year-old male with a locally advanced pancreatic head cancer (T4N1M0; stage III) received WT1 vaccine in combination with gemcitabine, and achieved PR (Fig. 4A). Five months after the beginning of the treatment, this patient underwent a complete surgical resection. Histopathologic examination of the resected specimen showed an invasive ductal adenocarcinoma with mononuclear cell infiltration around the cancer region and moderate to severe fibrotic change (Fig. 4B). This patient proved to be positive for DTH to WT1 peptide after 1 treatment course (Fig. 4C). The number of WT1-CTLs transiently decreased during the first 2–3 treatment courses but subsequently increased again, while the percentage of memory-phenotype WT1-CTLs remained high during the treatment courses (Fig. 4C). Of note, the percentage of WT1-CTLs in the tumor-infiltrating CD3<sup>+</sup> CD8<sup>+</sup> T lymphocytes was 2.48%, which was about 6 times higher than that in PB (0.39%) (Fig. 4D). This patient had been receiving monthly administration of WT1 vaccine in combination with gemcitabine for 3 years and has maintained a Karnofsky performance status of 100% with no evidence of disease recurrence.

DISCUSSION

This study was designed with a DLT target rate of 10% during the first 2 treatment courses, but only one of the 21 initial evaluable patients (4.8%) actually experienced DLT. These results confirmed that WT1 vaccine in combination with gemcitabine is acceptable for patients with

advanced pancreatic cancer. Cerebrovascular ischemia, reported here as a DLT, could be also caused by pancreatic cancer itself and/or the administration of gemcitabine, both of which are sometimes associated with a high risk of developing thrombotic disease.<sup>28,29</sup> Therefore, this adverse event was considered to be multifactorial and judged to be “possibly” related to treatment.

Except for skin reactions at the local injection sites, the toxicity profiles of WT1 vaccine in combination with gemcitabine were consistently similar to those of gemcitabine alone. As the *WT1* gene is physiologically expressed in hematopoietic progenitor cells,<sup>13</sup> damage to hematopoiesis is one of the major concerns in WT1-peptide-based immunotherapy. The incidence of hematological adverse events in our study, however, was similar to that observed for treatment with gemcitabine alone,<sup>30</sup> and these events were easily managed and reversible. These findings suggest that WT1 vaccine does not synergistically intensify hematological adverse events associated with gemcitabine. It seems unlikely that WT1-specific CTLs elicited by WT1 vaccine might damage normal WT1-expressing hematopoietic progenitor cells as well as WT-expressing tumor cells, as following reasons. First, in the previous clinical studies, we and others reported that WT1-specific CTLs elicited by WT1 vaccine decreased WT1-expressing leukemia cells and suppressed the disease progression of WT-expression cancer cells, but not significantly damaged normal hematopoiesis.<sup>19,23–26</sup> Second, it was demonstrated that, using mice in vivo experiments, WT1-targeting immunotherapy gave damage to tumor cells, but not WT1-expressing normal tissue, including hematopoietic cells.<sup>31,32</sup> The reason why the normal WT1-expressing hematopoietic cells are able to escape from the attack by WT1-specific CTLs is not well known. Further investigations should be required to address this issue.

The clinical efficacy of treatment with WT1 vaccine in combination with gemcitabine, especially in terms of survival, seemed to be better than of that with gemcitabine alone.<sup>1,2</sup> About half of patients who had been induced WT1-specific immunity after vaccination showed better clinical outcome with 12 months or longer survival time, suggesting additional or synergistic effects of WT1 vaccine in combination with gemcitabine. Furthermore, the former contributed to pain relief and thus to improvement of QOL. Recently, the result of the phase III study of gemcitabine plus S-1, S-1 alone, or gemcitabine alone in patients with locally advanced and metastatic pancreatic cancer (GEST study) conducted in Japan and Taiwan between 2007 and 2009 has been reported.<sup>33</sup> Median OS and OS rate at 12


TABLE 4. Association Between DTH and Survival

	Overall Survival				Total
	>12 mo (Superior)	≤12, >8 mo (Good)	≤8, >4 mo (Moderate)	≤4 mo (Poor)	
DTH positive	8*	3	6	0	17*
DTH negative	1	4	1	7	13
Total	9	7	7	7	30


$\chi^2 = 15.908$ ,  $P = 0.0012$ .  
\*One patient was excluded from this analysis.  
DTH indicates delayed-type hypersensitivity.

months in the gemcitabine alone group were 8.8 months and 35.4%, respectively. These results seemed a little better than those in our study. One reason for this may be the difference in the proportion of the patients with the locally advanced pancreatic cancer, in which survival data were apparently much better than those in metastatic ones. In our study, this proportion was 18.8%, which was lower than that in GEST study (23.8%). The other reason may be PS at baseline, which was also one of the important prognostic factors. The proportion of the patients with ECOG-

PS 0, 1, and 2 at baseline in our study were 46.9%, 31.3%, and 21.9%, respectively, whereas those in GEST study were 65.3%, 34.7%, and 0.0%, respectively. It is apparent that our patients are predicted to worse prognosis than those in GEST study. Despite lower proportion of locally advanced stage and worse PS, however, the survival data gained from the patients with DTH-positivity seemed to be better than those in GEST study. These results suggested additional or synergistic effects of WT1 vaccine. Although the number of patients in our present study was too small to reach any



**FIGURE 3.** Analysis of WT1-specific immune response. A, Immunologic monitoring of the phenotype analysis of WT1 tetramer<sup>+</sup> CD3<sup>+</sup>CD8<sup>+</sup> T lymphocytes (WT1-CTLs) in DTH-positive (light gray columns) and DTH-negative patients (dark gray columns). B, Immunologic monitoring of the phenotype analysis of WT1 tetramer<sup>+</sup>CD3<sup>+</sup>CD8<sup>+</sup> T lymphocytes (WT1-CTLs) in the patients of 4 groups classified according to overall survival time. The broken line represents the median percentage of memory-phenotype WT1-CTLs at baseline for all patients. WT1 tetramer=PE-conjugated WT1<sub>235</sub> tetramer [HLA-A\*24:02-restricted natural 9-mer WT1 peptide (CMTWNQMNL)], naïve (CD45RA<sup>+</sup>CCR7<sup>+</sup>), memory (CD45RA-CCR7<sup>+</sup> or CD45RA-CCR7<sup>-</sup>), and effector (CD45RA<sup>+</sup>CCR7<sup>-</sup>). 2nd indicates day 1 in the second course; 3rd, day 1 in the third course; B, baseline; DTH, delayed-type hypersensitivity.



**FIGURE 4.** Clinical course and immunologic monitoring of 1 patient. A, Abdominal computed tomography (CT) scan before and after treatment. Left: CT scan at baseline showed a large hypodense lesion in the head of the pancreas, which had also invaded the supramesenteric artery and portal vein. Right: 5 months after treatment (before operation), a follow-up CT scan showed >80% regression of the primary lesion. Gray arrows shows primary lesion of pancreas. B, Microscopic findings of the resected specimen (hematoxylin-eosin stain). C, Clinical course and immunologic monitoring. The black line represents the absolute number of WT1 tetramer<sup>+</sup>CD3<sup>+</sup>CD8<sup>+</sup> T lymphocytes (WT1-CTLs), and the gray column represents the percentage of memory-phenotype WT1-CTLs. D, Percentages of WT1-CTLs in the peripheral blood (PB) and tumor-infiltrating lymphocytes (TIL). CA19-9 indicates carbohydrate antigen 19-9; CTLs, cytotoxic T lymphocytes; DTH, delayed-type hypersensitivity; GEM, gemcitabine; KPS, Karnofsky performance status.

definitive conclusions about clinical efficacy, these findings have been sufficiently encouraging to prompt us to conduct a further clinical study to determine the potency of this combination therapy. No combination chemotherapy, with the exception of FOLFIRINOX,<sup>34</sup> has resulted in a significant improvement in survival of patients with pancreatic cancer although some combination therapies are thought to be more effective for several cancers than single-agent treatments.<sup>1</sup> The use of FOLFIRINOX, however, may have to be limited to patients with good performance status as this regimen has much higher toxicity that sometimes can impair QOL.<sup>34,35</sup> In contrast, as toxicities associated with cancer vaccines are generally mild and acceptable, combination therapies using chemotherapy and cancer vaccine can be

expected to exert their clinical benefits without worsening of QOL, which is often impaired by combination chemotherapy using several kinds of cytotoxic agents.

Immunologic monitoring is an important step in the development of evidence-based immunotherapy. Our data provided 2 useful prognostic markers of better clinical outcomes for the combination therapy used in our study. One is DTH to WT1 peptide and the other the frequency of memory-phenotype WT1-CTLs in PB although we did not find the correlation between clinical effects, including survival, and the frequency or absolute numbers of nonphenotypically divided WT1-specific CTLs statistically (data not shown). DTH-positive patients had a notably better prognosis than DTH-negative patients, and the OS curve for DTH-positive

patients showed a late separation beyond the median. As DTH has long been used for evaluation of antigen memory for bacterial, viral, and cancer antigens,<sup>36</sup> the occurrence of DTH to WT1 peptide may reflect the development and persistence of memory-phenotype WT1-CTLs. This can be inferred from our observation that DTH-positive patients showed a significantly higher frequency of memory-phenotype WT1-CTLs than did DTH-negative patients after WT1 vaccination. Furthermore, patients who survived 12 months or longer (superior responders) seemed to have the highest frequency of memory-phenotype WT1-CTLs in their PB although the number of patients in each subgroup was too small to make a statistically valid comparison. It was reported that long-term survivors who had been treated with mutant K-ras vaccine against pancreatic cancer showed the persistence of vaccinated peptide-recognizing T cells (long-term T-cell memory response) for many years after the last vaccination.<sup>37</sup> This report and our results suggest that the development and persistence of TAA-specific CTLs with memory-phenotype resulting from treatment with cancer vaccine contributed to the longer survival. Further investigations are needed to validate these findings in the larger-scale clinical trial.

Despite its potent cytotoxicity, gemcitabine reportedly has immune-modulating functions, such as increase in antigen cross-presentation,<sup>38</sup> and inhibition of B-cells,<sup>39</sup> myeloid-derived suppressive cells,<sup>40</sup> and regulatory T cells,<sup>41</sup> resulting in enhancement of the antigen-specific CTL function. Recently, we reported that gemcitabine enhanced the WT1 expression on human pancreatic cancer cells thus sensitizing the cancer cells to WT1-specific CTL.<sup>11</sup> Furthermore, it was reported that lymphopenia-induced memory-phenotype WT1-CTLs from naive-phenotype WT1-CTLs without self-antigen-induced tolerance.<sup>42</sup> Transient mild to moderate lymphopenia induced by gemcitabine and immediate recovery of T cells could thus promote both the differentiation of naive-phenotype WT1-CTLs into memory-phenotype WT1-CTLs and their proliferation in the clinical application of the combination therapy of gemcitabine and WT1 vaccine. In view of these immunostimulatory properties of gemcitabine, this combination therapy can be expected to generate additional or synergistic effects.

In conclusion, the combination of WT1 vaccine with the standard gemcitabine therapy was well tolerated for patients with advanced pancreatic cancer. WT1 vaccine might have additional effects on gemcitabine to improve survival benefit. An increase in memory-phenotype WT1-CTLs could be a useful predictive marker for a favorable clinical outcome. To determine the clinical efficacy of this combination therapy, we have started a phase 2 randomized clinical study (UMIN000005248).

#### ACKNOWLEDGMENTS

The authors thank the member of the nursing team for their care for the patients in this trial, Hiroko Nakajima, and Fumihiro Fujiki for their technical supports, Tomoe Umeda for her coordination of clinical research, Takushi Okusaka and Yuji Heike (National Cancer Center in Japan) for their advices about the planning of this study.

#### CONFLICTS OF INTEREST/ FINANCIAL DISCLOSURES

This study was supported by the Japanese Ministries of Education, Culture, Sports, Science and Technology, and of

Health, Labor and Welfare; the National Cancer Center Research and Development Fund; the Takeda Science Foundation; and Mitsui Life Social Welfare Foundation, Promotion of Cancer Research.

All authors have declared there are no financial conflicts of interest with regard to this work.

#### REFERENCES

1. Stathis A, Moore MJ. Advanced pancreatic carcinoma: current treatment and future challenges. *Nat Rev Clin Oncol*. 2010;7:163–172.
2. Burris HA III, Moore MJ, Andersen J, et al. Improvements in survival and clinical benefit with gemcitabine as first-line therapy for patients with advanced pancreas cancer: a randomized trial. *J Clin Oncol*. 1997;15:2403–2413.
3. Moore MJ, Goldstein D, Hamm J, et al. Erlotinib plus gemcitabine compared with gemcitabine alone in patients with advanced pancreatic cancer: a phase III trial of the National Cancer Institute of Canada Clinical Trials Group. *J Clin Oncol*. 2007;25:1960–1966.
4. Disis ML, Bernhard H, Jaffee EM. Use of tumour-responsive T cells as cancer treatment. *Lancet*. 2009;373:673–683.
5. Mellman I, Coukos G, Dranoff G. Cancer immunotherapy comes of age. *Nature*. 2011;480:480–489.
6. Rosenberg SA, Yang JC, Restifo NP. Cancer immunotherapy: moving beyond current vaccines. *Nat Med*. 2004;10:909–915.
7. Baxevas CN, Perez SA, Papamichail M. Combinatorial treatments including vaccines, chemotherapy and monoclonal antibodies for cancer therapy. *Cancer Immunol Immunother*. 2009;58:317–324.
8. Nowak AK, Lake RA, Robinson BWS. Combined chemotherapeutic of solid tumours: improving vaccines? *Adv Drug Deliv Rev*. 2006;58:975–990.
9. Zitvogel L, Apetoh L, Ghiringhelli F, et al. Immunological aspects of cancer chemotherapy. *Nat Rev Immunol*. 2008;8:59–73.
10. Ramakrishnan R, Assudani D, Nagaraj S, et al. Chemotherapy enhances tumor cell susceptibility to CTL-mediated killing during cancer immunotherapy in mice. *J Clin Invest*. 2010;120:1111–1124.
11. Takahara A, Koido S, Ito M, et al. Gemcitabine enhances Wilms' tumor gene WT1 expression and sensitizes human pancreatic cancer cells with WT1-specific T-cell-mediated antitumor immune response. *Cancer Immunol Immunother*. 2011;60:1289–1297.
12. Huff V. Wilms' tumours: about tumour suppressor genes, an oncogene and a chameleon gene. *Nature Rev Cancer*. 2011;11:111–121.
13. Sugiyama H. WT1 (Wilms' tumor gene 1): biology and cancer immunotherapy. *Jpn J Clin Oncol*. 2010;40:377–387.
14. Elisseeva OA, Oka Y, Tsuboi A, et al. Humoral immune responses against Wilms' tumor gene WT1 product in patients with hematopoietic malignancies. *Blood*. 2002;99:3272–3279.
15. Gaiger A, Carter L, Greinix H, et al. WT1-specific serum antibodies in patients with leukemia. *Clin Cancer Res*. 2001;7:761s–765s.
16. Oji Y, Kitamura Y, Kamino E, et al. WT1 IgG antibody for early detection of nonsmall cell lung cancer and as its prognostic factor. *Int J Cancer*. 2009;125:381–387.
17. Oka Y, Elisseeva OA, Tsuboi A, et al. Human cytotoxic T-lymphocyte responses specific for peptides of the wild-type Wilms' tumor gene (WT1) product. *Immunogenetics*. 2000;51:99–107.
18. Rezvani K, Brenchley JM, Price DA, et al. T-cell responses directed against multiple HLA-A\*0201-restricted epitopes derived from Wilms' tumor 1 protein in patients with leukemia and healthy donors: identification, quantification, and characterization. *Clin Cancer Res*. 2005;11:8799–8807.
19. Oka Y, Tsuboi A, Taguchi T, et al. Induction of WT1 (Wilms' tumor gene)-specific cytotoxic T lymphocytes by WT1 peptide vaccine and the resultant cancer regression. *Proc Natl Acad Sci USA*. 2004;101:13885–13890.



20. Morita S, Oka Y, Tsuboi A, et al. A phase I/II trial of a WT1 (Wilms' tumor gene) peptide vaccine in patients with solid malignancy: safety assessment based on the phase I data. *Jpn J Clin Oncol*. 2006;36:231–236.
21. Izumoto S, Tsuboi A, Oka Y, et al. Phase II clinical trial of Wilms tumor 1 peptide vaccination for patients with recurrent glioblastoma multiforme. *J Neurosurg*. 2008;108:963–971.
22. Miyatake T, Ueda Y, Morimoto A, et al. WT1 peptide immunotherapy for gynecologic malignancies resistant to conventional therapies: a phase II trial. *J Cancer Res Clin Oncol*. 2013;139:457–463.
23. Tsuboi A, Oka Y, Kyo T, et al. Long-term WT1 peptide vaccination for patients with acute myeloid leukemia with minimal residual disease. *Leukemia*. 2012;26:1410–1413.
24. Hashii Y, Sato-Miyashita E, Matsumura R, et al. WT1 peptide vaccination following allogeneic stem cell transplantation in pediatric leukemic patients with high risk for relapse: successful maintenance of durable remission. *Leukemia*. 2012;26:530–532.
25. Keilholz U, Letsch A, Busse A, et al. A clinical and immunologic phase 2 trial of Wilms tumor gene product 1 (WT1) peptide vaccination in patients with AML and MDS. *Blood*. 2009;113:6541–6548.
26. Rezvani K, Yong AS, Mielke S, et al. Leukemia-associated antigen-specific T-cell responses following combined PR1 and WT1 peptide vaccination in patients with myeloid malignancies. *Blood*. 2008;111:236–242.
27. Cella DF, Tulskey DS, Gray G, et al. The functional assessment of cancer therapy scale: development and validation of the general measure. *J Clin Oncol*. 1993;11:570–579.
28. Khorana AA, Fine RL. Pancreatic cancer and thromboembolic disease. *Lancet Oncol*. 2004;5:655–663.
29. Otten HM, Mathijssen J, ten Cate H, et al. Symptomatic venous thromboembolism in cancer patients treated with chemotherapy: an underestimated phenomenon. *Arch Intern Med*. 2004;164:190–194.
30. Okada S, Ueno H, Okusaka T, et al. Phase I trial of gemcitabine in patients with advanced pancreatic cancer. *Jpn J Clin Oncol*. 2001;31:7–12.
31. Gaiger A, Reese V, Disis ML, et al. Immunity to WT1 in the animal model and in patients with acute myeloid leukemia. *Blood*. 2000;96:1480–1489.
32. Gao L, Bellantuono I, Elsassner A, et al. Selective elimination of leukemic CD34 + progenitor cells by cytotoxic T lymphocytes specific for WT1. *Blood*. 2000;95:2198–2203.
33. Ueno H, Ioka T, Ikeda M, et al. Randomized phase III study of gemcitabine plus S-1, S-1 alone, or gemcitabine alone in patients with locally advanced and metastatic pancreatic cancer in Japan and Taiwan: GEST study. *J Clin Oncol*. 2013;31:1640–1648.
34. Conroy T, Desseigne F, Ychou M, et al. FOLFIRINOX versus gemcitabine for metastatic pancreatic cancer. *N Engl J Med*. 2011;364:1817–1825.
35. Gourgou-Bourgade S, Bascoul-Mollevi C, Desseigne F, et al. Impact of FOLFIRINOX compared with gemcitabine on quality of life in patients with metastatic pancreatic cancer: results from the PRODIGE 4/ACCORD 11 randomized trial. *J Clin Oncol*. 2013;31:23–29.
36. Disis ML. Immunologic biomarkers as correlates of clinical response to cancer immunotherapy. *Cancer Immunol Immunother*. 2011;60:433–442.
37. Wedén S, Klemp M, Gladhaug IP, et al. Long-term follow-up of patients with resected pancreatic cancer following vaccination against mutant K-ras. *Int J Cancer*. 2011;128:1120–1128.
38. Nowak AK, Lake RA, Marzo AL, et al. Induction of tumor cell apoptosis in vivo increases tumor antigen cross-presentation, cross-priming rather than cross-tolerizing host tumor-specific CD8 T cells. *J Immunol*. 2003;170:4905–4913.
39. Nowak AK, Robinson BWS, Lake RA. Gemcitabine exerts a selective effect on the humoral immune response: implications for combination chemo-immunotherapy. *Cancer Res*. 2002;62:2353–2358.
40. Suzuki E, Kapoor V, Jassar AS, et al. Gemcitabine selectively eliminates splenic Gr-1 + /CD11b + myeloid suppressor cells in tumor-bearing animals and enhances antitumor immune activity. *Clin Cancer Res*. 2005;11:6713–6721.
41. Rettig L, Seidenberg S, Parvanova I, et al. Gemcitabine depletes regulatory T-cells in human and mice and enhances triggering of vaccine-specific cytotoxic T-cells. *Int J Cancer*. 2010;129:832–838.
42. Pospori C, Xue SA, Holler A, et al. Specificity for the tumor-associated self-antigen WT1 drives the development of fully functional memory T cells in the absence of vaccination. *Blood*. 2011;117:6813–6824.

LETTER TO THE EDITOR

Maintenance of complete remission after allogeneic stem cell transplantation in leukemia patients treated with Wilms tumor 1 peptide vaccine

Blood Cancer Journal (2013) 3, e130; doi:10.1038/bcj.2013.29; published online 2 August 2013

The prognosis of patients after allogeneic hematopoietic stem cell transplantation (HSCT) is still not satisfactory because, while treatment-related mortalities have decreased, relapse after HSCT remains a major concern. The effectiveness of allogeneic HSCT for hematological malignancies is the result of immunologic rejection of recipient leukemia cells by donor T cells, known as the graft-versus-leukemia (GVL) effect.<sup>1</sup> It is thus obviously important to be able to exploit the GVL effect while minimizing graft-versus-host disease (GVHD). A targeted anti-leukemic immunotherapy, such as use of a leukemia vaccine,<sup>2</sup> is a promising strategy to boost the GVL effect.

Wilms tumor 1 (WT1) protein is one of the best targets for leukemia vaccines. Overexpression of the wild-type *WT1* gene has been detected in all types of human leukemia.<sup>3–5</sup> We performed a phase I clinical study of immunotherapy targeting the WT1 protein in patients with leukemia, and were able to show that WT1 vaccination was safe and could induce WT1-specific cytotoxic T lymphocyte (CTL).<sup>6</sup> Furthermore, reduction of minimal residual disease and long-lasting complete remission (CR) was observed in some leukemia patients who were given the WT1 vaccine.<sup>7</sup>


This report presents the results of phase I clinical study of WT1 vaccination for HLA-A\*2402-positive post-HSCT patients who were at high risk of relapse (HSCT in non-CR and 2nd HSCT for post-transplant relapse) or had already relapsed. The HLA-A\*2402-restricted modified 9-mer WT1 peptide (amino acids 235–243 CYTWNQMNL)<sup>8</sup> was emulsified with Montanide ISA51 adjuvant. Patients were intradermally injected with 1.0 mg (three patients: UPNs 1, 4 and 6) or 3.0 mg (other six patients) of WT1 peptide four times weekly. When no adverse effects and no obvious disease progression were observed after the fourth injection, further WT1 vaccinations at 2-week intervals were administered.

Nine patients (five with acute myeloid leukemia (AML), one each with acute lymphoblastic leukemia, chronic myelomonocytic leukemia, multiple myeloma and T-cell lymphoblastic lymphoma) were enrolled in this study (Supplementary Tables 1 and 2). Local inflammatory response was observed at the vaccine injection sites of all patients. One patient (UPN5) suffered mild hypoxia (PaO<sub>2</sub> 65 mm Hg at room air) and restrictive pulmonary dysfunction (FEV<sub>1.0</sub> 40%) 65 days after the start of WT1 vaccination (day 199 after HSCT; Figure 1a). He was diagnosed with bronchioleitis obliterans (BO), which was a symptom of chronic GVHD. The patient recovered soon after administration of inhaled steroids. While early and sudden discontinuation of prednisolone and tacrolimus (day 103 after HSCT) were considered to be the reason for development of BO, the possibility of an association between BO and WT1 vaccination cannot be entirely ruled out. In other eight patients, no severe toxicities related to WT1 vaccine were observed (Table1).

Three AML patients (UPN1–3), who had undergone HSCT in non-CR, started WT1 vaccine in CR (Supplementary Tables 1 and 2). They started WT1 vaccination on post-HSCT days 141, 76 and 93

and have remained in CR for 1038, 973 and 662 days, respectively (as of 8 April 2013; Table1), suggesting the potential of WT1 vaccination as a maintenance therapy after HSCT.

Six patients started WT1 vaccination in non-CR and two of them became CR after WT1 vaccination. One B-ALL patient (UPN4) with MLL-AF4 underwent bone marrow transplantation from an HLA-matched unrelated donor during the first CR. On post-HSCT day 111, MLL-AF4 and WT1 mRNA in peripheral blood (PB) had increased to 16 000 and 15 000 copies/μg RNA, indicating that the disease had relapsed. Tacrolimus and prednisolone doses were tapered off to induce GVL effects. The expression levels of MLL-AF4 and WT1 mRNA in PB had decreased to 2700 and 190



**Figure 1.** Clinical course of patients who attained CR after the start of WT1 peptide vaccination. (a) Clinical course of UPN5 who achieved CR after administration of WT1 vaccine but stopped vaccination because of the development of bronchioleitis obliterans. (b) Clinical course of UPN4. Residual leukemia cells that were detected by MLL/AF4 expression disappeared after the start of WT1 vaccination. In both cases, rapid tapering of immune-suppressive drugs preceded WT1 peptide vaccination.

**Table 1.** Patient outcomes

UPN	Disease	Status before vaccination	Adverse events	Number of vaccine doses	Outcome	Additional therapy	Survival	
							Post-HSCT	After start of vaccination
1	AML (M4)	CR	None	54	CR	—	1179 +	1038 +
2	AML(M4, DEK/CAN +)	CR	PLT↓	52	CR	—	1049 +	973 +
3	AML	CR	None	38	CR	—	759 +	662 +
4	B-ALL (MLL/AF4 +)	Molecular relapse	None	71	CR	—	1312 +	1179 +
5	AML (M4)	Relapse	Amylase↑, bronchileitis obliterans (cGVHD) <sup>a</sup>	2	CR	—	972 +	842 + <sup>b</sup>
6	CMMoL	Relapse	None	25	PD <sup>c</sup>	Chemo	2265	381
7	MM	PD	None	19	PD	Chemo	1301 +	804 +
8	T-LBL	Relapse	None	4	PD	Second transplant	955	656
9	AML (M2)	Relapse	None	17	PD	Second transplant	1544 +	749 +

Abbreviations: ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; CMMoL, chronic myelomonocytic leukemia; CR, complete remission; cGVHD, chronic graft-versus-host disease; HSCT, hematopoietic stem cell transplantation; MM, multiple myeloma; PD, progressive disease; T-LBL, T-cell lymphoblastic lymphoma. (8 April 2013). <sup>a</sup>A causal relationship between vaccination and this event was not strongly suspected, but could not be ruled out. <sup>b</sup>Vaccination was discontinued. (The last injection was on post-HSCT day 60). <sup>c</sup>Size of the subcutaneous tumor decreased, but the disease relapsed in axil lymph nodes and stomach.

copies/ $\mu$ g RNA by day 132, and WT1 vaccination was started on day 133. MLL-AF4 mRNA had become undetectable by day 146, and had never appeared until post-HSCT day 1312 (day 1179 after the start of WT1 vaccination as of 8 April 2013; Figure 1b).

Skin tumors appeared in UPN5 (AML-M5) on post-HSCT day 103 and was diagnosed by biopsy as leukemia relapse. Tacrolimus was discontinued on day103, and WT1 vaccination was started on day 130. Cutaneous tumors had regressed 2 weeks after the start of WT1 vaccination, but vaccination was terminated after the second injection because of the development of BO as described earlier (Figure 1a). This patient has been remained in CR until post-HSCT day 972 (day 842 after the start of WT1 vaccination at 8 April 2013). While the exact contribution of the vaccination effect to the disease remission in addition to the GVL effect was unclear, the fact that both of these two patients still have remained in CR until now is encouraging to continue this trial. In the following phase II trials, the enumeration of WT1-specific CTLs should be performed more frequently after the start of vaccination to clarify the relationship between the effect of WT1 peptide vaccination and leukemia regression.

WT1 (a natural 9-mer WT1 peptide) HLA-A\*2402 tetramer assays could be performed with peripheral blood mononuclear cell in seven of the nine patients to determine whether WT1<sub>235</sub> peptide-specific CD8<sup>+</sup> T cells had increased after WT1 vaccination. The gates for WT1 tetramer<sup>+</sup> cells were drawn as <0.1% of CD8<sup>+</sup> T cells were included in the tetramer-positive gate in multiple healthy individuals (Supplementary Figure 1A). WT1<sub>235</sub> tetramer<sup>+</sup> cells increased after the start of vaccination in three (UPNs1, 2 and 4) of the four patients who have remained in CR (Figure 1b and Supplementary Figure 1B). In the cases with progressive disease, continuous increase in the frequencies of WT1<sub>235</sub> tetramer<sup>+</sup> cells was not observed (Supplementary Figure 1B).

Our results suggest that WT1 vaccination should be started when the leukemia burden is minimal. The timing of the start of WT1 vaccination may be also important. For the cases with good outcomes, WT1 vaccination was started 76–140 days after transplantation (UPNs1–5), and at later times (days 299–1815) for PD cases (UPNs 6–9). A lymphopenic environment a few months after transplantation may be favorable for rapid and extensive expansion of tumor antigen-specific CTLs.

In summary, this report suggests that WT1 vaccine can be safely administrated for post-HSCT patients with hematological malignancies and has potential as a maintenance therapy. Clinical benefit of WT1 vaccination for post-HSCT patients will be evaluated in the subsequent phase II trials.

#### CONFLICT OF INTEREST

The authors declare no conflict of interest.

T Maeda<sup>1</sup>, N Hosen<sup>2,3</sup>, K Fukushima<sup>1</sup>, A Tsuboi<sup>4</sup>, S Morimoto<sup>2</sup>, T Matsui<sup>1</sup>, H Sata<sup>1</sup>, J Fujita<sup>1</sup>, K Hasegawa<sup>2</sup>, S Nishida<sup>4</sup>, J Nakata<sup>5</sup>, Y Nakae<sup>5</sup>, S Takashima<sup>5</sup>, H Nakajima<sup>6</sup>, F Fujiki<sup>6</sup>, N Tatsumi<sup>3</sup>, T Kondo<sup>7</sup>, M Hino<sup>8</sup>, Y Oji<sup>3</sup>, Y Oka<sup>5</sup>, Y Kanakura<sup>1</sup>, A Kumanogoh<sup>5</sup> and H Sugiyama<sup>2</sup>

<sup>1</sup>Department of Hematology and Oncology, Osaka University Graduate School of Medicine, Osaka, Japan;

<sup>2</sup>Department of Functional Diagnostic Science, Osaka University Graduate School of Medicine, Osaka, Japan;

<sup>3</sup>Department of Cancer Stem Cell Biology, Osaka University Graduate School of Medicine, Osaka, Japan;

<sup>4</sup>Department of Cancer Immunotherapy, Osaka University Graduate School of Medicine, Osaka, Japan;

<sup>5</sup>Department of Respiratory Medicine, Allergy and Rheumatic Diseases, Osaka University Graduate School of Medicine, Osaka, Japan;

<sup>6</sup>Department of Cancer Immunology, Osaka University Graduate School of Medicine, Osaka, Japan;

<sup>7</sup>Department of Hematology and Oncology, Kyoto University Graduate School of Medicine, Kyoto, Japan and

<sup>8</sup>Department of Hematology and Oncology, Osaka City University Graduate School of Medicine, Osaka, Japan  
E-mail: sugiyama@sahs.med.osaka-u.ac.jp

#### REFERENCES

- Weiden PL, Flournoy N, Thomas ED, Prentice R, Fefer A, Buckner CD *et al*. Antileukemic effect of graft-versus-host disease in human recipients of allogeneic-marrow grafts. *N Engl J Med* 1979; **300**: 1068–1073.
- Rezvani K, Yong AS, Mielke S, Savani BN, Musse L, Superata J *et al*. Leukemia-associated antigen-specific T-cell responses following combined PR1

- and WT1 peptide vaccination in patients with myeloid malignancies. *Blood* 2008; **111**: 236–242.
- 3 Cilloni D, Gottardi E, De Micheli D, Serra A, Volpe G, Messa F *et al.* Quantitative assessment of WT1 expression by real time quantitative PCR may be a useful tool for monitoring minimal residual disease in acute leukemia patients. *Leukemia* 2002; **16**: 2115–2121.
- 4 Miwa H, Beran M, Saunders GF. Expression of the Wilms' tumor gene (WT1) in human leukemias. *Leukemia* 1992; **6**: 405–409.
- 5 Inoue K, Sugiyama H, Ogawa H, Nakagawa M, Yamagami T, Miwa H *et al.* WT1 as a new prognostic factor and a new marker for the detection of minimal residual disease in acute leukemia. *Blood* 1994; **84**: 3071–3079.
- 6 Oka Y, Tsuboi A, Taguchi T, Osaki T, Kyo T, Nakajima H *et al.* Induction of WT1 (Wilms' tumor gene)-specific cytotoxic T lymphocytes by WT1 peptide

vaccine and the resultant cancer regression. *Proc Natl Acad Sci USA* 2004; **101**: 13885–13890.

- 7 Tsuboi A, Oka Y, Kyo T, Katayama Y, Elisseeva OA, Kawakami M *et al.* Long-term WT1 peptide vaccination for patients with acute myeloid leukemia with minimal residual disease. *Leukemia* 2012; **26**: 1410–1413.
- 8 Tsuboi A, Oka Y, Udaka K, Murakami M, Masuda T, Nakano A *et al.* Enhanced induction of human WT1-specific cytotoxic T lymphocytes with a 9-mer WT1 peptide modified at HLA-A\*2402-binding residues. *Cancer Immunol Immunother* 2002; **51**: 614–620.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>

Supplementary Information accompanies this paper on Blood Cancer Journal website (<http://www.nature.com/bcj>)

## Functional human Th17 clones with WT1-specific helper activity

Sho Tachino · Fumihiro Fujiki · Yoshihiro Oka · Akihiro Tsuboi · Soyoko Morimoto · Yu-Hung Lin · Taichi Tamanaka · Kenta Kondo · Hiroko Nakajima · Sumiyuki Nishida · Naoki Hosen · Yusuke Oji · Atsushi Kumanogoh · Haruo Sugiyama

Received: 3 September 2012 / Accepted: 7 December 2012 / Published online: 29 December 2012  
© Springer-Verlag Berlin Heidelberg 2012

**Abstract** Th17 plays important roles in the pathogenesis of various inflammatory and autoimmune diseases. Although the importance of Th17 in tumor immunity has also been suggested, precise roles of tumor-associated antigen-specific Th17 still remain poorly understood, especially in humans. We previously identified WT1<sub>332</sub>, a 16-mer helper epitope derived from tumor-associated antigen Wilms' tumor gene 1 (WT1) product, and WT1<sub>332</sub>-specific Th1 clones were established. In the present study, WT1-specific Th17 clones were established by the stimulation of

peripheral blood mononuclear cells with the WT1<sub>332</sub> helper peptide under human Th17-polarizing conditions. The WT1-specific Th17 clone exhibited the helper function for proliferation of conventional CD4<sup>+</sup> T cells in the antigenic stimulation-specific manner. This is the first report of establishment of functional Th17 clones with both antigen (WT1<sub>332</sub>) specificity and antigen-specific helper activity. Th17 clones established here and the method to establish antigen-specific Th17 clones should be a useful tool to further analyze the roles of human Th17 in tumor immunity.

S. Tachino · Y. Oka · Y.-H. Lin · T. Tamanaka · A. Kumanogoh  
Department of Respiratory Medicine, Allergy and Rheumatic Diseases, Osaka University Graduate School of Medicine, 2-2 Yamadaoka, Suita, Osaka, Japan  
e-mail: stachino@imed3.med.osaka-u.ac.jp

Y. Oka  
e-mail: yoshi@imed3.med.osaka-u.ac.jp

Y.-H. Lin  
e-mail: s3327501@gmail.com

T. Tamanaka  
e-mail: dtamanaka@imed3.med.osaka-u.ac.jp

A. Kumanogoh  
e-mail: kumanogo@imed3.med.osaka-u.ac.jp

F. Fujiki · H. Nakajima  
Department of Cancer Immunology, Osaka University Graduate School of Medicine, 2-2 Yamadaoka, Suita, Osaka, Japan  
e-mail: fu-fuji@sahs.med.osaka-u.ac.jp

H. Nakajima  
e-mail: hnakaji@cit.med.osaka-u.ac.jp

Y. Oka · A. Kumanogoh  
Department of Immunopathology, WPI Immunology Frontier Research Center, Osaka University, 3-1 Yamadaoka, Suita, Osaka, Japan

A. Tsuboi · S. Nishida  
Department of Cancer Immunotherapy, Osaka University Graduate School of Medicine, 3-1 Yamadaoka, Suita, Osaka, Japan  
e-mail: tsuboi@cit.med.osaka-u.ac.jp

S. Nishida  
e-mail: sumiyuki-n@imed3.med.osaka-u.ac.jp

S. Morimoto · K. Kondo · N. Hosen · H. Sugiyama (✉)  
Department of Functional Diagnostic Science, Osaka University Graduate School of Medicine, 1-7 Yamadaoka, Suita, Osaka, Japan  
e-mail: sugiyama@sahs.med.osaka-u.ac.jp

S. Morimoto  
e-mail: soyoko@imed3.med.osaka-u.ac.jp

K. Kondo  
e-mail: k.kondo@sahs.med.osaka-u.ac.jp

N. Hosen  
e-mail: hnaoki@imed3.med.osaka-u.ac.jp

Y. Oji  
Department of Cancer Stem Cell Biology, Osaka University Graduate School of Medicine, 1-7 Yamadaoka, Suita, Osaka, Japan  
e-mail: oji@sahs.med.osaka-u.ac.jp