

総説

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Oze T, Hiramatsu N, Yakushijin T, Miyazaki M, Yamada A, Oshita M, Hagiwara H, Mita E, Ito T, Fukui H, Inui Y, Hijioka T, Inada M, Katayama K, Tamura S, Yoshihara H, Inoue A, Imai Y, Hayashi E, Kato M, Miyagi T, Yoshida Y, Tatsumi T, Kasahara A, Hamasaki T, Hayashi N, Takehara T; the Osaka Liver Forum.	Post-treatment levels of α -fetoprotein predict Incidence of hepatocellular carcinoma after interferon therapy.	Clin Gastroenterol Hepatol.	In press.		2013 Dec

総説(和文)

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
豊田 秀徳、熊田 卓.	HCV 治療後発癌例の解析(C 型肝炎治療 2014:経口抗ウイルス薬時代の到来, HCV 駆除後肝発癌の対策).	肝胆膵.	67(6)	1035-1039	2013 Dec
髭 修平、木村 睦海、荒川 智宏、小関 至、狩野 吉康、豊田 成司.	肝線維化の定量的測定(検査値の読み方 連載).	臨床消化器内科.	28(13)	1783-1788	2013 Dec
多田 俊史、熊田 卓、豊田 秀徳.	超高齢肝細胞癌患者に対する治療法の選択と予後 -後期高齢者との比較検討-.	The Liver Cancer Journal.	5(4)	268-272	2013 Dec
山本 健太、熊田 卓、桐山 勢生、谷川 誠、久永康宏、豊田 秀徳、金森 明、多田 俊史、北畠 秀介、長谷川 綾平、伊藤 隆徳、安藤 祐資、颯田 祐介、田中 達也、曾根 康	肝細胞癌と鑑別が困難であった reactive lymphoid hyperplasia の 1 例.	肝臓.	54(11)	755-764	2013 Nov

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
博、福里 利夫.					
多田 俊史、熊田 卓、豊田 秀徳.	C型肝硬変例における分岐鎖アミノ酸製剤と肝発癌との関連 -Propensity score 分析を用いた検討-.	消化と吸収.	35(3)	290-294	2013 Jul
松田 厚志、久野 敦、成松 久、監修：第62回日本医学検査学会記念誌編集委員会.	グライコプロテオミクスによる臨床糖鎖腫瘍マーカー開発 -グライコームからグライコプロテオームへ.	検査診断学への展望 -臨床検査指針：測定とデータ判読のポイント-.			2013 May

IV.研究成果の刊行物・別刷

A novel serum marker, glycosylated *Wisteria floribunda* agglutinin-positive Mac-2 binding protein (WFA⁺-M2BP), for assessing liver fibrosis

Takeo Toshima · Ken Shirabe · Toru Ikegami · Tomoharu Yoshizumi ·
Atsushi Kuno · Akira Togayachi · Masanori Gotoh · Hisashi Narimatsu ·
Masaaki Korenaga · Masashi Mizokami · Akihito Nishie · Shinichi Aishima ·
Yoshihiko Maehara

Received: 1 November 2013 / Accepted: 7 February 2014
© Springer Japan 2014

Abstract

Background Recently, a novel marker, hyperglycosylated *Wisteria floribunda* agglutinin-positive Mac-2 binding protein (WFA⁺-M2BP), was developed for liver fibrosis using the glycan “sugar chain”-based immunoassay; however, the feasibility of WFA⁺-M2BP for assessing liver fibrosis has not been proven with clinical samples of hepatitis.

Methods Serum WFA⁺-M2BP values were evaluated in 200 patients with chronic liver disease who underwent histological examination of liver fibrosis. The diagnostic

accuracy of WFA⁺-M2BP values was compared with various fibrosis markers, such as ultrasound based-virtual touch tissue quantification (VTTQ), magnetic resonance imaging based-liver-to-major psoas muscle intensity ratio (LMR), and serum markers, including hyaluronic acid, type 4 collagen, and aspartate transaminase to platelet ratio index (APRI).

Results Serum WFA⁺-M2BP levels in patients with fibrosis grades F0, F1, F2, F3, and F4 had cutoff indices 1.62, 1.82, 3.02, 3.32, and 3.67, respectively, and there were significant differences between fibrosis stages F1 and F2, and between F2 and F3 ($P < 0.01$). The area under the receiver operating characteristic curves for the diagnosis of fibrosis ($F \geq 3$) using serum WFA⁺-M2BP values (0.812) was almost comparable to that using VTTQ examination (0.814), but was superior to the other surrogate markers, including LMR index (0.766), APRI (0.694), hyaluronic acid (0.683), and type 4 collagen (0.625) ($P < 0.01$ each). **Conclusions** Serum WFA⁺-M2BP values based on a glycan-based immunoassay is an accurate, reliable, and reproducible method for the assessment of liver fibrosis. This approach could be clinically feasible for evaluation of beneficial therapy through the quantification of liver fibrosis in hepatitis patients if this measurement application is commercially realized.

Electronic supplementary material The online version of this article (doi:10.1007/s00535-014-0946-y) contains supplementary material, which is available to authorized users.

T. Toshima · K. Shirabe (✉) · T. Ikegami · T. Yoshizumi ·
Y. Maehara

Department of Surgery and Science, Graduate School of Medical
Sciences, Kyushu University, 3-1-1 Maidashi, Higashi-ku,
Fukuoka 812-8582, Japan
e-mail: kshirabe@surg2.med.kyushu-u.ac.jp

A. Kuno · A. Togayachi · M. Gotoh · H. Narimatsu
Research Center for Medical Glycoscience (RCMG), National
Institute of Advanced Industrial Science and Technology
(AIST), Tsukuba 305-8568, Japan

M. Korenaga · M. Mizokami
The Research Center for Hepatitis and Immunology,
National Center for Global Health and Medicine,
Ichikawa 272-8516, Japan

A. Nishie
Department of Clinical Radiology, Graduate School of Medical
Sciences, Kyushu University, Fukuoka 812-8582, Japan

S. Aishima
Department of Anatomic Pathology, Graduate School of Medical
Sciences, Kyushu University, Fukuoka 812-8582, Japan

Keywords Liver fibrosis · Chronic hepatitis · M2BP ·
Mac-2 binding protein · VTTQ · Fibrosis marker

Abbreviations

ALT	Alanine aminotransferase
APRI	Aspartate transaminase-to-platelet ratio index
ARFI	Acoustic radiation force impulse
AST	Aspartate aminotransferase

COI	Cutoff index
Gd-EOB-DTPA	Gadolinium ethoxybenzyl diethylenetriamine pentaacetic acid
HBsAg	Hepatitis B virus surface antigen
HBV	Hepatitis B virus
HCV	Hepatitis C virus
HCVAb	Hepatitis C virus antibody
LMR	Liver-to-major psoas muscle intensity ratio
m/s	Meters per second
MRI	Magnetic resonance imaging
NPV	Negative predictive value
nonBnonC	Negative for hepatitis B virus surface antigen and hepatitis C virus antibody
PBC	Primary biliary cirrhosis
PPV	Positive predictive value
ROC	Receiver operating characteristic
VTTQ	Virtual Touch™ Tissue Quantification
WFA ⁺ -M2BP	<i>Wisteria floribunda</i> agglutinin-positive Mac-2 binding protein

Introduction

The management of chronic liver disease depends on the degree of liver fibrosis. Therefore, assessment of the degree of liver fibrosis is important for choosing a therapeutic strategy and for determining the prognosis [1, 2]. Liver biopsy is the gold standard method for evaluating the degree of liver fibrosis [3]. However, the invasiveness of liver biopsy, its potential for life-threatening complications, and sampling errors place a heavy burden on those patients with hepatitis who require follow-up [4–6]. Therefore, many reports have demonstrated non-invasive examination methods for assessing the degree of liver fibrosis, which might be alternatives to liver biopsy, such as new serum markers and transient elastography [7–9]. However, none of these studies developed a definitive method.

Recently, a novel marker for liver fibrosis was developed using the glycan “sugar chain”-based immunoassay, and the FastLec-Hepa system was used to determine serum values of “sweet-doughnut” hyperglycosylated *Wisteria floribunda* agglutinin-positive Mac-2 binding protein (WFA⁺-M2BP) for the assessment of liver fibrosis [10, 11]. This unique technical approach supported by multiple lectin-assisted glycan profiling may be applicable to the development pipeline for a wide variety of glycodiagnostic tools. However, the feasibility of WFA⁺-M2BP for assessing liver fibrosis has not been proven with clinical samples of hepatitis.

A method based on acoustic radiation force impulse (ARFI) imaging, with virtual touch tissue quantification (VTTQ), has been developed to evaluate liver fibrosis.

VTTQ measurements can be performed during observation of a particular liver lesion with an ultrasonic probe, and measurements are reproducible compared with transient elastography [12, 13]. For assessment of liver imaging, we have developed magnetic resonance imaging (MRI) using gadolinium ethoxybenzyl diethylenetriamine pentaacetic acid (Gd-EOB-DTPA) enhancement to assess liver function, which may vary inversely for liver fibrosis with the passing grade [14]. Indeed, the increase in rate of liver-to-major psoas muscle intensity ratio (LMR) in the hepatobiliary phase compared with the precontrast image is best correlated with the degree of liver fibrosis, and significantly decreases as liver fibrosis progresses (F0–F4).

This study aimed to compare the diagnostic accuracy of serum WFA⁺-M2BP values using the area under receiver operating characteristic (ROC) curves. The diagnostic performance of serum WFA⁺-M2BP values was compared with validated surrogate fibrosis markers, including VTTQ, the MRI-LMR index, and serum markers, such as hyaluronic acid, type 4 collagen, and the aspartate amino-transferase-to-platelet ratio index (APRI).

Methods

Patients

The study cohort consisted of 200 adults, including 40 healthy volunteers and 160 patients with or without hepatitis, who underwent hepatectomy or living donor liver transplantation, and whose serum WFA⁺-M2BP values and liver stiffness by VTTQ were measured at Kyushu University Hospital. Serum samples of inferior quality, which had the potential of measurement errors, such as those with hemolysis, milky fluid, or protein aggregation, were excluded [15]. Of the 160 patients, 106 were positive for antibody to hepatitis C virus (HCVAb), 21 were positive for hepatitis B virus (HBV) surface antigen (HBsAg), 12 had hepatitis due to alcohol, and 21 were negative for HBsAg and HCVAb. The study protocol conformed to the ethical guidelines of the 1975 Helsinki Declaration and was approved by our institutional review board.

Liver histology and quantification of liver fibrosis

All liver specimens were obtained by surgical resection and were fixed in formalin, embedded in paraffin wax, and stained with hematoxylin and eosin and Masson’s trichrome. The fibrosis staging in all surgical specimens was determined independently by two pathologists who did not know the VTTQ values. In case of discrepancies, histological sections were simultaneously reviewed using a multi-pipe microscope to reach a consensus. Fibrosis was

staged on a scale of 0–4 according to the METAVIR classification [18], with F0 indicating no fibrosis; F1, enlarged, fibrotic portal tracts; F2, periportal or portal–portal septa but intact architecture; F3, fibrosis with architectural distortion but no obvious cirrhosis; and F4, probable or definite cirrhosis.

Direct measurement of serum Mac-2 binding protein

The method of assessment of WFA⁺-M2BP was as follows [10, 11]. A glycan-based immunoassay, FastLec-Hepa, was developed as a simple and accurate system to automatically detect a unique fibrosis-related glyco-alteration in serum hyperglycosylated WFA⁺-M2BP. Briefly, the *Wisteria floribunda* agglutinin (WFA)-antibody immunoassay using the HISCL-2000i bedside clinical chemistry analyzer was developed to measure WFA⁺-M2BP values. These values were successfully adjusted every reaction condition during the automatic assay, and heat pretreatment of the serum was avoided to ensure binding avidity and a fast association rate within just 17 min. The measured values of WFA⁺-M2BP conjugated to WFA were indexed with the obtained values using the following equation:

$$\text{Cutoff index (COI)} = \frac{([\text{WFA}^+ - \text{M2BP}]_{\text{sample}} - [\text{WFA}^+ - \text{M2BP}]_{\text{NC}})}{([\text{WFA}^+ - \text{M2BP}]_{\text{PC}} - [\text{WFA}^+ - \text{M2BP}]_{\text{NC}})}$$

$[\text{WFA}^+ - \text{M2BP}]_{\text{sample}}$, WFA⁺-M2BP count of serum sample (PC, positive control; NC, negative control) [10, 11].

VTTQ and ARFI

The VTTQ system was installed on an ACUSON model S2000 ultrasound system (Siemens Medical Solutions, Inc., Ultrasound Division, Issaquah, WA, USA). The operators were surgeons trained by Siemens Medical Solutions, Inc. The VTTQ system uses an acoustic push pulse to generate shear waves, which pass through the liver parenchyma orthogonally to the acoustic push pulse, through a user-placed region of interest. When detection pulses interact with a passing shear wave, they reveal the wave's location at a specific time, allowing calculation of the shear wave speed. This absolute numerical value is related to the stiffness of the tissue within the region of interest [16, 17], and the results are expressed in meters per second (m/s). For each patient, seven successful measurements were performed several days before surgical operations during which histological specimens were obtained. The measurement of VTTQ in the right lobe of the liver was performed by placing the ultrasonic probe on the right intercostal space at a depth from 2 to 4 cm [9]. The median value of all measurements and the standard deviation of all

right and left VTTQ measurements for each patient were considered for analysis.

Analysis of the MRI-LMR index

The analysis system of the LMR in MRI using Gd-EOB-DTPA enhancement was as follows [14]. Briefly, the signal intensities were measured by placing as large a region of interest as possible on the left and right lobes of the liver parenchyma and major psoas muscle, avoiding vessels, tumors and artifacts, and one slice without significant artifacts for the precontrast image and hepatobiliary phase was selected. The increase in rate of the LMR in the hepatobiliary phase compared with the precontrast image was calculated using the following equation:

$$\frac{(\text{LMR in the hepatobiliary phase} - \text{LMR on the precontrast image})}{\text{LMR on the precontrast image}}$$

[14].

Surrogate serum markers

For all patients, blood samples were obtained on the same day that the VTTQ examination was performed, and they were examined in the same laboratory. The following parameters were determined: hyaluronic acid levels, type 4 collagen, platelet count, aspartate aminotransferase (AST) levels, alanine aminotransferase (ALT) levels, ICG-R15, and the APRI. The APRI was calculated as follows: AST level (per upper limit of normal; 33 U/L) \times 100/platelet count (10^9 /L) [18, 19].

Statistical analysis

Differences between quantitative variables for paired samples were analyzed using a nonparametric test (Wilcoxon rank sum test with Bonferroni's adjustment). The sensitivity, specificity, positive predictive value (PPV), and negative predictive value (NPV) of optimal cutoff values of liver surrogate fibrosis markers for the diagnosis of liver fibrosis were calculated as described [18, 19]. In addition, the diagnostic value of liver stiffness for predicting significant liver fibrosis (F1–F3) and cirrhosis (F4) was assessed by calculating the areas under the ROC curves. The ROC curve is a plot of sensitivity versus 1-specificity for all possible cutoff values. The most commonly used index of accuracy is the area under the ROC curve, where values close to 1.0 indicate high diagnostic accuracy, and 0.5 indicates a test of no diagnostic value. The optimal liver stiffness cutoff values used for the diagnosis of significant fibrosis and cirrhosis were selected based on the sensitivity, specificity, PPV, and NPV [18–20]. Statistical analysis of the differences between

the areas under the ROC curves was based on the theory of generalized U-statistics [21]. All of the differences were considered statistically significant at $P < 0.05$.

Results

Patients and liver specimens

Subject characteristics are summarized in Supplementary Table 1. The mean age of the 200 subjects (135 men and 65 women) was 64 ± 20 years. The 200 enrolled adults consisted of 40 healthy volunteers and 160 patients with or without hepatitis; of the latter, 106 were positive for HCVAb, 21 were positive for HBsAg, 12 had hepatitis due to alcohol, and 21 were negative for HBsAg and HCVAb (nonBnonC). Of the 200 surgical liver specimens, 84 had fibrosis grade F0; 45 were F1, 21 were F2, 16 were F3, and 34 were F4.

Liver fibrosis assessed by WFA⁺-M2BP values

Figure 1 shows box plots of serum WFA⁺-M2BP values for each fibrosis stage. Serum WFA⁺-M2BP values measured by the glycan-based immunoassay ranged from 0.22 to 8.69 (COI). WFA⁺-M2BP levels in patients with fibrosis grades F0 ($n = 84$), F1 ($n = 45$), F2 ($n = 21$), F3 ($n = 16$), and F4 ($n = 34$) had COIs of 1.62, 1.82, 3.02, 3.32, and 3.67, respectively. There were significant differences between fibrosis stages F1 and F2 ($P < 0.01$), and between fibrosis stages F2 and F3 ($P < 0.01$) (Fig. 1).

Correlation of WFA⁺-M2BP values with VTTQ, the LMR index, and serum markers of liver fibrosis

The log of WFA⁺-M2BP correlated with VTTQ results, a significant indicator of liver stiffness as liver fibrosis ($P = 0.0001$) (Fig. 2a). Log WFA⁺-M2BP values were also inversely related with the LMR index, a significant indicator of liver function that varies inversely with liver fibrosis ($P = 0.0001$) (Fig. 2b), as well as with type 4 collagen concentrations ($P = 0.0080$) (Fig. 2c) and the APRI ($P = 0.0001$) (Fig. 2d), but not with hyaluronic acid (Fig. 2e) and ICG-R15 concentrations (Fig. 2f).

APRI, aspartate transaminase-to-platelet ratio index; LMR, liver-to-major psoas muscle intensity ratio; VTTQ, Virtual Touch™ Tissue Quantification; WFA⁺-M2BP, *Wisteria floribunda* agglutinin-positive Mac-2 binding protein.

Diagnostic capability of WFA⁺-M2BP values for each type of liver fibrosis

The areas under the ROC curve for the diagnosis of types of fibrosis F1, F2, F3, and F4 with serum WFA⁺-M2BP values





Fig. 1 Box-and-whisker plot of serum WFA⁺-M2BP values for each fibrosis stage. The *tops* and *bottoms* of the boxes represent the first and third quartiles, respectively, with the *height* of the box represents the interquartile range, covering 50 % of the values. The *line* through the middle of each box represents the median. The *error bars* show the minimum and maximum values (range). Significant correlations were found between the stage of fibrosis and serum WFA⁺-M2BP values, and there were significant correlations between fibrosis stages F1 and F2 ($P < 0.01$) and between fibrosis stages F2 and F3 ($P < 0.01$). Asterisk Statistically significant by the Wilcoxon rank sum test with Bonferroni's adjustment; $P < 0.01$. WFA⁺-M2BP, *Wisteria floribunda* agglutinin-positive Mac-2 binding protein

were 0.686, 0.820, 0.817, and 0.806, respectively (Fig. 3). The optimal cutoff values were 1.00 m/s for $F \geq 1$, 1.86 m/s for $F \geq 2$, 2.21 m/s for $F \geq 3$, and 2.64 m/s for $F \geq 4$ (Table 1). The sensitivity of the serum WFA⁺-M2BP cutoffs for fibrosis grades $F \geq 1$, $F \geq 2$, $F \geq 3$, and $F \geq 4$ was 84.6, 84.7, 88.2, and 82.4 %, whereas the specificity was 47.6, 74.4, 78.7, and 71.9 %, respectively. The sensitivity of VTTQ cutoffs for these fibrosis grades was 81.9, 70.4, 72.6, and 82.4 %, respectively, while the specificity was 70.2, 82.2, 83.3, and 79.5 %, respectively (Table 1, Supplementary Table 2). Thus, WFA⁺-M2BP cutoffs were more sensitive, but less specific, in predicting liver fibrosis in each grade than the VTTQ cutoffs.

Comparison of WFA⁺-M2BP with other indicators for the diagnosis of fibrosis stage ≥ 3

We compared the area under the ROC curves of VTTQ, the LMR index, and serum markers (APRI, hyaluronic acid, and type 4 collagen) with that of WFA⁺-M2BP values. The cutoff values were determined, as described above. The area under the ROC curves for the diagnosis of fibrosis ($F \geq 3$) using serum WFA⁺-M2BP values (0.812) was comparable to that using VTTQ examination (0.814), but was significantly superior to the other surrogate markers, including LMR index (0.766), APRI (0.694), hyaluronic acid (0.683), and type 4 collagen (0.625) ($P = 0.0001$ each; Fig. 4). Serum WFA⁺-M2BP, VTTQ, LMR index, APRI, hyaluronic acid

Fig. 2 Correlation of WFA⁺-M2BP values with VTTQ, LMR index, and the other serum markers. The log of WFA⁺-M2BP correlated with VTTQ results, a significant indicator of liver stiffness as liver fibrosis ($P = 0.0001$) (a). Log WFA⁺-M2BP values were also inversely related with the LMR index, a significant indicator of liver function that varies inversely with liver fibrosis ($P = 0.0001$) (b), as well as with type 4 collagen concentrations ($P = 0.0080$) (c) and the APRI ($P = 0.0001$) (d), but not with hyaluronic acid (e) and ICG-R15 concentrations (f). APRI aspartate transaminase-to-platelet ratio index, LMR liver-to-major psoas muscle intensity ratio, VTTQ Virtual Touch™ Tissue Quantification, WFA⁺-M2BP *Wisteria floribunda* agglutinin-positive Mac-2 binding protein



concentration, and type 4 collagen concentration cutoffs had sensitivities for fibrosis grades $F \geq 3$ of 88.2, 72.6, 52.9, 66.7, 85.1, and 88.1 %, respectively; specificities of 78.7, 83.3, 78.7, 78.9, 40.8, and 40.5 %, respectively; PPVs of 58.9 %, 60.0 %/48.2, 59.6, 34.5, and 33.0 %, respectively; and NPVs of 94.5, 89.9, 83.4, 83.5, 96.2, and 91.1 %, respectively (Table 2).

The areas under the ROC curves for the diagnosis of fibrosis ($F \geq 4$) were comparable for serum WFA⁺-M2BP (0.806) and VTTQ (0.827), but significantly superior to the other surrogate markers, including LMR index (0.776), APRI (0.673), hyaluronic acid (0.657), and type 4 collagen (0.632) ($P = 0.0001$ each; Supplementary Figure 1 and Supplementary Table 3).

Diagnostic capability of WFA⁺-M2BP values for liver fibrosis in each groups

The area under the ROC curves for the diagnosis of fibrosis ($F \geq 3$) using serum WFA⁺-M2BP values was 0.797 in the

107 patients positive for HCVAb and 0.822 in the 21 patients negative for HBsAg and HCVAb, but was insufficient (0.620) in the 21 patients positive for HBsAg. The sensitivity, specificity, PPV, and NPV for the fibrosis grades $\geq F3$ in these three sets of patients using the serum WFA⁺-M2BP cutoff were 96.7 %/100.0 %/61.5 %, 63.6 %/73.3 %/87.5 %, 50.9 %/60.05/88.9 %, and 98.0 %/100.0 %/58.3 %, respectively.

Discussion

This is the first report to quantify liver fibrosis in a large population using serum WFA⁺-M2BP values by the glycan-based immunoassay, FastLec-Hepa, which was developed as a simple and accurate system for automatically detecting unique fibrosis-related glyco-alterations. The accuracy of WFA⁺-M2BP values for diagnosing liver




Fig. 3 Diagnostic ability of serum WFA⁺-M2BP values to assess stages of liver fibrosis. The areas under the ROC curves of serum WFA⁺-M2BP values for diagnosing liver fibrosis were **a** 0.698 for grade $F \geq 1$, **b** 0.837 for grade $F \geq 2$, **c** 0.812 for grade $F \geq 3$, and **d** 0.795 for grade $F \geq 4$. ROC receiver operating characteristic, WFA⁺-M2BP *Wisteria floribunda* agglutinin-positive Mac-2 binding protein

Table 1 WFA⁺-M2BP values for assessment of liver fibrosis

	Optimal cutoff (COI)	Sensitivity (%)	Specificity (%)	PPV (%)	NPV (%)
$F \geq 1$	1.00	84.6	47.6	69.2	69.0
$F \geq 2$	1.86	84.7	74.4	64.9	89.7
$F \geq 3$	2.21	88.2	78.7	58.9	94.5
$F \geq 4$	2.64	82.4	71.9	37.3	95.2

Optimal cutoff points gave the highest total sensitivity and specificity
 APRI Aspartate transaminase-to-platelet ratio index, COI cutoff index, WFA⁺-M2BP *Wisteria floribunda* agglutinin-positive Mac-2 binding protein, *m/s* meters per second NPV negative predictive value, PPV positive predictive value, VTTQ Virtual Touch™ Tissue Quantification

fibrosis grade $F \geq 3$, measured as sensitivity, specificity, PPV, and NPV, was better than that for other surrogate markers, such as the MRI-LMR index and other serum markers of liver fibrosis, including levels of hyaluronic acid, type 4 collagen, and the APRI. WFA⁺-M2BP could be an alternative non-invasive serum marker for liver biopsy for assessing liver fibrosis.

No reports have demonstrated the feasibility of serum WFA⁺-M2BP values as a predictor of liver fibrosis, and the function of WFA⁺-M2BP is unclear. Iacobelli et al. [22] identified WFA⁺-M2BP in 1986 as a tumor-associated antigen and detected it in culture media using a monoclonal antibody from CG-5 breast cancer cell lines. WFA⁺-M2BP

is a highly glycosylated secreted protein, a plant hemagglutinin Mac-2 (galectin-3) ligand, and has 90-kDa subunits, hence, the name 90K. WFA⁺-M2BP mainly mediates cell-to-cell and cell-to-matrix interactions, and is involved in cell proliferation and angiogenesis [23–25] by inducing the expression of cytokines, such as interleukin (IL)-1, IL-2, and IL-6 [23–25]. Indeed, endogenous WFA⁺-M2BP ligands laminin, fibronectin, and lysosome-associated membrane protein [26, 27], and enhances cell adhesion and the extracellular matrix to promote fibrosis. Therefore, expression of WFA⁺-M2BP levels may be proportional to the degree of liver fibrosis in patients with chronic liver diseases.

In our study, serum WFA⁺-M2BP values had better diagnostic ability for assessment of liver fibrosis than the other serum markers, such as the APRI, hyaluronic acid, and type 4 collagen, as evaluated by the area under the ROC curves. For diagnosing fibrosis stage ≥ 3 , the specificity and PPV using serum WFA⁺-M2BP were 78.7 and 58.9 %, respectively compared with 78.7 and 48.2 %, respectively, using APRI; 40.8 and 34.5 %, respectively, using hyaluronic acid, and 40.5 and 33.0 %, respectively, using type 4 collagen cutoffs. These results suggest that examination of serum WFA⁺-M2BP values is the most accurate diagnostic tool for liver fibrosis among the serum markers investigated in this study.

The diagnostic significance of surrogate markers in liver fibrosis has been evaluated distinguishing F3 from F2 fibrosis. The clinical significance of distinguishing F3 from F2 fibrosis has been widely accepted in the follow-up of patients with viral hepatitis, and also has been linked with hepatocarcinogenesis [28, 29]. The annual carcinogenesis rate was reported to be correlated with the stage of liver fibrosis in the study of 2,890 patients with hepatitis [28]. The annual incidence of hepatocellular carcinoma in patients with severe liver fibrosis of grade F3 is high (5.3 %), whereas the incidence in those with moderate liver fibrosis of grade F2 is low (1.9 %) [29]. Therefore, fibrotic change is closely correlated with hepatocarcinogenesis in patients with viral hepatitis, and it is critically important to distinguish between liver fibrosis of grades F3 and F2, which was the cause of the favorable point for the serum WFA⁺-M2BP examination superior to the other markers in our study. In addition, we found that the measurement time for our glycan-based immunoassay was only 17 min, which has practical implications. However, a limitation is that its measurement was just applied by the glycan “sugar chain”-based immunoassay, the FastLec-Hepa system, which is currently only available in a few selected institutions. Commercial application of this system is a primary goal for evaluation of a clinically beneficial therapy through quantification of liver fibrosis in hepatitis patients. This study, however, had one important limitation, in that




Fig. 4 Comparison of WFA⁺-M2BP with other indicators for the diagnosis of fibrosis stage $F \geq 3$ by areas under the receiver operating curves (ROC). The areas under the ROC curves for the diagnosis of grade $F \geq 3$ fibrosis were a 0.812 for serum WFA⁺-M2BP, b 0.814 for VTTQ, c 0.766 for LMR index, d 0.694 for APRI, e 0.683 for

hyaluronic acid, and f 0.625 for type 4 collagen. APRI aspartate transaminase-to-platelet ratio index, LMR liver-to-major psoas muscle intensity ratio, ROC receiver operating characteristic, VTTQ Virtual Touch™ Tissue Quantification, WFA⁺-M2BP *Wisteria floribunda* agglutinin-positive Mac-2 binding protein

Table 2 Diagnostic performance of indicators predicting liver fibrosis ($F \geq 3$)

	Optimal cutoff	Sensitivity (%)	Specificity (%)	PPV (%)	NPV (%)
WFA ⁺ -M2BP	2.21 (COI)	88.2	78.7	58.9	94.5
VTTQ	1.77 (m/s)	72.6	83.3	60.0	89.9
LMR index	1.15	66.7	78.9	59.6	83.5
APRI	3.33	52.9	78.7	48.2	83.4
Hyaluronic acid	58.0 (ng/mL)	85.1	40.8	34.5	96.2
Type 4 collagen	125.0 (ng/mL)	88.1	40.5	33.0	91.1

Optimal cutoff points gave the highest total sensitivity and specificity

APRI Aspartate transaminase-to-platelet ratio index, COI cutoff index, WFA⁺-M2BP *Wisteria floribunda* agglutinin-positive Mac-2 binding protein, m/s meters per second, NPV negative predictive value, PPV positive predictive value, VTTQ Virtual Touch™ Tissue Quantification

the number of patients with a fibrosis score of F3 was relatively low. Most patients who underwent hepatectomy had Child-Pugh grade A liver function and fibrosis scores of F0, F1, and F2, whereas most patients who underwent living donor liver transplantation had liver cirrhosis (F4). Accumulation of addition liver specimens by surgical resection may be necessary.

We have previously demonstrated that VTTQ values are correlated with liver fibrosis [9, 12, 13]. Compared with VTTQ, serum WFA⁺-M2BP values were almost equal for predicting liver fibrosis $\geq F3$, with similar areas under the ROC curves (0.812 vs 0.814) and similar sensitivity (88.2 vs 72.6 %), and specificity (78.7 vs 83.3 %). However, this VTTQ method has some limitations compared with serum WFA⁺-M2BP values. First, the diagnostic accuracy of

VTTQ values in the right and left lobes of the liver was significantly different; VTTQ values in the right lobe were more accurate for diagnosing liver fibrosis than those in the left lobe, as evaluated by the area under the ROC curves and the standard deviations of each VTTQ value [9]. The other limitation is the lower diagnostic accuracy of VTTQ values in fatty liver compared with other types of hepatitis for predicting liver fibrosis [30]. Therefore, the diagnostic accuracy for assessment of liver fibrosis needs to be compared between serum WFA⁺-M2BP and VTTQ values in these marginal clinical cases in the future. In addition, the VTTQ system installed on the ultrasound system by Siemens Medical Solutions is expensive [15–17]. Therefore, it may have the clinical significance of the WFA⁺-M2BP assessment for the medical economy to assess the