

- Chilamakuri CS, Lorenz S, Madoui MA, *et al.* (2014) Performance comparison of four exome capture systems for deep sequencing. *BMC Genomics* 15:449.
- Chin E, Zhang V, Wang J, *et al.* (2013) Frequently asked questions about the clinical utility of next-generation sequencing in molecular diagnosis of human genetic diseases. In: Wong L-JC (ed) *Next Generation Sequencing: Translation to Clinical Diagnostics*. Springer Science+Business Media, New York, pp 287–299.
- Chun S, Fay JC (2009) Identification of deleterious mutations within three human genomes. *Genome Res* 19:1553–1561.
- Cingolani P, Platts A, Wang le L, *et al.* (2012) A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w<sup>1118</sup>; iso-2; iso-3. *Fly* 6:80–92.
- Cooper DN, Ball EV, Krawczak M (1998) The human gene mutation database. *Nucleic Acids Res* 26:285–287.
- De Luca A, Bottillo I, Dasdia MC, *et al.* (2007) Deletions of *NF1* gene and exons detected by multiplex ligation-dependent probe amplification. *J Med Genet* 44:800–808.
- Fahsold R, Hoffmeyer S, Mischung C, *et al.* (2000) Minor lesion mutational spectrum of the entire *NF1* gene does not explain its high mutability but points to a functional domain upstream of the GAP-related domain. *Am J Hum Genet* 66:790–818.
- Frommolt P, Abdallah AT, Altmüller J, *et al.* (2012) Assessing the enrichment performance in targeted resequencing experiments. *Hum Mutat* 33:635–641.
- Gasparini P, D'Agruma L, Pio de Cillis G, *et al.* (1996) Scanning the first part of the neurofibromatosis type 1 gene by RNA-SSCP: identification of three novel mutations and of two new polymorphisms. *Hum Genet* 97:492–495.
- Han SS, Cooper DN, Upadhyaya MN (2001) Evaluation of denaturing high performance liquid chromatography (DHPLC) for the mutational analysis of the neurofibromatosis type 1 (*NF1*) gene. *Hum Genet* 109:487–497.
- Hattori M, Torii C, Yagihashi T, *et al.* (2009) Diagnosis of Russell-Silver syndrome by the combined bisulfite restriction analysis—denaturing high-performance liquid chromatography assay. *Genet Test Mol Biomarkers* 13:623–630.
- Japanese Genetic Variation Consortium (2013) Human genetic variation browser. Available at [www.genome.med.kyoto-u.ac.jp/SnpDB](http://www.genome.med.kyoto-u.ac.jp/SnpDB) (accessed March 18 2014).
- Jones K (2005) *Smith's Recognizable Patterns of Human Malformation*. Saunders, Philadelphia.
- Kosaki K, Udaka T, Okuyama T (2005) DHPLC in clinical molecular diagnostic services. *Mol Genet Metab* 86:117–123.
- Kumar P, Henikoff S, Ng PC (2009) Predicting the effects of coding non-synonymous variants on protein function using the SIFT algorithm. *Nat Protoc* 4:1073–1081.
- Lee MJ, Su YN, You HL, *et al.* (2006) Identification of forty-five novel and twenty-three known *NF1* mutations in Chinese patients with neurofibromatosis type 1. *Hum Mutat* 27:832.
- Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler Transform. *Bioinformatics* 25:1754–1760.
- Li Y, Bollag G, Clark R, *et al.* (1992) Somatic mutations in the neurofibromatosis 1 gene in human tumors. *Cell* 69:275–281.
- McKenna A, Hanna M, Banks E, *et al.* (2010) The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res* 20:1297–1303.
- Messiaen LM, Callens T, Mortier G, *et al.* (2000) Exhaustive mutation analysis of the *NF1* gene allows identification of 95% of mutations and reveals a high frequency of unusual splicing defects. *Hum Mutat* 15:541–555.
- Nemethova M, Bolcekova A, Ilencikova D, *et al.* (2013) Thirty-nine novel neurofibromatosis 1 (*NF1*) gene mutations identified in Slovak patients. *Ann Hum Genet* 77:364–379.
- Neurofibromatosis Conference Statement (1988) National Institutes of Health Consensus Development Conference. *Arch Neurol* 45:575–578.
- Purandare SM, Huntsman Breidenbach H, *et al.* (1995) Identification of neurofibromatosis 1 (*NF1*) homologous loci by direct sequencing, fluorescence *in situ* hybridization, and PCR amplification of somatic cell hybrids. *Genomics* 30:476–485.
- Purandare SM, Lanyon WG, Connor JM (1994) Characterisation of inherited and sporadic mutations in neurofibromatosis type-1. *Hum Mol Genet* 3:1109–1115.
- Richards CS, Bale S, Bellissimo DB, *et al.* (2008) ACMG recommendations for standards for interpretation and reporting of sequence variations: revisions 2007. *Genet Med* 10:294–300.
- Rozen S, Skaletsky H (2000) Primer3 on the www for general users and for biologist programmers. *Methods Mol Biol* 132:365–386.
- Sabbagh A, Pasmant E, Imbard A, *et al.* (2013) *NF1* molecular characterization and neurofibromatosis type I genotype-phenotype correlation: the French experience. *Hum Mutat* 34:1510–1518.
- Samejima H, Torii C, Kosaki R, *et al.* (2007) Screening for Alagille syndrome mutations in the *JAG1* and *NOTCH2* genes using denaturing high-performance liquid chromatography. *Genet Test Mol Biomarkers* 11:216–227.
- Schwarz JM, Rödelberger C, Schuelke M, *et al.* (2010) MutationTaster evaluates disease-causing potential of sequence alterations. *Nat Methods* 7:575–576.
- Siepel A, Pollard KS, Haussler D (2009) New methods for detecting lineage-specific selection. Proceedings of the 10th International Conference on Research in Computational. *Mol Biol* 3909:190–205.
- Takenouchi T, Hida M, Sakamoto Y, *et al.* (2013a) Severe congenital lipodystrophy and a progeroid appearance: mutation in the penultimate exon of *FBN1* causing a recognizable phenotype. *Am J Med Genet A* 161A:3057–3062.
- Takenouchi T, Matsuzaki Y, Torii C, *et al.* (2014) *SOX9* dimerization domain mutation mimicking type 2 collagen disorder phenotype. *Eur J Med Genet* 57:298–301.
- Takenouchi T, Shimizu A, Torii C, *et al.* (2013b) Multiple Café au Lait spots in familial patients with *MAP2K2* mutation. *Am J Med Genet A* 164A:392–396.
- Teer JK, Green ED, Mullikin JC, *et al.* (2012) VarSifter: visualizing and analyzing exome-scale sequence variation data on a desktop computer. *Bioinformatics* 28:599–600.
- Thorvaldsdóttir H, Robinson JT, Mesirov JP (2013) Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. *Brief Bioinform* 14:178–192.
- Udaka T, Imoto I, Aizu Y, *et al.* (2007) Multiplex PCR/liquid chromatography assay for screening of subtelomeric rearrangements. *Genet Test Mol Biomarkers* 11:241–248.
- Udaka T, Kurosawa K, Izumi K, *et al.* (2006) Screening for partial deletions in the *CREBBP* gene in Rubinstein-Taybi syndrome patients using multiplex PCR/liquid chromatography. *Genet Test Mol Biomarkers* 10:265–271.
- Udaka T, Torii C, Takahashi D, *et al.* (2005) Comprehensive screening of the thiopurine methyltransferase polymorphisms by denaturing high-performance liquid chromatography. *Genet Test Mol Biomarkers* 9:85–92.

- Upadhyaya M (2008) *NF1* gene structure and *NF1* genotype/phenotype correlations. In: Kaufmann D (ed) Neurofibromatosis. Karger, Basel, pp 46–62.
- Valero MC, Martín Y, Hernández-Imaz E, *et al.* (2011) A highly sensitive genetic protocol to detect NF1 mutations. *J Mol Diagn* 13:113–122.
- Yan D, Tekin M, Blanton SH, *et al.* (2013) Next-generation sequencing in genetic hearing loss. *Genet Test Mol Biomarkers* 17:581–587.

Address correspondence to:  
*Kenjiro Kosaki, MD, FACMG*  
*Center for Medical Genetics*  
*Keio University School of Medicine*  
*35 Shinanomachi, Shinjuku-ku*  
*Tokyo 160-8582*  
*Japan*  
*E-mail: kkosaki@z3.keio.jp*

