

研究成果の刊行に関する一覧表

雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Sakai T, Kusakabe T, Ebihara K, Aotani D, Yamamoto-Kataoka S, Zhao M, Gumbilai VM, Ebihara C, Aizawa-Abem M, Yamamoto Y, Noguchi M, Fujikura J, Hosoda K, Inagaki N, Nakao K	Leptin restores the insulinotropic effect of exenatide in a mouse model of type 2 diabetes with increased adiposity induced by streptozotocin and high-fat diet.	Am J Physiol Endocrinol Metab	307	E712-719	2014
Tanaka M, Yamaguchi S, Yamazaki Y, Kinoshita H, Kuwahara K, Nakao K, Jay PY, Noda T, Nakamura T.	Somatic chromosomal translocation between <i>Ewsr1</i> and <i>Fli1</i> loci leads to dilated cardiomyopathy in a mouse model	Sci Rep	5	7826	2015
Oshita K, Itoh M, Hirashima S, Kuwabara Y, Ishihara K, Kuwahara K, Nakao K, Kimura T, Nakamura K, Ushijima K, Takano M.	Ectopic automaticity induced in ventricular myocytes by transgenic overexpression of HCN2	J Mol Cell Cardiol	80	81-89	2015
Yamada Y, Kinoshita H, Kuwahara K, Nakagawa Y, Kuwabara Y, Minami T, Yamada C, Shibata J, Nakao K, Cho K, Arai Y, Yasuno S, Nishikimi T, Ueshima K, Kamakura S, Nishida M, Kiyonaka S, Mori Y, Kimura T, Kangawa K, Nakao K	Inhibition of N-type Ca ²⁺ channels ameliorates an imbalance in cardiac autonomic nerve activity and prevents lethal arrhythmias in mice with heart failure	Cardiovascular Research	104	1. 183-93	2014
Nakagawa Y, Nishikimi T, Kuwahara K, Yasuno S, Kinoshita H, Kuwabara Y, Nakao K, Minami T, Yamada C, Ueshima K, Ikeda Y, Okamoto H, Horii K, Nagata K, Kanagawa K, Minamino N, Nakao K.	The Effects of Super-Flux (High Performance) Dialyzer on Plasma Glycosylated Pro-B-Type Natriuretic Peptide (proBNP) and Glycosylated N-Terminal proBNP in End-Stage Renal Disease Patients on Dialysis	PLoS One	9	e92314	2014
Kuwabara T, Mori K, Mukoyama M, Kasahara M, Yokoi H, Nakao K	Macrophage-mediated glucolipototoxicity via myeloid-related protein 8/toll-like receptor 4 signaling in diabetic nephropathy	Clin Exp Nephrol	18	584-592	2014

<p>Kuwabara T, Mori K, Kasahara M, Yokoi H, Imamaki H, Ishii A, Koga K, Sugawara A, Yasuno S, Ueshima K, Morikawa T, Konishi Y, Imanishi M, Nishiyama A, Nakao K, Mukoyama M.</p>	<p>Predictive significance of kidney myeloid-related protein 8 expression in patients with obesity- or type 2 diabetes-associated kidney diseases.</p>	<p>PLoS One</p>	<p>9</p>	<p>e88942</p>	<p>2014</p>
<p>Kanda J, Mori K, Kawabata H, Kuwabara T, Mori KP, Imamaki H, Kasahara M, Yokoi H, Mizumoto C, Thoennissen NH, Koeffler HP, Barasch J, Takaori-Kondo A, Mukoyama M, Nakao K.</p>	<p>An AKI biomarker lipocalin 2 in the blood derives from the kidney in renal injury but from neutrophils in normal and infected conditions.</p>	<p>Clin Exp Nephrol</p>	<p>19</p>	<p>99-106</p>	<p>2015</p>
<p>Tomita T, Hosoda K, Fujikura J, Inagaki N, <u>Nakao K.</u></p>	<p>The G-Protein-Coupled Long-Chain Fatty Acid Receptor GPR40 and Glucose Metabolism.</p>	<p>Front Endocrinol</p>	<p>5</p>	<p>152</p>	<p>2014</p>
<p>Ebihara C, Ebihara K, Aizawa-Abe M, Mashimo T, Tomita T, Zhao M, Gumbilai V, Kusakabe T, Yamamoto Y, Aotani D, Yamamoto-Kataoka S, Sakai T, Hosoda K, Serikawa T, Nakao K.</p>	<p>Seipin is necessary for normal brain development and spermatogenesis in addition to adipogenesis.</p>	<p>Hum Mol Genet</p>		<p>In press</p>	<p>2015</p>