

- [36] S. Zhang, W.C. Huang, P. Li, H. Guo, S.B. Poh, S.W. Brady, Y. Xiong, L.M. Tseng, S.H. Li, Z. Ding, A.A. Sahin, F.J. Esteva, G.N. Hortobagyi, D. Yu, Combating trastuzumab resistance by targeting SRC, a common node downstream of multiple resistance pathways, *Nat. Med.* 17 (2011) 461–469.
- [37] J.O. Moore, M.A. Lemmon, K.M. Ferguson, Tie2 receptor dimerization mediated by its extracellular FNIII domains, *Biophysical Journal*, 2013, p. 608a.
- [38] M.A. Schumacher, N. Chinnam, T. Ohashi, R.S. Shah, H. Erickson, Structure of irisin reveals a novel intersubunit beta-sheet fibronectin (FNIII) dimer: implications for receptor activation, *J. Biol. Chem.* 288 (2013) 33738–33744.
- [39] L. Zabeau, D. Defeau, H. Iserentant, J. Vandekerckhove, F. Peelman, J. Tavernier, Leptin receptor activation depends on critical cysteine residues in its fibronectin type III subdomains, *J. Biol. Chem.* 280 (2005) 22632–22640.
- [40] G.M. Keating, Pertuzumab: in the first-line treatment of HER2-positive metastatic breast cancer, *Drugs* 72 (2012) 353–360.
- [41] S.M. Swain, S.B. Kim, J. Cortes, J. Ro, V. Semiglazov, M. Campone, E. Ciruelos, J.M. Ferrero, A. Schneeweiss, A. Knott, E. Clark, G. Ross, M.C. Benyunes, J. Baselga, Pertuzumab, trastuzumab, and docetaxel for HER2-positive metastatic breast cancer (CLEOPATRA study): overall survival results from a randomised, double-blind, placebo-controlled, phase 3 study, *Lancet Oncol.* 14 (2013) 461–471.
- [42] A. Freywald, N. Sharfe, C.M. Roifman, The kinase-null EphB6 receptor undergoes transphosphorylation in a complex with EphB1, *J. Biol. Chem.* 277 (2002) 3823–3828.
- [43] K. Nagano, S. Kanasaki, T. Yamashita, Y. Maeda, M. Inoue, K. Higashisaka, Y. Yoshioka, Y. Abe, Y. Mukai, H. Kamada, Y. Tsutsumi, S. Tsunoda, Expression of Eph receptor A10 is correlated with lymph node metastasis and stage progression in breast cancer patients, *Cancer Med.* 2 (2013) 972–977.

Regular Article

THROMBOSIS AND HEMOSTASIS

Anti-factor IXa/X bispecific antibody ACE910 prevents joint bleeds in a long-term primate model of acquired hemophilia A

Atsushi Muto,¹ Kazutaka Yoshihashi,¹ Minako Takeda,¹ Takehisa Kitazawa,¹ Tetsuhiro Soeda,¹ Tomoyuki Igawa,¹ Zenjiro Sampei,¹ Taichi Kuramochi,¹ Akihisa Sakamoto,¹ Kenta Haraya,¹ Kenji Adachi,¹ Yoshiki Kawabe,¹ Keiji Nogami,² Midori Shima,² and Kunihiro Hattori¹

¹Research Division, Chugai Pharmaceutical Co, Ltd, Gotemba, Shizuoka, Japan; and ²Department of Pediatrics, Nara Medical University, Kashihara, Nara, Japan

Key Points

- A long-term acquired hemophilia A model expressing spontaneous joint bleeds and other bleeds was newly established in nonhuman primates.
- Weekly SC dose of the anti-FIXa/X bispecific antibody ACE910 prevented joint bleeds and other bleeds in the primate hemophilia A model.

ACE910 is a humanized anti-factor IXa/X bispecific antibody mimicking the function of factor VIII (FVIII). We previously demonstrated in nonhuman primates that a single IV dose of ACE910 exerted hemostatic activity against hemophilic bleeds artificially induced in muscles and subcutis, and that a subcutaneous (SC) dose of ACE910 showed a 3-week half-life and nearly 100% bioavailability, offering support for effective prophylaxis for hemophilia A by user-friendly SC dosing. However, there was no direct evidence that such SC dosing of ACE910 would prevent spontaneous bleeds occurring in daily life. In this study, we newly established a long-term primate model of acquired hemophilia A by multiple IV injections of an anti-primate FVIII neutralizing antibody engineered in mouse-monkey chimeric form to reduce its antigenicity. The monkeys in the control group exhibited various spontaneous bleeding symptoms as well as continuous prolongation of activated partial thromboplastin time; notably, all exhibited joint bleeds, which are a hallmark of hemophilia. Weekly SC doses of ACE910 (initial 3.97 mg/kg followed by 1 mg/kg) significantly prevented these bleeding symptoms; notably, no joint bleeding symptoms were observed. ACE910 is expected to prevent spontaneous bleeds and joint damage in

hemophilia A patients even with weekly SC dosing, although appropriate clinical investigation is required. (*Blood*. 2014;124(20):3165-3171)

Introduction

Patients with severe hemophilia A (<1% of normal factor VIII [FVIII] level) typically suffer from recurrent bleeding episodes, primarily in the musculoskeletal system.^{1,2} Approximately 85% of the bleeding episodes are into joints,³ and repeated joint bleeding from early childhood results in a chronic degenerative arthritis. Although traditional on-demand treatment by a FVIII agent cannot prevent hemophilic arthropathy, routine prophylaxis with FVIII to maintain $\geq 1\%$ FVIII:C is beneficial in preventing it.^{4,5} However, the need for frequent IV injections of FVIII negatively affects patients' quality of life and their adherence to the routine prophylactic regimen, which is particularly problematic when treating pediatric patients at home.^{2,6}

Furthermore, ~30% of severe patients develop alloantibodies against FVIII (FVIII inhibitors),^{2,7} which largely restrict treatment with FVIII. FVIII inhibitors make hemorrhage more difficult to be controlled because alternative bypassing agents have shorter half-lives and are not always effective.^{7,8} Attempts to induce immune tolerance to FVIII inhibitors with high doses of FVIII are very expensive and do not always work.⁹

Therefore, a novel drug is needed: one that is long-lasting, subcutaneously injectable, effective regardless of FVIII inhibitors, and does not induce FVIII inhibitors.¹⁰⁻¹³ To achieve this desirable profile, we produced a series of humanized immunoglobulin G (IgG) antibodies bispecific to factors IXa and X (anti-FIXa/X antibodies) that mimic the FVIII cofactor function by binding and placing FIXa and FX into spatially appropriate positions (supplemental Figure 1, see supplemental Data available on the *Blood* Web site),¹⁴ and identified a clinical investigational drug termed ACE910.¹⁵ In a short-term primate model of acquired hemophilia A, ACE910 at a single IV dose of 1 or 3 mg/kg exerted hemostatic activity against artificial ongoing bleeds in muscles and subcutis to the same extent as recombinant porcine FVIII (rpoFVIII) at twice-daily IV doses of 10 U/kg.¹⁶ Furthermore, a multiple-dosing simulation calculated from the pharmacokinetic (PK) parameters of ACE910 in cynomolgus monkeys suggested that the plasma ACE910 concentration capable of stopping even ongoing bleeds would be maintained by once-weekly subcutaneous (SC) administration of 0.64 to 1.5 mg/kg ACE910.¹⁶

Submitted July 2, 2014; accepted September 14, 2014. Prepublished online as *Blood* First Edition paper, October 1, 2014; DOI 10.1182/blood-2014-07-585737.

Presented in abstract form at the World Federation of Hemophilia 2014 World Congress, Melbourne, Australia, May 11-15, 2014.

The online version of this article contains a data supplement.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2014 by The American Society of Hematology

Prevention of joint bleeding is of major importance in the care of hemophilia A patients.³ However, it remained unproven whether repeated SC dosing of ACE910 could actually prevent spontaneous bleeding episodes, including the joint bleeds that are a pathologic hallmark of hemophilia A. To address this question nonclinically, we required a primate model because ACE910 is highly species-specific in its FVIII-mimetic cofactor activity.¹⁶

In this study, we aimed first to establish a long-term acquired hemophilia A model expressing spontaneous bleeding episodes, including joint bleeds, in nonhuman primates, and second to evaluate the preventive effect of once-weekly SC dosing of ACE910 in this model for investigating the potential of a prophylactic treatment in hemophilia A patients.

Materials and methods

ACE910

ACE910 was expressed in HEK293 or CHO cells cotransfected with a mixture of plasmids encoding the anti-FIXa heavy chain, anti-FX heavy chain, and common light chain.¹⁵ ACE910 was purified by protein A and ion-exchange chromatography from the culture supernatants.

Anti-primate FVIII neutralizing antibodies

A mouse monoclonal anti-primate FVIII neutralizing antibody, termed VIII-2236, was prepared from hybridoma culture supernatants.^{14,16} A chimeric mouse-monkey anti-primate FVIII neutralizing antibody, termed cyVIII-2236, was constructed comprising the mouse variable region from VIII-2236 and the monkey constant region of IgG, which we originally cloned from cynomolgus monkey thymus. The cyVIII-2236 antibody was produced in HEK293 cells and isolated by protein A and gel permeation chromatography from the culture supernatants.

Comparison of cyVIII-2236 with VIII-2236 in an APTT assay

First, to compare the FVIII-neutralizing activity between cyVIII-2236 and VIII-2236, each was added to citrated plasma pooled from 3 normal male cynomolgus monkeys. Then, activated partial thromboplastin time (APTT) was measured with a standard method using Thrombocheck APTT-SLA (Sysmex) and coagulation analyzer KC4 Delta (Stago). Second, to compare the effect of cyVIII-2236 and VIII-2236 on the APTT-shortening activity of ACE910, each was added to FVIII-deficient human plasma (George King) at the final plasma concentration of 300 $\mu\text{g}/\text{mL}$ together with various concentrations of ACE910, then APTT was measured.

Animals

Ten male cynomolgus monkeys (2.35-4.25 kg, aged 3-4 years) were purchased from Hamri. All animal studies were approved by the institutional animal care and use committee of Chugai Pharmaceutical, and were conducted in accordance with the approved protocols and the Guidelines for the Care and Use of Laboratory Animals at the company. Chugai Pharmaceutical is fully accredited by the Association for Assessment and Accreditation of Laboratory Animal Care (AAALAC) International.

Long-term primate model of acquired hemophilia A

In the first part of this study, 2 cynomolgus monkeys received weekly IV injections (3 or 10 mg/kg each) of VIII-2236. Citrated blood was collected over time. Blood hemoglobin (Hgb) concentration was measured by hematology analyzer SF-3000 (Sysmex) to monitor hemorrhagic anemia. APTT was assessed with a standard method using Data-Fi APTT (Siemens) and coagulation analyzer AMAX CS-190 (Trinity). In addition, bleeding symptoms (bruises [dark-red areas on the body surface], hematuria, limping, and general condition including joint swelling) were monitored on 21

working days from day 0 until day 28. To observe bruises easily, the body hair was sheared on day 0, and the area of bruises and the number of days when bruises had been detected were assessed.

In the second part of this study, another 8 cynomolgus monkeys received weekly IV injections (10 mg/kg) of cyVIII-2236. Two groups ($n = 4$ each) were established: a control group (the vehicle group) treated with vehicle and a test group (the ACE910 group) treated with ACE910. ACE910 was administered as an initial 3.97 mg/kg SC dose 2 hours after cyVIII-2236 injection on day 0, and thereafter as weekly 1 mg/kg SC doses. The same dosing regimen of vehicle (histidine buffer containing a surfactant) was applied to the vehicle group. Citrated blood was collected over time. Blood Hgb concentration was measured; the change in blood Hgb level was expressed as a percentage of that on day 0 (2 hours after cyVIII-2236 injection). APTT was measured with Thrombocheck APTT-SLA and KC4 Delta. Bleeding symptoms were monitored on 41 working days until day 56. Necropsy was performed on day 56; organs and tissues from the whole body were macroscopically examined, and hemorrhagic findings in joints (shoulder, elbow, wrist, hip, knee, and ankle on both sides) were scored. Then, they were histopathologically examined.

We assessed cyVIII-2236 concentration, FVIII-neutralizing titer of cyVIII-2236, ACE910 concentration, and development of anti-ACE910 alloantibodies. The methods are described in detail in the supplemental Methods. Briefly, cyVIII-2236 concentration was determined with a sandwich enzyme-linked immunosorbent assay (ELISA) using recombinant human FVIII and an anti-human IgG antibody. FVIII-neutralizing titers of cyVIII-2236 on days 0 (just after cyVIII-2236 injection) and 56 were assessed with a modified Bethesda assay. ACE910 concentration was determined with a sandwich ELISA to quantify human IgG.¹⁶ Anti-ACE910 alloantibodies present at necropsy were examined with an electrochemiluminescent bridging immunoassay using labeled ACE910.¹⁶

In both parts, the monkeys were carefully monitored under the supervision of the attending veterinarian to determine the necessity of pain relief or other treatment. Consequently, no monkeys received an analgesic drug.

Statistical analysis and multiple-dosing simulation

The data are presented as individual values or as mean \pm standard deviation (SD). In the in vivo efficacy study, some were analyzed by 2-tailed Student *t* test (SAS Preclinical Package, version 5.00) between the vehicle and the ACE910 groups. A *P* value of .05 or less was considered statistically significant. Multiple doses of ACE910 were simulated on the basis of the PK parameters previously determined¹⁶ with SAAM II, version 1.2 software (SAAM Institute).

Results

Failure of long-term neutralization of endogenous FVIII by VIII-2236 in cynomolgus monkeys

First, we attempted to establish a long-term acquired hemophilia A model by weekly IV injection of VIII-2236 in cynomolgus monkeys. APTT prolongation was observed for the first 2 weeks (Figure 1), but gradually shortened after the third injection, and finally disappeared by the end of the fourth week, suggesting the development of alloantibodies to VIII-2236. VIII-2236 is a mouse IgG, and therefore entirely foreign to primates. In addition, no sign of joint bleeds was observed, although some bruises and a transient decrease in blood Hgb were detected (data not shown). Consequently, we sought an alternative way of establishing a longer-term primate model of acquired hemophilia A.

Chimerization of VIII-2236 to cyVIII-2236

As an attempt to avoid anti-VIII-2236 alloantibody development, we chimerized VIII-2236 (mouse IgG) by replacing its constant region with that of cynomolgus monkey IgG. The chimerized VIII-2236




Figure 1. Change in APTT after weekly IV injection of the mouse anti-primate FVIII neutralizing antibody VIII-2236 in cynomolgus monkeys. VIII-2236 was injected at 3 or 10 mg/kg IV doses to cynomolgus monkeys on days 0, 7, 14, and 21. Citrated blood was collected on days 0, 3, 7, 10, 14, 17, 21, 24, and 28 (before and 2 hours after VIII-2236 injection on days 0, 7, 14, and 21). The time course of APTT is shown for each monkey.

(cyVIII-2236) successfully prolonged APTT in cynomolgus monkey plasma in vitro with a similar concentration dependency to that of original VIII-2236 (Figure 2A). Additionally, cyVIII-2236 did not interfere with the APTT-shortening activity of ACE910, the same results were also found with VIII-2236 (Figure 2B). These results indicate that VIII-2236 was successfully chimerized to cyVIII-2236 while retaining the original characteristics.

Establishment of a long-term acquired hemophilia A model expressing spontaneous joint bleeds in cynomolgus monkeys

Using cyVIII-2236, we retried to establish long-term neutralization of endogenous FVIII, expecting reproducible development of spontaneous joint bleeds. The experimental protocol is illustrated in Figure 3A. In all the monkeys given vehicle (n = 4), APTT was prolonged to approximately twice the normal baseline for 8 weeks (Figure 3B). Plasma cyVIII-2236 concentration fluctuated, but remained over 58.6 μg/mL (nearly 400 nM) (supplemental Figure 2). As over 400 nM cyVIII-2236 prolonged APTT by twofold in vitro (Figure 2A), these observations seemed consistent. Thus, an acquired hemophilia A state was successfully maintained by using cyVIII-2236.

In this long-lasting acquired hemophilia A state, all the control monkeys given vehicle developed abnormal leg motion known as limping: a behavior to avoid using the affected leg (Figure 3C). The limping continued until the experiments ended, with days when limping was detected reaching 25.0 ± 8.3 days (61% ± 20%) of the 41 observation days. Two control monkeys (#3 and #4) developed joint swelling at the ankle in the limping limb from days 28 to 42 and from days 18 to 42, respectively. Macroscopic observation at necropsy on day 56 revealed intraarticular dark red areas (consistent with hemorrhagic findings in the histopathological examination) in joints (elbow, hip, or ankle) of all the control monkeys; the number of bleeding joints was 2.0 ± 0.8 per head (Figure 3D, Table 1). Bleeding was detected in the hip or ankle joint of the side that limped. Representative macroscopic findings of the hip joints are shown in Figures 3Ei-ii. Hemorrhagic findings or reactive findings associated with hemorrhage were histopathologically confirmed in the joints that limped: hemorrhage/hemosiderin deposition, mononuclear cell infiltration, and vascular proliferation in the synovial membranes;

synovial hyperplasia; granulation tissue; and destruction of articular cartilage or underlying bone (Figure 4A-C). Thus, a long-term acquired hemophilia A model expressing spontaneous joint bleeds was successfully established in cynomolgus monkeys.

Other spontaneous bleeding symptoms, that is, bruises and hematuria, were detected by daily observation in the vehicle group (data not shown). By macroscopic observation at necropsy, hemorrhagic findings were also observed in skeletal muscle (lower abdomen and femoral region), urinary bladder, seminal vesicle, rectum, and subcutis (back) in 1 or 2 of the 4 control monkeys (Table 1). Additionally, blood Hgb temporarily decreased during the observation period with the lowest values reaching 64% to 87% (Figure 5), suggesting substantial hemorrhagic blood loss.

Bleeding preventive effect of ACE910 in the long-term acquired hemophilia A model

The preventive effect of a weekly SC dose of ACE910 on spontaneous joint bleeds was evaluated in the model described in “Establishment of a long-term acquired hemophilia A model expressing spontaneous joint bleeds in cynomolgus monkeys.” The dosing regimen was simulated to maintain the plasma concentration trough at >30 μg/mL from the first week after administration using the PK parameters of ACE910 in cynomolgus monkeys previously obtained because around 26 μg/mL of plasma ACE910 would show hemostatic activity




Figure 2. Comparison of the mouse-monkey chimeric anti-primate FVIII neutralizing antibody cyVIII-2236 with the original mouse antibody VIII-2236 in an APTT assay. (A) Effects of cyVIII-2236 and VIII-2236 on APTT in normal cynomolgus monkey plasma. (B) Influence of 300 μg/mL cyVIII-2236 and VIII-2236 on APTT-shortening activity of ACE910 in FVIII-deficient human plasma. Data are expressed as means ± SD (n = 3). The bars depicting SD are shorter than the height of the symbols. The symbols for the group without anti-FVIII antibody are hidden behind the symbols for the other groups in panel B.

Table 1. Macroscopic hemorrhagic findings at necropsy on day 56 in a long-term primate model of acquired hemophilia A

Organ/tissue	Site	No. of animals with findings		
		Vehicle, n = 4	ACE910, n = 3	
Joint	Elbow	Left	1	0
		Right	1	0
	Hip	Left	4	0
		Right	1	0
	Ankle	Left	1	0
Skeletal muscle	Lower abdomen	2	0	
	Femur	1	0	
Urinary bladder	Wall	2	0	
Seminal vesicle	Serosa	1	0	
Rectum	Serosa	1	0	
Skin	Back		1	0
		Subcutis	1	0

In the other 3 monkeys, plasma ACE910 level remained around the target concentration (30 µg/mL) from day 4 to day 56 (Figure 6), demonstrating that the multiple-dosing simulation worked very well. Anti-ACE910 alloantibodies were examined in plasma collected at necropsy (day 28 for the #4 monkey; day 56 for the other 3 monkeys). In addition to the #4 monkey, 1 other monkey (ACE910 #3) had anti-ACE910 alloantibodies. In this monkey, however, the plasma ACE910 level slightly decreased only after day 49, and remained around 26 µg/mL, a level which is expected to show hemostatic activity.¹⁶

Regarding the joint bleeding symptoms, no limping and no macroscopic bleeding joints at necropsy were observed in any of the ACE910-treated monkeys, including monkey #4. In the ACE910 group (n = 3; excluding #4 statistically), the number of limping days (Figure 3C) and the number of bleeding joints at necropsy (Figure 3D, Table 1) significantly decreased (*P* < .01, 2-tailed Student *t* test), compared with those of the vehicle group (n = 4). In the ACE910 group, synovial hyperplasia and vascular proliferation in the synovial membranes of joints were histopathologically noted (Figure 4D-F). However, these findings were less severe and less frequent in the ACE910 group than those in the joints with neither macroscopic bleeding nor limping in the vehicle group.

Regarding the other spontaneous bleeding symptoms, bruises (the maximum value of areas and the number of days detected during the observation period) were ameliorated, and no hematuria and no organ bleeds at necropsy were found except for the oral cavity bleeds of the ACE910 #4 monkey (Table 1). Furthermore, monkeys #1 to 3

in the ACE910 group exhibited blood Hgb of 90% or above for 8 weeks (Figure 5). The minimum value of blood Hgb during the observation period was 94.9% ± 4.9% in the ACE910 group (n = 3, excluding monkey #4), which was significantly higher than 75.2% ± 11.3% in the vehicle group (n = 4) (*P* < .05, 2-tailed Student *t* test).

Discussion

Joint damage is one of the most problematic bleeding-related complications in hemophilia A patients. Manco-Johnson et al demonstrated in a prospective clinical study that the progression of joint damage could be decreased by routine alternate-day doses of FVIII from infancy.⁴ Although this dosing regimen could render a severe disease (<1% FVIII:C) into a moderate one (1%-5%), its preventive effect on joint bleeds was not perfect,^{4,17} and an analysis by den Uijl et al determined that the threshold FVIII:C level required to be free from joint bleeds would be 12%.¹⁸ Our previous study in a primate model of acquired hemophilia A predicted that ≥26 µg/mL ACE910 would exhibit hemostatic activity within the range of the mild phenotype (>5%) and that such a hemostatic level would be maintained by weekly SC doses of 0.64 mg/kg ACE910.¹⁶ Therefore, in the present study, we investigated weekly SC doses of 1 mg/kg ACE910 and demonstrated that such a regimen significantly prevented spontaneous joint bleeding, which also confirmed that the simulation of plasma ACE910 concentrations worked well. Because PK data of therapeutic antibodies from cynomolgus monkeys can be scaled to project human PK profiles,^{19,20} a similar PK profile and efficacy are expected for ACE910 in a clinical setting.

In this study, a long-term acquired hemophilia A model was newly established. The FVIII:C in this model was speculated to keep <4% and <3% in the vehicle and ACE910 groups, respectively, according to plasma cyVIII-2236 level and the cyVIII-2236 concentration vs FVIII:C correlation in the pooled cynomolgus monkey plasma (supplemental Figures 2 and 3). In the case of congenital hemophilia A, a small percentage of FVIII:C is considered to render the severity of bleeding; however, in acquired hemophilia A, measured FVIII:C does not necessarily correlate with the severity.²¹ Although cyVIII-2236 would not have necessarily decreased FVIII:C in the 1-stage clotting assay to <1% in the monkeys, the control monkeys actually presented severe bleeding symptoms. Thus, we considered that this model was a severe




Figure 4. Histopathological findings in a representative joint in a long-term primate model of acquired hemophilia A. Left hip joint with limping in the vehicle #1 monkey (A-C) and left hip joint without limping in the ACE910 #3 monkey (D-F) are shown at original magnification ×5 (A, D), ×20 (B, E), and ×400 (C, F). Hemorrhagic changes (arrowheads) including hemosiderin deposition (arrows) and destruction of articular cartilage/underlying bone (*) are detected in the joint with limping (A-C). No hemorrhagic changes are noted in the joint without limping (D-F). Hematoxylin and eosin stain. Aca, articular cartilage of acetabulum; Hb, hip bone; Sm, synovial membrane.




Figure 5. Relative blood Hgb concentrations in a long-term primate model of acquired hemophilia A. The time courses of blood Hgb concentration relative to concentration on day 0 (2 hours after cyVIII-2236 injection) are shown as individual values in respective cynomolgus monkeys (#1-4) of the vehicle and ACE910 groups.

acquired hemophilia A one. In an attempt to further validate this model by looking at reactivity to injected FVIII, we examined the effects of rpoFVIII, which is not neutralized by cyVIII-2236 (data not shown). Aiming to evaluate the marginal level of rpoFVIII efficacy, we administered rpoFVIII for 8 weeks with a twice-weekly 20 U/kg IV dosing regimen, which was set to keep $\geq 1\%$ rpoFVIII:C for 6 days a week (supplemental Figures 4 and 5A).²² The prolonged APTT was shortened immediately after the initial rpoFVIII injection (supplemental Figure 5B); however, in 3 of 4 monkeys, the APTT-shortening effect gradually disappeared in the middle of the dosing period, suggesting that anti-rpoFVIII alloantibodies had developed. In terms of bleeding symptoms, the number of days with limping was lower in the rpoFVIII group than in the vehicle group (supplemental Figure 5C); furthermore, the course of blood Hgb tended to worsen when anti-rpoFVIII alloantibodies presumably emerged (supplemental Figure 5D). In the monkey in which rpoFVIII maintained the APTT-shortening effect, limping and decrease in blood Hgb were not observed; nevertheless, 2 bleeding joints were noted by macroscopic observation at necropsy. Although it was difficult to fully evaluate the effect of rpoFVIII because anti-rpoFVIII alloantibodies developed, these results suggest that increasing FVIII activity could ameliorate the bleeding tendency in this model; however, the twice-weekly 20 U/kg IV dosing regimen was not sufficient to fully prevent joint bleeding.

ACE910 and rpoFVIII are proteins foreign to cynomolgus monkeys; therefore, the development of alloantibodies to these is theoretically inevitable.²³ Incidence rates of anti-humanized antibody alloantibodies in cynomolgus monkeys are reported to vary (0%-100%), and antigenicity in cynomolgus monkeys cannot predict that in humans.¹⁹ Although the antigenicity risk score of ACE910 in an *in silico* T-cell epitope prediction system was comparable to that of trastuzumab and palivizumab, which are nonimmunogenic in a clinical setting,¹⁵ the rate of anti-ACE910 alloantibody development must be evaluated in the actual clinical setting.

In this long-term acquired hemophilia A model, joint damage involving abnormal motion and histopathological features associated with intraarticular hemorrhage were similar to those of hemophilia A patients.^{3,24,25} Although a number of animal models have been reported for hemophilia A, it has been difficult to develop reproducible spontaneous joint bleeds.²⁶⁻²⁸ The FVIII-deficient mouse model of hemophilia A needs an artificial injuring procedure to induce joint bleeding.^{27,29} Although congenital hemophilia A models in dogs, sheep, rats, and pigs have been reported to develop

joint bleeds, it seems difficult to express reproducible joint bleeds in individuals and to efficiently evaluate drug efficacy in a practicable experimental period.²⁸⁻³⁴ Our model stably developed leg-joint damage within 8 weeks, which was possibly produced by the severe acquired hemophilic state and by bodyweight loading due to the bipedal motion of the monkeys. Thus, this model may be particularly useful in testing the efficacy of therapeutic agents from the orthopedic aspect of hemophilia A. In addition, this primate model should be useful for the many therapeutic antibodies that have poor interspecies cross-reactivity. Moreover, along with our model of hemophilia A, the strategy of using a mouse-host animal chimeric antibody may be beneficial for establishing long-term acquired animal models of other protein-deficiency diseases in the hematology research field.

Although limping and macroscopic intraarticular hemorrhage at necropsy were observed in all 4 control monkeys, joint swelling was noted in only 2 monkeys. Therefore, it was not so reproducible and may be inappropriate for quantitative assessment. The joint swelling was linked to limping in both monkeys. However, in one of the monkeys (vehicle #4), the ankle joint that had swollen did not present macroscopic intraarticular hemorrhage. We assume that the intraarticular hemorrhage may have been absorbed during 2 weeks after the remission of joint swelling.

In this study, although ACE910 completely prevented the macroscopic joint impairment, subclinical histopathological changes in the joint synovium were found, despite the presence of ACE910. We assume that the subclinical or small joint bleeds would occur in this model, and ACE910 prevented them from becoming larger and from leading to clinical symptoms, but did not provide a histopathological complete recovery at around 30 $\mu\text{g/mL}$ of plasma ACE910 concentration. An appropriate clinical investigation will be required to elucidate whether such an action of ACE910 can completely prevent macroscopic and clinical joint damage in the long run.

In conclusion, a long-term acquired hemophilia A model expressing reproducible spontaneous joint bleeds and other bleeds was newly established in cynomolgus monkeys, and weekly SC doses of ACE910 significantly prevented these bleeding symptoms. The difficulty in venous access negatively affects the adoption of and adherence to the routine prophylaxis regimen in home settings.³⁵




Figure 6. Plasma ACE910 concentrations in a long-term primate model of acquired hemophilia A. ACE910 was administered at an initial SC dose of 3.97 mg/kg on day 0 followed by weekly SC doses of 1 mg/kg on days 7, 14, 21, 28, 35, 42, and 49. The time courses of actual measured and simulated plasma concentrations of ACE910 are shown. The actual measured concentrations are presented as individual values for cynomolgus monkeys (#1-4) of the ACE910 group.

ACE910 is expected to provide hemophilia A patients, regardless of FVIII inhibitors, with a more effective and user-friendly way of bleeding prophylaxis. Clinical investigation of ACE910 is currently ongoing.

Acknowledgments

The authors thank their colleagues at Chugai Pharmaceutical Co, Ltd: T. Matsuura, M. Hiranuma, R. Takemoto, T. Koike, H. Kitamura, and T. Houjo for the *in vivo* pharmacologic experiments; T. Kojima, T. Wakabayashi, E. Tanaka, K. Esaki, Y. Kikuchi, M. Wada, M. Goto, H. Tsunoda, T. Suzuki, Y. Okuyama-Nishida, A. Harada, M. Funaki, S. Suzuki, T. Toyoda, and M. Ijiri for the preparation of test items and the *in vitro* experiments; Y. Sakamoto, T. Tachibana, and M. Nanami for the PK studies; and H. Saito for providing advice for the various experiments. The work was carried out at the Research Division, Chugai Pharmaceutical Co, Ltd.

This work was supported by research funding from Chugai Pharmaceutical Co, Ltd.

References

- Gringeri A, Muça-Perja M, Mangiafico L, von Mackensen S. Pharmacotherapy of haemophilia A. *Expert Opin Biol Ther*. 2011;11(8):1039-1053.
- Berntorp E, Shapiro AD. Modern haemophilia care. *Lancet*. 2012;379(9824):1447-1456.
- Roosendaal G, Lafeber FP. Blood-induced joint damage in hemophilia. *Semin Thromb Hemost*. 2003;29(1):37-42.
- Manco-Johnson MJ, Abshire TC, Shapiro AD, et al. Prophylaxis versus episodic treatment to prevent joint disease in boys with severe hemophilia. *N Engl J Med*. 2007;357(6):535-544.
- Oldenburg J. Prophylaxis in bleeding disorders. *Thromb Res*. 2011;127(suppl 1):S14-S17.
- Ragni MV, Fogarty PJ, Josephson NC, Neff AT, Raffini LJ, Kessler CM. Survey of current prophylaxis practices and bleeding characteristics of children with severe haemophilia A in US haemophilia treatment centres. *Haemophilia*. 2012;18(1):63-68.
- Astermark J, Donfield SM, DiMichele DM, et al; FENOC Study Group. A randomized comparison of bypassing agents in hemophilia complicated by an inhibitor: the FEIBA NovoSeven Comparative (FENOC) Study. *Blood*. 2007;109(2):546-551.
- Jimenez-Yuste V, Rodriguez-Merchan EC, Alvarez MT, Quintana M, Martin-Salces M, Hernandez-Navarro F. Experiences in the prevention of arthropathy in haemophilia patients with inhibitors. *Haemophilia*. 2008;14(suppl 6):28-35.
- Hay CR, DiMichele DM; International Immune Tolerance Study. The principal results of the International Immune Tolerance Study: a randomized dose comparison. *Blood*. 2012;119(6):1335-1344.
- Valentino LA, Ismael Y, Grygotis M. Novel drugs to treat hemophilia. *Expert Opin Emerg Drugs*. 2010;15(4):597-613.
- Lillicrap D. The future of hemostasis management. *Pediatr Blood Cancer*. 2013;60(suppl 1):S44-S47.
- Pipe SW. The hope and reality of long-acting hemophilia products. *Am J Hematol*. 2012;87(suppl 1):S33-S39.
- Kaufman RJ, Powell JS. Molecular approaches for improved clotting factors for hemophilia. *Blood*. 2013;122(22):3568-3574.
- Kitazawa T, Igawa T, Sampei Z, et al. A bispecific antibody to factors IXa and X restores factor VIII hemostatic activity in a hemophilia A model. *Nat Med*. 2012;18(10):1570-1574.
- Sampei Z, Igawa T, Soeda T, et al. Identification and multidimensional optimization of an asymmetric bispecific IgG antibody mimicking the function of factor VIII cofactor activity. *PLoS ONE*. 2013;8(2):e57479.
- Muto A, Yoshihashi K, Takeda M, et al. Anti-factor IXa/X bispecific antibody (ACE910): hemostatic potency against ongoing bleeds in a hemophilia A model and the possibility of routine supplementation. *J Thromb Haemost*. 2014;12(2):206-213.
- Collins PW. Personalized prophylaxis. *Haemophilia*. 2012;18(suppl 4):131-135.
- Den Uijl IE, Mauser Bunschoten EP, Roosendaal G, et al. Clinical severity of haemophilia A: does the classification of the 1950s still stand? *Haemophilia*. 2011;17(6):849-853.
- Ponce R, Abad L, Amaravadi L, et al. Immunogenicity of biologically-derived therapeutics: assessment and interpretation of nonclinical safety studies. *Regul Toxicol Pharmacol*. 2009;54(2):164-182.
- Deng R, Iyer S, Theil FP, Mortensen DL, Fielder PJ, Prabhu S. Projecting human pharmacokinetics of therapeutic antibodies from nonclinical data: what have we learned? *MABs*. 2011;3(1):61-66.
- Franchini M, Targher G, Montagnana M, Lippi G. Laboratory, clinical and therapeutic aspects of acquired hemophilia A. *Clin Chim Acta*. 2008;395(1-2):14-18.
- Srivastava A, Brewer AK, Mauser-Bunschoten EP, et al; Treatment Guidelines Working Group on Behalf of The World Federation Of Hemophilia. Guidelines for the management of hemophilia. *Haemophilia*. 2013;19(1):e1-e47.
- Dong JQ, Salinger DH, Endres CJ, et al. Quantitative prediction of human pharmacokinetics for monoclonal antibodies: retrospective analysis of monkey as a single species for first-in-human prediction. *Clin Pharmacokinet*. 2011;50(2):131-142.
- Roosendaal G, Vianen ME, Wenting MJ, et al. Iron deposits and catabolic properties of synovial tissue from patients with haemophilia. *J Bone Joint Surg Br*. 1998;80(3):540-545.
- Roosendaal G, van Rinsum AC, Vianen ME, van den Berg HM, Lafeber FP, Bijlsma JW. Haemophilic arthropathy resembles degenerative rather than inflammatory joint disease. *Histopathology*. 1999;34(2):144-153.
- Valentino LA, Hakobyan N, Kazarian T, Jabbar KJ, Jabbar AA. Experimental haemophilic synovitis: rationale and development of a murine model of human factor VIII deficiency. *Haemophilia*. 2004;10(3):280-287.
- Monahan PE. The expanding menagerie: animal models of hemophilia A. *J Thromb Haemost*. 2010;8(11):2469-2471.
- Nichols TC, Dillow AM, Franck HW, et al. Protein replacement therapy and gene transfer in canine models of hemophilia A, hemophilia B, von Willebrand disease, and factor VII deficiency. *ILAR J*. 2009;50(2):144-167.
- Hoffman M. Animal models of bleeding and tissue repair. *Haemophilia*. 2008;14(suppl 3):62-67.
- Nichols TC, Raymer RA, Franck HW, et al. Prevention of spontaneous bleeding in dogs with haemophilia A and haemophilia B. *Haemophilia*. 2010;16(suppl 3):19-23.
- Porada CD, Sanada C, Long CR, et al. Clinical and molecular characterization of a re-established line of sheep exhibiting hemophilia A. *J Thromb Haemost*. 2010;8(2):276-285.
- Booth CJ, Brooks MB, Rockwell S, et al. WAG-F8 (m1Ycb) rats harboring a factor VIII gene mutation provide a new animal model for hemophilia A. *J Thromb Haemost*. 2010;8(11):2472-2477.
- Lozier JN, Nichols TC. Animal models of hemophilia and related bleeding disorders. *Semin Hematol*. 2013;50(2):175-184.
- Kashiwakura Y, Mimuro J, Onishi A, et al. Porcine model of hemophilia A. *PLoS ONE*. 2012;7(11):e49450.
- Berntorp E. Joint outcomes in patients with haemophilia: the importance of adherence to preventive regimens. *Haemophilia*. 2009;15(6):1219-1227.

Authorship

Contribution: A.M., K.Y., M.T., T. Kitazawa, T.S., and Y.K. designed, performed, and analyzed the pharmacologic studies; T.I., Z.S., T. Kuramochi, and A.S. prepared the test items; K. Haraya conducted the PK studies; K.A. conducted the histopathological studies; K.N. and M.S. interpreted data and provided advice from the viewpoints of their medical expertise in hemophilia; K. Hattori provided direction and organized the program; and A.M. and T. Kitazawa wrote this manuscript.

Conflict-of-interest disclosure: A.M., K.Y., T. Kitazawa, T.S., T.I., Z.S., K. Hattori, M.T., T. Kuramochi, A.S., K. Haraya, K.A., and Y.K. are employees of Chugai Pharmaceutical, and the former 7 authors are inventors of the patents relating to anti-FIXa/X bispecific antibodies, of which all rights have been assigned to the company. K.N. and M.S. receive research support paid to their institution from Chugai Pharmaceutical, and have received consulting honoraria from Chugai Pharmaceutical and F. Hoffmann-La Roche.

Correspondence: Atsushi Muto, Research Division, Chugai Pharmaceutical Co, Ltd, 1-135 Komakado, Gotemba, Shizuoka 412-8513, Japan; e-mail: mutoats@chugai-pharm.co.jp.



blood

2014 124: 3165-3171
doi:10.1182/blood-2014-07-585737 originally published
online October 1, 2014

Anti-factor IXa/X bispecific antibody ACE910 prevents joint bleeds in a long-term primate model of acquired hemophilia A

Atsushi Muto, Kazutaka Yoshihashi, Minako Takeda, Takehisa Kitazawa, Tetsuhiro Soeda, Tomoyuki Igawa, Zenjiro Sampei, Taichi Kuramochi, Akihisa Sakamoto, Kenta Haraya, Kenji Adachi, Yoshiki Kawabe, Keiji Nogami, Midori Shima and Kunihiro Hattori

Updated information and services can be found at:
<http://www.bloodjournal.org/content/124/20/3165.full.html>

Articles on similar topics can be found in the following Blood collections
Free Research Articles (2875 articles)
Thrombosis and Hemostasis (788 articles)

Information about reproducing this article in parts or in its entirety may be found online at:
http://www.bloodjournal.org/site/misc/rights.xhtml#repub_requests

Information about ordering reprints may be found online at:
<http://www.bloodjournal.org/site/misc/rights.xhtml#reprints>

Information about subscriptions and ASH membership may be found online at:
<http://www.bloodjournal.org/site/subscriptions/index.xhtml>



Review

pH-dependent antigen-binding antibodies as a novel therapeutic modality[☆]T. Igawa^{*}, F. Mimoto, K. Hattori

Chugai Pharmaceutical Co. Ltd., Research Division, Japan

ARTICLE INFO

Article history:

Received 31 January 2014

Received in revised form 31 July 2014

Accepted 5 August 2014

Available online 12 August 2014

Keywords:

Monoclonal antibody

Antibody engineering

pH dependent

ABSTRACT

Monoclonal antibodies have become a general modality in therapeutic development. However, even with infinite binding affinity to an antigen, a conventional antibody is limited in that it can bind to the antigen only once, and this results in antigen-mediated antibody clearance when the a membrane-bound antigen is targeted, or in antibody-mediated antigen accumulation when a soluble antigen is targeted. Recently, a pH-dependent antigen-binding antibody that binds to an antigen in plasma at neutral pH and dissociates from the antigen in endosome at acidic pH has been reported to overcome this limitation and to reduce antigen-mediated antibody clearance and antibody-mediated antigen accumulation. A pH-dependent binding antibody against a soluble antigen can be further improved by Fc engineering to enhance the Fc receptor binding. Various approaches, including histidine-based engineering, direct cloning from immunized animals, and synthetic and combinatorial libraries, have been successfully applied to generate pH-dependent binding antibodies against various antigens. This review discusses the features, approaches, advantages, and challenges of developing a pH-dependent binding antibody as a novel therapeutic modality. This article is part of a Special Issue entitled: Recent advances in molecular engineering of antibody.

© 2014 Elsevier B.V. All rights reserved.

1. Introduction

With more than 30 monoclonal antibodies already approved for therapeutic use [1] and more than 300 under clinical development [2], monoclonal antibodies are clearly expected to play an important role in future therapeutics. Therefore, antibody engineering technologies that improve the therapeutic potency of monoclonal antibodies have been extensively studied in this decade. Such technologies include improving the binding affinity to a target, the specificity, the pharmacokinetics, and the effector function mediated by the Fc region of an antibody. Binding affinity can be improved by several methods, such as *in vitro* affinity maturation by library display [3] and computer-based *in silico* design [4]. For pharmacokinetics, lowering the isoelectric point of the variable region of an antibody is reported to prolong the antibody half-life by reducing the non-specific clearance of the antibody [5]. Fc engineering to enhance the binding affinity to FcRn at acidic pH also improves pharmacokinetics by increasing the recycling efficiency from the endosomes after non-specific uptake into cells [6]. Modulation of the antibody effector function by optimizing the interaction between the Fc region and an Fc gamma receptor is reported to improve the therapeutic efficacy of a monoclonal antibody [7]. Another noteworthy

technology that has recently been reported provides bispecific antibodies with a unique function that conventional monoclonal antibodies cannot achieve; they can neutralize two different disease-related cytokines [8]. In addition, bispecific antibodies can also redirect cytotoxic T cells to cancer cells by binding to CD3 and a tumor antigen [9], and they can mimic the function of coagulation factor VIII by binding to coagulation factor IXa and X [10].

Although these technologies have bestowed improved potency on therapeutic antibodies, it is now becoming more important to generate monoclonal antibodies that have further improved properties and are differentiated from conventional high-affinity monoclonal antibodies. In terms of modulating the interaction between an antigen and an antibody, achieving a high affinity binding to the antigen by affinity maturation has been the only approach for improvement. However, even with an infinite binding affinity to the antigen, a conventional antibody can bind to the antigen only once during its lifetime in plasma, and it is fundamentally limited by its consequent inability to neutralize further antigens, if the *in vivo* produced molar amount of antigen is larger than the amount of antibody injected.

If an antibody is targeting a membrane-bound antigen with a high rate of synthesis, such as IL-6 receptor (IL-6R), EGFR, CD4, or CD40 [11–14], the antibody is rapidly eliminated from plasma through antigen-mediated clearance. As a result, targeting these types of membrane antigen cannot be improved by affinity maturation and requires a high dose. Similarly, a high antibody dose is also required to neutralize soluble antigen at a high plasma concentration. The plasma

[☆] This article is part of a Special Issue entitled: Recent advances in molecular engineering of antibody.

^{*} Corresponding author. Tel.: +81 550 87 3029; fax: +81 550 87 5326.

E-mail address: igawatmy@chugai-pharm.co.jp (T. Igawa).

concentration of soluble antigen after antibody administration is determined by two factors: the baseline concentration and the antibody-mediated antigen accumulation. Some soluble antigens, such as IgE and C5, have a plasma baseline concentration that is already very high [15,16]; other soluble antigens without a high baseline concentration in plasma accumulate more than 1000-fold because the recycling function of an antibody inhibits degradation of the antigen and results in a very high concentration [17]. This accumulation after antibody administration occurs because an antigen in complex with an antibody has a longer half-life than the antigen alone [18,19]. The resulting high plasma antigen concentration requires a high antibody dose to neutralize the antigen. For both membrane-bound and soluble antigens, the limitation of conventional antibodies in only binding to the antigen once, even when the binding affinity is infinite, requires a high antibody dose.

Recently, we and others have reported that a pH-dependent antigen-binding property could overcome this limitation of conventional antibodies [20–22]. An antibody with a pH-dependent antigen-binding property dissociates the bound antigen in acidic endosomes after internalization into cells. Consequently, the dissociated antigen is trafficked to the lysosome and degraded, whereas the dissociated antibody, free of antigen, is recycled back to plasma by FcRn. The recycled free antibody can bind to another target antigen. By repeating this cycle, a pH-dependent antigen-binding antibody can bind to the target molecule more than once. Moreover, we have shown that the therapeutic potency of a pH-dependent antigen-binding antibody can be further enhanced by increasing its binding affinity to FcRn at neutral pH [23]. These studies demonstrate that pH-dependent antigen-binding antibodies can overcome the limitation of conventional antibodies and allow novel antibody therapeutics with differentiating properties to be generated. In this review, the features, advantages, and challenges of pH-dependent antigen-binding antibodies, and how they can be generated and optimized are discussed.

2. Effect of a pH-dependent antigen-binding antibody

2.1. Against a membrane-bound antigen

Fig. 1A describes the fate of a conventional antibody bound to a membrane-bound antigen. A conventional antibody bound to the cell surface antigen is internalized into cells, after which the antibody–antigen complex passes through the sorting endosome and is transferred to a lysosome, where eventually both the antibody and antigen are degraded by proteolysis. This means that, even with infinite binding

affinity to the antigen, a conventional antibody is limited to binding to an antigen only once. Conventional antibodies against membrane-bound antigens, such as IL-6R, EGFR, CD4, and CD40, exhibit non-linear clearance *in vivo* through antigen-mediated antibody clearance [11–14]. When a membrane-bound antigen is highly expressed in the body, the antibody is rapidly cleared from plasma, thereby requiring a high dose to neutralize this type of antigen over a long period.

Fig. 1B describes the fate of a pH-dependent antigen-binding antibody, which we also refer to as a recycling antibody, bound to a membrane-bound antigen. The recycling antibody bound to the cell surface antigen is internalized into the cells in the same way as a conventional antibody. However, in the sorting endosome, where the pH of the vesicle is acidic, the pH-dependent binding antibody is dissociated from the antigen. The antigen is transferred to the lysosome and degraded by proteolysis, while the dissociated antibody binds to FcRn in the endosome and is recycled back to the cell surface. Since FcRn does not bind to the antibody at neutral pH, the antibody dissociates from FcRn at the cell surface and returns back to plasma. A recycling antibody exhibits less non-linear antigen-mediated antibody clearance *in vivo* than conventional antibody and shows prolonged pharmacokinetics by avoiding antigen-mediated clearance. By repeating this cycle of binding at the cell surface and dissociating within the endosomes, a recycling antibody can bind to multiple antigens and overcomes the limitation of a conventional antibody.

The effect of a recycling antibody on a membrane-bound antigen was first demonstrated by our group using a monoclonal antibody against IL-6R [20]. A conventional antibody against IL-6R, tocilizumab, was engineered by a histidine mutagenesis approach (details described in section 4-1) to generate a pH-dependent binding antibody against IL-6R, named recycling tocilizumab. Importantly, tocilizumab and recycling tocilizumab have similar binding affinity to IL-6R at pH 7.4, while only recycling tocilizumab rapidly dissociates IL-6R at pH 6.0, allowing us to evaluate only the effect of pH-dependent antigen dissociation in the acidic endosomes. In a pharmacokinetic study in human IL-6R transgenic mice, the pharmacokinetics of recycling tocilizumab was better than that of tocilizumab, while in normal mice (whose IL-6R neither of the antibodies can bind) the pharmacokinetics of the two antibodies were similar, demonstrating that engineering the pH dependency reduced IL-6R-mediated clearance but did not influence non-specific clearance. By using a FcRn null variant of each antibody, we confirmed that the reduced IL-6R-mediated clearance by recycling tocilizumab was dependent on FcRn binding, indicating that recycling tocilizumab dissociated within the endosomes and




Fig. 1. Models showing the fate of (A) a conventional antibody and (B) a recycling antibody bound to a membrane-bound antigen. (A) A conventional antibody bound to a membrane-bound antigen is internalized into the cell, after which the antibody–antigen complex is trafficked to the sorting endosome and finally to lysosome, where eventually both the antibody and antigen are degraded by lysosomal protease. (B) A recycling antibody bound to a membrane-bound antigen is internalized into the cell in the same way as a conventional antibody. However, in the sorting endosome, where the pH of the vesicle is acidic, the recycling antibody is dissociated from the antigen by means of its pH-dependent antigen-binding property. The antigen is transferred to the lysosome and degraded by proteolysis, while the dissociated antibody is recycled back to the cell surface and plasma by utilizing FcRn.

was recycled back to plasma by FcRn (unpublished data). Chaparro-Riggers *et al.* also showed that FcRn is indispensable for a pH-dependent anti-PCSK9 antibody to be rescued from antigen-mediated clearance [21]. This was demonstrated by the fact that the half-life of the pH-dependent anti-PCSK9 antibody was the same as that of an antibody without pH-dependent binding (0.7 days) in FcRn knockout mice, while in wild-type mice, that of the pH-dependent antibody (14.4 days) was significantly longer than that of the antibody without pH-dependent binding (2.9 days).

A further engineered version of recycling tocilizumab, named SA237, which has a pH-dependent antigen-binding property and enhanced FcRn binding, demonstrated improved pharmacokinetics in cynomolgus monkeys compared to tocilizumab [20]. Whereas a subcutaneous injection of 2 mg/kg of tocilizumab inhibited C-reactive protein production for approximately 10 days, 2 mg/kg of SA237 was effective for more than 28 days. A Phase 1 clinical study of SA237 demonstrated significantly prolonged pharmacokinetics compared to tocilizumab, similar to that observed in cynomolgus monkeys (unpublished data). This study result clinically validated the concept of pH-dependent antigen-binding recycling antibodies.

2.2. Against a soluble antigen

Fig. 2A describes the fate of a conventional antibody bound to a soluble antigen. The conventional antibody bound to the soluble antigen in plasma is taken up into cells by non-specific endocytosis or pinocytosis. The antibody–antigen complex is transferred to a sorting endosome and binds to FcRn at acidic pH. FcRn recycles the antibody–antigen complex to the cell surface, where the complex is dissociated from FcRn back to plasma. Although the antibody is recycled, it is recycled in an antigen-bound form; therefore, the antibody cannot bind to the next antigen. This also limits a conventional antibody to binding only one soluble antigen, just as with a membrane-bound antigen. A soluble antigen in the absence of a conventional antibody is generally transferred to a lysosome and degraded, since most target antigens do not bind to FcRn (only IgG and albumin bind to FcRn). By binding to the soluble antigen, a conventional antibody inhibits the lysosomal degradation pathway of the soluble antigen. Thus, a conventional antibody against a soluble antigen *in vivo* generally results in an increase of total plasma antigen concentration or an accumulation of soluble antigen in plasma, an effect that has been reported for IL-6, MCP1, amyloid beta, hepcidin, and other soluble antigens [17,24–28]. When a soluble antigen is abundant in the

body, antigen accumulation results in the antigen binding sites of the antibodies becoming rapidly saturated, and free antibodies that are capable of binding to the antigens is lost.

Fig. 2B describes the fate of a recycling antibody bound to a soluble antigen. The antibody–antigen complex formed by the recycling antibody is also non-specifically taken up into cells in the same way as a conventional antibody. However, within the sorting endosome, the antibody pH-dependently dissociates from the soluble antigen, which is transferred to the lysosome and degraded by proteolysis. On the other hand, the free antibody is recycled back to plasma by FcRn. By repeating this cycle of binding in plasma and dissociating within the endosome, recycling antibodies against soluble antigens can also bind to multiple antigens and overcome the limitation of conventional antibodies. Since an antigen that forms a complex with a recycling antibody is degraded in the lysosome, recycling antibodies can degrade soluble antigens at the rate of non-specific uptake of the antibody–antigen complexes, and thus degrade the antigens more effectively than conventional antibodies. A recycling antibody exhibits reduced plasma antigen accumulation *in vivo* by enhancing soluble antigen clearance, and extends the half-life of the free antibodies.

The effect of a recycling antibody on a soluble antigen was also first demonstrated by the same monoclonal antibodies against IL-6R described above. Soluble IL-6R (sIL-6R) clearance was compared in normal mice that received sIL-6R alone, tocilizumab with sIL-6R, or recycling tocilizumab with sIL-6R. While sIL-6R alone was rapidly cleared from plasma, sIL-6R injected together with tocilizumab showed reduced clearance because it was bound to tocilizumab, which has a long half-life. In contrast, sIL-6R injected together with recycling tocilizumab showed accelerated sIL-6R clearance [20]. This antibody–antigen co-injection study can be generally applied to evaluate whether the clearance of a soluble antigen can be accelerated by pH-dependent antigen binding. However, this study does not mimic the clinical situation, where an antibody is injected to plasma at a steady-state soluble antigen concentration. To mimic the clinical situation, mice were implanted with an infusion pump filled with sIL-6R that continuously provided sIL-6R into plasma and maintained a steady-state sIL-6R concentration. In this model, an injection of tocilizumab resulted in approximately 20-fold increase of total plasma sIL-6R concentration, while recycling tocilizumab significantly reduced the accumulation and resulted in only a 2-fold increase of total plasma sIL-6R concentration [23].

Chaparro-Riggers *et al.* reported using a pH-dependent binding antibody against PCSK9 [21]. PCSK9 is a soluble antigen responsible for




Fig. 2. Models showing the fate of (A) a conventional antibody, (B) a recycling antibody and (C) a sweeping antibody bound to a soluble antigen. (A) A conventional antibody bound to a soluble antigen in plasma is taken up into cells by non-specific endocytosis or pinocytosis. The antibody–antigen complex is transferred to a sorting endosome, where it binds to FcRn at acidic pH and is recycled back to the cell surface and plasma by utilizing FcRn. Although the antibody is recycled, it is recycled in an antigen-bound form; therefore, the antibody cannot bind to the next antigen. (B) A recycling antibody bound to a soluble antigen in plasma is taken up into cells in the same way as a conventional antibody. However, within the sorting endosome, the recycling antibody dissociates the soluble antigen by utilizing its pH-dependent antigen-binding property. The dissociated antigen is transferred to a lysosome and degraded by lysosomal protease. On the other hand, the free antibody is recycled back to plasma by FcRn. (C) A sweeping antibody with enhanced FcRn binding at neutral pH binds to a soluble antigen, and the antibody–antigen complex is rapidly taken up into the cell in a FcRn-mediated manner. By accelerating the uptake of antibody–antigen complexes into cells, a sweeping antibody can degrade a larger amount of soluble antigen than a recycling antibody.

enhancing the degradation of LDL receptors. Inhibition of PCSK9 inhibits LDL receptor degradation and thereby reduces the LDL cholesterol level in plasma. A conventional antibody against PCSK9, J10, showed non-linear clearance in cynomolgus monkeys, which was presumably caused by an accelerated clearance of PCSK9–antibody complexes via an unknown pathway. To overcome the relatively short pharmacokinetics of J10 at low dosages, J17, a histidine-engineered version of J10 with a pH-dependent PCSK9-binding property, was generated. J17 demonstrated significantly improved pharmacokinetics compared to J10 in cynomolgus monkeys, and the LDL cholesterol-lowering effect was longer than that of J10 at the same dose. In the microscopic assay using HepG2 cells, J10 was more efficiently trafficked to lysosomes than J17 when co-incubated with PCSK9, while the trafficking of PCSK9 to lysosomes was comparable in the presence of J10 and J17. The authors demonstrated that a PCSK9-bound antibody shows accelerated lysosomal degradation by an unknown pathway, and this degradation can be avoided by a pH-dependent binding antibody that dissociates from PCSK9 in the endosomes. This study also confirmed the favorable effect of pH-dependent binding against a soluble antigen. Recently, Devanaboyina *et al.* reported a pH-dependent binding antibody against IL-6. In a mouse co-injection study, a conventional antibody against IL-6, clone 0218, significantly reduced the clearance of IL-6, while the pH-dependent binding antibody derived from clone 0218, VH4, accelerated the clearance of IL-6 from plasma [22]. In a microscopic assay using GFP-fused human FcRn variant-transfected HMEC-1 cells, they demonstrated that the conventional antibody against IL-6, clone 0218, colocalized with IL-6 in early endosomes, while the pH-dependent binding antibody, clone VH4, dissociated IL-6 in the early endosomes, and the dissociated IL-6 was subsequently trafficked to lysosomes.

These studies clearly demonstrated that recycling antibody against soluble antigen generally clears soluble antigen faster than conventional antibody and reduces antigen accumulation in plasma. Because administering recycling antibody reduces the total soluble antigen concentration compared with conventional antibody, the antibody dose required to neutralize the soluble antigen can be reduced.

3. Improving a recycling antibody against soluble antigens into a sweeping antibody by Fc engineering

A recycling antibody, with the capability of dissociating soluble antigens in the endosomes, can accelerate the clearance of the soluble antigens from plasma. However, the rate of antigen clearance by a recycling antibody is limited by the slow rate of non-specific uptake of antibody–antigen complexes into cells, so further enhancing the rate of soluble antigen degradation would be very beneficial. To further enhance the clearance of soluble antigens from plasma, we have recently developed a sweeping antibody, which is an Fc-engineered pH-dependent binding antibody [23]. Since uptake of antibody–antigen complexes was the rate-limiting step for antigen degradation, we engineered the Fc region of a pH-dependent antigen-binding antibody to increase the binding affinity to FcRn at neutral pH (wild-type IgG1 just marginally binds to FcRn at pH 7.4). This sweeping antibody, a pH-dependent binding antibody with engineered IgG1, can bind to FcRn at neutral pH on the cell surface, which causes antibody–antigen complexes to be rapidly taken up into the cells in an FcRn-mediated manner and, therefore, degrades more soluble antigens than a recycling antibody (Fig. 2C).

By moderately increasing the binding affinity to FcRn at neutral pH, an *in vivo* study using human FcRn transgenic mice demonstrated that the clearance of soluble antigens can be accelerated approximately 50-fold while the antibody maintains pharmacokinetics similar to a conventional antibody with wild-type IgG1 (Fig. 3). It is important to note that the antigen clearance, not the antibody clearance, was selectively enhanced by simultaneously engineering the pH-dependent antigen binding and moderately increasing the binding affinity to FcRn at neutral pH. For their microscopic assay, Devanaboyina *et al.* used human/mouse chimeric FcRn, which has increased affinity to wild-

type human IgG1. In the assay, they reported that, while a soluble antigen remained bound to a non-pH dependent binding antibody in the sorting endosomes, the antigen was dissociated from the pH-dependent binding antibody, and the dissociated antigen was finally transferred to lysosomes [22]. This result also suggests that both pH-dependent antigen binding and enhanced FcRn binding are required to increase antigen uptake into cells. We also observed the same phenomenon in our *in vitro* confocal microscopy assay using the human FcRn-transfected MDCK cell line. While a soluble antigen remained bound to a non-pH-dependent binding antibody in the sorting endosome (Fig. 4A), the soluble antigen was dissociated from a pH-dependent binding antibody with engineered IgG1 and enhanced FcRn binding (Fig. 4B).

It should be noted that previous Fc engineering to increase the binding affinity to FcRn at acidic pH aimed to improve antibody pharmacokinetics by increasing FcRn-mediated IgG recycling [29]. On the other hand, we engineered a pH-dependent antigen-binding antibody to moderately increase its binding affinity at neutral pH to generate a sweeping antibody, which selectively enhanced the clearance of the antigen while maintaining antibody pharmacokinetics similar to wild-type IgG1 (Fig. 3). However, consistent with the report from Vaccaro *et al.* in which they reported that engineering the Fc region of IgG1 to create a strong binding to FcRn at neutral pH modulated *in vivo* concentration of endogenous IgG antibodies [30], a sweeping antibody with strongly increased binding affinity to FcRn at neutral pH resulted in increased antibody clearance, although antigen clearance was more accelerated than that of a sweeping antibody with moderately increased binding affinity to FcRn at neutral pH [23]. An appropriate level of binding affinity to FcRn at neutral pH should be selected, depending on the characteristics of the target soluble antigen.

An alternative approach for enhancing the uptake of antibody–antigen complexes into cells is utilizing uptake mediated by Fc gamma receptor, which has been reported elsewhere as contributing to the clearance of antibody–antigen immune complexes *in vivo* [31]. Especially, the clearance of small immune complexes is dominantly supported by Fc gamma receptor IIb (or CD32b) expressed on liver sinusoidal endothelial cells [32]. Because Fc gamma receptor IIb is reported to be a recycling receptor, enhancing the binding affinity to this receptor could have a similar effect to that observed when the binding affinity to FcRn at neutral pH was enhanced [33]. Previously reported Fc engineering to enhance binding affinity to Fc gamma receptor IIb also enhanced the binding affinity to Fc gamma receptor IIa (or CD32a) with arginine 131 allotype, which represents approximately 70% of the Caucasian population [34]. However, since Fc gamma receptor IIa is not a recycling receptor and is expressed on platelets, enhancing the binding affinity also to Fc gamma receptor IIa could have both pharmacokinetic and safety issues. Recently, we have reported a novel engineered Fc variant that selectively enhances binding to Fc gamma receptor IIb over both the histidine and arginine 131 allotypes of Fc gamma receptor IIa [35]. By combining pH-dependent antigen binding with this novel Fc engineering, we may be able to accelerate the antigen clearance and degrade more soluble antigen than with a recycling antibody.

4. Approaches for generating recycling antibodies

4.1. Experimental approaches

Previously, monoclonal antibodies generated by either animal immunization or *in vitro* antibody library display technology have been selected based on their strong binding affinity to their target antigens at neutral pH and their binding to a functionally appropriate epitope. Therefore, the monoclonal antibodies obtained in such a way seldom had a pH-dependent antigen-binding property. Although most antibodies lose their binding affinity to the antigen at pH 3.0, they do not show




Fig. 3. The effect of a conventional antibody and a sweeping antibody against a soluble antigen in human FcRn transgenic mice. (A) shows the time course of plasma concentration of total soluble antigen, hsl-6R, after antibodies were administered. (B) shows the plasma antibody concentration. Blue lines indicate a conventional antibody, red lines a sweeping antibody (pH-dependent antigen-binding antibody with enhanced FcRn binding), and gray lines the baseline of the soluble antigen.

rapid enough dissociation of antigens at pH 5.5–6.0 in a sorting endosome.

In our first report of a pH-dependent binding antibody [20], we engineered a conventional antibody against IL-6R, tocilizumab, into a pH-dependent binding antibody, which was identified by mice immunization and subsequent screening of the binding affinity and IL-6-binding neutralizing capability [36]. We took advantage of a natural amino acid, histidine, which has pKa of around 6 (depending on the surrounding environment) and is utilized in naturally occurring pH-dependent protein-protein interactions [37]. We introduced several histidine residues in the CDR or FR of tocilizumab by site-directed mutagenesis. Because of the pKa of the imidazole group, histidine residues are protonated at endosomal acidic pH, which results in destabilizing the antibody-antigen interaction either directly, with histidine residues involved in interacting with the antigen, or indirectly, with histidine residues involved in maintaining the conformation of

the CDRs. A crystal structure of tocilizumab Fab fragment in complex with IL-6R suggested that a mutation of the tyrosine residue in the CDR of tocilizumab, which interacts with an arginine residue, into a histidine residue resulted in electrostatic repulsion between the arginine and the protonated histidine residue at acidic pH (unpublished data). A total of three histidine residues and several other mutations were introduced into tocilizumab to compensate for the reduced affinity at neutral pH or to improve its physicochemical properties, and finally histidine-based engineered tocilizumab that rapidly dissociates from IL-6R at pH 6.0 was generated, as we previously reported [20]. The kinetic parameters of tocilizumab and the histidine-based engineered antibody are described in Table 1.

We have applied the same histidine-based engineering approach to several other antibodies against different antigens, and in all cases we have successfully conferred a pH-dependent antigen-binding property. However, in some cases, although introducing histidine(s) in the




Fig. 4. The intracellular trafficking of an antigen in complex with either a conventional antibody or a pH-dependent antigen-binding antibody in MDCK transfected with human FcRn-EGFP (in green) and beta-2-microglobulin. (A–C) are images of representative endosomes after a pH-dependent antibody with enhanced FcRn binding was incubated in complex with its target antigen, hsl-6R, which was labeled with Alexa-555 (in red). (D–F) are images of representative endosomes after a conventional antibody with enhanced FcRn binding was incubated in complex with hsl-6R. (A) and (D) are images of FcRn-EGFP. (B) and (E) are images of hsl-6R-Alexa555. (C) and (F) are overlays of the FcRn and hsl-6R images. These are unpublished data.

appropriate position can provide pH dependency (i.e. reduction of the binding affinity at acidic pH), it simultaneously reduces the binding affinity to the antigen at neutral pH to some extent (but less than that at acidic pH). Although such an antibody would elicit a recycling property, the loss of binding affinity at neutral pH would reduce the biological activity of the antibody in plasma. If the biological activity does not meet the therapeutic goal, the binding affinity at neutral pH needs to be improved by the affinity maturation process.

Following our report, Chaparro-Riggers *et al.* reported their pH-dependent binding antibody against PCSK9, which was also generated through histidine-based engineering of the parent antibody [21]. Moreover, Devanaboyina *et al.* recently reported generating a pH-dependent binding antibody against IL-6 in the same way [22]. Changing a single aspartic acid to a histidine residue in HCDR3 conferred pH-dependent binding on IL-6, but the binding affinity to IL-6 at neutral pH was slightly reduced. Other histidine mutations tested in this report either did not display an increased pH dependency or resulted in a substantial loss of the binding affinity at neutral pH.

These reports verify that histidine-based engineering is a general approach for generating pH-dependent binding antibodies that demonstrate a recycling property. However, in some cases, a simultaneous loss of binding affinity at neutral pH may occur. Since the binding affinity at neutral pH is important for the biological activity of the antibody, this loss of binding affinity at neutral pH may need to be recovered by subsequent affinity maturation (discussed further in section 6).

The second approach for generating a pH-dependent binding antibody is to isolate the antibody directly from immunized animals. Our trial to screen a large number of antigen-binding antibodies suggested the feasibility of directly isolating rare pH-dependent antigen-binding antibodies (generally less than 5% of the binders) from immunized animals by using direct B cell-cloning technology, which allows antigen-binding antibodies to be identified more efficiently than conventional mouse hybridoma technology [38].

The third approach for generating a pH-dependent binding antibody is to isolate the antibody directly from a synthetic antibody phage library. Since histidine is rarely used in the CDR of human antibodies (0.5%–0.6%), the possibility of directly isolating a pH-dependent binding antibody from naïve or synthetic antibody libraries that mimic the human repertoire is low. Therefore, a library of synthetic antibodies that use more histidine in the CDR in combination with pH-dependent antigen panning (antigen binding at neutral pH and elution at acidic pH) can be used to directly isolate a pH-dependent antigen-binding antibody.

The fourth approach is to use a combinatorial antibody library to generate a pH-dependent binding antibody from a conventional antibody. Murtaugh *et al.* described a combinatorial histidine scanning library approach to engineer a conventional antibody into a pH-dependent binding antibody [39]. In their study, a VHH antibody against RNase A was used as a model antibody, and a combinatorial library was constructed that consisted of combinations of histidine and wild-type residues in the antibody–antigen interface. Antibodies displayed on phages were coselected for high-affinity binding and pH-dependent binding, which resulted in generating pH-dependent VHH antibody against RNase A. Although there was a slight loss in the binding affinity at neutral pH, this approach may be an efficient way to generate a pH-dependent binding antibody from a conventional antibody. Although a histidine-based pH-dependent binding antibody is a promising type of

recycling antibody, we are exploring an alternative approach to designing an antibody that dissociates the antigen within sorting endosomes. While the recycling antibodies discussed above take advantage of the pH difference between plasma and endosomes, another environmental difference between plasma and endosomes is calcium ion concentration. Calcium ion concentration in plasma is reported to be 1.2–2 mM in plasma which drops to 3–30 μ M in endosomes after endocytosis [40]. Therefore, a calcium ion-dependent binding antibody, which binds to the antigen at the calcium ion concentration in plasma and dissociates the antigen at the calcium ion concentration in endosomes, may be used to generate a recycling antibody.

4.2. Computational approaches

Significant progress has been achieved in the field of computational approaches that design functional proteins, so identifying a pH-dependent antigen-binding antibody by computational design could be an alternative approach. However, although computational design has already been applied to improve antibody–antigen interaction [4] and to generate pH-dependent binding proteins [41,42], there is no report to date of successfully designing a pH-dependent antigen-binding antibody. Strauch *et al.* computationally designed a pH-dependent protein that binds to the Fc region by utilizing histidine residues in the Fc region [41]. Sarkar *et al.* designed a pH-dependent granulocyte colony-stimulating factor (GCSF) that maintained the affinity to its receptor at neutral pH. To do this, they computationally identified negatively charged residues in the interface of GCSF and the GCSF receptor (GCSFR). If these negatively charged residues are replaced with histidine residues, the GCSFR binding affinity at neutral pH is expected to stay the same but the binding affinity at acidic pH will be reduced. They selected residues to mutate into histidine by comparing the free energy values at neutral and acidic pH of each histidine-mutated GCSF in complex with GCSFR [42]. A similar approach could be applied to provide antibodies with a pH-dependent antigen-binding property.

5. Advantages of pH-dependent antigen-binding antibodies and potential applications

pH-dependent antigen-binding antibodies have an advantage over conventional antibodies, especially when the target molecules are membrane-bound antigens or soluble antigens abundantly present in the body. Antibodies targeting these antigens require either a high dose that cannot be injected subcutaneously or frequent dosing, both of which are not patient-friendly, especially in chronic diseases.

When an antibody targets an antigen that is both membrane-bound and has a high rate of synthesis, such as IL-6R, EGFR, CD4, or CD40, the antibody is rapidly eliminated from plasma through antigen-mediated clearance [11–14]. In the case of tocilizumab, which targets IL-6R, neither improving the binding affinity to IL-6R nor reducing the non-specific clearance by increasing the binding affinity to FcRn at acidic pH significantly improved the duration of IL-6R neutralization *in vivo* [23]. This is because an antibody that targets IL-6R is mainly cleared from plasma by IL-6R-mediated clearance. Modeling and simulation suggested that the molar amount of IL-6R produced in the body within approximately 18 days is the same as 2 mg/kg of antibody. Since a conventional antibody can bind to the antigen only once, it is evident that,

Table 1
Kinetic parameters of parent antibody (tocilizumab) and histidine-engineered antibody (recycling tocilizumab).

	k_a at pH 7.4 ($s^{-1} M^{-1}$)	k_d at pH 7.4 (s^{-1})	K_D at pH 7.4 (M)	k_d at pH 6.0 (s^{-1})	pH dependency (k_d at pH 6.0/ k_d at pH 7.4)
Tocilizumab	2.2×10^5	1.1×10^{-3}	5.1×10^{-9}	2.3×10^{-3}	2.3
Recycling tocilizumab	5.3×10^5	1.0×10^{-3}	1.9×10^{-9}	1.5×10^{-2}	15

even if a conventional antibody has infinite binding affinity to IL-6R or very low non-specific clearance, a 2 mg/kg dose cannot be effective for more than 18 days. A pH-dependent binding antibody against IL-6R (SA237) that is recycled back to plasma after IL-6R-mediated uptake into cells, can bind to IL-6R multiple times, which allows 2 mg/kg of SA237 to remain effective for more than one month. Importantly, because the amount of an antibody that can be administered in a single subcutaneous injection is limited (generally 1 mL of 150 mg/mL IgG formulation), a pH-dependent binding antibody would be the only way to achieve monthly subcutaneous dosing for an antibody that targets IL-6R. The same strategy can be applied to other antibodies targeting membrane-bound antigens that are rapidly eliminated from plasma through antigen-mediated clearance.

An antibody that targets a soluble antigen with a high baseline plasma concentration, such as IgE or C5, would also require a high dose to neutralize the antigen [15,16]. To use omalizumab, an anti-IgE antibody, in patients with high plasma IgE concentration, subcutaneous injections at two or three sites are required, because plasma IgE concentration (in the range of sub $\mu\text{g/mL}$) is higher than the concentration of most cytokines (generally in the range of pg/mL) [16]. Similarly, eculizumab, an anti-complement C5 antibody, requires intravenous infusion, because plasma C5 concentration is reported to be 70–140 $\mu\text{g/mL}$ [15]. Although eculizumab requires chronic dosing, such a large amount of antigen cannot be neutralized by a subcutaneous dose. Administering an antibody against a soluble antigen with rapid clearance, such as amyloid beta, MCP1, hepcidin, IL-6, CD23, or VEGF, results in more than 1000-fold increase of plasma antigen concentration from the baseline by the antigen accumulation effect [17,24–28]. Notably, administering an anti-hepcidin antibody resulted in an accumulation of hepcidin by approximately 5000-fold, which required a 300 mg/kg dose to neutralize hepcidin, despite its picomolar binding affinity [17]. Since soluble antigens with a molecular weight below 60,000 g/L are generally very rapidly eliminated through renal excretion [43], an antibody targeting a soluble antigen with a small molecular weight would suffer from the same issue. Taken together, conventional antibodies targeting soluble antigens with a high baseline concentration and/or a large antigen-accumulation effect require a high dose. Since conventional antibodies can bind to antigens only once and cannot accelerate the antigen clearance, pH-dependent binding antibodies would provide a significant advantage over conventional antibodies against these soluble antigens. pH-dependent binding antibodies would enable subcutaneous dosing or less frequent dosing that cannot be achieved by simple high-affinity conventional antibodies with a long half-life.

Recycling antibodies can be engineered into sweeping antibodies by Fc engineering to enhance the binding to Fc gamma receptor or FcRn. The use of a sweeping antibody against a soluble antigen further reduces the dosage compared to a recycling antibody, providing another advantage over a conventional antibody. Moreover, since a sweeping antibody can actively eliminate the target soluble antigen from plasma, a sweeping antibody against soluble IL-6R that targeted a non-neutralizing epitope and showed no neutralizing activity *in vitro* was able to antagonize the activity of soluble IL-6R *in vivo* by directly removing soluble IL-6R from plasma [23]. Therefore, a sweeping antibody can target a soluble ligand with multiple target epitopes, through which the ligand binds to different receptors that cannot be neutralized by a single monoclonal antibody. Administering a conventional antibody may neutralize the binding of a ligand to its receptor through one epitope, but another epitope remains. In addition, the increase of antibody–antigen complexes in plasma may allow increased signaling to the second receptor through the epitope that is not neutralized. The sweeping antibody, on the other hand, could directly remove such antigens from plasma, allowing multiple epitopes to be antagonized *in vivo*. The same strategy could be applied to toxic antigens that do not have a functional epitope. Similarly to an LDL receptor removing LDL or an asialoglycoprotein receptor removing asialoglycoprotein from plasma, a sweeping antibody could directly remove a toxic antigen from plasma

to alleviate the disease condition [44,45]. Since these types of soluble antigen could not be therapeutic targets for conventional antibodies, sweeping antibodies may expand the target antigen space of antibody therapeutics.

6. Challenges in developing a pH-dependent antigen-binding antibody as a therapeutic

Although a pH-dependent antigen-binding antibody has attractive features, there are several challenges when generating the antibody for therapeutic application. The following challenges are specific to pH-dependent antigen-binding antibodies and need not be considered when developing conventional monoclonal antibodies.

First, histidine-based engineering to generate a pH-dependent antigen-binding antibody sometimes abrogates the binding affinity at neutral pH. In the case of the anti-IL-6 antibody, histidine mutagenesis conferred a recycling property onto the parental antibody, but at the same time it also reduced the binding affinity against IL-6 at pH 7.4 compared to the parent antibody [22]. If the binding affinity at neutral pH of a histidine-based recycling antibody is not sufficient to elicit the desired therapeutic activity, the binding affinity at neutral pH can be improved by the affinity maturation process, which is commonly applied to generate a potent therapeutic antibody [3,4]. Although a general affinity maturation strategy can be applied to recover the loss of binding affinity at pH 7.4, such mutations tend to also increase the binding affinity at acidic pH, which is undesirable. Nevertheless, we have been able to successfully generate high-affinity recycling antibodies with nanomolar K_D at pH 7.4 and simultaneously have sufficient pH dependency or large k_d at acidic pH by using histidine-based engineering and affinity maturation in all the antibodies targeting various antigens tested to date.

Second, the extent of pH-dependency required to dissociate the target antigens in acidic endosomes is not clear. An anti-IL-6R antibody with 20-fold larger K_D and 15-fold larger k_d at pH 6.0 than at pH 7.4 showed an improved pharmacokinetic property in cynomolgus monkeys and humans. PCSK9, an antibody with only 2.6-fold larger K_D and 12-fold larger k_d at pH 6.0 than at pH 7.4, showed a prolonged half-life in cynomolgus monkeys, although its binding affinity to cynomolgus monkey PCSK9 at pH 7.4 was also reduced. These results suggest that the pH-dependency in k_d is a more important factor than that in K_D for improving the pharmacokinetics of an antibody, but we assume that the absolute k_d value in acidic pH, not the relative pH-dependency of k_d (the ratio of k_d at pH 7.4 to that at pH 6.0), would also be important to achieve sufficient dissociation of the antigen. However, the absolute k_d value at pH 6.0 required to sufficiently dissociate an antigen from an antibody remains to be elucidated. The absolute K_D at pH 6.0 would also be important if we consider that the internalized antibody–antigen complexes reach a new equilibrium in the acidic endosomes. A detailed kinetic analysis of endosomal trafficking of antibody–antigen complexes would be required to identify the parameters necessary to sufficiently dissociate the antigens in the endosomes. Although these parameters could differ between the pH-dependent binding antibodies that target membrane-bound antigens, those that target soluble antigens, and the Fc-engineered sweeping antibodies, identifying these parameters would be important for generating pH-dependent binding antibodies with optimum *in vivo* properties.

Third, when considering therapeutic application, it would be necessary to predict antibody pharmacokinetics or the rate of antigen clearance in humans by preclinical study. This is often conducted in a pharmacokinetic study using cynomolgus monkeys; therefore, the antibody preferably needs to bind to a cynomolgus monkey antigen with a similar binding affinity as to the human antigen. Although cross-reactivity to cynomolgus monkey antigens is not so challenging for conventional antibodies, it could be more challenging for pH-dependent binding antibodies since the binding kinetics (both K_D and k_d) at pH 6.0 also need to be considered. In the case of the pH-dependent anti-PCSK9 antibody, since pH-dependency of K_D and k_d

and absolute k_d at pH 6.0 is somewhat different between humans and cynomolgus monkeys, predicting human PK/PD from a cynomolgus monkey study could be difficult. Moreover, with sweeping antibodies, since the Fc region is engineered to enhance the binding affinity to an Fc receptor, cross-reactivity of the Fc region also needs to be considered.

7. Conclusion

Recycling antibodies with pH-dependent antigen binding offer significant advantages over conventional antibodies by dissociating the antigens in endosomes and recycling free antibodies back to plasma. Although generating a high-affinity pH-dependent antigen-binding antibody is still challenging, various approaches are currently available and have been successfully applied to various antigens. Because affinity maturation has now become a basic technology for engineering improved antibody therapeutics, pH-dependent binding antibodies should play an important role as a novel modality for antibody therapeutics in the near future.

References

- J.G. Elvin, R.G. Couston, C.F. van der Walle, Therapeutic antibodies: market considerations, disease targets and bioprocessing, *Int. J. Pharm.* 440 (2013) 83–98.
- J.M. Reichert, Which are the antibodies to watch in 2013? *mAbs* 5 (2013) 1–4.
- A.R. Bradbury, S. Sidhu, S. Dubel, J. McCafferty, Beyond natural antibodies: the power of in vitro display technologies, *Nat. Biotechnol.* 29 (2011) 245–254.
- S.M. Lippow, K.D. Witttrup, B. Tidor, Computational design of antibody-affinity improvement beyond in vivo maturation, *Nat. Biotechnol.* 25 (2007) 1171–1176.
- T. Igawa, H. Tsunoda, T. Tachibana, A. Maeda, F. Mimoto, C. Moriyama, M. Nanami, Y. Sekimori, Y. Nabuchi, Y. Aso, K. Hattori, Reduced elimination of IgG antibodies by engineering the variable region, *Protein Eng. Des. Sel.* 23 (2010) 385–392.
- W.F. Dall'Acqua, P.A. Kiener, H. Wu, Properties of human IgG1s engineered for enhanced binding to the neonatal Fc receptor (FcRn), *J. Biol. Chem.* 281 (2006) 23514–23524.
- J.R. Desjarlais, G.A. Lazar, Modulation of antibody effector function, *Exp. Cell Res.* 317 (2011) 1278–1285.
- C. Spiess, J. Bevers III, J. Jackman, N. Chiang, G. Nakamura, M. Dillon, H. Liu, P. Molina, J.M. Elliott, W. Shatz, J.M. Scheer, G. Giese, J. Persson, Y. Zhang, M.S. Dennis, J. Giulianotti, P. Gupta, D. Reilly, E. Palma, J. Wang, E. Stefanich, H. Scheerens, G. Fuh, L.C. Wu, Development of a human IgG4 bispecific antibody for dual targeting of interleukin-4 (IL-4) and interleukin-13 (IL-13) cytokines, *J. Biol. Chem.* 288 (2013) 26583–26593.
- A. Satta, D. Mezzanzanica, F. Turatti, S. Canevari, M. Figini, Redirection of T-cell effector functions for cancer therapy: bispecific antibodies and chimeric antigen receptors, *Future Oncol.* 9 (2013) 527–539.
- T. Kitazawa, T. Igawa, Z. Sampei, A. Muto, T. Kojima, T. Soeda, K. Yoshihashi, Y. Okuyama-Nishida, H. Saito, H. Tsunoda, T. Suzuki, H. Adachi, T. Miyazaki, S. Ishii, M. Kamata-Sakurai, T. Iida, A. Harada, K. Esaki, M. Funaki, C. Moriyama, E. Tanaka, Y. Kikuchi, T. Wakabayashi, M. Wada, M. Goto, T. Toyoda, A. Ueyama, S. Suzuki, K. Haraya, T. Tachibana, Y. Kawabe, M. Shima, A. Yoshioka, K. Hattori, A bispecific antibody to factors IXa and X restores factor VIII hemostatic activity in a hemophilia A model, *Nat. Med.* 18 (2012) 1570–1574.
- Y. Ohsugi, T. Kishimoto, The recombinant humanized anti-IL-6 receptor antibody tocilizumab, an innovative drug for the treatment of rheumatoid arthritis, *Expert. Opin. Biol. Ther.* 8 (2008) 669–681.
- M.A. Tabrizi, C.M. Tseng, L.K. Roskos, Elimination mechanisms of therapeutic monoclonal antibodies, *Drug Discov. Today* 11 (2006) 81–88.
- M. Tabrizi, G.G. Bornstein, H. Suria, Biodistribution mechanisms of therapeutic monoclonal antibodies in health and disease, *AAPS J.* 12 (2010) 33–43.
- S.K. Kelley, T. Gelzleichter, D. Xie, W.P. Lee, W.C. Darbonne, F. Qureshi, K. Kissler, E. Ofiazoglu, I.S. Grewal, Preclinical pharmacokinetics, pharmacodynamics, and activity of a humanized anti-CD40 antibody (SGN-40) in rodents and non-human primates, *Br. J. Pharmacol.* 148 (2006) 1116–1123.
- N. Hayashi, Y. Tsukamoto, W.M. Sallas, P.J. Lowe, A mechanism-based binding model for the population pharmacokinetics and pharmacodynamics of omalizumab, *Br. J. Clin. Pharmacol.* 63 (2007) 548–561.
- K.M. Zareba, Eculizumab: a novel therapy for paroxysmal nocturnal hemoglobinuria, *Drugs Today (Barc)* 43 (2007) 539–546.
- J.J. Xiao, W. Krzyzanski, Y.M. Wang, H. Li, M.J. Rose, M. Ma, Y. Wu, B. Hinkle, J.J. Perez-Ruixo, Pharmacokinetics of anti-hepcidin monoclonal antibody Ab 12B9m and hepcidin in cynomolgus monkeys, *AAPS J.* 12 (2010) 646–657.
- F.D. Finkelman, K.B. Madden, S.C. Morris, J.M. Holmes, N. Boiani, I.M. Katona, C.R. Maliszewski, Anti-cytokine antibodies as carrier proteins. Prolongation of in vivo effects of exogenous cytokines by injection of cytokine-anti-cytokine antibody complexes, *J. Immunol.* 151 (1993) 1235–1244.
- Charles B. Davis, Leeann P. Tobia, Deborah C. Kwok, Christine M. Oishi, Neil Khetarpal, Timothy W. Hepburn, Lisa J. Benincosa, Fung-Sing Chow, William J. Jusko, Accumulation of antibody-target complexes and the pharmacodynamics of clotting after single intravenous administration of humanized anti-factor IX monoclonal antibody to rats, *Drug Deliv.* 6 (1999) 171–179.
- T. Igawa, S. Ishii, T. Tachibana, A. Maeda, Y. Higuchi, S. Shimaoka, C. Moriyama, T. Watanabe, R. Takubo, Y. Doi, T. Wakabayashi, A. Hayasaka, S. Kadono, T. Miyazaki, K. Haraya, Y. Sekimori, T. Kojima, Y. Nabuchi, Y. Aso, Y. Kawabe, K. Hattori, Antibody recycling by engineered pH-dependent antigen binding improves the duration of antigen neutralization, *Nat. Biotechnol.* 28 (2010) 1203–1207.
- J. Chaparro-Riggers, H. Liang, R.M. DeVay, L. Bai, J.E. Sutton, W. Chen, T. Geng, K. Lindquist, M.G. Casas, L.M. Boustany, C.L. Brown, J. Chabot, B. Gomes, P. Garzone, A. Rossi, P. Strop, D. Shelton, J. Pons, A. Rajpal, Increasing serum half-life and extending cholesterol lowering in vivo by engineering antibody with pH-sensitive binding to PCSK9, *J. Biol. Chem.* 287 (2012) 11090–11097.
- S.C. Devanaboyina, S.M. Lynch, R.J. Ober, S. Ram, D. Kim, A. Puig-Canto, S. Breen, S. Kasturirangan, S. Fowler, L. Peng, H. Zhong, L. Jermutus, H. Wu, C. Webster, E.S. Ward, C. Gao, The effect of pH dependence of antibody-antigen interactions on subcellular trafficking dynamics, *mAbs* 5 (2013) 851–859.
- T. Igawa, A. Maeda, K. Haraya, T. Tachibana, Y. Iwayanagi, F. Mimoto, Y. Higuchi, S. Ishii, S. Tamba, N. Hironiwa, K. Nagano, T. Wakabayashi, H. Tsunoda, K. Hattori, Engineered monoclonal antibody with novel antigen-sweeping activity in vivo, *PLoS One* 8 (2013) e63236.
- J.P. Davda, R.J. Hansen, Properties of a general PK/PD model of antibody-ligand interactions for therapeutic antibodies that bind to soluble endogenous targets, *mAbs* 2 (2010) 576–588.
- J.J. Haringman, D.M. Gerlag, T.J. Smeets, D. Baeten, F. van den Bosch, B. Bresnihan, F.C. Breedveld, H.J. Dinant, F. Legay, H. Gram, P. Loetscher, R. Schouder, T. Woodworth, P.P. Tak, A randomized controlled trial with an anti-CCl2 (anti-monocyte chemoattractant protein 1) monoclonal antibody in patients with rheumatoid arthritis, *Arthritis Rheum.* 54 (2006) 2387–2392.
- P.L. Martin, J. Cornacoff, U. Prabhakar, T. Lohr, G. Treacy, J.E. Sutherland, S. Hersey, E. Martin, Reviews preclinical safety and immune-modulating effects of therapeutic monoclonal antibodies to interleukin-6 and tumor necrosis factor- α in cynomolgus macaques, *J. Immunotoxicol.* 1 (2005) 131–139.
- J.C. Byrd, S. O'Brien, I.W. Flinn, T.J. Kipps, M. Weiss, K. Rai, T.S. Lin, J. Woodworth, D. Wynne, J. Reid, A. Molina, B. Leigh, S. Harris, Phase 1 study of lymfloximab with detailed pharmacokinetic and pharmacodynamic measurements in patients with relapsed or refractory chronic lymphocytic leukemia, *Clin. Cancer Res.* 13 (2007) 4448–4455.
- G.C. Jayson, C. Mulatero, M. Ranson, J. Zweit, A. Jackson, L. Broughton, J. Wagstaff, L. Hakansson, G. Groenewegen, J. Lawrance, M. Tang, L. Wauk, D. Levitt, S. Marreaud, F. F. Lehmann, M. Herold, H. Zwierzina, R. European Organisation for. C. Treatment of, Phase I investigation of recombinant anti-human vascular endothelial growth factor antibody in patients with advanced cancer, *Eur. J. Cancer* 41 (2005) 555–563.
- T. Olafsen, Fc engineering: serum half-life modulation through FcRn binding, *Methods Mol. Biol.* 907 (2012) 537–556.
- C. Vaccaro, J. Zhou, R.J. Ober, E.S. Ward, Engineering the Fc region of immunoglobulin G to modulate in vivo antibody levels, *Nat. Biotechnol.* 23 (2005) 1283–1288.
- R.J. Kurlander, D.M. Ellison, J. Hall, The blockade of Fc receptor-mediated clearance of immune complexes in vivo by a monoclonal antibody (2A2C2) directed against Fc receptors on murine leukocytes, *J. Immunol.* 133 (1984) 855–862.
- L.P. Ganesan, J. Kim, Y. Wu, S. Mohanty, G.S. Phillips, D.J. Birmingham, J.M. Robinson, C.L. Anderson, Fc γ RIIb on liver sinusoidal endothelium clears small immune complexes, *J. Immunol.* 189 (2012) 4981–4988.
- C.Y. Zhang, J.W. Booth, Divergent intracellular sorting of Fc γ RIIA and Fc γ RIIb, *J. Biol. Chem.* 285 (2010) 34250–34258.
- T. Lehrnbecher, C.B. Foster, S. Zhu, S.F. Leitman, L.R. Goldin, K. Huppi, S.J. Chanock, Variant genotypes of the low-affinity Fc γ RIIb receptors in two control populations and a review of low-affinity Fc γ RIIb receptor polymorphisms in control and disease populations, *Blood* 94 (1999) 4220–4232.
- F. Mimoto, H. Katada, S. Kadono, T. Igawa, T. Kuramochi, M. Muraoka, Y. Wada, K. Haraya, T. Miyazaki, K. Hattori, Engineered antibody Fc variant with selectively enhanced Fc γ RIIb binding over both Fc γ RIIA (R131) and Fc γ RIIb (H131), *Protein Eng. Des. Sel.* 26 (2013) 589–598.
- K. Sato, M. Tsuchiya, J. Saldanha, Y. Koishihara, Y. Ohsugi, T. Kishimoto, M.M. Bendig, Reshaping a human antibody to inhibit the interleukin 6-dependent tumor cell growth, *Cancer Res.* 53 (1993) 851–856.
- D.E. Vaughn, P.J. Bjorkman, Structural basis of pH-dependent antibody binding by the neonatal Fc receptor, *Structure* 6 (1998) 63–73.
- S. Tickle, R. Adams, D. Brown, M. Griffiths, D. Lightwood, A. Lawson, High-throughput screening for high affinity antibodies, *JALA* 14 (2009) 303–307.
- M.L. Murtaugh, S.W. Fanning, T.M. Sharma, A.M. Terry, J.R. Horn, A combinatorial histidine scanning library approach to engineer highly pH-dependent protein switches, *Protein Sci.* 20 (2011) 1619–1631.
- J.V. Gerasimenko, A.V. Tepikin, O.H. Petersen, O.V. Gerasimenko, Calcium uptake via endocytosis with rapid release from acidifying endosomes, *Curr. Biol.* 8 (1998) 1335–1338.
- E.M. Strauch, S.J. Fleishman, D. Baker, Computational design of a pH-sensitive IgG binding protein, *Proc. Natl. Acad. Sci. U. S. A.* 111 (2014) 675–680.
- C.A. Sarkar, K. Lowenhaupt, T. Horan, T.C. Boone, B. Tidor, D.A. Lauffenburger, Rational cytokine design for increased lifetime and enhanced potency using pH-activated "histidine switching", *Nat. Biotechnol.* 20 (2002) 908–913.
- C.E. Mogensen, The glomerular permeability determined by dextran clearance using Sephadex gel filtration, *Scand. J. Clin. Lab. Invest.* 21 (1968) 77–82.
- H. Feinberg, D. Torgersen, K. Drickamer, W.I. Weis, Mechanism of pH-dependent N-acetylgalactosamine binding by a functional mimic of the hepatocyte asialoglycoprotein receptor, *J. Biol. Chem.* 275 (2000) 35176–35184.
- T. Yamamoto, H.C. Chen, E. Guigard, C.M. Kay, R.O. Ryan, Molecular studies of pH-dependent ligand interactions with the low-density lipoprotein receptor, *Biochemistry* 47 (2008) 11647–11652.

ORIGINAL ARTICLE

Anti-factor IXa/X bispecific antibody (ACE910): hemostatic potency against ongoing bleeds in a hemophilia A model and the possibility of routine supplementation

A. MUTO,* K. YOSHIHASHI,* M. TAKEDA,* T. KITAZAWA,* T. SOEDA,* T. IGAWA,* Y. SAKAMOTO,* K. HARAYA,* Y. KAWABE,* M. SHIMA,† A. YOSHIOKA‡ and K. HATTORI*
 *Research Division, Chugai Pharmaceutical Co., Ltd, Gotemba, Shizuoka; †Department of Pediatrics, Nara Medical University; and ‡Nara Medical University, Kashihara, Nara, Japan

To cite this article: Muto A, Yoshihashi K, Takeda M, Kitazawa T, Soeda T, Igawa T, Sakamoto Y, Haraya K, Kawabe Y, Shima M, Yoshioka A, Hattori K. Anti-factor IXa/X bispecific antibody (ACE910): hemostatic potency against ongoing bleeds in a hemophilia A model and the possibility of routine supplementation. *J Thromb Haemost* 2014; 12: 206–13.

Summary. *Background:* We previously reported that a humanized anti-factor IXa/X bispecific antibody, hBS23, mimics the function of FVIII even in the presence of FVIII inhibitors, and has preventive hemostatic activity against bleeding in an animal model of acquired hemophilia A. After further molecular engineering of hBS23, we recently identified an improved humanized bispecific antibody, ACE910, for clinical investigation. *Objectives:* To elucidate the *in vivo* hemostatic potency of ACE910 by examining its effect against ongoing bleeds, and to determine its pharmacokinetic parameters for discussion of its potency for prophylactic use. *Methods:* A non-human primate model of acquired hemophilia A was established by injecting anti-primate FVIII neutralizing antibody. When bleeds emerged following an artificial bleed-inducing procedure, either ACE910 or recombinant porcine FVIII (rpoFVIII) was intravenously administered. rpoFVIII was additionally administered twice daily on the following 2 days. Bleeding symptoms were monitored for 3 days. A pharmacokinetic study and multiple-dosing simulations of ACE910 were also performed. *Results:* A single bolus of 1 or 3 mg kg⁻¹ ACE910 showed hemostatic activity comparable to that of 10 U kg⁻¹ (twice daily) rpoFVIII against ongoing bleeds. The determined

ACE910 pharmacokinetic parameters included a long half-life (3 weeks) and high subcutaneous bioavailability (nearly 100%). The simulation results based on pharmacokinetic parameters indicated that the above hemostatic level could be maintained with once-weekly subcutaneous administration of ACE910, suggesting the possibility of more effective prophylaxis. *Conclusions:* ACE910 may offer an alternative on-demand treatment option for patients with hemophilia A, as well as user-friendly and aggressive routine supplementation.

Keywords: antibodies, catalytic; factor VIII; hemophilia A; hemostasis; therapeutics.

Introduction

Hemophilia A is a bleeding disorder caused by an inherited deficiency of factor VIII. The severity is known to correlate with the plasma FVIII level: severe, moderate and mild phenotypes are defined by plasma FVIII levels of < 1, 1–5 and > 5 to < 40 U dL⁻¹, respectively. Severe cases have a high propensity to suffer bleeds, including joint bleeds, whereas moderate cases typically experience far fewer bleeding episodes, and mild cases rarely bleed spontaneously [1,2].

Patients are primarily treated with FVIII agents. However, as FVIII agents are exogenous for severely affected patients, ~ 30% of them develop alloantibodies against FVIII (FVIII inhibitors) [2]. FVIII inhibitors largely restrict treatment with FVIII agents, and consequently make it difficult to control bleeding, because alternative bypassing agents have shorter *in vivo* half-lives and are not always effective [3]. The eradication of FVIII inhibitors with high doses of FVIII is very expensive, and does not always work [4]. In patients with severe hemophilia A without FVIII inhibitors, routine prophylaxis with exoge-

Correspondence: Atsushi Muto, Research Division, Chugai Pharmaceutical Co., Ltd, 1-135 Komakado, Gotemba, Shizuoka 412-8513, Japan.

Tel.: +81 550 87 3411; fax: +81 550 87 1960.

E-mail: mutoats@chugai-pharm.co.jp

Institution where the work was carried out: Research Division, Chugai Pharmaceutical Co., Ltd.

Received 25 April 2013

Manuscript handled by: L. Aledort

Final decision: P. H. Reitsma, 27 November 2013

nous FVIII to maintain FVIII levels above 1 U dL^{-1} is beneficial to prevent bleeding [5,6]; however, the need for frequent intravenous injections negatively affects patients' quality of life [7].

In order to overcome these shortcomings, we previously created a humanized anti-FIXa/FX bispecific IgG antibody, termed hBS23, which replicated FVIII cofactor function by binding and placing FIXa and FX into spatially appropriate positions [8]. hBS23 had cofactor activity *in vitro*, even in the presence of FVIII inhibitors, and, in a non-human primate model of acquired hemophilia A, hBS23 at an intravenous dose of 0.3 mg kg^{-1} exerted hemostatic activity to prevent the progression of bleeding symptoms to the same extent as recombinant porcine FVIII (rpoFVIII) maintained at a plasma level of $\geq 1 \text{ U dL}^{-1}$. However, it remained unproven whether this bispecific antibody approach possessed the potency to ameliorate ongoing bleeds, which would require higher levels of FVIII, or how much hemostatic potency it had in comparison with FVIII. We recently modified hBS23 with a multidimensional optimization approach to improve the FVIII-mimetic cofactor activity, pharmacokinetic properties, immunogenicity, physicochemical stability, and ease of industrial manufacture for clinical application; we consequently identified an improved humanized anti-FIXa/FX bispecific IgG antibody, termed ACE910 [9]. In this study, we elucidate the *in vivo* hemostatic potency of ACE910, including that against ongoing bleeds as compared with rpoFVIII, by using a non-human primate model of acquired hemophilia A.

Furthermore, in order to elucidate the potency of ACE910 in routine supplementation, we performed a pharmacokinetic study of ACE910 in non-human primates to determine its pharmacokinetic parameters, and conducted multiple dosing simulations with those parameters. Routine supplementation with exogenous FVIII is aimed at keeping the FVIII level at 1 U dL^{-1} or above to convert a severe disease to a moderate one [2,5]. This strategy successfully reduces bleeding episodes and the risk of developing hemophilic arthropathy [5]. However, the effect is not necessarily perfect: a recent report suggested that the risk of joint damage remains until the baseline factor level is $10\text{--}15 \text{ U dL}^{-1}$ or higher [10]. In this study, we also discuss the possibility of once-weekly subcutaneous administration of ACE910 for routine supplementation that is more aggressive than the current one with exogenous FVIII; in other words, a regimen that can convert a severe disease to a mild disease.

Materials and methods

Materials

ACE910 was expressed in human embryonic kidney 293 or Chinese hamster ovary (CHO) cells, which were co-transfected with the mixture of plasmids encoding the

humanized anti-FIXa heavy chain, anti-FX heavy chain, and common light chain. ACE910 was purified by protein A and ion exchange chromatography from the culture supernatant [9]. B domain-deleted rpoFVIII [8] was prepared as described in the supporting information (Preparation and analysis of rpoFVIII). Briefly, we expressed it in CHO cells by stable transfection. Then, rpoFVIII was purified from the supernatant by using ion exchange and gel permeation chromatography. We confirmed its purity with SDS-PAGE under reduced conditions, and determined its activity (U dL^{-1}) with an activated partial thromboplastin time (APTT)-based one-stage coagulation assay. Recombinant human FVIII (rhFVIII) was purchased from Bayer HealthCare (Leverkusen, Germany). Anti-primate FVIII neutralizing antibody (VIII-2236) [8] and the other purchased reagents are described in the supporting information (Confirmation of non-reactivity of VIII-2236 to rpoFVIII and ACE910, and the other supporting methods).

Animals and ethics

We used 26 and 12 male cynomolgus monkeys for the *in vivo* hemostatic study (2.6–4.0 kg, aged 3 years; Hamri, Ibaraki, Japan) and for the pharmacokinetic study (2.9–5.0 kg, aged 4–5 years; Japan Laboratory Animals, Tokyo, Japan), respectively. We used 24 female mice (aged 5 weeks; Charles River, Yokohama, Japan, and SLC, Hamamatsu, Japan) and 24 female rats (aged 4 weeks; Charles River) for the immunization to generate anti-idiotypic antibodies against the variable region of ACE910. The details of anti-idiotypic antibodies are described in the supporting information (Generation and preparation of anti-idiotypic antibodies to each variable region of ACE910).

All animal studies were approved by the Institutional Animal Care and Use Committee of Chugai Pharmaceutical, and were conducted in accordance with the approved protocols and the Guidelines for the Care and Use of Laboratory Animals at the company. Chugai Pharmaceutical is fully accredited by the Association for Assessment and Accreditation of Laboratory Animal Care International.

APTT and thrombin generation (TG) assays

APTT and TG assays were performed with standard equipment. In the TG assay, we employed two kinds of triggering solutions that contained, respectively, FXIa and tissue factor (TF). The solution containing human FXIa (Enzyme Research Laboratories, Swansea, UK) was prepared in-house, and the solution containing TF, PPP-Reagent LOW (Thrombinoscope BV, Maastricht, the Netherlands) was purchased. The details are described in the supporting information (APTT and TG assays).

In vivo hemostatic study in an acquired hemophilia A model

As ACE910 is highly species-specific in its FVIII-mimetic cofactor activity, non-human primates were used. On day 0, the animals received an intravenous injection of VIII-2236 (10 mg kg⁻¹). Two hours later, the animals were anesthetized with isoflurane inhalation, and bruises on the body surface that might possibly have emerged because of FVIII neutralization were measured. Then, the following two surgical procedures were performed: a 18G-needle was inserted 1 cm deep into the muscle at 22 sites (two sites in each upper arm, three sites in each forearm, four sites on the inside of each thigh, and two sites on the outside of each thigh); and subcutaneous exfoliation was performed by inserting the tip of forceps beneath the abdominal skin to 3 cm at two sites. After administration of buprenorphine, an analgesic drug, the animals were allowed to recover from the anesthesia. (They received this analgesic treatment twice daily [morning and evening] from day 0 to day 2; six doses were given, and the condition of the animals was observed daily.) After recognition of bleeding, approximately 6–8 h after injury, the animals received intravenous ACE910 (0.3, 1 or 3 mg kg⁻¹; *n* = 4 for each group), rpoFVIII (3.4 or 10 U kg⁻¹; *n* = 4 for each group), or no test item (control; *n* = 6). In the rpoFVIII group, rpoFVIII was also intravenously administered in the morning and evening on days 1 and 2 (a total of five administrations). In the morning on days 1, 2, and 3, the animals were anesthetized for measurement of the bruised areas. After the evaluation on day 3, the animals were killed humanely. Citrated blood was collected before and 2 h after the VIII-2236 injection, ~ 10 min after the test item administration on day 0, and before measurement of the bruised area on days 1, 2, and 3. The change in blood hemoglobin level was expressed as a percentage of that on day 0 (2 h after the VIII-2236 injection). The plasma ACE910 concentration was determined with the method described in the supporting information (Measurement of plasma ACE910 concentration).

Pharmacokinetic study

Animals received intravenous ACE910 (6 mg kg⁻¹; *n* = 3) or subcutaneous ACE910 (0.06, 0.6 or 6 mg kg⁻¹; *n* = 3 for each group) in a single dose. For animals dosed intravenously, blood was sampled with a heparinized syringe at 0.25, 2, 8, 24, 48, 72 and 96 h postdose, as well as at 7, 14, 28, 42, 56, 70 and 84 days postdose. For animals dosed subcutaneously, blood was collected in the same way without sampling at 0.25 h. The plasma concentrations of ACE910 and anti-ACE910 alloantibodies were measured with the methods described in the supporting information (Measurement of plasma ACE910 concentration and detection of anti-ACE910 alloantibodies).

Statistical analysis, pharmacokinetic analysis, and multiple-dosing simulation

For the *in vivo* hemostatic study, data are presented as mean ± standard error. Other data are presented as mean ± standard deviation. The parametric Dunnett multiple comparison test (two-tailed) (SAS preclinical package version 5.00; SAS Institute Japan, Tokyo, Japan) was used to determine *P*-values. *P* < 0.05 was considered to be statistically significant.

In the pharmacokinetic study, the plasma ACE910 concentration data were analyzed by non-compartmental analysis with PHOENIX WINNONLIN version 6.2 (Pharsight, St Louis, MO, USA). Multiple-dosing simulations were performed with SAAM II version 1.2 (SAAM Institute, Seattle, WA, USA).

Results*In vitro cross-reactivity of ACE910 with cynomolgus monkeys*

First, we examined the cross-reactivity of ACE910 with cynomolgus monkeys by use of an APTT assay. ACE910 shortened APTT in FVIII-neutralized cynomolgus monkey plasma with a concentration dependency similar to that in human FVIII-deficient plasma (Fig. 1A,B).

We next examined the cross-reactivity of ACE910 with cynomolgus monkeys by use of one of the global assays, the TG assay [11]. Because we had not clearly detected rpoFVIII activity in the standard low-TF triggering condition in FVIII-neutralized cynomolgus monkey plasma, we instead employed FXIa as a trigger. Beforehand, we had confirmed, in human FVIII-deficient plasma, that the two triggering conditions produced similar peak heights for the purpose of comparing ACE910 with rhFVIII (Figs 1C and S1). ACE910 improved the peak height of the FXIa-triggered TG assay in FVIII-neutralized cynomolgus monkey plasma in a similar concentration-dependent manner to that in human FVIII-deficient plasma (Fig. 1C,D). ACE910 had a similar cofactor activity as rhFVIII or rpoFVIII in improving the peak height in each species. Beforehand, the rpoFVIII that we prepared had been analyzed for qualification (Fig. S2).

In vivo hemostatic study in an acquired hemophilia A model

To examine the *in vivo* hemostatic potency of ACE910, including that against ongoing bleeds, we modified the non-human primate model of acquired hemophilia A that we previously reported [8]. Briefly, more intense injury procedures were employed, and the dose timing of the test item was set to after the emergence of bleeding symptoms, so that the hemostatic action of 10 U kg⁻¹ rpoFVIII could be properly evaluated.