- Wiklund O, Agewall S, Alegria E, Chapman MJ, Durrington P, Erdine S, Halcox J, Hobbs R, Kjekshus J, Filardi PP, Riccardi G, Storey RF, Wood D; ESC Committee for Practice Guidelines (CPG) 2008–2010 and 2010–2012 Committees. ESC/EAS Guidelines for the management of dyslipidaemias: the Task Force for the Management of Dyslipidaemias of the European Society of Cardiology (ESC) and the European Atherosclerosis Society (EAS). *Eur Heart J* 2011; 32:1769–1818.
- 22. National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). Third report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report. Circulation 2002; 106:3143–3421.
- American Diabetes Association. Executive summary: standards of medical care in diabetes—2013. *Diabetes Care* 2013; 36:S4–S10.
- American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2013; 36:S67–S74.
- Walker AE, Robins M, Weinfeld FD. The National Survey of Stroke. Clinical Findings. Stroke 1981; 12:113–144.
- World Health Organization. Document for Meeting of MONICA Principal Investigators. WHO: Geneva, Switzerland, 1986.
- Woodward M. Epidemiology: Study Design and Data Analysis. 2nd ed. Chapman & Hall/CRC Press, Boca Raton, FL, 2005.

- Tanaka T, Okamura T. Blood cholesterol level and risk of stroke in community-based or worksite cohort studies: a review of Japanese cohort studies in the past 20 years. Keio J Med 2012; 61:79–88.
- 29. Amarenco P, Steg PG. The paradox of cholesterol and stroke. *Lancet* 2007; 370:1803–1804.
- 30. Kushiro T, Mizuno K, Nakaya N, Ohashi Y, Tajima N, Teramoto T, Uchiyama S, Nakamura H; Management of Elevated Cholesterol in the Primary Prevention Group of Adult Japanese Study Group. Pravastatin for cardiovascular event primary prevention in patients with mild-to-moderate hypertension in the Management of Elevated Cholesterol in the Primary Prevention Group of Adult Japanese (MEGA) Study. Hypertension 2009; 53:135–141.
- 31. Sever PS, Dahlöf B, Poulter NR, Wedel H, Beevers G, Caulfield M, Collins R, Kjeldsen SE, Kristinsson A, McInnes GT, Mehlsen J, Nieminen M, O'Brien E, Ostergren J; ASCOT investigators. Prevention of coronary and stroke events with atorvastatin in hypertensive patients who have average or lower-than-average cholesterol concentrations, in the Anglo-Scandinavian Cardiac Outcomes Trial-Lipid Lowering Arm (ASCOT-LLA): a multicentre randomised controlled trial. *Lancet* 2003; 361:1149–1158.
- MacMahon S, Peto R, Cutler J, Collins R, Sorlie P, Neaton J, Abbott R, Godwin J, Dyer A, Stamler J. Blood pressure, stroke, and coronary heart disease. Part 1, prolonged differences in blood pressure: prospective observational studies corrected for the regression dilution bias. *Lancet* 1990; 335:765–774.

