

細胞株より発現している MHC class I 遺伝子をクローニングしてクローンをとり、それぞれ塩基配列を決定した。各個体の B 細胞株からそれぞれ A ローカス、B ローカス 20 数個のクローンを取りシーケンスしたが、解析できたのが、表の数字である。それをそれぞれカニクイザル MHC のデータベースを検索して対応するクローンを同定した。取れてくる大腸菌のクローンと発現しているクローンが同じ(クローニング効率がどれも同じ)と仮定すると単一なものしか発現していないもの、2クローンが発現しているもの主要なものが 1 個でマイナーなものが発現しているものなど、個体によりまちまちであった。B 細胞株より発現している MHC class I A ローカス、B ローカスは、複数発現しており、これらの塩基配列を決定し、クラスタ解析を行った。SIV ウィルス感染抵抗性個体と感染感受性個体との MHC クラス I の特異的なタイプは、個体の n 数が少ないのでできなかった。ただし近縁のアカゲザルの報告から抵抗性を付与すると考えられる分子の発現がみられる個体(#018)があった。また A ローカスがホモであると思われる個体が3頭 (#022, #053, #053) いた。B ローカスについて、ホモと思われる個体はいなかった。

D. 考察

霊長類の実験動物として最もポピュラーであるカニクイザルについて、MHC のタイピングは非常に重要な情報であるが個体で発現している MHC の情報はクローンの塩基配列情報の登録は多くなされているが、その MHC 分子の発現により知られている感染症ウィルス感染に対して感受性かあるいは抵

抗性かは報告がない。これについては多くの個体で発現している MHC を解析し感染抵抗性の固体の数を増やして解析する必要がある。そしてデータを蓄積する必要がある。ヒトの MHC については、データベースが完備しルミネックス法で蛍光ビーズを使用し短時間で調べられる。ヒトとサルではニーズの要求も異なるので比較はできないが、できるだけ簡便で正確に同定できる方法の開発が待たれる。

またアカゲザルで感染抵抗性を付与すると報告のある Mamu-B*07:04 と相同性のある、Mafa-B*017:01 の発現が確認された。このようなデータの蓄積がヒトの医療への貢献ができると考えられるので、地道にデータを積み重ねていく努力は必要である。

E. 結論

MHC class I 抗原は感染症の抵抗性を付与することが知られており、あらかじめ抵抗性を付与する MHC class I 分子の発現を調べることができたとしたら実験動物としてのカニクイザル資源の高度化に役立つと考えられる。ただし MHC 分子の発現のデータの蓄積と解析が必要であると考えられる。

F. 健康危険情報

特になし。

G. 研究発表

なし。

H. 知的財産権の出願・登録状況 (予定を含む。)

なし。

個体	A locus		B locus	
#016	Mafa-A1*002:01:01	5	Mafa-B*121:01	3
	Mafa-A1*003:03	3	Mafa-B*136:02	2
			Mafa-B*007:01:01	2
			Mafa-B11L*01:04	2
			Mafa-B11L*01:01	1
			Mafa-B*050:01	1
			Mafa-B*050:04	1
#018	Mafa-A1*098:02	2	Mafa-B*017:01	4
	Mafa-A2*24:03	2	Mafa-B*007:01:02	3
	Mafa-A1*100:01	1		
#022	Mafa-A1*089:01	14	Mafa-B*104:01:02	3
			Mafa-B*073:01	2
			Mafa-B*064:01	1
			Mafa-B*144:02	1
#053	Mafa-A1*089:03	8	Mafa-B*099:01	3
			Mafa-B*114:01	1
#054	Mafa-A1*052:02	6	Mafa-B*068:02	3
	Mafa-A1*100:01	2	Mafa-B*037:02	2
			Mafa-B*033:02	2
			Mafa-B*068:03	1
#055	Mafa-A1*100:01	6	Mafa-B*108:01	3
			Mafa-B11L*01:04	2
			Mafa-B*007:01:02	1
			Mafa-B*033:03	1
#056	Mafa-A1*010:02:01	7	Mafa-B*058:01	5
	Mafa-A1*068:01	2	Mafa-B*153:01	3
	Mafa-A4*14:04	2	Mafa-B*043:01	2
#057	Mafa-A1*097:01	4	Mafa-B*007:01:01	4
	Mafa-A1*003:01	2	Mafa-B*121:01	3
			Mafa-B*011:01	1
			Mafa-B*004:03	1

図： MHC class I 遺伝子 A ローカス、B ローカスの発現パターン

研究成果の刊行に関する一覧表

(平成26年度)

研究成果の刊行に関する一覧表

書籍

著者氏名	論文タイトル名	書籍全体の編集者名	書 籍 名	出版社名	出版地	出版年	ページ
<u>Shimozawa Nobuhiro</u>	Cynomolgus monkey induced pluripotent stem cells generated by using allogeneic genes.	Andras Nagy	iPS cells: Generation characterization and differentiation –Methods and Protocols. Methods in Molecular Biology	Springer	Deutschland	2014	DOI 10.1007/7651_2014_137

雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Okabayashi S, <u>Shimozawa N</u> , <u>Yasutomi Y</u> , Yanagisawa K, Kimura N.	Diabetes mellitus accelerates A β pathology in brain accompanied by enhanced GA β generation in nonhuman primates.	PLoS ONE	10	e0117362	2015
Onishi M, Ozasa K, Kobiyama K, Ohata K, Kitano M, Taniguchi K, Homma T, Kobayashi M, Sato A, Katakai Y, <u>Yasutomi Y</u> , Wijaya E, Igarashi Y, Nakatsu N, Ise W, Inoue T, Yamada H, Vandenberg A, Standley DM, Kurosaki T, Coban C, Aoshi T, Kuroda E, Ishii KJ.	Hydroxypropyl- β -cyclodextrin spikes local inflammation that induce Th2 and Tfh responses to the coadministered antigen	J.Immunol			印刷中
Watanabe K., Matsubara A, Kawano M, Mizuno S, <u>Okamura T</u> , Tsujimura Y, Inada H, Noshika T, Matsuo K. and <u>Yasutomi Y</u> .	Recombinant Ag85B vaccine by taking advantage of characteristics of human parainfluenza type 2 virus vector showed Mycobacteria-specific immune responses by intranasal immunization.	Vaccine	32	1727-1735	2014
Kobiyama K., Aoshi T., Narita H., Kuroda E., Hayashi M., Tetsutani K., Koyama S., Mochizuki S., Sakurai K., Katakai Y., <u>Yasutomi Y</u> , Saijo S., Iwakura Y., Akira S., Coban C. and Ishii KJ.	A non-agonistic Dectin-1 ligand transforms CpG into a multitask nanoparticulate TLR9 agonist.	Proc. Natl. Acad. Sci. USA	111	3086-3091	2014

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Tsujimura Y, Inada H, Yoneda M, Fujita T, Matsuo K. and <u>Yasutomi Y.</u>	Effects of Mycobacteria major secretion protein, Ag85B, on allergic inflammation in the lung.	Plos One	E-pub		2014
Saito N, Chono H, Shibata H, <u>Ageyama N,</u> <u>Yasutomi Y,</u> Mineno J.	CD4(+) T Cells Modified by the Endoribonuclease MazF Are Safe and Can Persist in SHIV-infected Rhesus Macaques.	Mol Ther Nucleic Acids.	3	e168	2014
Machino-Ohtsuka T, Tajiri K, Kimura T, Sasaki S, Sato A, Yoshida T, Hiroe M, <u>Yasutomi Y,</u> Aonuma K, Imataka-Yoshida K.	Tenascin-C aggravates autoimmune myocarditis via dendritic cell activation and Th17 cell differentiation.	J.Am.Heart Assoc.	E-pub		2014
Tachibana SI, Kawai S, Katakai Y, Takahashi H, Nakade T, <u>Yasutomi Y,</u> Horii T, Tanabe K.	Contrasting infection susceptibility of the Japanese macaques and cynomolgus macaques to closely related malaria parasites, Plasmodium vivax and Plasmodium cynomolgi.	Parasitol. Int.	E-pub		2014
Fukuyama Y., Yuki Y., Katakai Y., Harada N., Takahashi H., Takeda S., Mjima M., Joo S., Kurokawa S., Sawada S., Shibata H., Park EJ., Fujihashi K., Briles DE., <u>Yasutomi Y.</u> , Tsukada H., Akiyoshi K. and Kiyono H	Nanogel-based pneumococcal surface protein A nasal vaccine induces microRNA-associated Th17 cell responses with neutralizing antibodies against Streptococcus pneumoniae in macaques.	Mucosal Immunology	E-pub		2015

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
L. Yasmin, J. Takano, Y. Nagai, J. Otsuki, <u>T. Sankai</u>	Detecting and quantifying male-specific fetal DNA in the serum of pregnant cynomolgus monkeys (<i>Macaca fascicularis</i>)	Com. Med.	65	70-76	2015
K. Honsho, M. Hirose, M. Hatori, L. Yasmin, H. Izu, S. Matoba, S. Togayachi, H. Miyoshi, <u>T. Sankai</u> , A. Ogura, A. Honda	Naïve-like conversion enhances the difference in innate <i>in vitro</i> differentiation capacity between rabbit ES cells and iPS cells	J. Reprod. Dev.	61	13-19	2015
A. Honda, M. Hirose, <u>T. Sankai</u> , L. Yasmin, K. Yuzawa, K. Honsho, H. Izu, A. Iguchi, M. Ikawa, A. Ogura	Single-step generation of rabbits carrying a targeted allele of <i>Tyrosinase</i> gene using CRISPR/Cas9	Exp.Anim.	64	31-37	2015
M. Hatori, <u>N. Shimozawa</u> , L. Yasmin, H. Suemori, N. Nakatsuji, A. Ogura, K. Yagami, <u>T. Sankai</u>	Role of retinoic acid and fibroblast growth factor 2 in neural differentiation from cynomolgus monkey (<i>Macaca fascicularis</i>) embryonic stem cells	Comp. Med.	64	1-8	2014

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Kumagai, A., Fujita, A., Yokoyama, T., Nonobe, Y., Hasaba, Y., Sasaki, T., Itoh, Y., Koura, M., <u>Suzuki, O.</u> , Adachi, S., Ryo, H., Kohara, A., Tripathi, L., Sanosaka, M., Fukushima, T., Takahashi, H., Kitagawa, K., Nagaoka, Y., Kawahara, H., Mizuguchi, K., Nomura, T., Matsuda, J., Tabata, T., and Takemori, H.	Altered Actions of Memantine and NMDA- Induced Currents in a New Grid2-Deleted Mouse Line.	Genes	5(4)	1095-1114	2014
鈴木治, 松田潤一郎	心筋症のマウスモデル： 拡張型心筋症マウスモデル の心機能解析と遺伝子 解析	医学のあゆみ	252 (10)	1027-1031	2015

RESEARCH ARTICLE

Diabetes Mellitus Accelerates A β Pathology in Brain Accompanied by Enhanced GA β Generation in Nonhuman Primates

Sachi Okabayashi^{1,2}, Nobuhiro Shimozawa¹, Yasuhiro Yasutomi¹, Katsuhiko Yanagisawa³, Nobuyuki Kimura^{1,3*}

1 Tsukuba Primate Research Center, National Institute of Biomedical Innovation, 1–1 Hachimandai, Tsukuba-shi, Ibaraki, 305–0843, Japan, **2** The Corporation for Production and Research of Laboratory Primates, 1–1 Hachimandai, Tsukuba-shi, Ibaraki, 305–0843, Japan, **3** Section of Cell Biology and Pathology, Department of Alzheimer's Disease Research, Center for Development of Advanced Medicine for Dementia, National Center for Geriatrics and Gerontology (NCGG), Gengo 35, Morioka, Obu, Aichi, 474–8511, Japan

* kimura@ncgg.go.jp



OPEN ACCESS

Citation: Okabayashi S, Shimozawa N, Yasutomi Y, Yanagisawa K, Kimura N (2015) Diabetes Mellitus Accelerates A β Pathology in Brain Accompanied by Enhanced GA β Generation in Nonhuman Primates. PLoS ONE 10(2): e0117362. doi:10.1371/journal.pone.0117362

Academic Editor: Yoshitaka Nagai, National Center of Neurology and Psychiatry, JAPAN

Received: July 20, 2014

Accepted: December 21, 2014

Published: February 12, 2015

Copyright: © 2015 Okabayashi et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This study was supported by The Research Funding for Longevity Sciences (25-20) from National Center for Geriatrics and Gerontology (NCGG), Japan. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Abstract

Growing evidence suggests that diabetes mellitus (DM) is one of the strongest risk factors for developing Alzheimer's disease (AD). However, it remains unclear why DM accelerates AD pathology. In cynomolgus monkeys older than 25 years, senile plaques (SPs) are spontaneously and consistently observed in their brains, and neurofibrillary tangles are present at 32 years of age and older. In laboratory-housed monkeys, obesity is occasionally observed and frequently leads to development of type 2 DM. In the present study, we performed histopathological and biochemical analyses of brain tissue in cynomolgus monkeys with type 2 DM to clarify the relationship between DM and AD pathology. Here, we provide the evidence that DM accelerates A β pathology *in vivo* in nonhuman primates who had not undergone any genetic manipulation. In DM-affected monkey brains, SPs were observed in frontal and temporal lobe cortices, even in monkeys younger than 20 years. Biochemical analyses of brain revealed that the amount of GM1-ganglioside-bound A β (GA β)—the endogenous seed for A β fibril formation in the brain—was clearly elevated in DM-affected monkeys. Furthermore, the level of Rab GTPases was also significantly increased in the brains of adult monkeys with DM, almost to the same levels as in aged monkeys. Intraneuronal accumulation of enlarged endosomes was also observed in DM-affected monkeys, suggesting that exacerbated endocytic disturbance may underlie the acceleration of A β pathology due to DM.

Introduction

Alzheimer's disease (AD) is a progressive neurological disorder that is histopathologically characterized by the formation of senile plaques (SPs) and neurofibrillary tangles (NFTs) [1, 2]. It is widely accepted that β -amyloid protein (A β), the major component of SPs, is a key molecule

underlying AD pathogenesis [3, 4]. Several epidemiological/clinical studies have shown that diabetic mellitus (DM) patients are significantly more likely to develop cognitive dysfunction and exhibit increased susceptibility to AD [5–9], in consistent with the original Rotterdam study [10]. Recent findings also showed that there are several pathogenic connections between AD and DM patient brains, such as brain inflammation, mitochondrial dysfunction, and defective neuronal insulin signaling [11]. Insulin resistance causes alteration in GSK3 β kinase signaling pathway as observed in AD brains, and it is also associated with an AD-like pattern of reduced cerebral glucose metabolic rate in brain [12, 13]. Moreover, accumulating evidences showed that the experimental induction of DM enhanced AD pathology even in rodents [14–25]. However, it remains unclear how DM physiologically accelerates AD pathology in the brain.

With advancing age, both SPs and NFTs occur spontaneously in brains of cynomolgus monkeys [26, 27]. In addition, the amino acid sequence of A β of cynomolgus monkeys is completely consistent with that of humans [28]. These advantages make this species a useful model to study age-dependent AD pathophysiology. As with humans, obesity occasionally occurs in adult, middle-aged monkeys, and it can result in the development of type 2 DM [29, 30]. Similar to the case of humans, these monkeys have a period of insulin resistance and hyperinsulinemia before developing overt DM, which is then accompanied by deficiency in pancreatic insulin production [29–31]. The pathological changes that occur in the pancreatic islets of aged monkeys are also similar to those seen in human diabetics, including the deposition of islet amyloid polypeptide (IAPP) [29–31]. In addition, gestational diabetes has been also reported in female cynomolgus monkeys [29–31]. Thus, cynomolgus monkeys are a useful species to investigate not only age-dependent AD lesions but also the relationship between DM and AD pathology.

Here, we report that DM enhances the generation of GM1-ganglioside-bound A β (GA β) to accelerate SP deposition in cynomolgus monkey brains. GA β was previously identified as the endogenous seed for A β fibril formation in the brain, and its generation is enhanced by endocytic disturbance, which is considered to be involved in early-stage AD pathology [32–34]. In DM-affected adult monkeys, the level of Rab GTPases in the brain was obviously increased as compared to normal adult monkeys, and intraneuronal endosomes were apparently enlarged. These findings suggest that DM exacerbates age-dependent endocytic disturbance, which then may lead to accelerate A β pathology via enhanced GA β generation.

Materials and Methods

Animals

Forty-one cynomolgus monkey (*Macaca fascicularis*) brains were used in this study. Of these, six brains were from young monkeys (age: 6 and 7 years); six were from normal adult monkeys (age: 17, 18, 19, and 20 years); nine were from DM-affected adult monkeys (age: 17, 18, 19, and 20 years); ten were from normal aged monkeys (age: 24, 25, 26, and 28 years); and ten were from DM-affected aged monkeys (age: 24, 25, 26, and 28 years). The frontal and temporal lobes were used for immunohistochemical studies. The cerebral cortices of 12 monkeys were used for dot blot analyses. Of these 12, 3 were from young monkeys (age: 6 years [N = 2] and 7 years [N = 1]); 3 were from normal adult monkeys (age: 18 years, 19 years, and 20 years [N = 1 each]); 3 were from DM-affected adult monkeys (age: 18 years, 19 years, and 20 years [N = 1 each]); and 3 were from normal aged monkeys (age: 26 years [N = 2] and 28 years [N = 1]). The cerebral cortices of 12 female monkeys were used for Western blot analyses. Of these 12, 3 were from young monkeys (age: 7 years [N = 3]); 3 were from normal adult monkeys (age: 18 years, 19 years, and 20 years [N = 1 each]); 3 were from DM-affected adult monkeys (age: 18 years, 19 years, and 20 years [N = 1 each]); and 3 were from normal aged monkeys (age: 25 years and 26 years [N = 2]). The cerebral cortices of 14 female monkeys were used for

A β ELISA. Of these 12, 4 were from young monkeys (age: 7 years [N = 4]); 3 were from normal adult monkeys (age: 18 years, 19 years, and 20 years [N = 1 each]); 3 were from DM-affected adult monkeys (age: 18 years, 19 years, and 20 years [N = 1 each]); and 4 were from normal aged monkeys (age: 24 years [N = 3], and 25 years [N = 1]). All brains were obtained from the Tsukuba Primate Research Center (TPRC), National Institute of Biomedical Innovation (NIBIO), Japan. Monkeys in the TPRC are reared in individual cages (0.5 m wide \times 0.8 m high \times 0.9 m deep; stain-less steel mesh). The breeding rooms are rectangular, and the individual cages are installed on the long sides of the room. Each room contains at least 90 cages. Therefore, monkeys can always make visual, auditory and olfactory contact with their roommates. Ambient temperature in the rooms is kept about 25°C, and humidity is set at 50% to 60%. The air is replaced 12 times hourly. Monkeys are provided with 100 g of apples in the morning, and 70 g of commercial food (Type AS; Oriental Yeast Co., Ltd., Tokyo, Japan) is given to them twice in the afternoon. Water is available ad libitum. Every morning their health status (e.g., viability, appetite, fur-coat appearance) was monitored by experienced animal technicians. When any abnormality is found, a veterinarian examines the monkey promptly and applies the appropriate treatment. Moreover, the monkeys are medically examined under anesthesia (ketamine hydrochloride) at least once every 2 y. The medical examination consists of body weight measurement, tuberculin test, blood sample, stool test, examination of the fundus, and a medicated bath. The maintenance of animals was conducted according to the rules for animal care of the TPRC at NIBIO for the care, use, and biohazard countermeasures of laboratory animals [35]. This study was carried out in strict accordance with the recommendations in the Animal Care and Use Committee of the NIBIO, Japan. The protocol was approved by the Committee on the Ethics of Animal Experiments of the NIBIO (DS17-001R1). When the monkey presents clinical symptoms by injury or illness and it could not expect recovery from pain or morbidity, it is judged as poor prognosis. In the present study, the animals used in this study died of natural causes were euthanized when they reached endpoints determined as poor prognosis. All DM-affected monkeys were also sacrificed because of poor prognosis. For euthanasia, the monkeys were deeply anesthetized with a lethal dose of pentobarbital, and all efforts were made to minimize suffering.

Antibodies

In this study, we used the following antibodies: mouse monoclonal anti-dynein heavy chain (DHC) (Sigma, Saint Louis, MO); mouse monoclonal anti-dynein intermediate chain (DIC) (Millipore, Temecula, CA); mouse monoclonal anti- GA β antibody (4396C) (22); mouse monoclonal kinesin light chain (KLC) (Santa Cruz Biotechnology, Santa Cruz, CA); mouse monoclonal anti-nephrilysin (NEP; DAKO, Glostrup, Denmark); mouse monoclonal anti-phosphorylated tau antibody (AT8; Innogenetics, Gent, Belgium); mouse monoclonal anti-Rab5 antibody (Rab5m; Santa Cruz Biotechnology); rabbit polyclonal anti-A β antibody (IBL); rabbit polyclonal anti-full-length β -amyloid precursor protein (APP) antibody (IBL); rabbit polyclonal anti-Cathepsin D (CatD; Cell Signaling Technology, Danvers, MA); rabbit polyclonal anti-kinesin heavy chain (KHC) (Sigma); rabbit polyclonal anti-LC3 (Novus Biologicals, Littleton, CO); rabbit polyclonal anti-Rab5 antibody (Santa Cruz Biotechnology); rabbit polyclonal anti-Rab7 antibody (Sigma); rabbit polyclonal anti-Rab11 antibody (Santa Cruz Biotechnology); and rabbit polyclonal anti-synaptophysin (DAKO).

Histopathological analyses for DM-associated pathology

Most of the tissues were fixed in 10% neutral buffered formalin, processed routinely and stained with hematoxylin and eosin (HE) for histopathological analyses. The pancreases were also stained with direct fast scarlet (DFS; Muto) for identification of islet amyloid.

Immunohistochemistry

Brain samples were immersion-fixed in 10% neutral buffered formalin, embedded in paraffin, and cut into 4 μ m-thick sections. For immunohistochemical analyses with anti-GA β antibody, brain samples were fixed in 4% paraformaldehyde. Sections were deparaffinized by pretreatment with 0.5% periodic acid, and then incubated overnight at 4°C free floating in the following primary antibody solutions: (1) A β (1:100); (2) APP (1:100); (3) Rab5 (1:100); (4) GA β (1:20); or (5) AT8 (1:100). Following brief washes with buffer, the sections were sequentially incubated with biotinylated goat anti-mouse IgG (1:200) or goat anti-rabbit IgG (1:200), followed by streptavidin-biotin-horseradish peroxidase complex (DAKO). Immunoreactive elements were visualized by treating the sections with 3–3' diaminobenzidine tetrahydrochloride (Dojin Kagaku). The sections were then counterstained with hematoxylin. The immunoreactivity of A β , GA β , APP or Rab5 in a given cortical area was quantified with computer software Image J 1.49i (National Institute of Health). For quantification analyses, the sections were not counterstained with hematoxylin.

Biochemical analyses of monkey brains

Frozen monkey brain tissue (wet weight 0.2 g) was homogenized in a glass homogenizer with 4 ml of homogenate buffer solution (0.32 M sucrose, 10 mM Tris-HCl [pH 7.6], 0.25 mM PMSF, and 1 mM EDTA, and Complete Mini proteinase inhibitor cocktail), and then centrifuged at 100,000 \times g for 20 min to obtain the supernatant fraction. The proteins in the fraction were subjected to dot blot analyses. Brain homogenates were also centrifuged at 1,000 \times g for 10 min to remove the nuclear fraction, and then the supernatant was centrifuged at 105,000 \times g for 60 min to obtain the microsomal fraction. The proteins in the microsomal fraction were subjected to Western blot analyses. For A β ELISA, brain homogenates were incubated with Triton-X100 at a final concentration of 1% for 15 min at 37°C, and spun at 100,000 \times g for 20 min at 20°C. The pellet was homogenized in homogenate buffer containing 1% sarkosyl, incubated for 15 min at 37°C, and spun at 100,000 \times g for 20 min at 20°C. The sarkosyl-insoluble pellet was sonicated in 70% folic acid, cleared by centrifugation at 100,000 \times g for 20 min at 20°C. The supernatant was evaporated, and then resuspended in dimethyl sulfoxide (Sigma).

Dot blot analyses

Dot blot analyses were performed to assess age- and DM-related changes in GA β formation. Brain samples were adjusted to 1 mg, 2.5 mg, and 5 mg, and then applied onto nitrocellulose membranes and dried. The membranes were blocked with 5% nonfat dried milk in 20 mM PBS (pH 7.0) and 0.1% Tween-20 for 1 h at room temperature, and then incubated in the anti-GA β antibody solution overnight at 4°C. They were then incubated with horseradish peroxidase-conjugated goat anti-mouse IgG (1:10000; Cell Signaling Technology) for 1 h at room temperature. Immunoreactive elements were visualized using enhanced chemiluminescence (Luminata Forte Western HRP Substrate, Millipore).

Western blot analyses

Western blot analyses were performed to assess age- and DM-related changes in APP, Rab5, Rab7, Rab11, DHC, DIC, KHC, KLC, NEP, Cathepsin D heavy chain (CatD HC), and LC3 expression. Each microsome fraction prepared as described above was adjusted to 5 mg, and then analyzed using SDS-polyacrylamide gel electrophoresis using 14% (for LC3) or 12.5% (other proteins) acrylamide gels. Separated proteins were blotted onto polyvinylidene fluoride membranes (Immobilon P; Millipore). The membranes were blocked with 5% nonfat dried milk in

20 mM PBS (pH 7.0) and 0.1% Tween-20 for 1 h at room temperature, and then incubated overnight at 4°C in the following primary antibody solutions: (1) synaptophysin (1:20000); (2) APP (1:2000); (3) Rab5m (1:2000); (4) Rab7 (1:10000); (5) Rab11 (1:2000); (6) DHC (1:1000); (7) DIC (1:20,000); (8) KHC (1:5,000); (9) KLC (1:1000); (10) NEP (1:4,000); (11) CatD (1:2,000); or (12) LC3 (1:2,000). They were then incubated with either horseradish peroxidase-conjugated goat anti-mouse IgG or goat anti-rabbit IgG (1:10,000; Cell Signaling Technology) for 1 h at room temperature. Immunoreactive elements were visualized using enhanced chemiluminescence. To confirm reproducibility, immunoreactive bands obtained from the Western blots were quantified using commercially available software (Quantity One; PDI, Inc.).

A β ELISA

Total A β levels in young monkey, normal adult monkey, DM-affected monkey, and normal aged monkey brains were determined using a sandwich ELISA. The kit for A β (1-X) was obtained from IBL (Gunma, Japan). The ELISA assay was carried out according to the instruction manual. All samples were measured in duplicate.

Data analyses

Data obtained from quantitative analyses of immunohistochemistry and biochemical analyses are shown as means \pm SD. For statistical analyses, one-way ANOVAs were performed, followed by the Fisher's post hoc test.

Results

Clinical background and DM-associated pathology in cynomolgus monkeys

Tsukuba Primate Research Center (TPRC) maintains a large breeding and rearing colony of cynomolgus monkeys for high-quality production of nonhuman primate models and biomedical investigations. In the TPRC colony, some adult monkeys are spontaneously affected with type 2 DM for various reasons, such as pregnancy history and environmental factors. The clinical background of all monkeys used for this study is shown in (S1 Table). TPRC has accumulated clinical data for more than 40 years. Based on these data, the normal blood glucose level for female monkeys is in the range of 24 to 74 mg/dL, and for male monkeys the range is 24 to 76 mg/dL. Normal blood triglyceride levels are in the range of 8 to 85 mg/dL for females, and 6 to 52 mg/dL for males. The blood glucose and triglyceride levels of normal adult monkeys used in the present study were all in the normal range, and their body weights were in the range of historical TPRC data for normal monkeys (S1 Table). All DM-affected monkeys, on the other hand, exhibited hyperlipidemia and hyperglycemia (S1 Table). These DM-affected monkeys were severely obese; however, their body weight at acquisition of their brains for this study had decreased to about less than half the maximum body weights they attained during their lifetime (S1 Table).

Histopathologically, islet amyloid was found in the pancreases of all DM-affected adult monkeys (Fig. 1A). In eight DM cases, most hyalinized islets were replaced with severe amyloid deposits, with the remaining islet cells being severely degenerated or decreased in apparent number (Fig. 1A-C). The hyalinized islets with severe amyloid deposits stained positive for direct fast scarlet (DFS) (Fig. 1D). Furthermore, two DM cases had fatty degeneration of the liver, and one DM case had mild atheromatosis. Thus, histopathological analyses demonstrated that these adult monkeys clearly had DM. On the other hand, we did not observe apparent vascular lesions in all DM monkey brains.

DM accelerates A β pathology in cynomolgus monkey brain

Next, we conducted immunohistochemical analyses to assess whether DM affects AD pathology in these cynomolgus monkey brains. As previously reported [26, 36], we observed SP depositions in the brains of aged monkeys (Fig. 2A), but not in those of normal adult monkeys younger than 20 years of age (Fig. 2B-D). Strikingly, we apparently observed diffuse A β -immunopositive SPs in brains of six DM-affected adult monkeys younger than 20 years old, even though they were very small quantities as compared to aged monkey brains (Fig. 2E-I). In aged cynomolgus monkey brains, cerebral amyloid angiopathy (CAA) lesions are also observed, and NFTs are observed over 32-year-old monkey brains [26, 27]. Although abnormally phosphorylated tau accumulation was not observed, we found much severe CAA in the brains of aged monkeys with DM (Fig. 3).

DM enhances GA β generation in adult monkey brains

To assess whether DM affects A β level in monkey brains, we examined A β ELISA analyses. In aged monkey brains, A β level was significantly increased, suggesting that the increase of A β level correlates with age-dependent SP depositions (Fig. 4A). However, surprisingly, A β level was just slightly increased in DM-affected adult monkey brains as compared to normal adult monkey brains (Fig. 4A). We previously identified a unique A β species, called GA β , characterized by its binding to GM1 ganglioside; GA β was demonstrated in brain tissue along with early pathological changes characteristic of AD [32]. Accumulating evidence suggests that GA β accelerates A β fibril formation by acting as a seed molecule [37–40]. We have also confirmed age-dependent GA β generation in cynomolgus monkey brains [32, 41].

To test our hypothesis that DM enhances GA β generation in the brain, we performed additional immunohistochemical analyses using an anti-GA β -specific antibody. In the brains of normal adult monkeys, we observed little, if any, immunoreactivity with anti-GA β antibody (Fig. 4B). In contrast, in the brains of DM-affected adult monkeys, we frequently observed neurons clearly immunopositive for anti-GA β antibody (Fig. 4B). Quantitative analyses also confirmed that the immunoreactivity of GA β was significantly increased in DM-affected adult monkey brains (Fig. 4C).

For biochemical analyses, we carried out dot blot analyses on monkey brain samples. Because of its conformational features, GA β is not recognized by western blot analyses. Our dot blot analyses showed that GA β generation increased in an age-dependent manner in cynomolgus monkey brains (Fig. 4D). GA β immunoreactivity was much stronger in aged monkey brains, indicating that the amount of GA β generated generally parallels age-dependent SP deposition as well as A β level. In normal adult monkey brains, the amount of GA β generation was not much different from that in young monkey brains (Fig. 4D). It is noteworthy that GA β generation was significantly increased in DM-affected adult monkey brains (Fig. 4E).

DM exacerbates endocytic disturbance with significant increase of Rab GTPases

In the brains of early-stage AD patients, neuronal endocytic pathology, such as intracellular accumulation of abnormally enlarged endosomes, is frequently observed [42–44]. We previously demonstrated that aging causes endocytic pathology along with a significant increase in Rab GTPases, resulting in the intracellular accumulation of APP [45]. This endocytic disturbance was observed in brains from cynomolgus monkeys almost 10 years before SP deposition begins [46]. We also reported that GA β accumulates in enlarged endosomes of neurons in aged cynomolgus monkey brains [33]. Moreover, endocytic disturbance induces GA β generation [34].




Fig 1. Histopathology of adult monkeys with type 2 diabetes mellitus. (A) Hematoxylin-eosin (HE)-stained section of the pancreas from a 17-year-old cynomolgus monkey with type 2 diabetes mellitus (DM) (S1 Table, Number 1). Most of the islets were replaced by abundant amyloid deposits. (B) HE-stained section of the pancreas from an 18-year-old cynomolgus monkey with DM (S1 Table, Number 3). Most of the islets were replaced with severe amyloid deposits. (C) HE-stained section of the pancreas from a 19-year-old cynomolgus monkey with DM (S1 Table, Number 7) showing hyalinized islets. Very few

islet cells remain. (D) Direct fast scarlet-stained section of pancreas from an 18-year-old cynomolgus monkey with DM (S1 Table, Number 5). Hyalinized islets with severe amyloid deposition were positive for direct fast scarlet staining. (E) HE-stained section of the liver from an 18-year-old cynomolgus monkey with DM (S1 Table, Number 3). Marked fatty degeneration was observed in the liver. (F) HE-stained section of the aorta from an 18-year-old cynomolgus monkey with DM (S1 Table, Number 5). Mild atheromatosis with foam cells and sterol clefts was observed in the aorta. Scale bars for a-f, 100 μ m.

doi:10.1371/journal.pone.0117362.g001

Taken together, these findings suggest that endocytic disturbance is involved in age-dependent A β pathology. Thus, to assess whether DM enhances endocytic disturbance, we investigated endocytic pathology in monkey brains.

In the brains of normal adult monkeys, APP and Rab5-positive early endosomes were observed as small granules in neurons (Fig. 5A, B). By contrast, in the brains of DM-affected adult monkeys, the immunoreactivity of APP and Rab5 was significantly stronger and the immunopositive granules were larger (Fig. 5C, D). Quantitative analyses confirmed that both APP- and Rab5-immunopositive granules were significantly increased in DM-affected adult monkey brains as compared to normal adult monkey brains (Fig. 5E). Western blot analyses showed that Rab5, Rab7 (late endosome-associated GTPase), and Rab11 (recycling endosome-associated GTPase) were increased in aged monkey brains, corroborating that age-dependent endocytic disturbance does occur in monkey brains, as previously reported (Fig. 5F, G) [45]. The amount of APP was also significantly increased in aged monkey brains as previously reported (Fig. 5F, G) [45, 47]. In the brains of DM-affected adult monkeys, Rab GTPases and APP levels were apparently increased compared to the brains of normal adult monkeys, being almost same as in aged monkey brains (Fig. 5F, G).

DM affects cathepsin D level and autophagosome clearance

Endosome trafficking is mediated by axonal transport motor proteins [48], and a recent study showed that the experimental induction of DM alters axonal motor protein levels in rodent model [49]. In the present study, we did not find apparent differences in axonal motor protein levels between normal and DM-affected adult monkey brains (Fig. 6). Endocytic disturbance is also induced by the breakdown in lysosomal degradation [50]. In DM-affected adult monkey brains, the level of cathepsin D (CatD) heavy chain increased (Fig. 6). On the other hand, we observed the significant increase in autophagosome marker LC3-II level without any alterations in LC3-I level (Fig. 6). A recent study also showed that DM-associated down-regulation of insulin signals reduces the level of A β degrading enzymes such as neprilysin (NEP) [51]. However, we did not find a clear reduction tendency in NEP levels in the brains of DM-affected adult monkeys (Fig. 6).

Discussion

Here, we present the evidence that DM accelerates A β pathology in the brain parenchyma of nonhuman primates, which have not undergone any genetic manipulation. We demonstrated that DM does so by enhancing the generation of GA β , the endogenous seed for A β fibril formation in the brain. The brains of DM-affected adult monkeys contained robust endocytic pathology, such as a significant increase in Rab GTPases and intraneuronal accumulation of enlarged endosomes. Endocytic disturbance is a cellular pathological characteristic of neurons of AD patients and enhances GA β generation [33, 34]. Thus, our present findings suggest that DM exacerbates age-dependent endocytic disturbance, which in turn enhance GA β generation resulting in accelerated A β pathology.

Recent epidemiological/clinical studies suggest that DM is a major risk factor for developing AD [5–9]. However, the underlying mechanisms for this association remain unclear. Thus, in the present study, we performed histopathological and biochemical analyses using brains from




Fig 2. Senile plaques in the brains of adult monkeys with DM. Images of temporal lobe (TL) and frontal lobe (FL) sections from normal cynomolgus monkeys (A-D) and cynomolgus monkeys with DM (E-H). Sections were immunostained with anti-A β antibody and counterstained with hematoxylin. In aged monkey brains, we observed SPs immunostained with anti-A β antibody, as previously reported (A). In contrast, we did not observe Ab-immunopositive structures in the normal adult monkey brains (B-D). However, we did observe small but obvious A β -immunopositive senile plaques (SPs) in the frontal and temporal cortices of DM-affected adult monkeys (E-H). Scale bars, 100 μ m. (I) Quantitative image analysis of Ab-immunopositive area in the sections obtained from female normal adult monkey, DM-affected adult monkey, and normal aged monkey brains. Data obtained from normal aged monkey brains were set as standards. Y-axes show the mean values of the quantified data. CT, normal cynomolgus monkeys. DM, DM-affected monkeys.

doi:10.1371/journal.pone.0117362.g002




Fig 3. Cerebral amyloid angiopathy in the brains of aged monkeys with DM. Images of temporal lobe (TL) sections from normal cynomolgus monkeys (A, B) and cynomolgus monkeys with DM (C, D). Sections were immunostained with anti-A β antibody and counterstained with hematoxylin. In the brains of DM-affected aged monkeys, we observed very severe CAA lesions (arrowheads) as compared to normal aged monkeys. CT, normal aged monkeys. DM, DM-affected aged monkeys. Scale bars, 100 μ m.

doi:10.1371/journal.pone.0117362.g003

DM-affected cynomolgus monkeys in order to assess the relationship between DM and AD pathology. As previously reported, SPs spontaneously form in the brains of aged monkeys over the age of 25 years, but never in the brains of normal young monkeys and adult monkey younger than 20 years (Fig. 2A-D) [26]. Strikingly, our immunohistochemical analyses revealed SP depositions in the brains of DM-affected adult monkeys as young as 18 years (Fig. 2E-H). To our knowledge, this is the first study to show that DM enhances A β pathology even in nonhuman primate brains without genetic manipulation. We also observed much severe CAA lesions in the brains of DM-affected aged monkeys than in those of normal aged monkeys (Fig. 3). These findings are consistent with the previous studies showing that DM-related conditions induce amyloidogenesis and A β pathology in rodent models [14–25]. Although additional




Fig 4. The analyses of A β and GA β in the brains of normal and DM-affected monkeys. (A) A β level in young monkey, normal adult monkey, DM-affected monkey, and normal aged monkey brains were assessed with sandwich ELISA. A β level was significantly increased in normal aged monkey brains. In DM-affected monkey brains, A β level seemed unchanged. Data obtained from young monkey brains were set as standards. Y-axes show the mean values of the quantified data. Values are means \pm SD. *P < 0.02. (B) Image of temporal lobe sections from a 19-year-old normal adult monkey and an 18-year-old cynomolgus monkey with DM. Sections were immunostained with the anti-GA β -specific antibody 4396C and counterstained with hematoxylin. In the brain of the normal adult monkey, we observed little, if any, immunoreactivity for anti-GA β antibody. By contrast, in the brain of the DM-affected adult monkey, we observed several neurons immunopositive for anti-GA β antibody. Scale bars, 100 μ m. (C) Quantitative image analysis of GA β -immunopositive area in the sections obtained from female normal adult monkey and DM-affected adult monkey brains. Data obtained from normal adult monkey brains were set as standards. Y-axes show the mean values of the quantified data. Values are means \pm SD. *P < 0.02. (D) Dot blots showing the amount of GA β generated in brains of cynomolgus monkeys of different ages. The blot samples were adjusted to 1 μ g, 2.5 μ g, or 5 μ g of total protein. Dot blot analyses showed that GA β generation increased in an age-dependent manner. (E) Dot blots showing the amount of GA β generated in the brains of normal adult monkeys and DM-affected adult monkeys. The blot samples were adjusted to 1 μ g, 2.5 μ g, or 5 μ g of total protein. The amount of GA β in brains samples from DM-affected monkeys was significantly increased compared to those from normal adult monkeys. CT, normal adult monkeys; DM, DM-affected adult monkeys.

doi:10.1371/journal.pone.0117362.g004




Fig 5. Immunohistochemistry and Western blot analyses of Rab5, Rab7, Rab11, and APP in the brains of normal and DM-affected monkeys. Images of temporal lobe sections from a 19-year-old normal cynomolgus monkey (A, B) and an 18-year-old cynomolgus monkey with DM (C, D). Sections were immunostained with anti-APP antibody (A, C) or anti-Rab5 antibody (B, D), and then counterstained with hematoxylin. Scale bars, 100 μ m. In the brains of normal adult monkeys, APP and Rab5 were observed as small granules in neurons (A, B). By contrast in the brains of DM-affected adult monkeys, APP- and Rab5-immunopositive granules were enlarged, and APP and Rab5 immunoreactivity was significantly more robust (C, D). (E) Quantitative image analysis of APP or Rab5-immunopositive area in the sections obtained from female normal adult monkey and DM-affected adult monkey brains. Data obtained from normal adult monkey brains were set as standards respectively. Y-axes show the mean values of the quantified data. Values are means \pm SD. * $P < 0.01$. (F) Western blots showing the amounts of APP, Rab5, Rab7, Rab11, and synaptophysin in the brains of normal monkeys and DM-affected monkeys of different ages. Western blot analyses showed that APP and Rab GTPases were significantly increased in both DM-affected adult and aged monkey brains. In the brains of DM-affected adult monkeys, APP and Rab GTPases levels were obviously increased compared to those of normal adult monkeys. Lanes contained microsome fractions derived from the brains of young monkeys, normal adult monkeys, DM-affected adult monkeys, and aged