

treatment status, and disease stage. The FACT-O, particularly the physical well-being, functional well-being, and ovarian subscales were sensitive to changes in performance status over a two-three month period. To assess the effects of bevacizumab-related side effects on QOL, questions from the FACIT measurement system have been added related to rash, concerns about appearance, diarrhea, fatigue, and appetite (labeled “Additional Concerns (TSE)”).

In order to evaluate the effect of surgery on quality of life, patients will complete the Physical Functioning Subscale of the Rand 36-Item Short Form Health Survey (Rand SF-36). The Physical Functioning (PF) Subscale is a 10-item subscale of the Rand SF-36a global quality of life questionnaire, designed to assess quality of life of patients across all medical conditions<sup>54-56</sup>.

The PF Subscale consists of items concerning activities of daily living: walking, climbing stairs, bathing, dressing, and performance of physical activities, with each item rated on a three-point scale of limitation of activity due to the patients' health, from "not limited" to "limited a lot." Internal consistency of the PF subscale is excellent, with an alpha co-efficient ranging from 0.89 to 0.92.<sup>56</sup> The PF subscale has been found to significantly correlate with other physical functioning measures (Sickness Impact Profile [SIP],  $r=.67-.78$ ; shortened Arthritis Impact Measurement Scale (sAIMS,  $r=.60$ ). Further evidence of validity was provided by the PF subscale distinguishing between patients with serious and mild medical conditions.<sup>57</sup> Furthermore, the PF subscale has been found to be responsive to changes in functioning after surgical procedures (thoracic surgery for treatment of non-small-cell lung cancer, abdominal aortic aneurysm repair, and total hip arthroplasty<sup>58</sup>), and sensitive to differences in quality of life between laparoscopic and open surgical procedures<sup>59,60</sup> and between epidural and patient-controlled analgesia after colonic surgery.<sup>57</sup> Norms have been developed for all subscales of the SF-36, by gender and age groups, based upon 2,474 respondents, as well as for patients with physical limitations.<sup>58, 59</sup>

Eight questions will be included to measure specific quality of life problems after surgery (labeled “Additional Concerns (S)” in). These questions will address issues such as pain, fatigue, problems with the surgical incision, and ostomy appliances. Similar questions have been used in GOG-0152 (A Phase III Randomized Study of Cisplatin And Taxol<sup>®</sup> with Interval Secondary Cytoreduction versus Cisplatin and Paclitaxel in Patients with Suboptimal Stage III Epithelial Ovarian Carcinoma). Several of the questions were taken from questionnaires in the FACIT quality of life measurement system.<sup>61</sup> others were drafted to be similar in format to FACIT questions.

## 2.10 Background and Rationale for Translational Research(08/04/08)

The translational research component of this protocol will focus on the molecular and biochemical phenotype of recurrent ovarian cancer. It is well known that the vast majority of patients with advanced ovarian cancer who respond to initial

therapy will recur. However, these recurrent tumors remain essentially a molecular enigma because of their general unavailability for analysis. A brief review of the GOG Tissue Bank demonstrated that less than 5% of ovarian cancer specimens are from sources other than the primary tumor. Further, only 22 specimens of recurrent ovarian cancer with attached clinical data have been banked.

This protocol provides an extraordinary opportunity to study these tumors, characterize them on a molecular basis, compare them to the original primary tumor, and determine the basis for disease recurrence and altered drug sensitivities. In the past five years, over 600 manuscripts on expression profiling of cancers using microarray technology have been published, illustrating the recognized utility of this approach in exploring questions of tumor biology and clinical correlates. The principles of class prediction and class discovery as they apply to the molecular classification of human cancers were exemplified by Golub et al., who used oligonucleotide microarrays to monitor gene expression in acute leukemias as a test case.<sup>62</sup> Class prediction identified and validated a subset of informative genes whose expression was highly correlated with previously defined classes. Further, subsequent studies have utilized these approaches to provide proof of the "molecular profiling principle" as well as to gain novel insights into clinical cancer problems. Using a specialized, lymphoid cell-specific cDNA microarray, Alizadeh et al. performed expression profiling of diffuse large B-cell lymphomas and identified two molecularly distinct forms of this malignancy that correlated with overall survival.<sup>63</sup>

Further, recent work on the problem of drug resistance has detailed multiple potential biochemical mechanisms, which may be critical for the development of drug resistance in ovarian cancer. For instance the expression level of DNA repair enzymes and membrane transporters have been implicated in cisplatin resistance while microtubule mutations have been shown to affect paclitaxel sensitivity.<sup>64, 65</sup> These *in vitro* determined mechanisms require testing and validation on *in vivo* derived tumor specimens.

GOG-0213 patients with platinum-sensitive, recurrent epithelial ovarian, peritoneal primary or Fallopian tube carcinoma undergoing secondary cytoreduction will be able to provide archival formalin-fixed and paraffin-embedded primary or metastatic tumor, a pre-op serum specimen, a pre-op plasma specimen, formalin-fixed recurrent tumor, frozen recurrent tumor, formalin-fixed normal tissue and/or frozen normal tissue to establish an enduring resource for defining the molecular and biochemical phenotype of recurrent ovarian cancer. The pre-op serum and plasma will be prepared from blood drawn prior to secondary cytoreductive surgery. The exact choice of the biomarkers and profiles to be evaluated and the assays to be performed in the tissue, serum and plasma specimens submitted for the GOG-0213 patients undergoing secondary cytoreduction will be reevaluated based on evolving data in the field.

### 2.11 Rationale for Banking DNA from Whole Blood for Research (06/22/09)

The National Cancer Institute is encouraging Cooperative Clinical Trial Groups including the Gynecologic Oncology Group to bank whole blood from women participating in clinical trials such that the blood specimens will be linked to clinical outcome data (progression-free survival, overall survival, response and adverse effects) and information regarding treatment. The purpose of this effort is to support research including pharmacogenomic and pharmacogenetic research.

Women who are candidates for this clinical trial or who have already been enrolled on GOG-0213 will be asked to give permission for 10 ml of their blood to be collected for this research study and for future research. No matter what the women decide to do, it will not affect their care. The women can still participate in this GOG study even if they do not allow their blood to be collected and used for this research study and/or for future research. Women already enrolled on GOG-0213 will need to be re-consented for this collection.

### 2.12 Single Nucleotide Polymorphisms (SNPs) and SNP Profiling(06/22/09)

It is well known that individual single nucleotide polymorphisms (SNPs) and SNP profiles are associated with many clinical aspects of cancer. This includes risk of developing invasive cancer, risk of recurrence of cancer, patient survival and chemotherapy toxicity. We propose to use genome wide SNP-association studies and individual SNP analyses to identify SNPs which correlate with a variety of clinical measures including but not limited to patient survival, recurrence of disease, response, and toxicity

### 2.13 Rationale for the inclusion of fallopian tube carcinoma (FTCA)

Primary carcinoma of the fallopian tube is among the rarest malignancies of the female genital tract accounting for approximately 3.3/1,000,000 women annually. Despite its rarity, the disease shares many features of ovarian and primary peritoneal cancer including, risk factors (age and nulliparity), genomic alterations (LOH 3q and 8q, 1q, 5p, 7q, 12p and 20q), genetic abnormalities (Her 2-neu, P53, and k-ras mutations), natural history (local followed peritoneal metastases), response to chemotherapy, and anticipated survival by stage.<sup>66-68</sup> The latter feature is modeled after primary ovarian cancer as well. Most strikingly though is the relationship between BRCA mutation and the attendant increased risk of fallopian tube cancer over baseline. A life-time risk increase of 120 fold over background has been reported for women who harbor BRCA mutation. In fact, women diagnosed with FTCA may be at greater risk for harboring a BRCA mutation than women diagnosed with ovarian cancer. As such, women undergoing risk-reducing bilateral salpingo-oophorectomy (RRBSO) are recommended to have as much of the fallopian tube resected as possible and undergo step-sectioning as is performed for the ovary.

Since there appears to be a common set of environmental and genetic risk factors for FTCA and ovarian cancer, it is not surprising that the clinical approach for these two neoplasms is similar including primary surgical resection and debulking or staging, adjuvant platinum- and taxane-based chemotherapy and surveillance protocols (including CA-125). Based on these features and the lack of consensus as to the precise diagnostic criteria separating primary entities of the ovary, fallopian tube and peritoneum it is appropriate to consider FTCA within this spectrum of disease.

#### 2.14 Inclusion of Women and Minorities

The Gynecologic Oncology Group and GOG participating institutions will not exclude potential subjects from participating in this or any study solely on the basis of ethnic origin or socioeconomic status. Every attempt will be made to enter all eligible patients into this protocol and therefore address the study objectives in a patient population representative of the entire ovarian, fallopian tube and peritoneal primary cancer population treated by participating institutions.

### 3.0 PATIENT ELIGIBILITY AND EXCLUSIONS

#### 3.1 Eligible Patients

- 3.10 Patients enrolled after August 28, 2011 must be candidates for cytoreductive surgery and consent to have their surgical treatment determined by randomization.(08/29/11)(12/19/11)
- 3.11 Patients must have histologic diagnosis of epithelial ovarian carcinoma, peritoneal primary or Fallopian tube carcinoma, which is now recurrent.
- 3.12 Patients with the following histologic epithelial cell types are eligible: Serous adenocarcinoma, endometrioid adenocarcinoma, mucinous adenocarcinoma, undifferentiated carcinoma, clear cell adenocarcinoma, mixed epithelial carcinoma, transitional cell carcinoma, malignant Brenner's Tumor, or adenocarcinoma not otherwise specified (N.O.S.).
- 3.13 Patients must have had a complete response to front-line platinum-taxane therapy (at least three cycles).(08/04/08)
- 3.131 A complete response to front-line chemotherapy must include: negative physical exam, negative pelvic exam and normalization of CA125, if elevated at baseline. Although not required, any radiographic assessment of disease status (e.g. CT, MRI, PET/CT, etc) obtained following the completion of primary therapy (defined in 3.133) should be considered negative for disease.
- 3.132 All patients must have also had a treatment-free interval without clinical evidence of progressive disease of at least 6 months from completion of front-line chemotherapy (both platinum and taxane). Front-line therapy may have included a biologic agent (i.e. bevacizumab).
- 3.133 Front-line treatment may include maintenance therapy following complete clinical or pathological response. However, maintenance cytotoxic chemotherapy must be discontinued for a minimum of 6 months prior to documentation of recurrent disease. Patients receiving maintenance biological therapy **or hormonal therapy** are ELIGIBLE provided their recurrence is documented more than 6 months from primary cytotoxic chemotherapy completion (includes maintenance chemotherapy) AND a minimum 4 weeks has elapsed since their last infusion of biological therapy.(06/22/09)

3.14 Patients must have clinically evident recurrent disease for the purpose of this study, **(08/29/11)**

3.142 *Measurable disease* (RECIST) is defined as at least one lesion that can be accurately measured in at least one dimension (longest dimension to be recorded). Each lesion must be more than or equal to 20 mm when measured by conventional techniques, MRI or CT, or more than or equal to 10 mm when measured by spiral CT.

3.15 Patients must have adequate:

3.151 Bone marrow function: Absolute neutrophil count (ANC) greater than or equal to 1,500/mm<sup>3</sup>, equivalent to Common Toxicity Criteria for Adverse Events v3.0 (CTCAE) Grade 1.

3.152 Platelets greater than or equal to 100,000/mm<sup>3</sup>. (CTCAE Grade 0-1).

3.153 Renal function: Creatinine (non-IDMS)  $\leq 1.5 \times$  institutional upper limit normal (ULN), CTCAE Grade 1 **(03/15/10) (08/23/10)**

3.154 Hepatic function:

3.1541 Total bilirubin  $\leq 1.5$  ULN (CTCAE Grade 1).

3.1542 SGOT/AST and Alkaline Phosphatase  $\leq 2.5$  times the upper limit of normal in the absence of liver metastasis.  
SGOT/AST and Alkaline Phosphatase  $< 5.0$  times ULN in the presence of liver metastasis.

3.155 **This criterion applies only to the patients enrolled before August 29, 2011 and those enrolled after this date electing to receive Bevacizumab.**

Patients must have a urine protein-to-creatinine ratio (UPCR)  $< 1.0$  mg/dL. The UPCR has been found to correlate directly with the amount of protein excreted in a 24 hr urine collection. Specifically, a UPCR of 1.0 is equivalent to 1.0 gram of protein in a 24-hour urine collection. Obtain at least 4 ml of a random urine sample in a sterile container (does not have to be a 24 hour urine). Send the sample to the lab with a request for urine protein and creatinine levels [separate requests]. The lab will measure protein concentration (mg/dL) and creatinine concentration (mg/dL). The UPCR is derived as: protein concentration (mg/dL) / creatinine concentration (mg/dL).

- 3.16 (This eligibility criterion does not apply to patients enrolled after August 28, 2011).**(08/29/11)(12/19/11)**Patients who are not candidates for surgical cytoreduction are eligible for the chemotherapy randomization. Patients are not considered candidates for surgical cytoreduction if complete cytoreduction in the estimation of the investigator is impossible or a medical infirmity precludes exploration and debulking.
- 3.17 Patients must have met the pre-entry requirements specified in Section 7.0.
- 3.18 Patients must have signed an approved informed consent and authorization permitting release of personal health information.
- 3.19 Patients must have a GOG Performance Status of 0, 1, or 2.
- 3.110 Patients must be at least 18 years old.

### 3.2 Ineligible Patients

- 3.21 Patients who have received more than one previous regimen of chemotherapy (maintenance is not considered a second regimen).
- 3.22 Patients receiving concurrent immunotherapy, or radiotherapy.
- 3.23 Patients who have received prior radiotherapy to any portion of the abdominal cavity or pelvis are excluded.
- 3.24 Patients whom have already undergone secondary cytoreduction for recurrent disease are excluded.**(08/29/11)**
- 3.25 Patients with a prior histologic diagnosis of borderline, low malignant potential (grade 0) epithelial carcinoma that was surgically resected and who subsequently developed an unrelated, new invasive epithelial ovarian or peritoneal primary cancer are eligible provided that they meet the criteria listed in Section 3.12.
- 3.26 Patients who require parenteral hydration or nutrition and have evidence of partial bowel obstruction or perforation.
- 3.27 Patients who have received prior chemotherapy for any abdominal or pelvic tumor (other than ovarian, fallopian tube, and primary peritoneal) are excluded. **(06/22/09) (03/15/10)**
- 3.28 Patients with synchronous primary endometrial cancer, or a past history of primary endometrial cancer, are excluded, unless all of the following conditions are met: Stage not greater than I-B; no more than superficial myometrial invasion, without vascular or lymphatic invasion; no poorly

differentiated subtypes, including papillary serous, clear cell or other FIGO Grade 3 lesions.

- 3.29 Patients with uncontrolled infection.
- 3.30 Patients with concurrent severe medical problems unrelated to the malignancy that would significantly limit full compliance with the study or expose the patient to extreme risk or decreased life expectancy.
- 3.31 Patients with  $\geq$  grade 2 peripheral neuropathy
- 3.32 Patients with a history of allergic reactions to carboplatin and/or paclitaxel or chemically similar compounds. Patients with allergic (hypersensitivity) reactions to these chemotherapeutic agents are **NOT** excluded **IF** they were successfully retreated following a desensitization program or protocol.
- 3.33 **This criterion applies only to the patients enrolled before August 29, 2011 and those enrolled after this date electing to receive Bevacizumab.(10/01/12)**

Patients with known hypersensitivity to Chinese hamster ovary cell products or other recombinant human or humanized antibodies.

- 3.34 Patients of childbearing potential, not practicing adequate contraception, patients who are pregnant or patients who are nursing are not eligible for this trial. To date, no fetal studies in animal or humans have been performed. The possibility of harm to a fetus is likely. Bevacizumab specifically inhibits VEGF, which is responsible for the formation of new blood vessels during development, and antibodies can cross the placenta. Therefore, bevacizumab should not be administered to pregnant women. In addition, there are unknown immediate and long-term consequences of chemotherapy administration to these women. In addition, surgical exploration as mandated by randomization during pregnancy may cause imminent mortal consequences. Further, it is not known whether bevacizumab is excreted in human milk. Because many drugs are excreted in human milk, bevacizumab should not be administered to nursing women. Subjects will be apprised of the large potential risk to a developing fetus.
- 3.35 Patients with other invasive malignancies, with the exception of non-melanoma skin cancer, who had (or have) any evidence of the other cancer present within the last 5 years or whose previous cancer treatment contraindicates this protocol therapy.



- 3.36 **This criterion applies only to the patients enrolled before August 29, 2011 and those enrolled after this date electing to receive Bevacizumab.(10/01/12)**

Patients with active bleeding or pathologic conditions that carry high risk of bleeding such as a known bleeding disorder, coagulopathy, or tumor involving major vessels.

- 3.37 **This criterion applies only to the patients enrolled before August 29, 2011 and those enrolled after this date electing to receive Bevacizumab.(10/01/12)**

Patients with a history or evidence upon physical examination of CNS disease, including primary brain tumor, seizures not controlled with standard medical therapy, any brain metastases or a history of stroke within 5 years of the first date of treatment on this study.

- 3.38 **This criterion applies only to the patients enrolled before August 29, 2011 and those enrolled after this date electing to receive Bevacizumab. (10/01/12)**

Patients with clinically significant cardiovascular disease. This includes:

- 3.381 Patients with significant cardiac conduction abnormalities, i.e. PR interval > 0.24 sec or 2nd or 3rd degree AV block.
- 3.382 Uncontrolled hypertension, defined as systolic > 150 mm Hg or diastolic > 90 mm Hg.
- 3.383 Myocardial infarction, cardiac arrhythmia or unstable angina < 6 months prior to registration.
- 3.384 New York Heart Association (NYHA) Grade II or greater congestive heart failure.
- 3.385 Serious cardiac arrhythmia requiring medication.
- 3.386 Grade II or greater peripheral vascular disease (exception: episodes of ischemia < 24 hrs in duration, that are managed non-surgically and without permanent deficit).(03/15/10)
- 3.387 History of CVA within six months.

- 3.39 **This criterion applies only to the patients enrolled before August 29, 2011 and those enrolled after this date electing to receive Bevacizumab.(10/01/12)**

Patients who have had a major surgical procedure, open biopsy, dental extractions or other dental surgery/procedure that results in an open wound, or significant traumatic injury within 28 days prior to the first date of treatment on this study, or anticipation of need for major surgical procedure during the course of the study; patients with placement of vascular access device or core biopsy within 7 days prior to the first date of treatment on this study.

3.391 Patients undergoing pre-treatment secondary cytoreduction will undergo therapy with bevacizumab on cycle #2 (See Section 5.234).

3.392 Patients undergoing pre-treatment surgery for purposes other than cytoreduction may also participate provided they meet eligibility in Section 3.1. Patients randomized to arms containing bevacizumab must wait a minimum of 28 days since that procedure to begin protocol treatment. Patients who undergo an uncomplicated port placement must wait a minimum of 7 days to begin protocol treatment. **(03/15/10)**

## 4.0 STUDY MODALITIES

### 4.1 Carboplatin (Paraplatin®, NSC # 241240)

4.11 Formulation: Carboplatin is supplied as a sterile lyophilized powder available in single-dose vials containing 50 mg, 150 mg and 450 mg of carboplatin for administration by intravenous infusion. Each vial contains equal parts by weight of carboplatin and mannitol.

4.12 Solution Preparation: Immediately before use, the content of each vial must be reconstituted with either sterile water for injection, USP, 5% dextrose in water, or 0.9% sodium chloride injection, USP, according to the following schedule:

<u>Vial Strength</u>	<u>Diluent Volume</u>
50 mg	5 ml
150 mg	15 ml
450 mg	45 ml

These dilutions all produce a carboplatin concentration of 10 mg/ml.

NOTE: Aluminum reacts with carboplatin causing precipitate formation and loss of potency. Therefore, needles or intravenous sets containing aluminum parts that may come in contact with the drug must not be used for the preparation or administration of carboplatin.

4.13 Storage: Unopened vials of carboplatin are stable for the life indicated on the package when stored at controlled room temperature and protected from light.

4.14 Stability: When prepared as directed, carboplatin solutions are stable for eight hours at room temperature. Since no antibacterial preservative is contained in the formulation, it is recommended that carboplatin solutions be discarded eight hours after dilution.

4.15 Supplier: Commercially available from Bristol-Myers Squibb Company.

4.16 Administration: See Section 5.2.

4.17 Adverse effects:

Hematologic: Myelosuppression

Gastrointestinal: Nausea, vomiting, diarrhea, abdominal pain, constipation

Neurologic: Peripheral neuropathy, ototoxicity, visual disturbances, change in taste, central nervous system symptoms

Renal: Abnormal renal function test results including serum creatinine, blood urea nitrogen, and creatinine clearance

Hepatic: Abnormal liver function tests including bilirubin, SGOT, and alkaline phosphatase

Electrolyte Changes: Abnormally decreased serum electrolyte values reported for sodium, potassium, calcium, and magnesium.

Allergic Reactions: Rash, urticaria, erythema, pruritus, and rarely bronchospasm and hypotension.

Injection Site Reactions: Redness, swelling, pain; necrosis associated with extravasation has been reported.

Other: Pain, asthenia, alopecia. Cardiovascular, respiratory, genitourinary, and mucosal side effects have occurred in 6% or less of the patients.

Cardiovascular events (cardiac failure, embolism, cerebrovascular accidents) were fatal in less than 1% of patients and did not appear to be related to chemotherapy. Cancer-associated hemolytic-uremic syndrome has been reported rarely. Malaise, anorexia, and hypertension have been reported as part of post-marketing surveillance.

\*See FDA-approved package insert for a comprehensive list of adverse events associated with carboplatin.

## 4.2 Paclitaxel (Taxol®, NSC #673089)

- 4.21 Formulation: Paclitaxel is a poorly soluble plant product from the western yew, *Taxus brevifolia*. Improved solubility requires a mixed solvent system with further dilutions of either 0.9% sodium chloride or 5% dextrose in water.

Paclitaxel is supplied as a sterile solution concentrate, 6 mg/ml in 5 ml vials (30 mg/vial) in polyoxyethylated castor oil (Cremophor EL) 50% and dehydrated alcohol, USP, 50%. The contents of the vial must be diluted just prior to clinical use. It is also available in 100 and 300 mg vials.

- 4.22 Solution Preparation: Paclitaxel, at the appropriate dose, will be diluted in 500-1000 ml of 0.9% Sodium Chloride injection, USP or 5% Dextrose injection, USP (D5W) (500 ml is adequate if paclitaxel is a single agent). Paclitaxel must be prepared in glass or polyolefin containers due to leaching of diethylhexylphthalate (DEHP) plasticizer from polyvinyl chloride (PVC) bags and intravenous tubing by the Cremophor vehicle in which paclitaxel is solubilized.

NOTE: Formation of a small number of fibers in solution (within acceptable limits established by the USP Particulate Matter Test for LVPs) has been observed after preparation of paclitaxel. Therefore, in-line filtration is necessary for administration of paclitaxel solutions. In-line filtration should be accomplished by incorporating a hydrophilic, microporous filter of pore size not greater than 0.22 microns (e.g.: IVEX-

II, IVEX-HP or equivalent) into the IV fluid pathway distal to the infusion pump. Although particulate formation does not indicate loss of drug potency, solutions exhibiting excessive particulate matter formation should not be used.

- 4.23 Storage: The intact vials can be stored in a temperature range between 2-25° C (36-77°F).
- 4.24 Stability: Commercially available paclitaxel will be labeled with an expiration date. All solutions of paclitaxel exhibit a slight haziness directly proportional to the concentration of drug and the time elapsed after preparation, although when prepared as described above, solutions of paclitaxel (0.3-1.2 mg/ml) are physically and chemically stable for 27 hours.
- 4.25 Supplier: Commercially available from Bristol-Myers Squibb Company.
- 4.26 Administration: Paclitaxel, at the appropriate dose and dilution, will be given as a 3-hour continuous IV infusion. Paclitaxel will be administered via an infusion control device (pump) using non-PVC tubing and connectors, such as the IV administration sets (polyethylene or polyolefin) that are used to infuse parenteral Nitroglycerin. Nothing else is to be infused through the line where paclitaxel is being administered. See section 5.2.
- 4.27 Adverse Effects: Hematologic: Myelosuppression  
Gastrointestinal: Nausea and vomiting, diarrhea, stomatitis, mucositis, pharyngitis, typhlitis, ischemic colitis, neutropenic enterocolitis  
Heart: Arrhythmia, heart block, ventricular tachycardia, myocardial infarction (MI), bradycardia, atrial arrhythmia  
Pulmonary: Pneumonitis  
Blood Pressure: Hypotension, hypertension (possibly related to concomitant medication--Dexamethasone)  
Neurologic: Sensory (taste), peripheral neuropathy, seizures, mood swings, hepatic encephalopathy, encephalopathy  
Skin: Infiltration: erythema, induration, tenderness, rarely ulceration, injection-recall reactions, erythema multiforme (e.g., Stevens-Johnson syndrome, toxic epidermal necrolysis)  
Allergy: Anaphylactoid and urticarial reactions (acute), flushing, rash, pruritus  
Liver: Increased SGOT, SGPT, bilirubin, alkaline phosphatase and triglycerides, hepatic failure, hepatic necrosis  
Other: Alopecia, fatigue, arthralgia, myalgia, light-headedness, myopathy, headaches

Other, Vision: Sensation of flashing lights, blurred vision, scintillating scotomata

\*See FDA- approved package insert for a comprehensive list of adverse events associated with paclitaxel.

#### 4.3 Bevacizumab (NSC #704865, IND #113912) (08/04/08) (12/19/11)

All investigators who receive a copy of the protocol should also obtain a copy of the Investigator's Brochure (IB). IB's are available from the Pharmaceutical Management Branch, CTEP, DCTD, NCI and may be obtained by emailing the IB Coordinator ([ibcoordinator@mail.nih.gov](mailto:ibcoordinator@mail.nih.gov)) or by calling the IB Coordinator at 301-496-5725.

4.31 Description: Bevacizumab is a recombinant humanized anti-VEGF monoclonal antibody, consisting of 93% human and 7% murine amino acid sequences. The agent is composed of human IgG framework and murine antigen-binding complementarity-determining regions. Bevacizumab blocks the binding of vascular endothelial growth factor (VEGF) to its receptors resulting in inhibition of angiogenesis.

4.32 How Supplied: Bevacizumab is supplied as a clear to slightly opalescent, sterile liquid ready for parenteral administration in a 400 mg (25mg/ml – 16 mL) fill

glass vial containing bevacizumab with phosphate, trehalose, polysorbate 20, and Sterile Water for Injection, USP.

4.33 Storage and Stability: Bevacizumab is shipped on blue ice for next day delivery. On receipt, bevacizumab should be stored in the refrigerator (2° to 8°C) and should remain refrigerated until just prior to use. Do not freeze. Do not shake. Shelf-life studies of bevacizumab are continuing. Investigators will be notified when lots have expired. The sterile single use vials contain no antibacterial preservatives; therefore, vials should be discarded eight hours after initial entry.

4.34 Preparation: Vials contain no preservative and are intended for single use only. Place the calculated dose in 100 mL of 0.9% Sodium Chloride for injection. Once diluted in 0.9% Sodium Chloride for injection, the bevacizumab solution must be administered within 8 hours.

4.35 Administration: Bevacizumab is administered intravenously as a continuous infusion. The initial dose should be administered over a minimum of 90 minutes. If no adverse reactions occur after the initial dose, the second dose should be administered over a minimum of 60 minutes. If no adverse reactions occur after the second dose, all

subsequent doses should be administered over a minimum of 30 minutes. If infusion-related adverse reactions occur, all subsequent infusions should be administered over the shortest period that was well tolerated.

To insure complete delivery of bevacizumab, flush the IV infusion line with 0.9% sodium chloride. The following are two recommended methods for flushing the bevacizumab IV infusion line:

1. When the bevacizumab infusion is complete, add an additional 50mL of 0.9% sodium chloride for injection to the bevacizumab infusion bag. Continue the infusion until a volume equal to that of the volume contained in the tubing has been administered.
2. Replace the empty bevacizumab infusion bag with a 50mL bag of 0.9% sodium chloride for injection and infuse a volume equal to the volume contained in the tubing.

Please note: the flush is not included in the total recommended infusion times.

#### 4.36 Comprehensive Adverse Events and Potential Risks List (CAEPR) for Bevacizumab (NSC #704865) (08/23/10) (12/19/11) (09/29/14)

The Comprehensive Adverse Event and Potential Risks list (CAEPR) provides a single list of reported and/or potential adverse events (AE) associated with an agent using a uniform presentation of events by body system. In addition to the comprehensive list, a subset, the Specific Protocol Exceptions to Expedited Reporting (SPEER), appears in a separate column and is identified with bold and italicized text. This subset of AEs (SPEER) is a list of events that are protocol specific exceptions to expedited reporting to NCI (except as noted below). Refer to the 'CTEP, NCI Guidelines: Adverse Event Reporting Requirements' [http://ctep.cancer.gov/protocolDevelopment/electronic\\_applications/docs/aeguidelines.pdf](http://ctep.cancer.gov/protocolDevelopment/electronic_applications/docs/aeguidelines.pdf) for further clarification. *Frequency is provided based on 3540 patients.* Below is the CAEPR for bevacizumab (rhuMAb VEGF).

**NOTE:** Report AEs on the SPEER **ONLY IF** they exceed the grade noted in parentheses next to the AE in the SPEER. If this CAEPR is part of a combination protocol using multiple investigational agents and has an AE listed on different SPEERs, use the lower of the grades to determine if expedited reporting is required.

Version 2.3, August 1, 2013<sup>1</sup>

Adverse Events with Possible Relationship to Bevacizumab (rhuMAb VEGF) (CTCAE 4.0 Term) [n= 3540]			Specific Protocol Exceptions to Expedited Reporting (SPEER)
Likely (>20%)	Less Likely (<=20%)	Rare but Serious (<3%)	
<b>BLOOD AND LYMPHATIC SYSTEM DISORDERS</b>			
	Anemia		<b><i>Anemia (Gr 3)</i></b>
		Blood and lymphatic system disorders - Other (renal thrombotic microangiopathy)	
	Febrile neutropenia		<b><i>Febrile neutropenia (Gr 3)</i></b>
<b>CARDIAC DISORDERS</b>			
		Acute coronary syndrome <sup>2</sup>	

	Cardiac disorders - Other (supraventricular arrhythmias) <sup>3</sup>		<b>Cardiac disorders - Other (supraventricular arrhythmias)<sup>3</sup> (Gr 3)</b>
		Heart failure	
		Left ventricular systolic dysfunction	
		Myocardial infarction <sup>2</sup>	
		Ventricular arrhythmia	
		Ventricular fibrillation	
<b>GASTROINTESTINAL DISORDERS</b>			
	Abdominal pain		<b>Abdominal pain (Gr 3)</b>
	Colitis		<b>Colitis (Gr 3)</b>
	Constipation		<b>Constipation (Gr 3)</b>
	Diarrhea		<b>Diarrhea (Gr 3)</b>
	Dyspepsia		<b>Dyspepsia (Gr 2)</b>
		Gastrointestinal fistula <sup>4</sup>	
	Gastrointestinal hemorrhage <sup>5</sup>		<b>Gastrointestinal hemorrhage<sup>5</sup> (Gr 2)</b>
	Gastrointestinal obstruction <sup>6</sup>		
		Gastrointestinal perforation <sup>7</sup>	
		Gastrointestinal ulcer <sup>8</sup>	
	Ileus		
	Mucositis oral		<b>Mucositis oral (Gr 3)</b>
	Nausea		<b>Nausea (Gr 3)</b>
	Vomiting		<b>Vomiting (Gr 3)</b>
<b>GENERAL DISORDERS AND ADMINISTRATION SITE CONDITIONS</b>			
	Fatigue		<b>Fatigue (Gr 3)</b>
	Infusion related reaction		<b>Infusion related reaction (Gr 2)</b>
	Non-cardiac chest pain		<b>Non-cardiac chest pain (Gr 3)</b>
	Pain		<b>Pain (Gr 3)</b>
<b>IMMUNE SYSTEM DISORDERS</b>			
	Allergic reaction		<b>Allergic reaction (Gr 2)</b>
		Anaphylaxis	
<b>INFECTIONS AND INFESTATIONS</b>			
	Infection <sup>9</sup>		<b>Infection<sup>9</sup> (Gr 3)</b>
		Infections and infestations - Other (necrotizing fasciitis)	
	Infections and infestations - Other (peri-rectal abscess)		
<b>INJURY, POISONING AND PROCEDURAL COMPLICATIONS</b>			
		Injury, poisoning and procedural complications – Other (anastomotic leak) <sup>10</sup>	
	Wound complication		<b>Wound complication (Gr 2)</b>
	Wound dehiscence		<b>Wound dehiscence (Gr 2)</b>
<b>INVESTIGATIONS</b>			
	Alanine aminotransferase increased		<b>Alanine aminotransferase increased (Gr 3)</b>
	Alkaline phosphatase increased		<b>Alkaline phosphatase increased (Gr 3)</b>
	Aspartate aminotransferase increased		<b>Aspartate aminotransferase increased (Gr 3)</b>
	Blood bilirubin increased		<b>Blood bilirubin increased (Gr 2)</b>
	Cardiac troponin I increased		
Neutrophil count decreased			<b>Neutrophil count decreased (Gr 3)</b>
	Platelet count decreased		<b>Platelet count decreased (Gr 4)</b>



	Weight loss		<b>Weight loss (Gr 3)</b>
	White blood cell decreased		<b>White blood cell decreased (Gr 3)</b>
<b>METABOLISM AND NUTRITION DISORDERS</b>			
	Anorexia		<b>Anorexia (Gr 3)</b>
	Dehydration		<b>Dehydration (Gr 3)</b>
<b>MUSCULOSKELETAL AND CONNECTIVE TISSUE DISORDERS</b>			
	Arthralgia		<b>Arthralgia (Gr 3)</b>
	Musculoskeletal and connective tissue disorder - Other (bone metaphyseal dysplasia) <sup>11</sup>		
	Myalgia		<b>Myalgia (Gr 3)</b>
	Osteonecrosis of jaw <sup>12</sup>		
<b>NERVOUS SYSTEM DISORDERS</b>			
	Dizziness		<b>Dizziness (Gr 2)</b>
	Headache		<b>Headache (Gr 3)</b>
		Intracranial hemorrhage	
		Ischemia cerebrovascular <sup>2</sup>	
	Peripheral sensory neuropathy <sup>13</sup>		
		Reversible posterior leukoencephalopathy syndrome	
	Syncope		
<b>RENAL AND URINARY DISORDERS</b>			
		Acute kidney injury	
	Hematuria		<b>Hematuria (Gr 3)</b>
	Proteinuria		<b>Proteinuria (Gr 2)</b>
		Renal and urinary disorders - Other (Nephrotic Syndrome)	
		Urinary fistula	
<b>REPRODUCTIVE SYSTEM AND BREAST DISORDERS</b>			
Reproductive system and breast disorders - Other (ovarian failure) <sup>14</sup>			
		Vaginal fistula	
	Vaginal hemorrhage		<b>Vaginal hemorrhage (Gr 3)</b>
<b>RESPIRATORY, THORACIC AND MEDIASTINAL DISORDERS</b>			
	Allergic rhinitis		<b>Allergic rhinitis (Gr 3)</b>
		Bronchopleural fistula	
		Bronchopulmonary hemorrhage	
	Cough		<b>Cough (Gr 3)</b>
	Dyspnea		<b>Dyspnea (Gr 2)</b>
	Epistaxis		<b>Epistaxis (Gr 3)</b>
	Hoarseness		<b>Hoarseness (Gr 3)</b>
		Respiratory, thoracic and mediastinal disorders - Other (nasal-septal perforation)	
		Respiratory, thoracic and mediastinal disorders - Other (tracheo-esophageal fistula)	
<b>SKIN AND SUBCUTANEOUS TISSUE DISORDERS</b>			
	Pruritus		<b>Pruritus (Gr 2)</b>
	Rash maculo-papular		<b>Rash maculo-papular (Gr 2)</b>

	Urticaria		<b>Urticaria (Gr 2)</b>
<b>VASCULAR DISORDERS</b>			
Hypertension			<b>Hypertension (Gr 3)</b>
	Thromboembolic event		<b>Thromboembolic event (Gr 3)</b>
		Vascular disorders - Other (arterial thromboembolic event) <sup>2,15</sup>	

<sup>1</sup>This table will be updated as the toxicity profile of the agent is revised. Updates will be distributed to all Principal Investigators at the time of revision. The current version can be obtained by contacting [PIO@CTEP.NCI.NIH.GOV](mailto:PIO@CTEP.NCI.NIH.GOV). Your name, the name of the investigator, the protocol and the agent should be included in the e-mail.

<sup>2</sup>The risks of arterial thrombosis such as cardiac or CNS ischemia are increased in elderly patients and in patients with a history of diabetes.

<sup>3</sup>Supraventricular arrhythmias may include supraventricular tachycardia, atrial fibrillation and atrial flutter.

<sup>4</sup>Gastrointestinal fistula may include: Anal fistula, Colonic fistula, Duodenal fistula, Esophageal fistula, Gastric fistula, Gastrointestinal fistula, Rectal fistula, and other sites under the GASTROINTESTINAL DISORDERS SOC.

<sup>5</sup>Gastrointestinal hemorrhage may include: Colonic hemorrhage, Duodenal hemorrhage, Esophageal hemorrhage, Esophageal varices hemorrhage, Gastric hemorrhage, Hemorrhoidal hemorrhage, Intra-abdominal hemorrhage, Oral hemorrhage, Rectal hemorrhage, and other sites under the GASTROINTESTINAL DISORDERS SOC.

<sup>6</sup>Gastrointestinal obstruction may include: Colonic obstruction, Duodenal obstruction, Esophageal obstruction, Ileal obstruction, Jejunal obstruction, Rectal obstruction, Small intestinal obstruction, and other sites under the GASTROINTESTINAL DISORDERS SOC.

<sup>7</sup>Gastrointestinal perforation may include: Colonic perforation, Duodenal perforation, Esophageal perforation, Gastric perforation, Jejunal perforation, Rectal perforation, Small intestinal perforation, and other sites under the GASTROINTESTINAL DISORDERS SOC.

<sup>8</sup>Gastrointestinal ulcer may include: Duodenal ulcer, Esophageal ulcer, Gastric ulcer, and other sites under the GASTROINTESTINAL DISORDERS SOC.

<sup>9</sup>Infection may include any of the 75 infection sites under the INFECTIONS AND INFESTATIONS SOC.

<sup>10</sup>Anastomotic leak may include Gastric anastomotic leak; Gastrointestinal anastomotic leak; Large intestinal anastomotic leak; Rectal anastomotic leak; Small intestinal anastomotic leak; Urostomy leak; Vaginal anastomotic leak

<sup>11</sup>Metaphyseal dysplasia was observed in young patients who still have active epiphyseal growth plates.

<sup>12</sup>Cases of osteonecrosis of the jaw (ONJ) have been reported in cancer patients in association with bevacizumab treatment, the majority of whom had received prior or concomitant treatment with i.v. bisphosphonates.

<sup>13</sup>Increased rate of peripheral sensory neuropathy has been observed in trials combining bevacizumab and chemotherapy compared to chemotherapy alone.

<sup>14</sup>*Ovarian failure, defined as amenorrhea lasting 3 or more months with follicle-stimulating hormone (FSH) elevation ( $\geq 30$  mIU/mL), was increased in patients receiving adjuvant bevacizumab plus mFOLFOX compared to mFOLFOX alone (34% vs. 2%). After discontinuation of bevacizumab, resumption of*

*menses and an FSH level <30 mIU/mL was demonstrated in 22% (7/32) of these women. Long term effects of bevacizumab exposure on fertility are unknown.*

<sup>15</sup>Arterial thromboembolic event includes visceral arterial ischemia, peripheral arterial ischemia, heart attack and stroke.

**Also reported on bevacizumab (rhuMAb VEGF) trials but with the relationship to bevacizumab (rhuMAb VEGF) still undetermined:**

**BLOOD AND LYMPHATIC SYSTEM DISORDERS** - Blood and lymphatic system disorders - Other (idiopathic thrombocytopenia purpura); Bone marrow hypocellular; Disseminated intravascular coagulation; Hemolysis

**CARDIAC DISORDERS** - Atrioventricular block complete; Atrioventricular block first degree; Cardiac arrest; Myocarditis; Pericardial effusion; Restrictive cardiomyopathy; Right ventricular dysfunction

**EAR AND LABYRINTH DISORDERS** - Ear and labyrinth disorders - Other (tympanic membrane perforation); Hearing impaired; Tinnitus; Vertigo

**ENDOCRINE DISORDERS** - Hyperthyroidism; Hypothyroidism

**EYE DISORDERS** - Blurred vision; Cataract; Dry eye; Extraocular muscle paresis; Eye disorders - Other (blindness); Eye disorders - Other (conjunctival hemorrhage); Eye disorders - Other (corneal epithelial defect); Eye disorders - Other (floaters); Eye disorders - Other (ischemic CRVO); Eye disorders - Other (macular pucker); Eye disorders - Other (transient increased IOP > or =30 mm Hg); Eye disorders - Other (vitreous hemorrhage); Eye pain; Keratitis; Optic nerve disorder; Photophobia; Retinal detachment; Retinal tear; Retinopathy; Watering eyes

**GASTROINTESTINAL DISORDERS** - Ascites; Chelitis; Colonic stenosis; Dry mouth; Dysphagia; Enterocolitis; Esophageal pain; Esophageal stenosis; Flatulence; Gastrointestinal disorders - Other (peritonitis); Oral pain; Pancreatitis; Proctitis; Rectal mucositis; Rectal stenosis; Typhlitis

**GENERAL DISORDERS AND ADMINISTRATION SITE CONDITIONS** - Death NOS; Edema face; Edema limbs; Edema trunk; Facial pain; Fever; Flu like symptoms; Gait disturbance; Injection site reaction; Localized edema; Multi-organ failure; Sudden death NOS

**HEPATOBIILIARY DISORDERS** - Cholecystitis; Gallbladder necrosis; Gallbladder obstruction; Hepatic failure; Hepatic necrosis

**INFECTIONS AND INFESTATIONS** - Infections and infestations - Other (aseptic meningitis)

**INJURY, POISONING AND PROCEDURAL COMPLICATIONS** - Arterial injury; Bruising; Burn; Dermatitis radiation; Fracture

**INVESTIGATIONS** - Activated partial thromboplastin time prolonged; Blood antidiuretic hormone abnormal; CD4 lymphocytes decreased; CPK increased; Carbon monoxide diffusing capacity decreased; Electrocardiogram QT corrected interval prolonged; Forced expiratory volume decreased; GGT increased; INR increased; Lipase increased; Lymphocyte count decreased; Serum amylase increased; Weight gain

**METABOLISM AND NUTRITION DISORDERS** - Acidosis; Hypercalcemia; Hyperglycemia; Hyperkalemia; Hypermagnesemia; Hyponatremia; Hypertriglyceridemia; Hyperuricemia; Hypoalbuminemia; Hypocalcemia; Hypokalemia; Hypomagnesemia; Hyponatremia; Hypophosphatemia

**MUSCULOSKELETAL AND CONNECTIVE TISSUE DISORDERS** - Arthritis; Back pain; Bone pain; Chest wall pain; Fibrosis deep connective tissue; Generalized muscle weakness; Head soft tissue necrosis; Joint effusion; Muscle weakness lower limb; Muscle weakness upper limb; Musculoskeletal and connective tissue disorder - Other (aseptic necrotic bone); Musculoskeletal and connective tissue disorder - Other (myasthenia gravis); Musculoskeletal and connective tissue disorder - Other (polymyalgia rheumatica); Neck pain; Pain in extremity; Pelvic soft tissue necrosis; Soft tissue necrosis lower limb

**NEOPLASMS BENIGN, MALIGNANT AND UNSPECIFIED (INCL CYSTS AND POLYPS)** - Tumor pain  
**NERVOUS SYSTEM DISORDERS** - Arachnoiditis; Ataxia; Central nervous system necrosis; Cerebrospinal fluid leakage; Cognitive disturbance; Depressed level of consciousness; Dysesthesia; Dysgeusia; Dysphasia; Encephalopathy; Extrapyramidal disorder; Facial nerve disorder; Hydrocephalus; Leukoencephalopathy; Memory impairment; Nervous system disorders - Other (increased intracranial pressure); Paresthesia; Peripheral motor neuropathy; Pyramidal tract syndrome; Seizure; Somnolence; Tremor; Vasovagal reaction

**PSYCHIATRIC DISORDERS** - Agitation; Anxiety; Confusion; Depression; Insomnia; Libido decreased; Psychosis

**RENAL AND URINARY DISORDERS** - Bladder spasm; Chronic kidney disease; Cystitis noninfective; Renal and urinary disorders - Other (dysuria); Renal and urinary disorders - Other (ureterolithiasis); Renal hemorrhage; Urinary frequency; Urinary incontinence; Urinary retention; Urinary tract obstruction; Urinary tract pain

**REPRODUCTIVE SYSTEM AND BREAST DISORDERS** - Breast pain; Erectile dysfunction; Irregular menstruation; Pelvic pain; Vaginal discharge

**RESPIRATORY, THORACIC AND MEDIASTINAL DISORDERS** - Adult respiratory distress syndrome; Atelectasis; Hypoxia; Nasal congestion; Pulmonary fibrosis; Pulmonary hypertension; Respiratory failure; Respiratory, thoracic and mediastinal disorders - Other (dry nares); Respiratory, thoracic and mediastinal disorders - Other (pulmonary infarction)

**SKIN AND SUBCUTANEOUS TISSUE DISORDERS** - Alopecia; Dry skin; Hyperhidrosis; Nail loss; Pain of skin; Palmar-plantar erythrodysesthesia syndrome; Photosensitivity; Purpura; Rash acneiform; Skin and subcutaneous tissue disorders - Other (diabetic foot ulcer); Skin and subcutaneous tissue disorders - Other (skin breakdown/ decubitus ulcer); Skin hyperpigmentation; Skin induration; Skin ulceration; Stevens-Johnson syndrome

**VASCULAR DISORDERS** - Flushing; Hot flashes; Hypotension; Lymphocele; Phlebitis; Vasculitis

**Note:** Bevacizumab (rhuMAb VEGF) in combination with other agents could cause an exacerbation of any adverse event currently known to be caused by the other agent, or the combination may result in events never previously associated with either agent.

#### 4.37 General Information on Adverse Effects of Bevacizumab (06/22/09)

Based on clinical trials with bevacizumab as monotherapy or in combination with chemotherapy, the most common adverse events of any severity include asthenia, pain, headache, hypertension, diarrhea, stomatitis, constipation, epistaxis, dyspnea, dermatitis and proteinuria. The most common grade 3-4 adverse events were asthenia, pain, hypertension, diarrhea and leukopenia. The most serious AEs include life-threatening or fatal hemorrhage, arterial thromboembolic events, gastrointestinal perforation and wound dehiscence; these events were uncommon but occurred at an increased frequency compared to placebo or chemotherapy controls in randomized studies.

The following is a description of major adverse events associated with bevacizumab therapy. A list of Comprehensive Adverse Events and Potential Risks (CAEPR) in NCI-CTCAE v3.0 terms is included above. Reference may also be made to the Investigators' Brochure and the FDA package insert ([www.fda.gov/cder/foi/label/2004/1250851bl.pdf](http://www.fda.gov/cder/foi/label/2004/1250851bl.pdf)).

**Infusion-Related Reactions:** Infusion reactions with bevacizumab were uncommon (<3%) and rarely severe (0.2%). Infusion reactions may include rash, urticaria, fever, rigors, hypertension, hypotension, wheezing, or hypoxia. Currently, there is no adequate information on the safety of retreatment with bevacizumab in patients who have experienced severe infusion-related reactions.