


15. Millward MJ, Boyer MJ, Lehnert M, *et al*: Docetaxel and carboplatin is an active regimen in advanced non-small-cell lung cancer: a phase II study in Caucasian and Asian patients. *Ann Oncol* 14: 449-454, 2003.
16. Zhang X, Hunt JL, Landsittel DP, *et al*: Correlation of protease-activated receptor-1 with differentiation markers in squamous cell carcinoma of the head and neck and its implication in lymph node metastasis. *Clin Cancer Res* 10: 8451-8459, 2004.
17. Muller S, Su L, Tighiouart M, *et al*: Distinctive E-cadherin and epidermal growth factor receptor expression in metastatic and nonmetastatic head and neck squamous cell carcinoma: predictive and prognostic correlation. *Cancer* 113: 97-107, 2008.
18. Belani CP, Einzig A, Bonomi P, *et al*: Multicenter phase II trial of docetaxel and carboplatin in patients with stage IIIB and IV non-small-cell lung cancer. *Ann Oncol* 11: 673-678, 2000.
19. Jahanzeb M, Sarna G, Hirsch R, *et al*: Docetaxel and carboplatin as first-line therapy in advanced non-small cell lung carcinoma: a phase II study. *Anticancer Res* 24: 1239-1242, 2004.
20. Schuette W, Nagel S, von Weikersthal LF, *et al*: Randomized phase III trial of docetaxel plus carboplatin with or without levofloxacin prophylaxis in elderly patients with advanced non-small cell lung cancer: the APRONTA trial. *J Thorac Oncol* 6: 2090-2096, 2011.
21. Groen HJ, Sietsma H, Vincent A, *et al*: Randomized, placebo-controlled phase III study of docetaxel plus carboplatin with celecoxib and cyclooxygenase-2 expression as a biomarker for patients with advanced non-small-cell lung cancer: the NVALT-4 study. *J Clin Oncol* 29: 4320-4326, 2011.
22. Koch A, Bergman B, Holmberg E, *et al*: Swedish Lung Cancer Study Group: Effect of celecoxib on survival in patients with advanced non-small cell lung cancer: a double blind randomised clinical phase III trial (CYCLUS study) by the Swedish Lung Cancer Study Group. *Eur J Cancer* 47: 1546-1555, 2011.
23. Edelman MJ, Watson D, Wang X, *et al*: Eicosanoid modulation in advanced lung cancer: cyclooxygenase-2 expression is a positive predictive factor for celecoxib+chemotherapy - Cancer and Leukemia Group B Trial 30203. *J Clin Oncol* 26: 848-855, 2008.
24. Wang Z, Duan J, Guo Q, *et al*: A phase II clinical trial of celecoxib combined with platinum-based chemotherapy in the treatment of patients with advanced NSCLC as first-line treatment. *Chin J Lung Cancer* 11: 425-430, 2008 (In Chinese).
25. Mendes RT, Stanczyk CP, Sordi R, Otuki MF, dos Santos FA and Fernandes D: Selective inhibition of cyclooxygenase-2: risks and benefits. *Rev Bras Reumatol* 52: 767-782, 2012.
26. Matsui Y, Amano H, Ito Y, *et al*: Thromboxane A₂ receptor signaling facilitates tumor colonization through P-selectin-mediated interaction of tumor cells with platelets and endothelial cells. *Cancer Sci* 103: 700-707, 2012.
27. Honn KV: Inhibition of tumor cell metastasis by modulation of the vascular prostacyclin/thromboxane A₂ system. *Clin Exp Metastasis* 1: 103-114, 1983.
28. Wang ZL, Fan ZQ, Jiang HD and Qu JM: Selective Cox-2 inhibitor celecoxib induces epithelial-mesenchymal transition in human lung cancer cells via activating MEK-ERK signaling. *Carcinogenesis* 34: 638-646, 2013.
29. Kozak KR, Milne GL, Bentzen SM and Yock TI: Elevation of prostaglandin E₂ in lung cancer patients with digital clubbing. *J Thorac Oncol* 7: 1877-1878, 2012.
30. Sorenson S, Fohlin H, Lindgren A, *et al*: Predictive role of plasma vascular endothelial growth factor for the effect of celecoxib in advanced non-small cell lung cancer treated with chemotherapy. *Eur J Cancer* 49: 115-120, 2013.
31. Lynch TJ, Bell DW, Sordella R, *et al*: Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib. *N Engl J Med* 350: 2129-2139, 2004.
32. Soda M, Choi YL, Enomoto M, *et al*: Identification of the transforming EML4-ALK fusion gene in non-small-cell lung cancer. *Nature* 448: 561-566, 2007.
33. Cebola I and Peinado MA: Epigenetic deregulation of the COX pathway in cancer. *Prog Lipid Res* 51: 301-313, 2012.
34. de Maat MF, van de Velde CJ, Umetani N, *et al*: Epigenetic silencing of cyclooxygenase-2 affects clinical outcome in gastric cancer. *J Clin Oncol* 25: 4887-4894, 2007.
35. Suzuki R, Yamamoto M, Saka H, *et al*: A phase II study of carboplatin and paclitacel with meloxicam. *Lung Cancer* 63: 72-76, 2009.
36. O'Donnell PH and Dolan ME: Cancer pharmacoethnicity: ethnic differences in susceptibility to the effects of chemotherapy. *Clin Cancer Res* 15: 4806-4814, 2009.



Contents lists available at ScienceDirect

Respiratory Investigation

journal homepage: www.elsevier.com/locate/resinv

Original article

Phase II study of amrubicin combined with carboplatin for refractory relapsed small-cell lung cancer: North Japan Lung Cancer Group Trial 0802



Yosuke Kawashima^a, Akira Inoue^{b,*}, Shunichi Sugawara^a, Satoshi Oizumi^c, Makoto Maemondo^d, Koichi Okudera^e, Toshiro Suzuki^f, Kazuhiro Usui^g, Masao Harada^h, Naoto Morikawaⁱ, Yukihiro Hasegawa^j, Ryota Saito^b, Osamu Ishimoto^a, Tomohiro Sakakibara^b, Hajime Asahina^c, Toshihiro Nukiwa^k

ARTICLE INFO

Article history:

Received 10 June 2013

Received in revised form

17 October 2013

Accepted 9 December 2013

Available online 28 January 2014

Keywords:

Amrubicin

Small-cell lung cancer

Refractory relapse

Phase II trial

ABSTRACT

Background: Amrubicin (AMR), a new anthracycline agent, has shown promising results for advanced small-cell lung cancer (SCLC), although the efficacy of AMR alone against refractory relapsed SCLC is insufficient. This study was conducted to evaluate the safety and efficacy of the combination of AMR and carboplatin (CBDCA) in patients with refractory relapsed SCLC.

Methods: Patients with advanced SCLC who relapsed within 90 days after the completion of first-line chemotherapy received AMR (30 mg/m², days 1–3) and CBDCA (area under the curve 4.0 mg mL⁻¹ min⁻¹, day 1) every 3 weeks. The primary endpoint of this study was the overall response rate (ORR), and the secondary endpoints were progression-free survival (PFS), overall survival, and the toxicity profile. Assuming that an ORR of 45% in eligible patients would indicate potential usefulness and an ORR of 20% would be the lower limit of interest, with $\alpha=0.10$ and $\beta=0.10$, at least 24 patients were required.

Results: Among 29 eligible patients, the ORR was 34% (90% confidence interval, 20–48). The median PFS was 3.5 months, whereas the median survival time was 7.3 months. The most common grade 3–4 toxicity was neutropenia (79%), although only one patient (3%) suffered from febrile neutropenia. Non-hematological toxicities were of moderate severity and no treatment-related death was observed.

Conclusions: This is the first prospective study of AMR combined with CBDCA for refractory relapsed SCLC, which was effective and well tolerated. However, further investigation of this regimen is warranted.

© 2014 The Japanese Respiratory Society. Published by Elsevier B.V. All rights reserved.

*Corresponding author. Tel.: +81 22 717 8539; fax: +81 22 717 8549.

E-mail addresses: "affiliation_string.xml"yousukekawashima3@yahoo.co.jp (Y. Kawashima), akinoue@idac.tohoku.ac.jp (A. Inoue), swara357@cat-v.ne.jp (S. Sugawara), soizumi@med.hokudai.ac.jp (S. Oizumi), maemondo-ma693@miyagi-pho.jp (M. Maemondo), hppokudera@gmail.com (K. Okudera), toshi@isawa-hosp.mizusawa.iwate.jp (T. Suzuki), usui@east.ntt.co.jp (K. Usui), mharada@sap-cc.go.jp (M. Harada), carcinoma@nifty.com (N. Morikawa), yukihiro_hasegawa@med.pref.aomori.jp (Y. Hasegawa), beambitious0529@yahoo.co.jp (R. Saito), ishimoto@crest.ocn.ne.jp (O. Ishimoto), sakatomo@idac.tohoku.ac.jp (T. Sakakibara), asahinah@huhp.hokudai.ac.jp (H. Asahina), toshinkw47@gmail.com (T. Nukiwa).

^aDepartment of Pulmonary Medicine, Sendai Kousei Hospital, 4-15 Hirose-machi, Aoba-ku, Sendai 980-0873, Japan

^bDepartment of Respiratory Medicine, Tohoku University Hospital, 1-1 Seiryu-machi, Aoba-ku, Sendai 980-8574, Japan

^cFirst Department of Medicine, Hokkaido University School of Medicine, Kita 15-jo Nishi 7-chome, Kita-ku, Sapporo 060-8638, Japan

^dDepartment of Respiratory Medicine, Miyagi Cancer Center, 47-1 Nodayama, Medeshima-shiote, Natori 981-1293, Japan

^eDepartment of Respiratory Medicine, Hirosaki Central Hospital, 3-1 Yoshino-machi, Hirosaki 036-8188, Japan

^fDepartment of Respiratory Medicine, Isawa Hospital, 61 Ryugababa, Mizusawa-ku, Oshu 023-0864, Japan

^gDivision of Respiratory, NTT Medical Center Tokyo, 5-9-22 Higashi Gotanda, Shinagawa-ku, Tokyo 141-8625, Japan

^hDepartment of Respiratory Medicine, Hokkaido Cancer Center, 2-3-54 Kikusui 4-jo, Shiroishi-ku, Sapporo 003-0804, Japan

ⁱDivision of Pulmonary Medicine, Allergy, and Rheumatology, Department of Internal Medicine, Iwate Medical University School of Medicine, 19-1 Uchimarui, Morioka 020-8505, Japan

^jDepartment of Respiratory Medicine, Aomori Prefectural Central Hospital, 2-1-1 Higashitukurimichi, Aomori 030-8553, Japan

^kSouth Miyagi Medical Center, 38-1 Nishi, Ogawara, Shibata 989-1253, Japan

1. Introduction

Lung cancer is currently the leading cause of cancer death in many countries, and small-cell lung cancer (SCLC) accounts for 12–15% of all lung cancer cases [1]. SCLC is chemosensitive, and the standard first-line chemotherapy for advanced SCLC is platinum-doublet regimens such as cisplatin (CDDP) plus etoposide (ETP) or CDDP plus irinotecan (CPT) [2,3]. Despite high response rates to first-line chemotherapy, most patients experience SCLC relapse. The efficacy of second-line chemotherapy differ according to the relapse type (sensitive relapse, defined as relapse after >90 days from the completion of first-line chemotherapy or refractory relapse, defined as relapse during first-line chemotherapy or within 90 days after completion of first-line chemotherapy). There has been no standard treatment for patients with refractory relapsed SCLC, and few single agents have shown a response rate of >10% [4].

Amrubicin (AMR), a new anthracycline agent, has shown some promising results for advanced SCLC. A Japanese phase II study of the intravenous administration of single-agent first-line AMR therapy (45 mg/m²) for 3 consecutive days demonstrated a high overall response rate (ORR) (75.8%) and long median survival time (MST) (11.7 months) [5]. AMR was also more effective than topotecan (TOP) for chemosensitive relapsed SCLC in our previous phase II trial (response rates, 38% and 13%, respectively), although the response rate of AMR for refractory relapsed SCLC was only 17% (that of TOP was 0%) [6], a finding compatible with the result of AMR in a similar population in a subsequent large phase II study by Ettinger [7].

Since some of the patients with refractory relapsed SCLC did not receive a sufficient dose of platinum agent during first-line chemotherapy, we thought that second-line chemotherapy consisting of AMR combined with platinum might be worth investigating. Thus, we conducted this phase II study to evaluate the safety and efficacy of the combination of AMR and CBDCA in patients with refractory relapsed SCLC.

2. Patients and methods

2.1. Patient selection

This multicenter phase II trial was conducted in accordance with the principles outlined in the Helsinki Declaration of the World

Medical Association, and the protocol was approved by the institutional review board of each participating institution (Approval date: December 15, 2008; Approved No: 2008-365). Patients >20 years of age with histologically or cytologically confirmed SCLC who had progressed during first-line chemotherapy or had relapsed within 90 days after the completion of first-line chemotherapy were enrolled in this study. Other eligibility criteria included an Eastern Cooperative Oncology Group performance status (PS) of 0–2, measurable lesions according to Response Evaluation Criteria in Solid Tumors (RECIST), an estimated life expectancy \geq 3 months, and adequate organ function (white blood cell count \geq 4000/mm³, absolute neutrophil count \geq 2000/mm³, platelet count \geq 100,000/mm³, hemoglobin \geq 9.0 g/dL, serum bilirubin \leq 1.5 mg/dL, aspartate aminotransferase and alanine aminotransferase \leq 100 IU/L, creatinine level \leq 1.5 mg/dL, and arterial oxygen pressure \geq 60 mmHg). Written informed consent was obtained from all enrolled patients. Patients with symptomatic brain metastasis, interstitial lung disease, massive effusion requiring drainage, or severe comorbidities such as uncontrolled diabetes or cardiac disease were excluded. This trial was registered at UMIN (ID: R000001597).

2.2. Treatment schedule

The AMR was diluted in 50 mL of normal saline and administered by 10-min intravenous infusion at a dose of 30 mg/m² on days 1–3 of each treatment cycle. CBDCA was diluted in 250 mL of 5% glucose solution or normal saline and administered at infusion intervals of \geq 30 min at a dose of area under the curve (AUC) 4.0 mg mL⁻¹ min⁻¹ after AMR on day 1. The doses of both agents were determined according to our previous phase I study of this combination for patients with untreated SCLC [8]. The treatment was repeated every 21 days. Premedication with corticosteroids and an antiemetic 5-HT₃ antagonist was recommended. The dose of AMR was reduced by 5 mg/m² each in the subsequent cycle in cases of severe toxic effects such as grade 3 or more non-hematological toxicities, thrombocytopenia \leq 20,000/mm³, grade 4 neutropenia lasting \geq 4 days, or febrile neutropenia in the previous cycle. Use of granulocyte colony-stimulating factor (G-CSF) was permitted for neutropenia but not for prophylaxis. No prophylactic antibiotic support was planned. All patients were scheduled to receive at least three cycles of treatment unless their disease progressed, unacceptable toxicity occurred, the patient refused further treatment, or the physician

decided to discontinue the treatment. Subsequent chemotherapy after disease progression was not limited.

2.3. Patient assessment

Patient assessments, including a physical examination, a complete blood count, and biochemistry analysis, were repeated once a week after the initial evaluation. Tumor measurement was performed during the baseline assessment by computed tomography (CT) and was repeated every month until the best response to the protocol treatment was identified. Complete response (CR), partial response (PR), stable disease (SD), and progressive disease (PD) were determined based on RECIST version 1.0. CR and PR were confirmed by re-assessment performed at least 4 weeks after the first observation. SD was confirmed by re-assessment performed at least 6 weeks after registration. After confirmation, CT scans were acquired every 2 months until PD was observed. The CT scans of all patients were extramurally reviewed to confirm the response and progression-free survival (PFS). PFS was defined as the time from the date of registration to the date of the first observation of PD or death. Overall survival (OS) was defined as the time from the date of registration to the date of death or the latest follow-up (censored case). Toxicities were evaluated according to Common Terminology Criteria for Adverse Events version 3.0.

2.4. Statistical analysis

The primary endpoint of this study was the overall response rate (ORR), and secondary endpoints were PFS, OS, and the toxicity profile. Assuming that an ORR of 45% in eligible patients would indicate potential usefulness while an ORR of 20% would be the lower limit of interest, with $\alpha=0.10$ and $\beta=0.10$, at least 24 patients were required. Survival estimation was performed using the Kaplan–Meier method.

3. Results

3.1. Patient characteristics and treatment delivery

Between September 2008 and May 2011, 30 patients were enrolled from 10 institutions. One patient was excluded because of ineligible histology. Most of patients were male with a good PS (Table 1). Most patients received a CBDCA-based regimen as first-line chemotherapy, with a median of 4 cycles (range, 2–11 cycles). The median number of treatment cycles in the current study was 4 (range, 1–7), and 83% (24 of 29) of patients received three or more cycles.

3.2. Efficacy

All 29 patients were evaluable for response. The ORR was 34% (90% confidence interval, 20–48) and the disease-control rate was 83% (Table 2). The response rate of patients treated with CBDCA-based first-line chemotherapy was 40%, whereas that of patients treated with CDDP-based first-line chemotherapy was 22%, although the difference was not statistically significant. The response rates of patients treated with ETP and

Table 1 – Patient characteristics.

Number of patients	29
Gender	
Male	26
Female	3
Age (years)	
Median	67
Range	50–81
Performance status	
0	9
1	16
2	4
Prior chemotherapy	
Cisplatin+etoposide	2
Carboplatin+etoposide	15
Cisplatin+irinotecan	7
Carboplatin+irinotecan	5

Table 2 – Response.

Response	Number of patients	%	90% CI
Complete response	0	0	
Partial response	10	34	
Stable disease	14	48	
Progressive disease	5	17	
Overall response rate	10	34	20–48
Disease control rate	24	83	

CI, confidence interval.

of those treated with CPT as first-line chemotherapy were 35% and 33%, respectively. At the data cut-off point in September 2013, the median PFS was 3.5 months and the median survival time was 7.3 months (Fig. 1).

3.3. Safety

The toxicities (>grade 2) are summarized in Table 3. The most common adverse event in this study was neutropenia (79%), although only one patient (3%) suffered from febrile neutropenia. Thirteen patients (45%) required G-CSF support, the median duration of which was 4 days (range, 1–11). Two patients (7%) received a blood transfusion. Eight patients (28%) required AMR dose reduction due to hematological toxicity. Non-hematological toxicities were moderate. One patient died only 5 days after the initiation of protocol treatment. The attending physician reported that the cause of death was rapid progression of SCLC, and the independent data and safety monitoring committee of this study reviewed the clinical course and accepted the physician's decision. No treatment-related death was observed.

4. Discussion

This study met its primary endpoint. Since there have been few promising monotherapy options for refractory relapsed




Fig. 1 – (A) Progression-free survival and (B) overall survival.

SCLC, the combination of AMR and CBDCA is worth investigating. Contrary to our expectations, most patients in this study received sufficient cycles of platinum-doublet therapy as first-line chemotherapy. The ORR might have increased if more patients had been treated with insufficient first-line chemotherapy. According to subgroup analysis, this regimen might be suitable for patients treated with CBDCA as first-line chemotherapy. The efficacy of CBDCA plus AMR was not different in patients treated with ETP or CPT as first-line chemotherapy with platinum, which was consistent with our previous result of AMR as second-line chemotherapy [6]. Although the sample size was too small, the above-mentioned results require further validation.

In another Japanese study, even AMR alone demonstrated a quite high response rate (40%) in refractory relapsed SCLC [9], although the result might be biased due to its small sample size ($n=16$), considering the result of a subsequent larger study [7]. Other studies have used combined regimens for relapsed SCLC, some of which suggested high efficacy. However, most of those studies included both sensitive and refractory relapse patterns [4]; thus, their usefulness in refractory relapsed SCLC was unclear.

Toxicity is another important issue for such combination regimens. The above-mentioned previous regimens for relapsed SCLC were generally very toxic. For example, Kubota reported that dose-intensive CODE (CDDP, vincristine, doxorubicin, and ETP) could result in an ORR of approximately 80% in patients with refractory relapsed SCLC; however, that regimen required prophylactic G-CSF support due to severe

Table 3 – Toxicity profile.

Toxicity (\geq grade 2)	Grade (CTCAE)			Grade 3/4 (%)
	Number of patients			
	2	3	4	
Hematological				
Neutropenia	0	10	13	23 (79%)
Decreased hemoglobin	11	6	1	7 (24%)
Thrombocytopenia	6	4	3	7 (24%)
Febrile neutropenia	–	1	0	1 (3%)
Non-hematological				
Infection	4	2	0	2 (6%)
Nausea	2	0	0	0
Fatigue	1	0	0	0
Mucositis oral	1	0	0	1 (3%)
Stomach pain	1	0	0	0
Phlebitis	1	0	0	0
Hiccups	1	0	0	0
Pain	1	0	0	0
Interstitial lung disease	0	1	0	1 (3%)
Hyponatremia	0	2	0	2 (6%)
Hypoglycemia	0	0	1	1 (3%)

CTCAE, Common terminology criteria for adverse events.

neutropenia [10]. In contrast, AMR combined with CBDCA showed moderate toxicity in this study, which might be attributable to the dose of CBDCA being AUC 4. We reported this regimen in another study, where toxicity profiles tended to be similar and the efficacy for SCLC was sufficient (ORR was 89% as first-line treatment) [11]. Regarding the AMR dose, the current dose was one level lower than the recommended dose in our phase I and phase II studies of patients with chemotherapy-naïve SCLC because we considered that previously treated patients would be at a higher risk of myelosuppression. Although we believe this combination with the current dosage would be worth investigating in the second-line setting in terms of the risk–benefit balance, there might be scope for increasing the AMR dose to increase its efficacy.

This study has a few limitations. First, the sample size was too small to draw definite conclusions, the efficacy of this combination needs to be confirmed in a future phase III study in which the current regimen could be compared with AMR alone. Second, the drug dose might be insufficient for refractory relapsed cases. Considering that the toxicity of the current dose was moderate, there might be scope to increase the CBDCA or AMR dosage. In addition, the patients that would benefit most from the re-administration of platinum during second-line chemotherapy should be identified.

In conclusion, AMR combined with CBDCA was effective for refractory relapsed SCLC and demonstrated acceptable toxicity. Since treatment options for patients with refractory relapsed SCLC remain limited, further investigation of this regimen is warranted.

Conflict of interest

Akira Inoue received honoraria and research funding from AstraZeneca; Satoshi Oizumi received honoraria from AstraZeneca and research funding from Eli Lilly; Toshihiro Nukiwa received honoraria from Boehringer Ingelheim.

REFERENCES

- [1] Govindan R, Page N, Morgensztern D, et al. Changing epidemiology of small-cell lung cancer in the United States over the last 30 years: analysis of the surveillance, epidemiologic, and end results database. *J Clin Oncol* 2006;24:4539-44.
- [2] Roth BJ, Johnson DH, Einhorn LH, et al. Randomized study of cyclophosphamide, doxorubicin, and vincristine versus etoposide and cisplatin versus alternation of these two regimens in extensive small-cell lung cancer: a phase III trial of the Southeastern Cancer Study Group. *J Clin Oncol* 1992;10:282-91.
- [3] Noda K, Nishiwaki Y, Kawahara M, et al. Irinotecan plus cisplatin compared with etoposide plus cisplatin for extensive small-cell lung cancer. *N Engl J Med* 2002;346:85-91.
- [4] Tiseo M, Ardizzoni A. Current status of second-line treatment and novel therapies for small cell lung cancer. *J Thorac Oncol* 2007;2:764-72.
- [5] Yana T, Neguro S, Takada M, et al. Phase II study of amrubicin in previously untreated patients with extensive-disease small cell lung cancer: West Japan Thoracic Oncology Group (WJTOG) study. *Invest New Drugs* 2007;25:253-8.
- [6] Inoue A, Sugawara S, Yamazaki K, et al. Randomized phase II trial comparing amrubicin with topotecan in patients with previously treated small-cell lung cancer: North Japan Lung Cancer Study Group Trial 0402. *J Clin Oncol* 2008;26:5401-6.
- [7] Ettinger DS, Jotte R, Lorigan P, et al. Phase II study of amrubicin as second-line therapy in patients with platinum-refractory small-cell lung cancer. *J Clin Oncol* 2010;28:2598-603.
- [8] Inoue A, Yamazaki K, Maemondo M, et al. A phase I study of amrubicin combined with carboplatin for elderly patients with small-cell lung cancer. *J Thorac Oncol* 2006;1:551-5.
- [9] Onoda S, Masuda N, Seto T, et al. Phase II trial of amrubicin for treatment of refractory or relapsed small-cell lung cancer: Thoracic Oncology Research Group Study 0301. *J Clin Oncol* 2006;24:5448-53.
- [10] Kubota K, Nishiwaki Y, Kakinuma R, et al. Dose-intensive weekly chemotherapy for treatment of relapsed small-cell lung cancer. *J Clin Oncol* 1997;15:292-6.
- [11] Inoue A, Ishimoto O, Fukumoto S, et al. A phase II study of amrubicin combined with carboplatin for elderly patients with small-cell lung cancer: North Japan Lung Cancer Study Group Trial 0405. *Ann Oncol* 2010;21:800-3.

Phase II trial of carboplatin and pemetrexed as first-line chemotherapy for non-squamous non-small cell lung cancer, and correlation between the efficacy/toxicity and genetic polymorphisms associated with pemetrexed metabolism: Hokkaido Lung Cancer Clinical Study Group Trial (HOT) 0902

Kenya Kanazawa · Hiroshi Yokouchi · Xintao Wang · Takashi Ishida · Yuka Fujita · Satoru Fujiuchi · Toshiyuki Harada · Masao Harada · Kei Takamura · Satoshi Oizumi · Ichiro Kinoshita · Yutaka Katsuura · Osamu Honjo · Tetsuya Kojima · Hirotohi Dosaka-Akita · Hiroshi Isobe · Mitsuru Munakata · Masaharu Nishimura · Hokkaido Lung Cancer Clinical Study Group

Received: 8 May 2014 / Accepted: 8 September 2014 / Published online: 8 October 2014
© Springer-Verlag Berlin Heidelberg 2014

Abstract

Purpose This phase II study evaluated the response rate (RR) and safety of combination therapy with carboplatin (CBDCA) and pemetrexed (PEM) in Japanese patients with non-squamous non-small cell lung cancer (non-sq NSCLC). Further, the relationship between therapy efficacy/toxicity and genetic polymorphisms associated with PEM metabolism was analyzed.

Methods Forty-one patients received CBDCA at a dose targeting an area under the concentration–time curve of

5 mg/mL × min and PEM of 500 mg/m² on day 1 every 3 weeks. Single-nucleotide polymorphisms of the thymidylate synthase (*TYMS*) coding gene, the variable number of tandem repeat (VNTR) in the *TYMS*, and the methyl-ene-tetrahydrofolate reductase (*MTHFR*) coding gene were analyzed.

Results The overall RR was 36.6 %. Median progression-free survival and median survival time were 4.7 months [95 % confidence interval (CI) 3.9–5.6 months] and 16.2 months (95 % CI 6.1–26.2 months), respectively.

K. Kanazawa (✉) · H. Yokouchi · X. Wang · T. Ishida · M. Munakata
Department of Pulmonary Medicine, Fukushima Medical University School of Medicine, 1 Hikarigaoka, Fukushima 960-1295, Japan
e-mail: k-kenya@fmu.ac.jp

T. Ishida
Clinical Oncology Center, Fukushima Medical University Hospital, 1 Hikarigaoka, Fukushima 960-1295, Japan

Y. Fujita · S. Fujiuchi
Department of Respiratory Medicine, National Hospital Organization Asahikawa Medical Center, 7-4048 Hanasaki-cho, Asahikawa, Hokkaido 070-8644, Japan

T. Harada
Center for Respiratory Disease, Japan Community Health Care Organization Hokkaido Hospital, 1-8-3-18 Nakanoshima, Toyohira-ku, Sapporo, Hokkaido 062-8618, Japan

M. Harada
Department of Respiratory Medicine, National Hospital Organization Hokkaido Cancer Center, 3-54, Kikusui 4-jo 2-chome, Shiroishi-ku, Sapporo, Hokkaido 003-0804, Japan

K. Takamura
First Department of Medicine, Hokkaido P. W. F. A. C. Obihiro-Kosei General Hospital, 1-West-8-South-6, Obihiro, Hokkaido 080-0016, Japan

S. Oizumi · M. Nishimura
First Department of Medicine, Hokkaido University School of Medicine, Kita 8, Nishi 5, Kita-ku, Sapporo, Hokkaido 060-0808, Japan

I. Kinoshita · H. Dosaka-Akita
Department of Medical Oncology, Hokkaido University Graduate School of Medicine, Kita 8, Nishi 5, Kita-ku, Sapporo, Hokkaido 060-0808, Japan

Y. Katsuura
Department of Pulmonary Medicine, Saiseikai Fukushima General Hospital, 4-15 Sakuragi-cho, Fukushima 960-8133, Japan

O. Honjo
Department of Respiratory Medicine, Hokkaido P. W. F. A. C. Sapporo-Kosei General Hospital, 5, Kita 3-jo Higashi 8-chome, Chuo-ku, Sapporo, Hokkaido 003-0804, Japan

Epidermal growth factor receptor gene mutations were detected in 6 patients (14.6 %). The VNTR in the *TYMS* significantly correlated with anemia ($p = 0.047$) and thrombocytopenia ($p = 0.038$).

Conclusions This combination therapy was effective and tolerable in patients with advanced non-sq NSCLC. The VNTR in the *TYMS* appears to be a predictive factor for anemia and thrombocytopenia in patients treated with this regimen.

Keywords Non-squamous non-small cell lung cancer · Carboplatin · Pemetrexed · Genetic polymorphisms · Methylenetetrahydrofolate reductase (MTHFR) · Thymidylate synthase (TS)

Introduction

The majority of patients with non-small cell lung cancer (NSCLC) are diagnosed at inoperable stages, and platinum-based chemotherapy remains a key strategy for the management of patients with advanced NSCLC [1–3]. Overall survival of front-line chemotherapy using pemetrexed (PEM) in combination with cisplatin (CDDP) was superior to that with other platinum doublets, particularly for patients with advanced non-squamous (non-sq) NSCLC [4–7].

PEM-based regimens have a mild toxicity profile and can improve patients' quality of life (QOL) [4, 6, 8, 9]. Further, carboplatin (CBDCA)-based regimens are also widely used and are associated with relatively mild toxicity [9–11]. However, the efficacy and safety of PEM combined with CBDCA have not been well established in Japanese patients with NSCLC.

Thymidylate synthetase (TS) is one of the main targets of PEM [12], and methylenetetrahydrofolate reductase (MTHFR) is an enzyme indispensable for folate metabolism. Both enzymes are strongly associated with cell proliferation and efficacy of pyrimidine-antagonist chemotherapies, such as the one with PEM [13, 14], and studies have demonstrated a correlation between clinical efficacy of various anticancer agents and the polymorphisms of these genes [15]. Patients with homozygous mutations for the MTHFR coding gene (*MTHFR*) C677T had a significantly increased progression-free survival (PFS) when compared to patients with wild-type or heterozygous mutations [16].

Similarly, TS is a critical target for various chemotherapies [17, 18], including those used for the treatment of NSCLC. One study reported that TS expression correlated with PEM sensitivity in NSCLC cell lines [19]. Tanaka et al. [20] conducted a large-scale study of the Japanese population showing that TS expression was lower in adenocarcinoma than in squamous cell carcinoma of the lung.

Thus, the goal of the present phase II study was to evaluate the response rate (RR) and safety of CBDCA and PEM in Japanese patients with non-sq NSCLC. Further, the relationship between therapy efficacy/toxicity and genetic polymorphisms of folate metabolism-associated enzyme coding genes, the TS coding gene (*TYMS*) and *MTHFR*, or the variable number of tandem repeat (VNTR) in the *TYMS* in peripheral blood cells was examined.

Patients and methods

Eligibility criteria

Eligibility criteria were as follows: cytologically or histologically confirmed diagnosis of non-sq NSCLC; patients without prior systemic chemotherapy, including ones with postoperative recurrence; stage IIIB or IV disease according to the 7th edition of TNM criteria; no indications for curative chemoradiotherapy; age between 20 and 74 years; Eastern Cooperative Oncology Group (ECOG) performance status 0–1; measurable disease (Response Evaluation Criteria in Solid Tumors, ver. 1.0); and normal organ function (as defined by absolute white blood cell count $\geq 4.0 \times 10^9/L$ or neutrophil count $\geq 2.0 \times 10^9/L$; hemoglobin ≥ 9.0 g/dL; platelets $\geq 100 \times 10^9/L$; alanine aminotransferase [ALT] and aspartate aminotransferase [AST] ≤ 100 IU/L [ALT and AST ≤ 150 IU/L was acceptable if liver metastasis was present]; serum creatinine ≤ 1.2 mg/dL; calculated creatinine clearance using Cockcroft-Gault formula ≥ 60 ml/min; arterial partial pressure of oxygen [PaO_2] ≥ 60 Torr or arterial hemoglobin oxygen saturation by pulse oximetry [SpO_2] ≥ 90 % at ambient air); and projected life expectancy ≥ 12 weeks. The main exclusion criteria were as follows: active infection; temperature ≥ 38 °C; severe complications, such as heart failure, renal failure, liver dysfunction, uncontrolled diabetes mellitus, and hypertension; a concomitant malignancy within the last 5 years; central nervous system metastases with symptoms; uncontrolled pleural effusion or ascites; interstitial pneumonia or pulmonary fibrosis on chest X-ray; history of severe hypersensitivity to drug components; required concurrent treatment with systemic steroid; and pregnancy. The protocol was approved by each institutional review board. The study was performed in accordance with the ethics principles of the Declaration of Helsinki. Written informed

T. Kojima · H. Isobe
Department of Medical Oncology and Respiratory Medicine,
KKR Sapporo Medical Center, 3-40, Hiragishi 1-jo 6-chome,
Toyohira-ku, Sapporo, Hokkaido 062-0931, Japan

Hokkaido Lung Cancer Clinical Study Group
Sapporo, Hokkaido, Japan

consent was obtained from all patients. This study was registered with the University Hospital Medical Information Network (UMIN) [UMIN000002846].

Treatment

PEM at a dose of 500 mg/m² on day 1 and CBDCA at a dose calculated to produce an area under the concentration–time curve (AUC) of 5.0 mg/mL × min on day 1 were administered intravenously every 3 weeks. The treatment was discontinued in the case of any of the following: disease progression; unacceptable toxicity; patient refusal; death during treatment; and investigator's decision. All patients received oral folic acid (500 μg) daily and a vitamin B12 injection (1,000 μg) every 9 weeks, beginning one or more weeks before the first dose and continuing until three weeks after the last dose of study treatment. Any treatment was permitted after protocol discontinuation. Dose adjustment and cycle delay of 21 days or less were permitted to allow for resolution of toxic effects.

Assessment of toxicity and response

Toxicities or adverse events were graded according to the Common Terminology Criteria for Adverse Events (CTCAE), version 3.0. Tumor responses were assessed using chest X-ray, computed tomography, or magnetic resonance imaging (when clinically indicated), before and during treatment. Assessments were repeated at least every month unless progression was detected. Responses were recorded as complete response (CR), partial response (PR), stable disease (SD), or progressive disease (PD) in accordance with Response Evaluation Criteria in Solid Tumors (RECIST) guidelines version 1.0. Disease control rate (DCR) was defined as the sum of the objective response (CR or PR) rate and the rate of SD. Clinical response data were all confirmed by central review.

Endpoints and statistical analysis

The primary endpoint was overall RR. Secondary endpoints were safety, PFS, and overall survival (OS). Duration of tumor response was defined as the time between the date of the first objective assessment of CR or PR and the date when PD or death was recorded from any cause. PFS was defined as the time between the date of registration and the date of PD or death from any cause. OS was defined as the time between the date of registration and the date of death from any cause. The Kaplan–Meier method was used to estimate PFS and OS.

On the assumption that threshold RR and expected RR would, respectively, be 20 and 40 %, a sample size of 36 patients was required by the Simon's and Fleming's

designs with a one-sided α error of 0.05 and a β error of 0.20. A total of 40 patients were planned to enroll considering later exclusion of patients. All analyses were based on the intent-to-treat population.

Genetic analyses

Analyses of genetic variants were performed with the investigator blinded to patient characteristics and clinical outcome. Five milliliters of peripheral blood was taken from each patient who had enrolled in this study and who had consented to the genetic analysis. DNA was extracted from each blood sample for analysis of the *MTHFR* single-nucleotide polymorphisms (SNPs), C677T and A1298C. Then, each extracted DNA was used to determine the TS genotypes of the VNTR in the five prime untranslated region (5'-UTR) of *TYMS*, two tandem repeat (2R)/3R/4R, and the 3R-SNP, G/C, in *TYMS* by polymerase chain reaction (PCR) and PCR-restriction fragment length polymorphism (PCR–RFLP), using the forward primer, 5'-AAAAG GCGCGCGGAAGGGGTCCT-3', and reverse primer, 5'-TCCGAGCCGCCCACAGGCAT-3'. DNA amplification was performed three times per sample, using the GeneAmp[®] PCR System 9700 (Applied Biosystems, Life Technologies Japan, Inc., Tokyo, Japan). PCR with the genomic DNA template was performed in reaction mixtures containing 1 × PCR buffer II without MgCl₂, MgCl₂ solution, 200 μM of deoxyribonucleoside triphosphates, 500 nM of each primer, 0.5 units of AmpliTaq[®] DNA Polymerase, and 100 ng of genomic DNA (all of these reagents were obtained from Applied Biosystems). The cycling conditions were one cycle of 94 °C for 5 min, 35 cycles of 94 °C for 40 s, 62 °C for 60 s, 72 °C for 40 s, with a final extension at 72 °C for 5 min. Aliquots of amplified fragments were separated on 4 % agarose gels, and the TS VNTR genotype was determined, with 2R 116 bp, 3R 144 bp, and 4R 172 bp. Samples showing the 3R genotype were analyzed further for G/C polymorphism by RFLP. PCR products were digested with *Hae*III (TaKaRa Bio, Inc., Shiga, Japan) followed by electrophoresis in 4 % agarose gel and ethidium bromide stain. The 3R fragments of 66, 37, 28, and 10 bp were classified into the 3G allele, and the 3R fragments of 94, 37, and 10 bp were classified into the 3C allele, as previously reported [21]. Analysis was performed at least three times to confirm the genotype. *TYMS* genotype was categorized into a high-expression genotype (2R/3G, 3C/3G, 3G/3G, 3G/4R) and a low-expression genotype (2R/2R, 2R/3C, 3C/3C), depending on the 5'-UTR VNTR polymorphism and the C/G polymorphism within the third VNTR.

The association between polymorphisms and RR or chemotherapy-related toxicity was analyzed by the χ^2 test or Fisher's exact test. The correlation between

polymorphisms and PFS/OS was analyzed by the log-rank test. For each test, patients were compared among each genotype, such as wild type, heterozygous, and homozygous. Statistical significance was established at $p < 0.05$.

Results

Patients characteristics

From November 2009 to November 2010, 41 patients were enrolled (Table 1). Twenty-four patients (58.5 %) died during the follow-up, mostly as a result of disease progression (23 of 24 patients). The median age of the enrolled patients was 63 years (range of 43–73 years), and 28 patients (68.3 %) were male. Of the 41 patients, 27 (65.9 %) had an ECOG performance status of 1, 36 (87.8 %) had stage IV disease, and 39 (95.1 %) were diagnosed with adenocarcinoma.

Epidermal growth factor receptor (EGFR) gene mutations were investigated in 40 of 41 patients. The mutation was not searched for one patient because diagnostic yield from the tissue sample for detection was not enough. The positive mutations were detected in six patients, four of whom had an exon 19 deletion mutation, while the other two had an L858R mutation in exon 21.

Treatment administered

The median number of treatment cycles delivered was four (range of 1–6 cycles), with 33 patients (80.5 %) completing more than three cycles.

The dose of agents was reduced in three patients (7.3 %) because of adverse events, including grade 3 general fatigue, grade 4 hematologic toxicity, and grade 4 anaphylaxis. Protocol treatment was terminated in eight patients (19.5 %) before completion of three cycles.

At the time of final analysis, 23 patients (56.2 %) received second- or third-line treatment, following the initial therapy. Sixteen patients (39.0 %) received cytotoxic chemotherapies as a second-line treatment and 3 of them received EGFR tyrosine kinase inhibitors (EGFR-TKIs) as a third-line treatment. Seven patients (17.1 %) received EGFR-TKIs as a second-line treatment and 2 of them received cytotoxic chemotherapies as a third-line treatment. Among the 10 patients (24.4 %) who were treated with an EGFR-TKI in the second- or third-line, five had an EGFR gene mutation. One patient with EGFR gene mutation did not receive an EGFR-TKI.

Response to treatment

All 41 patients were assessable for tumor responses. Fifteen patients exhibited PR, 20 patients exhibited SD,

Table 1 Patient demographics and baseline characteristics

Parameters	(N = 41) n (%)
Age, median (years) (range)	
Gender	
Male	28 (68.3)
Female	13 (31.7)
ECOG performance status	
0	14 (34.1)
1	27 (65.9)
Disease stage	
IIIB	5 (12.2)
IV	36 (87.8)
Histology	
Adenocarcinoma	39 (95.1)
Large cell carcinoma	2 (4.9)
EGFR gene mutation	
Wild type	34 (82.9)
Exon 19 deletion	4 (9.8)
Exon 21 L858R	2 (4.9)
Unknown	1 (2.4)
No. of chemotherapy cycle	
1	1 (2.4)
2	7 (17.1)
3	6 (14.6)
4	11 (26.8)
5	3 (7.3)
6	13 (31.7)
≥3 cycles	33 (80.5)

Table 2 Treatment efficacy

Best response	(N = 41) n (%)
CR	0 (0.0)
PR	15 (36.6)
SD	20 (48.8)
PD	6 (14.6)
Overall response rate (RR)	36.6 % [95 % CI 22.1–53.1 %]
Disease control rate (DCR)*	85.4 % [95 % CI; 70.8–94.4 %]

CI confidence interval

* DCR = {CR + PR + SD}/(CR + PR + SD + PD)

and disease progressed in six cases, resulting in a RR of 36.6 % [95 % confidence interval (CI) 22.1–53.1 %] and a DCR of 85.4 % (95 % CI, 70.8–94.4 %) (Table 2). The lower limit of the 95 % CI of the RR exceeded the threshold RR of 20 %; thus, the primary endpoint was achieved.




Fig. 1 Kaplan–Meier curves for (a) progression-free survival (PFS) and (b) overall survival (OS)

The final survival assessment was conducted in July 2014, 4 years and 8 months after the last patient's enrollment. With a median follow-up time of 16.2 months (range of 2.4–54.1 months), median PFS and median survival time (MST) were 4.7 months (95 % CI 3.9–5.6 months) and 16.2 months (95 % CI 6.1–26.2 months), respectively (Fig. 1). The one-year survival rate was 53.6 % (95 % CI 37.4–69.3 %).

Retrospective subanalysis of PFS and OS of patients with or without an EGFR gene mutation showed that EGFR gene mutation status did not influence the PFS (a median of 5.9 months for the mutation-positive group, as compared to 4.6 months for the mutation-negative group; $p = 0.738$), but patients with an EGFR gene mutation were associated with a trend of longer OS than those without an EGFR gene mutation (a median of 16.2 months for the mutation-positive group, as compared to 11.7 months for the mutation-negative group; $p = 0.06$).

Adverse events

All 41 eligible patients were also evaluable for toxicity analysis. The rates of grade 3 and 4 hematologic and non-hematologic adverse events during the treatment were as follows: eight patients (19.5 %) had grade ≥ 3 leukopenia; 12 patients (29.3 %) had grade ≥ 3 neutropenia; 14 patients (34.1 %) had grade ≥ 3 anemia; seven patients (17.1 %) had grade ≥ 3 thrombocytopenia. Only one patient (2.4 %) experienced grade 3 febrile neutropenia. Granulocyte colony stimulating factors (G-CSF) were given to three patients (7.3 %). The most common non-hematologic adverse event was anorexia (29/41 patients, 70.7 %); other

Table 3 Toxicity profile

Toxicity	Grade, <i>n</i> (<i>N</i> = 41)				Percentage of grade 3 or 4 (%)
	1	2	3	4	
<i>Hematologic</i>					
Leukopenia	4	9	7	1	19.5
Neutropenia	3	8	9	3	29.3
Febrile neutropenia	–	–	1	0	2.4
Anemia	9	11	11	3	34.1
Thrombocytopenia	20	6	5	2	17.1
<i>Non-hematologic</i>					
Hypoalbuminemia	3	2	0	0	0.0
Liver dysfunction	16	2	1	0	2.4
High LDH	5	0	0	0	0.0
Hyponatremia	10	0	1	0	2.4
Hypopotassemia	1	2	0	0	0.0
Infection	0	2	1	0	2.4
Anorexia	23	3	3	0	7.3
Nausea	15	6	2	0	4.9
Vomiting	3	4	0	0	0.0
Allergic reaction	0	1	0	1	2.4
Stomatitis	2	0	0	0	0.0
Skin rash	1	1	1	0	2.4
Pigmentation	2	0	0	0	0.0
Alopecia	2	0	0	0	0.0
Fatigue	7	0	0	0	0.0
Constipation	2	1	0	0	0.0

non-hematologic adverse events were rare (Table 3). No interstitial lung disease was reported. No treatment-related death was observed.

Genetic polymorphisms and clinical indices

Blood samples were collected from 37 patients (90.2 %). Gene polymorphisms of C677T and A1298C in the *MTHFR* were in Hardy–Weinberg equilibrium according to Pearson's Chi-square test (C677T: $\chi^2 = 0.182 < \chi^2 [0.05] = 3.84$; A1298C: $\chi^2 = 0.946 < \chi^2 [0.05]$) (Table 4). Patients with 3R/3R and 3R/4R of the tandem repeat in 5'-UTR of *TYMS* had experienced significantly more grade 3/4 anemia ($p = 0.047$) or grade 3/4 thrombocytopenia ($p = 0.038$) than those with 2R/3R. Other variants explored in this study did not significantly correlate with hematologic toxicities (Table 5). Non-hematologic toxicities were mild, and the relationship between those events and gene variants was not evaluated. Gene variants did not significantly correlate with RR, DCR (Table 5), PFS, or OS (data not shown).

Discussion

Although CDDP and CBDCA have substantially different toxicity profiles, a meta-analysis comparing CDDP- and CBDCA-based chemotherapy failed to establish which regimen is associated with superior survival [5, 22–24]. CBDCA was used in this study, as non-hematologic toxicity and strong subjective symptoms (e.g., nausea, vomiting, and general fatigue) predominate in patients treated with CDDP, whereas hematologic toxicity with relatively less symptoms is observed more commonly in patients treated with CBDCA [20, 23, 24]. In a phase III study conducted in Norway, an AUC of 5.0 mg/mL \times min of CBDCA was selected for patients in the CBDCA plus PEM arm [9] and resulted in good clinical efficacy with a tolerable toxicity profile. We also adopted the same dose of CBDCA in this study, and the treatment efficacy of our CBDCA plus PEM regimen was favorable. Further, the toxicity profile was mild and tolerable. The PFS in this study seemed to be shorter than that in the other studies for Japanese patients [25, 26]. A continuation maintenance therapy with PEM was not adopted in our protocol due to the fact that the maintenance strategy [27] had not been established when our trial was launched. Compared to other Japanese phase II trials (from 5–6 cycles) [25, 26], there were fewer chemotherapy cycles in the current study, while Norwegian study reported even fewer (a median of 3.3 cycles) [9]. In our study, PD was the major cause (13 patients, 52 %) of undergoing 4 or less cycles of the chemotherapy. The dose of CBDCA may be associated with the fewer cycle of chemotherapy, providing another reason for the shorter PFS.

PEM targets folate-dependent reactions and acts on TS, a key enzyme for DNA synthesis [28, 29]. Alteration of TS activity due to polymorphisms in the cognate coding gene

Table 4 Number of cases with *MTHFR* and *TYMS* variants

Gene	Gene variants	(<i>N</i> = 41) <i>n</i> (%)
<i>MTHFR</i> -C677T	C/C	12 (32.4)
	C/T	17 (45.9)
	T/T	8 (21.6)
<i>MTHFR</i> -A1298C	A/A	26 (70.3)
	A/C	9 (24.3)
	C/C	2 (5.4)
<i>TYMS</i> -VNTR	2R/3R	7 (18.9)
	3R/3R	29 (78.4)
	3R/4R	1 (2.7)
SNP of the VNTR (3R)	3G (–)	2 (5.4)
	3G (+)	35 (94.6)

MTHFR methylenetetrahydrofolate reductase coding gene, *TYMS* thymidylate synthase coding gene, *VNTR* variable number of tandem repeat, 2R 2 repeats, 3R 3 repeats, 4R 4 repeats, *SNP* single-nucleotide polymorphism

Table 5 χ^2 test of correlation between the treatment efficacy/toxicities and SNPs of *MTHFR/TS* gene

Indexes	<i>MTHFR</i>		<i>TYMS</i>	
	C677T	A1298C	VNTR	SNP of the VNTR (3R)
	<i>p</i>	<i>p</i>	<i>p</i>	<i>p</i>
Response rate	0.757	0.307	0.699	0.902
Hematologic toxicities				
Leukopenia	0.703	0.393	0.372	0.149*
Neutropenia	0.565	0.684	0.194	0.220*
Anemia	0.237	0.217	0.047	0.237*
Thrombocytopenia	0.598	0.393	0.038	0.598*

VNTR variable number of tandem repeat

* Fisher's exact test

influences outcomes in patients with NSCLC [13, 30, 31]. *MTHFR* is an essential enzyme for one-carbon metabolism needed for DNA synthesis, repair, and methylation [32]. The alteration of *MTHFR* activity plays a role in carcinogenesis [33, 34], which supports the notion that *MTHFR* polymorphisms may affect patient outcomes [13, 15, 32]. Only an increasing repeat number of VNTR in 5'-UTR of *TYMS* correlated with anemia and thrombocytopenia, suggesting that this genetic marker might be useful for the prediction of hematologic toxicity in response to PEM. In order to avoid the severe hematologic toxicities in patients with the 3R/3R variant, we should consider switching the regimen, reducing the initial dose of PEM, or upwardly adjusting the minimum number of red blood cells or platelets required for starting the treatment.

A significant correlation between the *MTHFR*-C677T allele and improved clinical outcome has been found in another study [16]. However, we were not able to show the same result. One of the reasons appears to be race/ethnicity in those variants. In breast cancer, race/ethnicity has been reported to modify the association between the two SNPs of *MTHFR* and breast cancer survival [35]. It is known that PEM targets another enzyme associated with folate metabolism other than TS, such as dihydrofolate reductase (DHFR), which inhibits a cytotoxic effect of antifolates, thereby reducing treatment efficacy [36]. Indeed, a previous study showed the association between PFS and either of TS or of DHFR [37]. Collectively, a comprehensive analysis of polymorphisms of all enzymes associated with folate metabolism is required.

In terms of toxicity profile, the two Japanese studies of CBDCA (AUC = 6.0 mg/mL × min) plus PEM (500 mg/m²) followed by maintenance PEM (500 mg/m²) conducted by the Kyoto Thoracic Oncology Research Group (KTOGT0902) [25] and by Okamoto et al. [26], demonstrated that grade 3/4 neutropenia, anemia, and thrombocytopenia were seen in 33, 31, and 18 % and 56, 29.4, and 41.3 %, respectively. In the former study, red blood cell and platelet transfusions were required for 6.1 and 4.1 %, respectively. In our study, grade 3/4 neutropenia, anemia, and thrombocytopenia were seen, respectively, in 29.3, 34.1, and 17.1 %, whereas packed red blood cell transfusions were given to three patients (7.3 %) and platelet concentrate given to one patient (2.4 %). Non-hematologic toxicities were also mild, and our study achieved a good treatment completion rate over three courses. AUC of 5 or 6 mg/mL × min of CBDCA should be adjusted for individual patients in terms of the balance between efficacy and toxicity. In clinical practice, we assume that CBDCA plus PEM regimen can be selected for patients with non-squamous and EGFR wild-type NSCLC as a first-line therapy, particularly for those unfit for CDDP or with an ECOG performance status (PS) of two as demonstrated in a previous study [38] or elderly patients with good PS who have also a benefit of CBDCA-based platinum doublet [39, 40].

Recent phase III trials have shown favorable efficacies of EGFR-TKI in NSCLC patients with active EGFR gene mutation [41, 42]. There was no significant difference of either PFS or OS according to EGFR gene mutation status in our study, but OS of the patients with EGFR gene mutation tended to be longer than those without the mutation, presumably due to the post-treatment therapy using EGFR-TKIs. The smaller size of patients with EGFR mutation (6/40, 15 %) in our study was due to the fact that several institutes were also participating in another ongoing trial, which was recruiting NSCLC patients with EGFR gene mutation. However, our data implicate the importance of the use of EGFR-TKIs for treating patients with an active

EGFR gene mutation. CBDCA plus PEM regimen seems also appropriate for those with EGFR mutant-positive NSCLC, who failed the initial EGFR-TKI therapy.

Although the development of effective treatment strategies against advanced NSCLC has progressed swiftly over the last decade, minimization of toxicities remains important for QOL purposes. The toxicity profile of chemotherapy can vary according to race, ethnicity, and genetic makeup. Gene analyses in this study are the first to demonstrate a possibility of correlation between genetic polymorphisms and hematologic toxicity in non-squamous NSCLC patients treated with PEM. Further studies to confirm this evidence are warranted.

Acknowledgments We thank all the patients who participated in this study and also thank all institutions for recruiting patients.

Conflict of interest Satoshi Oizumi received an honorarium from AstraZeneca and a grant from Eli Lilly. Others had no conflict of interest to declare.

References

- Jemal A, Bray F, Center MM, Farley J, Ward E, Forman D (2011) Global cancer statistics. *CA Cancer J Clin* 61:69–90
- NSCLC Meta-Analyses Collaborative Group (2008) Chemotherapy in addition to supportive care improves survival in advanced non-small-cell lung cancer: a systematic review and meta-analysis of individual patient data from 16 randomized controlled trials. *J Clin Oncol* 26:4617–4625
- Hotta K, Fujiwara Y, Matsuo K, Suzuki T, Kiura K, Tabata M, Takigawa N, Ueoka H, Tanimoto M (2007) Recent improvement in the survival of patients with advanced non-small-cell lung cancer enrolled in phase III trials of first-line systemic chemotherapy. *Cancer* 109:939–948
- Scagliotti GV, Parikh P, von Pawel J, Biesma B, Vansteenkiste J, Manegold C, Serwatowski P, Gatzemeier U, Digumarti R, Zukin M, Lee JS, Mellemegaard A, Park K, Patil S, Rolski J, Goksel T, de Marinis F, Simms L, Sugarman KP, Gandara D (2008) Phase III study comparing cisplatin plus gemcitabine with cisplatin plus pemetrexed in chemotherapy-naïve patients with advanced-stage non-small-cell lung cancer. *J Clin Oncol* 26:3543–3551
- Manegold C, Gatzemeier U, von Pawel J, Pirker R, Malayeri R, Blatter J, Krejcy K (2000) Front-line treatment of advanced non-small-cell lung cancer with MTA (LY231514, pemetrexed disodium, ALIMTA) and cisplatin: A multicenter phase II trial. *Ann Oncol* 11:435–440
- Treat J, Scagliotti GV, Peng G, Monberg MJ, Obasaju CK, Socinski MA (2012) Comparison of pemetrexed plus cisplatin with other first-line doublets in advanced non-small cell lung cancer (NSCLC): a combined analysis of three phase 3 trials. *Lung Cancer* 76:222–227
- Al-Saleh K, Quinton C, Ellis PM (2012) Role of pemetrexed in advanced non-small-cell lung cancer: meta-analysis of randomized controlled trials, with histology subgroup analysis. *Curr Oncol* 19:e9–e15
- Shepherd FA, Dancey J, Arnold A, Neville A, Rusthoven J, Johnson RD, Fisher B, Eisenhauer E (2001) Phase II study of pemetrexed disodium, a multitargeted antifolate, and cisplatin as first-line therapy in patients with advanced nonsmall cell lung

- carcinoma: A study of the National Cancer Institute of Canada Clinical Trials Group. *Cancer* 92:595–600
9. Grønberg BH, Bremnes RM, Fløtten O, Amundsen T, Brunsvig PF, Hjelde HH, Kaasa S, von Plessen C, Stornes F, Tollåli T, Wammer F, Aasebø U, Sundstrøm S (2009) Phase III study by the Norwegian Lung Cancer Study Group: Pemetrexed plus carboplatin compared with gemcitabine plus carboplatin as first-line chemotherapy in advanced non-small-cell lung cancer. *J Clin Oncol* 27:3217–3224
 10. Zinner RG, Fossella FV, Gladish GW, Glisson BS, Blumenschein GR Jr, Papadimitrakopoulou VA, Pisters KM, Kim ES, Oh YW, Peeples BO, Ye Z, Curiel RE, Obasaju CK, Hong WK, Herbst RS (2005) Phase II study of pemetrexed in combination with carboplatin in the first-line treatment of advanced non-small cell lung cancer. *Cancer* 104:2449–2456
 11. Hotta K, Matsuo K, Ueoka H, Kiura K, Tabata M, Tanimoto M (2004) Meta-analysis of randomized clinical trials comparing cisplatin to carboplatin in patients with advanced non-small-cell lung cancer. *J Clin Oncol* 22:3852–3859
 12. Muhsin M, Gricks C, Kirkpatrick P (2004) Pemetrexed disodium. *Nat Rev Drug Discov* 3:825–826
 13. Takehara A, Kawakami K, Ohta N, Oyama K, Ota Y, Oda M, Watanabe G (2005) Prognostic significance of the polymorphisms in thymidylate synthase and methylenetetrahydrofolate reductase gene in lung cancer. *Anticancer Res* 25:4455–4462
 14. Maring JG, Groen HJ, Wachters FM, Uges DR, de Vries EG (2005) Genetic factors influencing pyrimidine-antagonist chemotherapy. *Pharmacogenomics J* 5:226–243
 15. Ishihama H, Chida M, Araki O, Karube Y, Seki N, Tamura M, Umezumi H, Honma K, Masawa N, Miyoshi S (2008) Comparison of 5-fluorouracil-related gene expression levels between adenocarcinomas and squamous cell carcinomas of the lung. *Jpn J Clin Oncol* 39:33–36
 16. Smit EF, Burgers SA, Biesma B, Smit HJ, Eppinga P, Dingemans AM, Joerger M, Schellens JH, Vincent A, van Zandwijk N, Groen HJ (2009) Randomized phase II and pharmacogenetic study of pemetrexed compared with pemetrexed plus carboplatin in pre-treated patients with advanced non-small-cell lung cancer. *J Clin Oncol* 27:2038–2045
 17. Rose MG, Farrell MP, Schmitz JC (2002) Thymidylate synthase: a critical target for cancer chemotherapy. *Clin Colorectal Cancer* 1:220–229
 18. Okamoto I, Hirabayashi N, Kitano M, Nakagawa K (2011) Thymidylate synthase and dihydropyrimidine dehydrogenase expression levels are associated with response to S-1 plus carboplatin in advanced non-small cell lung cancer. *Lung Cancer* 73:103–109
 19. Ozasa H, Oguri T, Uemura T, Miyazaki M, Maeno K, Sato S, Ueda R (2010) Significance of thymidylate synthase for resistance to pemetrexed in lung cancer. *Cancer Sci* 101:161–166
 20. Tanaka F, Wada H, Fukui Y, Fukushima M (2011) Thymidylate synthase (TS) gene expression in primary lung cancer patients: a large-scale study in Japanese population. *Ann Oncol* 22:1791–1797
 21. Kawakami K, Watanabe G (2003) Identification and functional analysis of single nucleotide polymorphism in the tandem repeat sequence of thymidylate synthase gene. *Cancer Res* 63:6004–6007
 22. Go RS, Adjei AA (1999) Review of the comparative pharmacology and clinical activity of cisplatin and carboplatin. *J Clin Oncol* 17:409–422
 23. Jiang J, Liang X, Zhou X, Huang R, Chu Z (2007) A meta-analysis of randomized controlled trials comparing carboplatin-based to cisplatin-based chemotherapy in advanced non-small cell lung cancer. *Lung Cancer* 57:348–358
 24. Ardizzoni A, Boni L, Tiseo M, Fossella FV, Schiller JH, Paesmans M, Radosavljevic D, Paccagnella A, Zatloukal P, Mazzanti P, Bisset D, Rosell R, CISCA (CISplatin versus CARboplatin) Meta-analysis Group (2007) Cisplatin- versus carboplatin-based chemotherapy in first-line treatment of advanced non-small-cell lung cancer: an individual patient data meta-analysis. *J Natl Cancer Inst* 99:847–857
 25. Kim YH, Hirabayashi M, Togashi Y, Hirano K, Tomii K, Masago K, Kaneda T, Yoshimatsu H, Otsuka K, Mio T, Tomioka H, Suzuki Y, Mishima M (2012) Phase II study of carboplatin and pemetrexed in advanced non-squamous, non-small-cell lung cancer: Kyoto Thoracic Oncology Research Group Trial 0902. *Cancer Chemother Pharmacol* 70:271–276
 26. Okamoto I, Aoe K, Kato T, Hosomi Y, Yokoyama A, Imamura F, Kiura K, Hirashima T, Nishio M, Nogami N, Okamoto H, Saka H, Yamamoto N, Yoshizuka N, Sekiguchi R, Kiyosawa K, Nakagawa K, Tamura T (2013) Pemetrexed and carboplatin followed by pemetrexed maintenance therapy in chemo-naïve patients with advanced nonsquamous non-small-cell lung cancer. *Invest New Drugs* 31:1275–1282
 27. Paz-Ares L, de Marinis F, Dediu M, Thomas M, Pujol JL, Bidoli P, Molinier O, Sahoo TP, Laack E, Reck M, Corral J, Melemed S, John W, Chouaki N, Zimmermann AH, Visseren-Grul C, Gridelli C (2012) Maintenance therapy with pemetrexed plus best supportive care versus placebo plus best supportive care after induction therapy with pemetrexed plus cisplatin for advanced non-squamous non-small-cell lung cancer (PARAMOUNT): a double-blind, phase 3, randomized controlled trial. *Lancet Oncol* 13:247–255
 28. Takimoto CH (1996) New antifolates: pharmacology and clinical applications. *Oncologist* 1:68–81
 29. Hanauske AR, Chen V, Paoletti P, Niyikiza C (2001) Pemetrexed disodium: a novel antifolate clinically active against multiple solid tumors. *Oncologist* 6:363–373
 30. Hashimoto H, Ozeki Y, Sato M, Obara K, Matsutani N, Nakagishi Y, Ogata T, Maehara T (2006) Significance of thymidylate synthase gene expression level in patients with adenocarcinoma of the lung. *Cancer* 106:1595–1601
 31. Kaira K, Ohde Y, Nakagawa K, Okumura T, Murakami H, Takahashi T, Kondo H, Nakajima T, Endo M, Yamamoto N (2012) Thymidylate synthase expression is closely associated with outcome in patients with pulmonary adenocarcinoma. *Med Oncol* 29:1663–1672
 32. Lee MS, Asomaning K, Su L, Wain JC, Mark EJ, Christiani DC (2012) MTHFR polymorphisms, folate intake and carcinogen DNA adducts in the lung. *Int J Cancer* 131:1203–1209
 33. Suzuki T, Matsuo K, Hiraki A, Saito T, Sato S, Yatabe Y, Mitsudomi T, Hida T, Ueda R, Tajima K (2007) Impact of one-carbon metabolism-related gene polymorphisms on risk of lung cancer in Japan: a case control study. *Carcinogenesis* 28:1718–1725
 34. Liu H, Jin G, Wang H, Wu W, Liu Y, Qian J, Fan W, Ma H, Miao R, Hu Z, Sun W, Wang Y, Jin L, Wei Q, Shen H, Huang W, Lu D (2008) Association of polymorphisms in one-carbon metabolizing genes and lung cancer risk: a case-control study in Chinese population. *Lung Cancer* 61:21–29
 35. Martin DN, Boersma BJ, Howe TM, Goodman JE, Mechanic LE, Chanock SJ, Ambs S (2006) Association of MTHFR gene polymorphism with breast cancer survival. *BMC Cancer* 6:257
 36. Askari BS, Krajcinovic M (2010) Dihydrofolate reductase gene variations in susceptibility to disease and treatment outcomes. *Curr Genomics* 11:578–583
 37. Chen CY, Chang YL, Shih JY, Lin JW, Chen KY, Yang CH, Yu CJ, Yang PC (2011) Thymidylate synthase and dihydrofolate reductase expression in non-small cell lung cancer: the association with treatment efficacy of pemetrexed. *Lung Cancer* 74:132–138
 38. Zukin M, Barrios CH, Pereira JR, Ribeiro RDA, Beato CA, do Nascimento YN, Murad A, Franke FA, Precivale M, Araujo LH,

- Baldotto CS, Vieira FM, Small IA, Ferreira CG, Lilenbaum RC (2013) Randomized phase III trial of single-agent pemetrexed versus carboplatin and pemetrexed in patients with advanced non-small-cell lung cancer and Eastern Cooperative Oncology Group performance status of 2. *J Clin Oncol* 31:2849–2853
39. Quoix E, Zalcman G, Oster JP, Westeel V, Pichon E, Lavolé A, Dauba J, Debieuvre D, Souquet PJ, Bigay-Game L, Dansin E, Poudenx M, Molinier O, Vaylet F, Moro-Sibilot D, Herman D, Bennouna J, Tredaniel J, Ducoloné A, Lebitasy MP, Baudrin L, Laporte S, Milleron B, Intergroupe Francophone de Cancérologie Thoracique (2011) Carboplatin and weekly paclitaxel doublet chemotherapy compared with monotherapy in elderly patients with advanced non-small-cell lung cancer: IFCT-0501 randomised, phase 3 trial. *Lancet Oncol* 378:1079–1088
40. Gervais R, Robinet G, Clément-Duchêne C, Denis F, El Kouri C, Martin P, Chouaki N, Bourayou N, Morère JF (2013) Pemetrexed and carboplatin, an active option in first-line treatment of elderly patients with advanced non-small cell lung cancer (NSCLC): a phase II trial. *Lung Cancer* 80:185–190
41. Maemondo M, Inoue A, Kobayashi K, Sugawara S, Oizumi S, Isobe H, Gemma A, Harada M, Yoshizawa H, Kinoshita I, Fujita Y, Okinaga S, Hirano H, Yoshimori K, Harada T, Ogura T, Ando M, Miyazawa H, Tanaka T, Saijo Y, Hagiwara K, Morita S, Nukiwa T, North-East Japan Study Group (2010) Gefitinib or chemotherapy for non-small-cell lung cancer with mutated EGFR. *Engl J Med* 362:2380–2388
42. Mitsudomi T, Morita S, Yatabe Y, Negoro S, Okamoto I, Tsurutani J, Seto T, Satouchi M, Tada H, Hirashima T, Asami K, Katakami N, Takada M, Yoshioka H, Shibata K, Kudoh S, Shimizu E, Saito H, Toyooka S, Nakagawa K, Fukuoka M, West Japan Oncology Group (2010) Gefitinib versus cisplatin plus docetaxel in patients with non-small-cell lung cancer harbouring mutations of the epidermal growth factor receptor (WJTOG3405): An open label, randomised phase 3 trial. *Lancet Oncol* 11:121–128

Clinical Trials. gov number : NCT00016211 ; PCI 試験今昔

Prophylactic cranial irradiation in extensive small cell lung cancer

赤松 弘朗¹ / 山本 信之²

Hiroaki Akanatsu Nobuyuki Yamamoto

和歌山県立医科大学医学部内科学第三講座¹ / 教授²

Clinical Trials. gov number : NCT00016211 の詳細

Slotman らによる本試験(以下, Slotman 試験)についての論文は「Prophylactic cranial irradiation in extensive small-cell lung cancer」として, 2007年の『The New England Journal of Medicine』誌に発表された¹⁾。周知のとおり, その後各国のガイドラインに採用されるに至った重要な論文である。

対象は初回化学療法になんらかの反応をみせた進展型小細胞肺癌(extensive disease-small cell lung cancer; ED-SCLC)で, 経過観察群と, 試験治療として予防的全脳照射(prophylactic cranial irradiation; PCI)を行う PCI 群とに 1 : 1 に無作為化した(図 1)。主要評価項目は, 症候性脳転移発症までの期間とされた。計286名の患者が参加し, 予想どおり PCI は症候性脳転移発症までの期間を有意に抑え(ハザード比(HR)0.27(95%信頼区間(CI) : 0.16~0.44), $p < 0.001$), それだけでなく無増悪生存期間(PFS)(12.0週 vs. 14.7週), 全生存期間(OS)(5.4ヵ月 vs. 6.7ヵ月)ともに PCI 群で有意な延長が示された。

Slotman 試験の問題点

以上が本論文の抄録から得られる情報である。しかしな




図1 Slotman 試験のシエーマ

がら本論文を読み込むと(読み込まなくても?)不十分なデザインに基づいた試験であることが明らかであり, 論文発表当初からいくつかの疑義が寄せられていた²⁾³⁾。なかでも最大の弱点は, 登録時に頭部画像検査が必須でなかったことである。つまり, 試験治療が本当に「予防的」全脳照射であったのか誰も確認できていないという, 今から考えると少しいかげんな試験であった。そのほかにも主要評価項目が OS でないこと, 化学療法が現在の標準治療であるプラチナ併用療法でないこと, 全脳照射の方法が一律でないこと, 無作為化後の頭部画像検査が定期的に行われず, 転移を示唆する状況になってはじめて行うことが規定されていることなどの問題点が指摘された。

検証試験の結果

このような背景をもとに, 考えるかぎり十分堅固な試験デザインをもって PCI の有効性を再検証したのが, 2014年の米国臨床腫瘍学会(ASCO)において日本から報告された第Ⅲ相臨床試験である(図 2)⁴⁾。本検証試験では無作為化前の頭部 MRI を必須とし, PCI の線量もこれまでで最もエビデンスのある 25Gy/10frs に統一した。もちろん, 主要評価項目は OS とされた。結果, 初回の中間解析において PCI 群が経過観察群に対して OS が上回る可能性がきわ




図2 日本からの検証試験のシエーマ(Slotman 試験との主な違いを青字で示す)

めて乏しい(PCI群が有意に上回る確率は $p < 0.001$ だが, OSにおける p 値は0.091であった)との結果が判明し, 本検証試験は無効中止となった(図3)。

改めて両試験の結果を比較してみる(表1)。両試験間でPFSにはそれほど差はないが, OSは大きく異なっている。PFS中央値である2~3ヵ月時点において, 脳転移増悪頻度がそれほど高くないため, 後治療における全身化学療法の導入割合がこうした差異をもたらした可能性が高い。実際, Slotman 試験において後治療(放射線治療も含む)の導入割合が45~68%であったのに対し, 検証試験では両群とも80%以上の患者に対して2次化学療法が行われている。それでは, なぜ後治療にこのような差が生じたのであるのか。両試験における脳転移発症割合は検証試験においておおむね高いが, これはデザインの問題でSlotman 試験では症候性脳転移が, 検証試験では無症候性脳転移が多いためと思われる。つまり検証試験においては定期的なMRIによって無症候性脳転移を早期に検出したことが, 高い後治療の導入割合につながり, ひいては良好な生存期間をもたらしたと考えられる。

最後に

検証試験の結果は2014年のASCOで九州がんセンターの瀬戸貴司先生より発表されたが, くしくも前の演者がBen Slotman先生であったため新旧PCI試験の発表者が壇上で対峙することになった。(Ben Slotman先生には気の毒であったが)ディスカッサントは検証試験の結果について非常に好意的で, 日本からのエビデンスがこのようなかたちで認められたことを非常に誇らしく思った。本検証試験の結果は, ED-SCLCのガイドラインを塗り替えることになると思われる。




図3 日本より発表された第Ⅲ相臨床試験におけるOS

(文献4)より引用・一部改変

表1 2つのPCI試験のまとめ

試験	治療	患者数	6ヵ月時点の	12ヵ月時点の	mPFS (月)	MST (月)
			脳転移増悪頻度 (%)	脳転移増悪頻度 (%)		
Slotman 5 ¹⁾	経過観察群	143	32	40.4	2.8	5.4
	PCI群	143	4.4	14.6	3.4	6.7
Seto 5 ⁴⁾	経過観察群	79	38*	58.0	2.4	15.1
	PCI群	84	12*	32.4	2.2	10.1

mPFS: PFS中央値, MST: 生存期間中央値

*: ASCO発表スライドより推定

文献

- 1) Slotman B, Faivre-Finn C, Kramer G, et al: Prophylactic cranial irradiation in extensive small-cell lung cancer. *N Engl J Med* 357: 664-672, 2007
- 2) Shivnani AT: Prophylactic cranial irradiation in small-cell lung cancer. *N Engl J Med* 357: 1977, 2007; author reply 1978
- 3) Fujiwara Y, Hotta K, Kiura K: Prophylactic cranial irradiation in small-cell lung cancer. *N Engl J Med* 357: 1977-1978, 2007; author reply 1978
- 4) Seto T, Takahashi T, Yamanaka T, et al: Prophylactic cranial irradiation (PCI) has a detrimental effect on the overall survival (OS) of patients (pts) with extensive disease small cell lung cancer (ED-SCLC); Results of a Japanese randomized phase III trial. *J Clin Oncol* 32(Suppl.): 2014 (ASCO 2014, Abstr. 7503)



話題

小細胞肺がんに対する 予防的全脳照射*

赤松 弘朗** 山本 信之**

Key Words : prophylactic cranial irradiation, small cell lung cancer

はじめに

小細胞肺がんは肺がんの約15%を占め、転移をきたしやすく悪性度の高い疾患である¹⁾。診断時に切除可能であることは非常に少なく、化学療法²⁾や放射線療法との同時併用^{3,4)}が選択される。しかし治療にいったんは奏効を示すものの再発も多く、なかでも脳転移の制御は大きな課題である。小細胞肺がんにおける脳転移は初診時約20%にみられるだけでなく、経過中にも50~65%で発症するとされている⁵⁾。これは疾患としての性質だけでなく、血液脳関門の存在により全身治療である抗がん剤の効果が十分得られないことも一因とされており、実際脳転移に対する化学療法の効果は非常に乏しいことが示されている⁶⁾。

予防的全脳照射(prophylactic cranial irradiation ; PCI)はそのような背景からでてきた非常にユニークな治療戦略であり、多くのがん腫の中でも小細胞肺がんにおいてのみ採用される手法である。

本稿では小細胞肺がんに対するPCIの科学的根拠についてこれまでの知見を概説し、日常臨床への応用についてもふれる。

初回治療に完全奏効を示した 小細胞肺がんに対するPCI

初回治療(化学療法・放射線療法)でいったん完全奏効(complete response ; CR)が得らるもの




図1 初回治療に完全奏効を示した小細胞肺がんに対するPCIのメタ解析;初めてPCIによる生存期間延長が示された (文献¹⁰⁾より引用一部改変)

の、初再発として脳転移を呈することが多く経験されたことから、このような症例を対象としたPCIの有効性を検証する臨床試験が1980年代から本邦も含めて複数行われた⁷⁻⁹⁾。その結果、PCIが脳転移再発を有意に減少させることが示されたものの、生存期間延長に結びつくかについてははっきりしなかった。Auperinらはこれら7つの臨床試験をもとに987名の個々のデータを用いたメタ解析を行い¹⁰⁾、これによってPCIによる生存期間の延長(3年生存割合15.3% vs. 20.7%, $P=0.01$)が初めて証明された(図1)。その後、Arriagataらがこのメタ解析で用いられた試験のうち主要な2報について長期成績を報告している(観察期間中央値11年)。結果、5年生存割合はPCI群18%に対し経過観察群15%と有意差はわず

* Prophylactic cranial irradiation for small cell lung cancer.

** Hiroaki AKAMATSU, M.D. & Nobuyuki YAMAMOTO, M.D., Ph.D.: 和歌山県立医科大学内科学第三講座(〒641-8509 和歌山県和歌山市紀三井寺811-1); Third Department of Internal Medicine, Wakayama Medical University, Wakayama 641-8509, JAPAN




図2 進展型小細胞肺癌に対するPCIの第III相試験
左：脳転移発生割合，右：全生存期間（文献¹³⁾より引用改変）

かに認められなかったものの($P=0.06$)，PCI群で良好な傾向が示された。生存曲線においてもAuperinらの報告と同様，常に一定の割合をもってPCI群が上回っているという結果であった。これらの報告により，初回治療にCRとなった症例に対してはPCIが標準治療として認識され，その後各国のガイドラインにも反映されることになっている¹¹⁾¹²⁾。

進展型小細胞肺癌に対するPCI

進展型小細胞肺癌は初回治療でCRに至ることは稀であり，通常は化学療法と緩和治療を中心に行う。しかし経過中の脳転移発生割合が高く，その場合の化学療法は前述したように効果が期待できない。そこで限局型小細胞肺癌におけるPCIの報告を受け，進展型に対してもPCIの導入が検討された。Slotmanらは初回化学療法によってなんらかの効果が得られた進展型小細胞肺癌を対象としてPCIと経過観察の第III相試験を行った¹³⁾。結果，1年以内の脳転移発生割合はPCI群14.6%に対し経過観察群40.4%と実に約1/3に抑えられ，かつ生存期間中央値についても前者6.7か月に対し後者5.4か月と有意な延長が示された(図2)。ただし，これについては後述するような問題点があり，本邦ではより厳密な対象選択をもとにした第III相試験が厚生労働省科学研究(研究班長 山本信之)において行われている。

至適線量・有害事象に関する報告

PCIの有効性が示されるとともに，至適線量や有害事象に関しても以下のようにさまざまな報告が行われてきた。

1. 至適線量

Auperinらのメタ解析では総線量の増加に従って脳転移発生が抑えられることが示されたものの，これが生存には関与しないことが報告された。これを確認すべく，PechouxらはCRが得られた限局型を対象に25 Gy/10 frsと36 Gy[18 frsもしくは24 frs(1日2回)]の第III相試験を行った¹⁴⁾。結果，主要評価項目であった2年時点の脳転移再発に差はなく，生存についてはむしろ高線量群が下回っていた[図3，37% vs. 42%(HR 1.00~1.44, $P=0.05$)]。高線量群で生存が劣っていた理由についてははっきりしないが，この結果をもって25 Gy/10 frsが標準線量と考えられるようになった。

2. 有害事象

全脳照射による晩期毒性についても，これまでに議論が行われてきた。無治療群との比較についてはGregorら⁷⁾・Arriagataら¹⁵⁾の試験で検討されているものの，半数近くの症例でPCI開始前に神経症状の出現が認められており，PCIの影響に関して厳密な判断は難しいと考えられた。しかし，がん腫は異なるものの非小細胞肺癌に対して行われたPCIと経過観察の比較試験¹⁶⁾では，




図3 高線量によるPCIは生存期間を延長しなかった
(文献¹⁴⁾より引用改変)

PCI群において1年時点の認知機能低下や記憶障害が指摘されている¹⁷⁾。また無治療との比較ではないが、前述したPechouxらの検討でも中央値3年以上の観察期間において、25 Gy・36 Gyの両群で下肢筋力・記憶力・知的能力などの経時的な低下が認められている¹⁸⁾。限局型の場合、根治に至る可能性もあるため、こうした晩期毒性について事前に十分説明しておく配慮が必要と思われる。

日常臨床へ外挿する前に…

ここまでPCIのエビデンスについて述べてきたが、実はこれらの試験が行われた20年近く前の背景を考えると、われわれの日常臨床に応用するにあたっては、いくつかの注意が必要である。

最も注意すべき事項は対象患者の適切さである。これらの試験では初回治療については化学療法単独のみのものが含まれており、CRの確認についても多くが胸部X線のみでなされていた。PCI直前の脳転移の有無について確認されていない試験も多い。つまり対象患者における真のCR割合が低い可能性があるため、これらの試験で行われたPCIが実は「無症候性脳転移に対する全脳照射であった」可能性が潜んでいることは否定できない。近年ではCT・PETの普及によって病期診断・効果判定がより正確になされるようになってきている。1999年に報告されたAuperinらによるメタ解析におけるPCI群の3年生存割合が20%




図4 限局型小細胞肺癌におけるPETの有無； stagingにおけるPETの有無が生存期間に影響する
(文献¹⁹⁾より引用改変)

であるのに対し、Pechouxらによる第III相試験(2009年)における3年生存割合が35%と大きな開きがあるのは、こうした患者選択の正確性に影響されている可能性がある。後方視的研究ではあるが、Xanthopoulosらは54例の限局型小細胞肺癌について、病期診断におけるPETの有無がもたらす有効性について報告しており¹⁹⁾、stagingにおけるPETの有無によって生存期間に有意な差が生じており(図4)、多変量解析においても全身状態(performance status)・N因子と並んでPETの有無が有意な予後因子であった(HR 0.24, $P=0.04$)としている。

特に脳転移の頻度が増す進展型においては、こうした背景に注意が必要と考えられる。Slotmanらの報告では登録時の脳転移検索が行われた症例は30%に満たなかったとされているが、本邦のようにMRIの普及が進んでいる国においては、定期的な頭部検索により迅速に無症候性脳転移を治療することが可能かもしれない。PCIより生存期間や有害事象で優れる可能性がある。こうした考えをもとに、本邦では厳密な対象選択を行った上での進展型小細胞肺癌に対するPCIの意義を問う第III相試験を行っている(シエマを図5に示す)。本試験ではPCI前の頭部MRIを必須とし、また照射線量もこれまでで最もエビデンスのある25 Gy/10 frsに統一した。本研究の結果は2014年の米国臨床腫瘍学会(ASCO)で報告される予定であり、詳細な結果報告が非常に待たれる。