

# Endogenous Mesenchymal Stromal Cells in Bone Marrow Are Required to Preserve Muscle Function in mdx Mice

RYO FUJITA,<sup>a</sup> KATSUTO TAMAI,<sup>b</sup> ERIKO AIKAWA,<sup>b</sup> KEISUKE NIMURA,<sup>a</sup> SAKI ISHINO,<sup>c</sup> YASUSHI KIKUCHI,<sup>b</sup> YASUFUMI KANEDA<sup>a</sup>

**Key Words.** Mesenchymal stromal cells • Duchenne muscular dystrophy • TNF- $\alpha$ -stimulated gene/protein-6 • TSG-6 • Muscle satellite cells • Bone marrow cells

<sup>a</sup>Division of Gene Therapy Science, <sup>b</sup>Department of Stem Cell Therapy Science, and <sup>c</sup>Center for Medical Research and Education, Graduate School of Medicine, Osaka University, Osaka, Japan

Correspondence: Katsuto Tamai, M.D., Ph.D., Department of Stem Cell Therapy Science, Graduate School of Medicine, Osaka University, 2-2, Yamada-oka, Suita, Osaka 565-0871, Japan. Telephone: 81-6-6879-3901; Fax: 81-6-6879-3909; e-mail: tamai@gts.med.osaka-u.ac.jp; or Yasufumi Kaneda, M.D., Ph.D., Division of Gene Therapy Science, Graduate School of Medicine, Osaka University, 2-2, Yamada-oka, Suita, Osaka 565-0871, Japan. Telephone: 81-6-6879-3901; Fax: 81-6-6879-3909; e-mail: kaneday@gts.med.osaka-u.ac.jp

Received July 9, 2014; accepted for publication October 31, 2014; first published online in *STEM CELLS EXPRESS* November 19, 2014.

© AlphaMed Press  
1066-5099/2014/\$30.00/0

<http://dx.doi.org/10.1002/stem.1900>

## ABSTRACT

The physiological role of “endogenous” bone marrow (BM) mesenchymal stromal cells (MSCs) in tissue regeneration is poorly understood. Here, we show the significant contribution of unique endogenous BM-MSC populations to muscle regeneration in Duchenne muscular dystrophy (DMD) mice (mdx). Transplantation of BM cells (BMCs) from 10-week-old mdx into 3–4-week-old mdx mice increased inflammation and fibrosis and reduced muscle function compared with mdx mice that received BMCs from 10-week-old wild-type mice, suggesting that the alteration of BMC populations in mdx mice affects the progression of muscle pathology. Two distinct MSC populations in BM, that is, hematopoietic lineage (Lin<sup>−</sup>/ckit<sup>−</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> and Lin<sup>−</sup>/ckit<sup>−</sup>/CD106<sup>+</sup>/CD44<sup>−</sup> cells, were significantly reduced in 10-week-old mdx mice in disease progression. The results of a whole-transcriptome analysis indicated that these two MSC populations have distinct gene expression profiles, indicating that the Lin<sup>−</sup>/ckit<sup>−</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> and Lin<sup>−</sup>/ckit<sup>−</sup>/CD106<sup>+</sup>/CD44<sup>−</sup> MSC populations are proliferative- and dormant-state populations in BM, respectively. BM-derived Lin<sup>−</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> MSCs abundantly migrated to damaged muscles and highly expressed tumor necrosis factor- $\alpha$ -stimulated gene/protein-6 (TSG-6), an anti-inflammatory protein, in damaged muscles. We also demonstrated that TSG-6 stimulated myoblast proliferation. The injection of Lin<sup>−</sup>/ckit<sup>−</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> MSCs into the muscle of mdx mice successfully ameliorated muscle dysfunction by decreasing inflammation and enhancing muscle regeneration through TSG-6-mediated activities. Thus, we propose a novel function of the unique endogenous BM-MSC population, which countered muscle pathology progression in a DMD model. *STEM CELLS* 2015;33:962–975

## INTRODUCTION

Bone marrow (BM) is a source of hematopoietic stem cells (HSCs) and also contains nonhematopoietic stromal cell populations known as mesenchymal stromal cells (MSCs). Conventionally, MSCs have been defined as colony-forming unit-fibroblasts (CFU-Fs) with the capacity to differentiate into adipocytes, chondrocytes, and osteocytes in vitro [1, 2]. Culture-expanded MSCs have been used to treat various tissue injuries, such as myocardial infarction and lung and skin injuries [3–5]. A focus of great interest regarding the use of culture-expanded MSCs in clinical applications is their anti-inflammatory and regeneration activities, which render the microenvironment more efficient for promoting tissue regeneration [3, 6].

Our previous studies revealed that, similar to culture-expanded BM-MSCs, circulating

BM-derived osteoblast progenitor cells were attracted by bone morphogenetic protein-2 to contribute to ectopic bone formation [7, 8]. We also found previously that necrotic skin-derived high-mobility group box 1 (HMGB1) mobilizes platelet-derived growth factor receptor  $\alpha$ -positive (PDGFR $\alpha$ <sup>+</sup>) mesenchymal cells from the BM into circulation [9]. These data suggest that injury stimuli attract endogenous BM-MSC populations via the circulation to support tissue regeneration.

However, the characteristics of these endogenous BM-MSC populations mobilized from BM into circulation and damaged tissues have not been established because of their low number and high heterogeneity. In addition, another endogenous function of MSCs in BM is to maintain HSC quality and quantity as a niche by secreting stem cell factor and/or C-X-C type chemokine ligand 12 (CXCL12), known as stromal cell-derived factor 1 $\alpha$  (SDF-1 $\alpha$ )

[10–14]. Together, the results of our previous investigations along with those of other seminal studies suggest that multiple MSC populations exist in BM, possibly with distinct functions, such as supporting HSCs and contributing to tissue regeneration [15, 16].

Although information regarding the functions and surface markers of culture-expanded MSCs has accumulated, the molecular characteristics of endogenous BM-MSCs are still poorly understood [17, 18]. It is thus necessary to explore BM-MSC populations contributing to the tissue regeneration processes and their molecular signatures—such as gene expression profiles—to elucidate the specific functions of each MSC population *in vivo*. In previous research, we investigated the dynamics of BM-MSCs in acute injury and regeneration models [8, 9], but it remains unclear whether endogenous BM-MSCs also support tissue regeneration in chronic tissue damage and regeneration processes, such as those in Duchenne muscular dystrophy (DMD).

DMD is characterized primarily by progressive muscle degeneration and weakness due to mutations in the dystrophin gene [19–21]. Muscle cells without dystrophin are vulnerable to damage, resulting in cycles of degeneration and regeneration, chronic inflammation, severe fibrosis, and reduced muscle contractility. Currently, therapies for DMD can be divided into two groups: treatments targeting the restoration of dystrophin expression by cell therapies [22–26] and treatments for tilting the balance of overall muscle condition in favor of regeneration by the activation of satellite cells and the suppression of inflammation and fibrosis [27, 28]. However, there is no curative treatment for DMD at present.

In addition to these therapies, boosting endogenous muscle repair mechanisms is an alternative and feasible approach to treat muscle diseases. In light of the accumulated findings, the activation of muscle repair mechanisms by endogenous BM-MSCs could be an attractive novel approach to slow the progression of muscle dysfunction in DMD. For this purpose, a better understanding of the roles of endogenous BM-MSCs in the pathology of DMD as well as the identification of endogenous molecular characteristics *in vivo* is urgently required.

In this study, we investigated the contribution of “endogenous” BM-MSCs to muscle regeneration in a DMD mouse model (mdx). Using a bone marrow transplantation (BMT) model, we found that alterations of the endogenous BM-MSC population affect the muscle pathological conditions in mdx mice. In an evaluation of an endogenous BM-MSC population by a whole-genome transcriptome analysis and flow cytometry analysis, we characterized the heterogeneity of endogenous BM-MSCs and identified the dominantly recruited BM-MSC population into damaged muscles in mdx mice. The treatment of this specific BM-MSC population alone stimulated the muscle regeneration processes. We believe that these findings provide new insight into muscle repair mechanisms with specific endogenous BM-MSC populations, and our results contribute to the understanding of BM-MSC-mediated muscle repair mechanisms that can slow disease progression in DMD.

## MATERIALS AND METHODS

### Mice

All animals were handled according to approved protocols and the guidelines of the Animal Committee of the Osaka

University Graduate School of Medicine. C57BL/6, C57BL/10ScSn/J, and C57BL/10ScSn-Dmd<sup>mdx</sup>/J (referred to as mdx) mice were purchased from CLEA Japan (Tokyo). C57BL/6 mice that ubiquitously expressed enhanced green fluorescent protein (GFP, referred to as GFP mice) were kindly provided by Masaru Okabe (Osaka University, Osaka, Japan). B6.129S4-Pdgfra<sup>tm11(EGFP)Sor</sup>/J mice (referred to as PDGFR $\alpha$ -H2BGFP) were purchased from The Jackson Laboratory (West Grove, PA). PDGFR $\alpha$ -H2BGFP mice were heterozygous knockin mice in which the histone H2B-GFP fusion gene was inserted into the locus of the PDGFR $\alpha$  gene.

### Muscle Endurance Test and Grip-Strength Test

To determine the muscle endurance in mice, a mouse was allowed to grasp the bar or metallic mesh and remain suspended (Supporting Information Fig. S1F, S1G). The time until the mouse released its grip was recorded. Each mouse was subjected to six tests, and the results from these tests were averaged. The final outcome value for the muscle endurance was determined using the holding impulse (second  $\times$  body weight, g) to correct for the negative effects of body mass on the hang time. The whole-limb grip strength was measured using a Grip Strength Meter for mice (Muromachi Kikai, Tokyo). The mouse was allowed to grasp a horizontal mesh platform with all four limbs, and then, the mouse's tail was pulled back parallel to the mesh platform. The peak tension (gram; g) was recorded when the mouse released its grip. Six sets of measurements were performed for each mouse, and the average values were defined as the mouse whole-limb grip strength.

### Isolation of Bone Marrow Cells and Bone Marrow Transplantation

The isolation of BM cells (BMCs) and BMT was performed as described [7, 9]. Briefly, under sterile conditions, BMCs were isolated by flushing the femurs and tibiae with 2% fetal bovine serum (FBS)/phosphate-buffered saline (PBS). For the total BMT,  $5 \times 10^6$  BMCs were injected into the tail veins of mice irradiated with 10 Gy. Six weeks after reconstitution, the chimerisms were examined, and the results confirmed that more than 90% of the total BMCs were donor-derived cells (Supporting Information Fig. S2A, S2B). For the preparation of each mouse with reduced MSCs in BM, GFP<sup>+</sup>/Lin<sup>−</sup>/ckit<sup>+</sup> BMCs, which include HSC but not MSC populations, were sorted with the FACSARIA II cell sorter (BD Biosciences, San Jose, CA; Supporting Information Fig. S3A). Each irradiated mouse received  $1 \times 10^5$ – $2 \times 10^5$  GFP<sup>+</sup>/Lin<sup>−</sup>/ckit<sup>+</sup> BMCs, the number of which was almost identical to that of the GFP<sup>+</sup>/Lin<sup>−</sup>/ckit<sup>+</sup> BMCs for the total BMT. After reconstitution, the numbers and percentages of donor-derived-MSC populations in BM were examined (Supporting Information Fig. S3B–S3F).

### Preparation of Conditioned Media

For the preparation of conditioned media, culture-expanded BM-MSCs at passage 3 were used. The BM-MSCs were stimulated with 10% wild-type (WT) or mdx mouse serum for 2 days. The medium was then changed to fresh  $\alpha$ -MEM, and another 2 days of incubation was performed. The conditioned media of these cells were collected and used to stimulate C2C12 cells.



### Isolation of Primary Satellite Cells and C2C12 Cells

For the isolation of single myofibers, the extensor digitorum longus muscles were isolated and digested in 0.2% collagenase for 90 minutes at 37°C, as described [29]. Primary myoblasts were obtained from all hind limb muscles, according to the procedure of Musaro and Barberi [30]. C2C12 cells were obtained from the American Type Culture Collection (Rockville, MD). For antibody treatment, C2C12 cells were preincubated for 1 hour with 2.5 µg/ml of control IgG (rat IgG) or blocking antibody for CD44 (clone KM81; Abcam, Cambridge, MA) with or without BM-MSC-derived condition media. Recombinant mouse tumor necrosis factor- $\alpha$  (TNF- $\alpha$ )-stimulated gene/protein-6 (TSG-6) (0.5 mg/ml) was applied to stimulate the C2C12 cells for 48 hours.

### Colony-Forming Unit Assay

Lin<sup>−</sup> BMCs were collected to exclude all hematopoietic lineage cells from total BMCs by the MACS system (Miltenyi Biotec, Bergisch Gladbach, Germany). Isolated Lin<sup>−</sup> BMCs were then seeded in a six-well plate. The adherent cells were stained as described [31]. The numbers of colonies containing more than 50 cells were counted under a light microscope.

### In Vivo Migration Assay

Silicon tubes containing either PBS or SDF-1 $\alpha$  (100 ng/ml; Peprotech, Rocky Hill, NJ) were implanted under the skin of mdx mice. After 2 days, the tubes were collected and cultured for 2 days and analyzed by immunocytochemistry.

### Muscle Injury and Treatment

To induce muscle injury, a cardiotoxin (CTX; *Naja mossaibica* venom; Sigma, St. Louis, MO) was used [32]. For cell treatment, freshly isolated BMCs ( $1 \times 10^5$ ) were injected directly into both the superior and inferior regions of CTX-damaged WT or mdx muscles. The day after treatment, 10 µg of CD44 antibody (clone KM81, Abcam) and 5 µg of TSG-6 antibody (clone MAB2104, R&D Systems, Minneapolis, MN) were administered to the muscles. As a control, the same amounts of rat or mouse IgG (R&D Systems) were used. Recombinant mouse TSG-6 (2 µg in 10 µl of PBS; R&D Systems) was administered to the muscles. The same volume of PBS was injected into the muscle as a control.

### Histology and Immunostaining

Hematoxylin and eosin (H&E) staining was performed as described [32]. For collagen staining, a picrosirius red staining kit was used (Polysciences, Warrington, PA). The average cross-sectional area (CSA) of muscle fibers was measured using NIH ImageJ software. Only regenerating muscle fibers with central nuclei (100–300 fibers per muscle) were measured. The antibodies used for immunostaining are listed in Supporting Information Table S1. For the nuclear staining, prolong gold antifade reagent containing DAPI (Life Technologies, Carlsbad, CA) was used. The MyoD-, Pax7-, and embryonic myosin heavy chain (eMyHC)-positive cells and the area in three regions of cross-section per mouse were determined by NIH ImageJ software.

### Evans Blue Dye Uptake

First, 10 mg/ml of Evans blue dye (EBD) stock solution was prepared in PBS. A single dose (50 mg/g of b.wt.) was administered to the mouse by intraperitoneal injection. The mice

were sacrificed 20 hours later, and the muscles were dissected. The EBD<sup>+</sup> cells per muscle were counted in five regions using  $\times 20$  magnification sections.

### Fluorescence-Activated Cell Sorting Analysis and Sorting of Bone Marrow Cells

Isolated BMCs containing  $2 \times 10^6$  cells were suspended in 200 µl of staining buffer (2% FBS in PBS), followed by an incubation with purified rat anti-mouse CD16/CD32 (Mouse BD Fc Block, BD Pharmingen, San Diego, CA) for 10 minutes at 4°C. The fluorescence-conjugated antibodies were incubated for 30 minutes at 4°C. The antibodies used in the flow cytometry analysis are listed in Supporting Information Table S1. Fluorescence was measured by the FACSCanto II (BD Biosciences). Gates were defined based on the isotype control staining. The fluorescence-activated cell sorting data were analyzed using FlowJo software ver. 6.3.3 (Tree Star, Ashland, OR).

### Muscle Digestion for the Flow Cytometry Analysis

The harvested muscles (soleus, tibialis anterior [TA], gastrocnemius, and plantaris) were minced into small pieces and digested using 0.2% collagenase A (Roche, Mannheim, Germany) for 40 minutes at 37°C with frequent triturating by pipette. The digested muscles were centrifuged at 1,200 rpm for 5 minutes and then further incubated in 0.2% collagenase/dispase (Roche) for 1 hour at 37°C. Following digestion, the cells were filtered using a sequence of 70-µm and 40-µm nylon mesh strainers, followed by centrifugation at 1,500 rpm for 10 minutes. The pellets were then suspended in the staining buffer and stained with specific antibodies, and the BMC populations in damaged muscles were analyzed and isolated by the BD FACSCanto II and BD FACSAria II, respectively.

### Western Blot Analysis

Western blot analyses were performed as described [32]. The antibodies used in the Western blot analysis are listed in Supporting Information Table S1. All bands were visualized by Chemi-Lumi One (Nacalai Tesque, Kyoto, Japan) with ImageQuant LAS 4000mini software (GE Healthcare, Buckinghamshire, U.K.). The band density was analyzed using NIH ImageJ software.

### RNA Extraction and Real-Time Polymerase Chain Reaction

Total RNA was extracted from muscle and BMCs using ISOGEN (Nippon Gene, Toyama, Japan) according to the manufacturer's instructions. cDNA was synthesized from an equal quantity of total RNA (2 µg for muscle and 500 ng for sorted BMCs) by reverse transcription using the High Capacity RNA-to-cDNA kit (Applied Biosystems, Foster City, CA). A real-time polymerase chain reaction (PCR) was performed with SYBR PremixEX Taq (Takara Bio, Shiga, Japan) using oligonucleotide primers. The sequences of oligonucleotide primers used for the real-time PCR are listed in Supporting Information Table S2. The quantitative data were obtained in triplicate within a single experiment on a 384-well plate based on the standard curve method using CFX manager software (BioRad, Hercules, CA). All data were normalized to GAPDH levels as an internal control.

### RNA Sequencing

RNA was extracted from sorted Lin<sup>−</sup>/ckit<sup>−</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> and Lin<sup>−</sup>/ckit<sup>−</sup>/CD106<sup>+</sup>/CD44<sup>−</sup> BMCs using ISOGEN (Nippon

Gene) according to the manufacturer's instructions. Strand-specific sequencing libraries from two biological replicate RNA samples were prepared according to the Life Technologies protocol, as described [33]. Briefly, digested poly-A-tailed RNA was ligated to the SOLiD Adaptor Mix and then reverse-transcribed using the SOLiD Total RNA-Seq Kit (Life Technologies). Size-selected first-strand cDNA was amplified by SOLiD 5' PCR primers and barcoded SOLiD 3' PCR primers (Life Technologies). RNA-seq libraries were sequenced with the 5500xl Genetic Analyzer (Life Technologies). The resulting reads were mapped using Lifescope (Life Technologies) and analyzed using Cufflinks and CummeRbund [34, 35]. The enrichment of gene ontology was calculated by gene ontology tool Database for Annotation, Visualization, and Integrated Discovery [36, 37].

### Statistical Analyses

The values are expressed as the mean  $\pm$  SEM. Statistical analyses were performed with Student's *t* test using the SPSS ver. 10.0 (Japan, Inc., Tokyo). For all statistical tests, significance was accepted at  $p < .05$ .

### Accession Numbers

The accession number of RNA seq data is DRA002440.

## RESULTS

### WT and mdx Mouse Bone Marrow Transplantations Showed Different Effects on mdx Muscle Pathology

We first examined whether the transplantation of WT BMCs in mdx mice would ameliorate muscle pathology by restoring dystrophin expression. To address this question, we transplanted BMCs derived from either 10-week-old WT mice (or 10-week-old GFP mice) or mdx mice into lethally irradiated 3–4-week-old mdx mice to generate BMT mice (Fig. 1A; Mdx-wt vs. Mdx-mdx, Supporting Information Fig. S4A; Mdx-GFP). To clarify the influence of BMT itself on muscle pathology, we used age-matched non-BMT mdx mice as controls. The body weights and muscle wet weights were not significantly different between the groups during the experimental period (Supporting Information Fig. S1A, S1B).

We did not detect any BM-derived (GFP<sup>+</sup>) myofibers in the soleus muscles of the Mdx-GFP mice (Supporting Information Fig. S4A, S4B) or in the cardiotoxin (CTX)-damaged TA muscles (Supporting Information Fig. S4C, S4D), indicating that BM-derived myofibers are negligible, if any, in the mdx mouse after BMT. However, we found significantly increased collagen deposition, which is a marker of fibrosis, in soleus muscles of the Mdx-mdx mice compared with those of Mdx-wt and Mdx (age-matched non-BMT) mice 7 weeks after BMT (Fig. 1B). The aged-matched, non-BMT mdx mice showed less collagen staining compared with the Mdx-wt mice (Fig. 1B), suggesting certain adverse effects of the lethal-dose irradiation in the BMT procedure on muscle pathology in mdx mice.

To obtain quantitative data for fibrosis [27], we measured the levels of fibronectin and MMP13 in soleus muscles 7 weeks after BMT and found that these fibrosis markers were also significantly higher in the Mdx-mdx mice than in Mdx-wt and Mdx mice (Fig. 1C, 1D; Supporting Information Fig. S1C).

MMP13 expression is regulated by TNF- $\alpha$ . TNF- $\alpha$  expression tended to be higher in Mdx-mdx mice than in Mdx-wt and Mdx mice (Supporting Information Fig. S1D), suggesting that TNF- $\alpha$  and MMP13 participate, at least in part, in fibrotic deposition.

In contrast, the expression of TNF- $\alpha$ -stimulated gene/protein-6 (TSG-6), which is an anti-inflammatory mediator, was slightly higher in Mdx-wt mice compared with Mdx-mdx mice (Supporting Information Fig. S1E). EBD staining of soleus muscles 7 weeks after BMT showed greatly enhanced necrotic tissue damage in Mdx-mdx mice compared with Mdx-wt and Mdx mice (Fig. 1E, 1F), suggesting increased necrosis by excess inflammation or decreased overall repair mechanisms in Mdx-mdx mice.


We next analyzed the muscle strength and endurance of those BMT mice from 4 to 7 weeks after BMT using grip strength (Fig. 1G) and two different hang wire tests (Supporting Information Fig. S1F, S1G) [28]. Consistent with the deterioration of muscle pathology observed in the Mdx-mdx mice, the muscle functions in Mdx-mdx mice were significantly lower than those of Mdx-wt and Mdx mice (Fig. 1G; Supporting Information Fig. S1F, S1G). These results suggest that mdx-related alterations in BM have significant adverse effects on the progression of muscular degeneration, whereas normal BMT in mdx mice can rescue this mechanism without dystrophin supplementation.

### Chronic Muscle Injury in mdx Mice Induced a Reduction of MSC Populations in BM

In the mdx mouse, massive muscle degeneration starts at approximately 3–4 weeks of age, and the degeneration/regeneration cycle continues throughout the lifespan, although muscle degeneration is milder after 12 weeks of age [38]. We thus examined whether the numbers of BM-MSCs in 10-week-old mdx mice were altered as a result of the systemic effect of the chronic muscle degeneration/regeneration cycle. We used a CFU-F assay to determine the total numbers of MSCs in the BM of 10-week-old mdx mice.

Strikingly, the number of BM CFU-Fs in 10-week-old mdx mice was significantly decreased compared with those in the WT mice (Fig. 2A). BM-MSCs have been shown to be highly heterogeneous. We thus investigated the subpopulations of BM-MSCs in 10-week-old mdx mice by conducting flow cytometry analysis. Both CD106 and CD44 have been reported as being expressed in cultured BM mesenchymal "stem" cells [6]. However, a recent report suggests that freshly isolated BM mesenchymal stem cells do not express CD44, and therefore, the expression of CD44 on MSCs in BM remains controversial [18].

Interestingly, in this study, both the Lin<sup>−</sup>/ckit<sup>−</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> and Lin<sup>−</sup>/ckit<sup>−</sup>/CD106<sup>+</sup>/CD44<sup>−</sup> subpopulations in BM were greatly reduced in 10-week-old mdx mice (Fig. 2B, 2C), but the percentage of the Lin<sup>+</sup>/ckit<sup>−</sup> population in BM, which includes mature hematopoietic lineage cells [24], was not significantly different between the 10-week-old mdx and WT mice (Supporting Information Fig. S5A). These differences between WT and mdx mice were not observed at 3 weeks of age, before or immediately after the onset of DMD (Supporting Information Fig. S5B–S5F). The results thus suggest that the reduction of these BM subpopulations depends on DMD progression.



**Figure 1.** Effects of 10-week-old WT or mdx bone marrow cell (BMC) transplantation on Duchenne muscular dystrophy muscle pathology. **(A):** Schematic illustration of the experimental models. Either 10-week-old WT mouse BMCs or mdx mouse BMCs were transplanted into 3–4-week-old mdx mice after lethal-dose irradiation (Mdx-wt and Mdx-mdx). **(B):** Representative soleus muscle sections of Mdx (age-matched non-BMT control), Mdx-wt, and Mdx-mdx mice stained by picrosirius red 7 weeks after BMT. Scale bar = 200  $\mu$ m. **(C):** Western blot analyses of fibronectin and Pan-actin (loading control) of soleus muscle 7 weeks after BMT. **(D):** Quantified band densities of fibronectin/actin. The values are the mean  $\pm$  SEM; \*,  $p < .05$ . **(E):** Representative soleus muscle sections of Mdx (age-matched non-BMT control), Mdx-wt, and Mdx-mdx mice visualized by Evans blue dye (EBD) 7 weeks after BMT. Scale bar = 100  $\mu$ m. **(F):** The number of EBD<sup>+</sup> fibers was quantified. The values are the mean  $\pm$  SEM ( $n = 4$ ); \*,  $p < .05$ . **(G):** Measurement of grip strength in Mdx (age-matched non-BMT control), Mdx-wt, and Mdx-mdx mice from 4 to 7 weeks after BMT. The values are the mean  $\pm$  SEM ( $n = 7$ –8/group); \*,  $p < .05$ . Abbreviation: BMT, bone marrow transplantation.


To exclude the possibility that these reductions of specific BM subpopulations were simply the result of different degrees of cellularity in dystrophin-deficient BMCs, we established three types of BMT models (Fig. 2D). BMCs from 3-week-old WT mice (donor) were transplanted into either 3-week-old WT (WT-wt) or mdx mice (Mdx-wt) after a lethal dose of irradiation, which replaces the host BMCs with donor BMCs (Supporting Information Fig. S2A, S2B). As another control, BMCs from a 3-week-old mdx mouse (donor), which had BM subpopulations and CFU-F numbers identical to those of WT mice (Supporting Information Fig. S5B–S5F), were transplanted into 3-week-old WT mice (WT-mdx). Both the  $\text{Lin}^-/\text{ckit}^+/\text{CD106}^+/\text{CD44}^+$  and  $\text{Lin}^-/\text{ckit}^+/\text{CD106}^+/\text{CD44}^-$

subpopulations in the Mdx-wt mice were significantly lower than those in the WT-wt and WT-mdx mice (Fig. 2E, 2F). These results suggest that both the  $\text{Lin}^-/\text{ckit}^+/\text{CD106}^+/\text{CD44}^+$  and  $\text{Lin}^-/\text{ckit}^+/\text{CD106}^+/\text{CD44}^-$  BM subpopulations were gradually reduced by chronic exposure to a muscle injury environment after the onset of muscular dystrophy.

#### Whole-Transcriptome Analysis of the Reduced MSC Populations in mdx Mice

We further investigated the precise properties of endogenous  $\text{Lin}^-/\text{ckit}^+/\text{CD106}^+/\text{CD44}^+$  and  $\text{Lin}^-/\text{ckit}^+/\text{CD106}^+/\text{CD44}^-$  BM populations. PDGFR $\alpha$  has been reported to be expressed in mesenchymal lineage cells [39, 40]. We therefore examined






**Figure 2.** Reduction of mesenchymal stromal cell populations in bone marrow (BM) by chronic muscle injury in mdx mice. **(A):** The numbers of Lin<sup>−</sup> cells in the BM of 10-week-old WT and Mdx mice were analyzed in a CFU-F assay. The frequency of CFU-Fs is expressed as the mean ± SEM (*n* = 6/group); \*, *p* < .05. **(B):** Representative fluorescence-activated cell sorting (FACS) profiles of BMCs in 10-week-old WT and Mdx mice BM, stained with Lin, ckit, CD106, and CD44. FACS charts gated within Lin<sup>−</sup>/ckit<sup>−</sup> cells are shown. **(C):** The percentages of CD106<sup>+</sup>/CD44<sup>+</sup> and CD106<sup>+</sup>/CD44<sup>−</sup> cells gated within Lin<sup>−</sup>/ckit<sup>−</sup> cells were quantified. The values are the mean ± SEM (*n* = 6/group); \*, *p* < .05. **(D):** Scheme of the bone marrow transplantation (BMT) models, which indicates the following: (1) 3-week-old WT BMCs were transplanted into preirradiated 3-week-old WT mice (WT-wt); (2) 3-week-old WT BMCs were transplanted into preirradiated 3-week-old Mdx mice (Mdx-wt); and (3) 3-week-old Mdx BMCs were transplanted into preirradiated 3-week-old WT mice (WT-mdx). **(E):** Representative FACS profiles of BMCs 7 weeks after BMT, stained with Lin, ckit, CD106, and CD44. FACS plot gated within Lin<sup>−</sup>/ckit<sup>−</sup> cells is shown. **(F):** The percentages of CD106<sup>+</sup>/CD44<sup>+</sup> and CD106<sup>+</sup>/CD44<sup>−</sup> cells gated within Lin<sup>−</sup>/ckit<sup>−</sup> cells were quantified. The values are the mean ± SEM (*n* = 4/group); \*, *p* < .05. Abbreviation: BMC, bone marrow cell.

PDGFR $\alpha$  expression in these BM populations using PDGFR $\alpha$ -H2B GFP mice (see Materials and Methods). Lin<sup>−</sup>/ckit<sup>−</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> and Lin<sup>−</sup>/ckit<sup>−</sup>/CD106<sup>+</sup>/CD44<sup>−</sup> BM populations comprised 19.9% and 71.9% of the PDGFR $\alpha$ -expressing cells, respectively (Fig. 3A, 3B), suggesting that both of these BM populations contain PDGFR $\alpha$ <sup>+</sup> mesenchymal lineage cells.

Next, we conducted a whole-transcriptome analysis by RNA sequencing (RNA seq) analyses to further investigate the molecular properties of these populations. We sorted Lin<sup>−</sup>/ckit<sup>−</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> and Lin<sup>−</sup>/ckit<sup>−</sup>/CD106<sup>+</sup>/CD44<sup>−</sup> BMCs. The transcript of CD106 mRNA was enriched in the Lin<sup>−</sup>/ckit<sup>−</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> and Lin<sup>−</sup>/ckit<sup>−</sup>/CD106<sup>+</sup>/CD44<sup>−</sup> populations, but was not detected in any other populations (Fig. 3C), indicating a successful enrichment of these two populations. Our comparison of the whole-genome transcriptome revealed the enrichment of different functional clusters in each population (Fig. 3D, 3E). The Lin<sup>−</sup>/ckit<sup>−</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> population showed significantly higher expression levels of

genes, including DNA replication- and cell cycle-related genes (Fig. 3D), such as Myc and Mcm2 (Fig. 3F). In contrast, these genes were not expressed in the Lin<sup>−</sup>/ckit<sup>−</sup>/CD106<sup>+</sup>/CD44<sup>−</sup> population. The Lin<sup>−</sup>/ckit<sup>−</sup>/CD106<sup>+</sup>/CD44<sup>−</sup> population showed significantly higher expression levels of genes, including cell adhesion- and extracellular matrix receptor interaction-related genes (Fig. 3E). CXCL12 (SDF-1 $\alpha$ ), Col1A, and FGFR1, which have been reported to be expressed in HSC niche cells [10, 14, 41] (Fig. 3G; Supporting Information Fig. S6).

One of the features associated with stem cell populations is cell-cycle quiescence under physiological conditions. Therefore, to further investigate the differences in the expression of cell cycle-related genes between the two cell populations observed in the RNA seq analysis, we analyzed the expression of the nuclear proliferation marker Ki67 in a flow cytometry analysis. We found that approximately 50% of the Lin<sup>−</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> population was Ki67<sup>+</sup>, whereas only 10% of the Lin<sup>−</sup>/CD106<sup>+</sup>/CD44<sup>−</sup> population was Ki67<sup>+</sup>



**Figure 3.** Prospective characterization of freshly isolated mesenchymal stromal cell populations in bone marrow. **(A):** Representative fluorescence-activated cell sorting (FACS) profiles of total BMCs in PDGFR $\alpha$ -H2BGFP knockin mouse. The percentage of PDGFR $\alpha$ -H2BGFP $^{+}$  cells in total BMCs is shown (left). CD106 and CD44 expression profiles in PDGFR $\alpha$ -GFP $^{+}$  cells (right). **(B):** The percentages of CD106 $^{+}$ /CD44 $^{+}$  and CD106 $^{+}$ /CD44 $^{-}$  cells in PDGFR $\alpha$ -GFP $^{+}$  cells were quantified. The values are the mean  $\pm$  SEM ( $n = 3$ /group); \*,  $p < .05$ . **(C):** CD106 mRNA expressions in sorted BMCs. The values are the mean  $\pm$  SEM. The data are for three independent sorting experiments. **(D, E):** Gene ontology (GO) analysis of RNA seq data from sorted Lin $^{-}$ /ckit $^{-}$ /CD106 $^{+}$ /CD44 $^{+}$  and Lin $^{-}$ /ckit $^{-}$ /CD106 $^{+}$ /CD44 $^{-}$  BMCs. Genes that are significantly upregulated in Lin $^{-}$ /ckit $^{-}$ /CD106 $^{+}$ /CD44 $^{+}$  BMCs (197 genes) or Lin $^{-}$ /ckit $^{-}$ /CD106 $^{+}$ /CD44 $^{-}$  cells (342 genes) were categorized by GO annotations. **(F):** RNA-seq results showing the expression of cell cycle-associated genes (Myc and Mcms) in sorted Lin $^{-}$ /ckit $^{-}$ /CD106 $^{+}$ /CD44 $^{+}$  and Lin $^{-}$ /ckit $^{-}$ /CD106 $^{+}$ /CD44 $^{-}$  BMCs. **(G):** RNA-seq results showing the expressions of CXCL12 (SDF-1 $\alpha$ ) and Col1a1 in sorted Lin $^{-}$ /ckit $^{-}$ /CD106 $^{+}$ /CD44 $^{+}$  and Lin $^{-}$ /ckit $^{-}$ /CD106 $^{+}$ /CD44 $^{-}$  BMCs. The RNA-seq data were obtained from two biological replicates. **(H):** Representative FACS profiles of Ki67 expression on Lin $^{-}$ /CD106 $^{+}$ /CD44 $^{+}$  and Lin $^{-}$ /CD106 $^{+}$ /CD44 $^{-}$  BMCs. **(I):** The percentages of Ki67 $^{+}$  cells on each population were quantified. The values are the mean  $\pm$  SEM ( $n = 3$ ); \*,  $p < .05$ . **(J, K):** The expressions of cdk6 (J) and p21 (K). The data are for three independent sorting experiments. The values are the mean  $\pm$  SEM; \*,  $p < .05$ . Abbreviations: BMC, bone marrow cell; FSC, forward scatter; GFP, green fluorescent protein.

(Fig. 3H, 3I). Real-time PCR also revealed that p21, a cell cycle inhibitor gene, showed reduced expression in the  $\text{Lin}^-/\text{ckit}^-/\text{CD106}^+/\text{CD44}^+$  population than in the  $\text{Lin}^-/\text{ckit}^-/\text{CD106}^+/\text{CD44}^-$  population (Fig. 3J). In contrast, cyclin-dependent kinase 6 (cdk6) showed higher expression in the  $\text{Lin}^-/\text{ckit}^-/\text{CD106}^+/\text{CD44}^+$  population than in the  $\text{Lin}^-/\text{ckit}^-/\text{CD106}^+/\text{CD44}^-$  population (Fig. 3K). Together, these data suggest that  $\text{Lin}^-/\text{ckit}^-/\text{CD106}^+/\text{CD44}^+$  and  $\text{Lin}^-/\text{ckit}^-/\text{CD106}^+/\text{CD44}^-$  BM populations, which were decreased in 10-week-old mdx mice, were a proliferative and a slow-dividing mesenchymal lineage, respectively.

### Endogenous Migration of BM-Derived MSC Population in Injured Muscles

We then investigated the recruitment of BM-derived mesenchymal populations into damaged muscles. GFP BMCs were transplanted in 3–4-week-old mdx mice (Mdx-GFP), and we analyzed the characteristics of the BM-derived cells recruited to the damaged muscles 4 weeks after BMT by flow cytometry. We found that the  $\text{Lin}^-/\text{GFP}^+$  cells, which include BM-derived MSC populations (Supporting Information Fig. S7A, S7B), were substantially recruited into the damaged muscle of the mdx mice compared with WT muscles (red squares in Fig. 4A, 4B). Importantly, most of the  $\text{Lin}^-/\text{GFP}^+$  cells in the damaged muscles expressed both CD106 and CD44 on their surface (red squares in Fig. 4C; Supporting Information Fig. S7C), suggesting that BM-derived  $\text{Lin}^-/\text{CD106}^+/\text{CD44}^+$  cells predominantly accumulated into damaged muscle in mdx mice compared with other  $\text{Lin}^-/\text{GFP}^+$  populations.

These results led us to examine the possible recruitment mechanism of this population. We found that CXCR4 was highly expressed in freshly isolated  $\text{Lin}^-/\text{ckit}^-/\text{CD106}^+/\text{CD44}^+$  BMCs by the RNA seq (Fig. 4E) and flow cytometry analyses (Supporting Information Fig. S7D). CXCR4 is a receptor for SDF-1 $\alpha$  and has been found to be involved in the migration of stem/progenitor cells in various tissues. To elucidate whether the SDF-1 $\alpha$ /CXCR4 axis plays a role in recruiting  $\text{Lin}^-/\text{ckit}^-/\text{CD106}^+/\text{CD44}^+$  BMCs into damaged muscles, we first compared CXCR4 expression in the  $\text{Lin}^-/\text{ckit}^-/\text{CD106}^+/\text{CD44}^+$  BMCs of WT and mdx mice. We found that CXCR4 expression was slightly but not significantly elevated in the  $\text{Lin}^-/\text{ckit}^-/\text{CD106}^+/\text{CD44}^+$  BM populations of mdx mice compared with those of the WT mice (Supporting Information Fig. S7E), suggesting that elevated CXCR4 levels are not responsible for the specific migration of  $\text{Lin}^-/\text{ckit}^-/\text{CD106}^+/\text{CD44}^+$  BMCs into damaged muscles.

Next, we assessed SDF-1 $\alpha$  expression in mdx soleus muscles. SDF-1 $\alpha$  was specifically detected in mdx muscles but not WT muscles, and CD68 $^+$  cells primarily colocalized with SDF-1 $\alpha$  signals (Supporting Information Fig. S7F). We also detected significant increases in  $\text{Lin}^-/\text{ckit}^-/\text{CD106}^+/\text{CD44}^+/\text{CXCR4}^+$  cells in the peripheral blood of mdx mice (Fig. 4F). In addition, we implanted silicon tubes containing recombinant SDF-1 $\alpha$  under the skin of mdx mice. Two days later, we collected the tubes and counted the numbers and percentages of adherent  $\text{CD106}^+/\text{CD44}^+$  tube-trapped cells. As shown in Supporting Information Figure S7G, S7H, significantly more  $\text{CD106}^+/\text{CD44}^+$  cells migrated into the SDF-1 $\alpha$ -containing tubes compared with the tubes containing PBS. These results suggest that the migration of  $\text{Lin}^-/\text{ckit}^-/\text{CD106}^+/\text{CD44}^+$  BMCs into damaged muscles although the circulation may be

partially explained, if not exclusively so, by the CXCR4-SDF-1 $\alpha$  pathway.

### Function of the BM-Derived MSC Population in Damaged Muscles

Because the TSG-6-mediated immunosuppressive function of cultured expanded MSCs has been documented [42], we next investigated the expression of TSG-6 in the BM-derived  $\text{Lin}^-/\text{CD106}^+/\text{CD44}^+$  population in the damaged muscles of Mdx-GFP mice. We found that BM-derived ( $\text{GFP}^+$ )/ $\text{Lin}^-/\text{CD106}^+/\text{CD44}^+$  cells significantly and preferentially expressed TSG-6 in damaged muscles compared with other BM-derived populations (Fig. 5A). However, interleukin-10, which has also been reported to be expressed in BM-MSCs, was not detected (data not shown). It is well known that TSG-6 suppresses inflammation by interacting with CD44 on inflammatory cells, but we also observed the expression of CD44 on the C2C12 myoblasts and primary satellite cells (Fig. 5B; Supporting Information Fig. S8A, S8B). Recombinant TSG-6 (rTSG-6) significantly augmented the proliferation of C2C12 myoblasts (Fig. 5C), suggesting that TSG-6 from BM-MSCs also activated myoblasts.


A previous study suggested that damage signals enhanced TSG-6 expression in BM-MSCs [3], and thus, we investigated whether the treatment of primary BM-MSCs with mdx mouse serum (which contains damage signals) would enhance TSG-6 expression (Fig. 5D). As shown in Figure 5E, elevated TSG-6 expression in mdx serum-treated BM-MSCs was observed, suggesting that the recruited BM-MSC population exposed to damage signals in mdx mice likely elevates TSG-6 expression to suppress inflammation and to activate myoblasts in the damaged muscles.

We also cocultured C2C12 myoblasts with conditioned medium (CM) from BM-MSCs treated with mdx mouse serum for 48 hours (Fig. 5D). We observed a significant proliferation of C2C12 myoblasts following CM treatment (Fig. 5F). However, the CM-mediated C2C12 proliferation was moderately but significantly blocked in the presence of a CD44-neutralizing antibody (Fig. 5F). These data suggest that the TSG-6 secreted from activated BM-MSCs stimulates myoblast activation.

### BM-Derived $\text{Lin}^-/\text{ckit}^-/\text{CD106}^+/\text{CD44}^+$ Cells Accelerated Muscle Regeneration

Lastly, we examined whether the  $\text{Lin}^-/\text{ckit}^-/\text{CD106}^+/\text{CD44}^+$  BMCs, the main source of TSG-6 in damaged muscles, could support muscle regeneration. To minimize the effect of the endogenous BM-MSCs population during muscle regeneration, we generated mice with reduced MSCs in BM ( $\text{ckit}^+$  BMT) by transplanting a sorted  $\text{Lin}^-/\text{ckit}^+$  population (including HSCs but not MSCs) after lethal-dose irradiation (Supporting Information Fig. S3A). As shown in Supporting Information Figure S3B, S3C, the percentage of the  $\text{Lin}^-/\text{ckit}^-$  population in the BM, which includes an MSC population, was significantly less in the  $\text{ckit}^+$  BMT mice compared with the total BMT mice after 6 weeks of reconstitution. Furthermore, fewer donor BM-derived fibroblastic cells were observed in the  $\text{ckit}^+$  BMT mice compared with the total BMT mice (Supporting Information Fig. S3D–S3F). These data indicate that  $\text{ckit}^+$  BMT mice have few endogenous MSCs in their BM.






**Figure 4.** Identification of bone marrow (BM)-mesenchymal stromal cell populations in mdx muscles. **(A):** Flow cytometry analysis in muscles of WT-GFP (left) and Mdx-GFP (right) bone marrow transplantation (BMT) mice performed 4 weeks after BMT. The red squares indicate Lin<sup>-</sup>/GFP<sup>+</sup> cells (BM-derived nonhematopoietic cells). **(B):** The percentages of Lin<sup>-</sup>/GFP<sup>+</sup> cells were quantified. The values are the mean  $\pm$  SEM ( $n = 4$ ); \*,  $p < .05$ . **(C):** CD106 and CD44 expression in Lin<sup>-</sup>/GFP<sup>+</sup> cells in the muscles of Mdx-GFP BMT mice. Most of the Lin<sup>-</sup>/GFP<sup>+</sup> cells were positive for both CD106 and CD44 (red squares). The percentages of CD106<sup>+</sup>/CD44<sup>+</sup> cells in Lin<sup>-</sup>/GFP<sup>+</sup> cells in mdx muscles are shown with red squares. **(D):** The percentages of Lin<sup>-</sup>/GFP<sup>+</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> cells in mdx muscles were evaluated. The values are the mean  $\pm$  SEM ( $n = 4$ ); \*,  $p < .05$ . **(E):** CXCR4 expression from the RNA seq analysis of sorted Lin<sup>-</sup>/GFP<sup>+</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> and Lin<sup>-</sup>/GFP<sup>+</sup>/CD106<sup>+</sup>/CD44<sup>-</sup> bone marrow cells. **(F):** The percentages of Lin<sup>-</sup>/GFP<sup>+</sup>/CD106<sup>+</sup>/CD44<sup>+</sup>/CXCR4<sup>+</sup> cells in the peripheral blood of WT and mdx mice were quantified by flow cytometry. The values are the mean  $\pm$  SEM ( $n = 6-8$ ); \*,  $p < .05$ . Abbreviation: GFP, green fluorescent protein.

Using the ckit<sup>+</sup> BMT model, we investigated whether an intramuscular injection of Lin<sup>-</sup>/GFP<sup>+</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> BMCs would enhance muscle regeneration in CTX-induced damaged muscles of WT-ckit<sup>+</sup> BMT mice (Supporting Information Fig. S9). At 14 days post-treatment, H&E staining showed well-arranged

myofibers with fewer mononuclear cells in damaged muscles treated with Lin<sup>-</sup>/GFP<sup>+</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> BMCs compared with PBS-treated or Lin<sup>-</sup>/GFP<sup>+</sup>/CD106<sup>+</sup>/CD44<sup>-</sup> BMCs-treated damaged muscles, which were the second-most recruited BMCs into damaged muscles (Supporting Information Fig. S9A).



**Figure 5.** Effect of TSG-6 secreted from BM-MSC populations on myoblasts. **(A):** TSG-6 expression in sorted BM-derived cells from mdx muscles. The values are the mean  $\pm$  SEM. The data are for three independent sorting experiments; \*,  $p < .05$ . **(B):** Representative fluorescence-activated cell sorting profiles of CD44 expression on C2C12 myoblasts (red line). Isotype control staining is represented by the black line. **(C):** The number of C2C12 myoblasts treated with TSG-6 after 48 hours. **(D):** Strategy for the preparation of CM from primary BM-MSCs. **(E):** Western blot analyses of TSG-6 and Pan-actin (loading control) expression in primary BM-MSCs prepared as in panel (D). **(F):** The number of C2C12 myoblasts treated with CM containing either control IgG (rat IgG) or CD44-blocking antibody (KM81) for 48 hours. The values are the mean  $\pm$  SEM ( $n = 3$ ); \*,  $p < .05$ . Abbreviations: BM-MSCs, bone marrow-mesenchymal stromal cells; CM, conditioned medium; FBS, fetal bovine serum; PBS, phosphate-buffered saline.

We quantified newly regenerating myofibers or myotubes by counting the desmin<sup>+</sup> cells with central nuclei, and the results indicated that the Lin<sup>-</sup>/ckit<sup>-</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> BMCs significantly increased the number of desmin<sup>+</sup> cells with central nuclei compared with the number in PBS-treated damaged muscles (Supporting Information Fig. S9B, S9C). We also investigated eMyHC<sup>+</sup> cell areas in damaged muscle sections. The percentage area of the eMyHC<sup>+</sup> cell area was increased by Lin<sup>-</sup>/ckit<sup>-</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> BMC treatment compared with PBS- or Lin<sup>-</sup>/ckit<sup>-</sup>/CD106<sup>-</sup>/CD44<sup>+</sup> BMC-treated damaged muscle (Supporting Information Fig. S9D, S9E). These data suggest that Lin<sup>-</sup>/ckit<sup>-</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> BMCs accelerate muscle regeneration processes in CTX-induced muscle damage.

We next investigated whether Lin<sup>-</sup>/ckit<sup>-</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> BMC treatment supports muscle regeneration in Mdx mice. Using Mdx-ckit<sup>+</sup> BMT mice reconstituted for 6 weeks (10-week-old mice), we first assessed the effect of Lin<sup>-</sup>/ckit<sup>-</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> BMC treatment on satellite cell activation 7 days after cell injection. The injection of Lin<sup>-</sup>/ckit<sup>-</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> BMCs increased the number of activated satellite cells, as indicated by MyoD staining (Fig. 6A, 6B). We also observed slightly elevated MyoD expression after Lin<sup>-</sup>/ckit<sup>-</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> BMC treatment by Western blotting (Fig. 6C). However, the elevation of the number of MyoD<sup>+</sup> satellite cells was partially inhibited by the administration of a CD44-neutralizing antibody or a TSG-6 antibody (Fig. 6A, 6B). MyoD expression also tended to be decreased by the CD44-neutralizing antibody or TSG-6 antibody (Fig. 6C). The number of Pax7<sup>+</sup> cells and the expression level of Pax7, which is a quiescent satellite cell marker, were not significantly changed

by Lin<sup>-</sup>/ckit<sup>-</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> BMC treatment (Supporting Information Fig. S10A–S10C).

We also observed that Lin<sup>-</sup>/ckit<sup>-</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> BMC treatment suppressed the infiltration of inflammatory cells and enhanced the overall muscle repair mechanisms in Mdx-ckit<sup>+</sup> BMT mice, as assessed by the average muscle fiber CSA at day 7 post-treatment (Fig. 6D, 6E). However, again, the effects of Lin<sup>-</sup>/ckit<sup>-</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> BMC treatment on muscle regeneration were partially blocked by a CD44-neutralizing antibody or a TSG-6 antibody (Fig. 6D, 6E). Of note, recombinant TSG-6 treatment also enhanced muscle regeneration with Lin<sup>-</sup>/ckit<sup>-</sup>/CD106<sup>-</sup>/CD44<sup>+</sup> BMC treatment (Fig. 6D, 6E). These data may indicate that the BM-derived Lin<sup>-</sup>/ckit<sup>-</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> population slowed muscle disease progression by the activation of satellite cells and the suppression of inflammation to enhance the overall repair mechanisms, in part via the TSG-6/CD44-mediated pathway.

## DISCUSSION

In this study, we demonstrated for the first time that injured muscular tissues in DMD mice recruit a Lin<sup>-</sup>/ckit<sup>-</sup>/CD106<sup>+</sup>/CD44<sup>+</sup> mesenchymal population from the BM to suppress inflammatory and fibrotic reactions and activate a muscular regeneration mechanism. Such chronic injury/regeneration cycles in DMD mice were shown to eventually decrease the MSC populations in BM, resulting in a disruption of the intrinsic anti-inflammatory and regeneration-promoting activities of BM-MSCs to exacerbate muscular dystrophy (Fig. 7).