

別紙4

研究成果の刊行に関する一覧表

雑誌

発表者氏名	論文タイトル名	発表誌名	出版年	巻号	ページ
Azuma K, Okamoto I, Kawahara A, Taira T, Nakashima K, Hattori S, Kinoshita T, Takeda M, <u>Nakagawa K</u> , Takamori S, Kuwano M, Ono M, Kage M.	Association of the Expression of Mutant Epidermal Growth Factor Receptor Protein as Determined with Mutation-Specific Antibodies in Non-small Cell Lung Cancer with Progression-Free Survival after Gefitinib Treatment.	J Thorac Oncol	2012	7(1)	122-127
Okamoto K, Okamoto I, Hatashita E, Kuwata K, Yamaguchi H, Kita A, Yamanaka K, Ono M, <u>Nakagawa K</u> .	Overcoming erlotinib resistance in EGFR mutation-positive non-small cell lung cancer cells by targeting survivin	Mol Cancer Ther	2012	11(1)	204-213
Takeda M, Okamoto I, Tsurutani J, Oiso N, Kawada A, <u>Nakagawa K</u> .	Clinical Impact of Switching to a Second EGFR-TKI After a Severe AE Related to a First EGFR-TKI in EGFR-mutated NSCLC.	Jpn J Clin Oncol	2012	42(6)	528-533
Nakagawa K, Kudoh S, Ohe Y, Johkoh T, Ando M, Yamazaki N, Seki A, Takemoto S, Fukuoka M.	Postmarketing surveillance study of erlotinib in Japanese patients with non-small-cell lung cancer (NSCLC): an interim analysis of 3488 patients (POLARSTAR).	J Thorac Oncol	2012	7(8)	1296-1303
Aoyama Y, Nishimura T, Sawamoto T, Satoh T, Katashima M, <u>Nakagawa K</u> .	Pharmacokinetics of sepantromium bromide (YM155), a small-molecule suppressor of survivin, in Japanese patients with advanced solid tumors: dose proportionality and influence of renal impairment.	Cancer Chemother Pharmacol	2012	70(3)	373-380
Hayashi H, Kurata T, Fujisaka Y, Kawakami H, Tanaka K, Okabe T, Takeda M, Satoh T, Yoshida K, Tsunoda T, Arao T, Nishio K, <u>Nakagawa K</u> .	Phase I trial of OTS11101, an anti-angiogenic vaccine targeting Vascular Endothelial Growth Factor Receptor 1 in solid tumor.	Cancer Science	2013	104(1)	98-104
Kiyota H, Okamoto I, Takeda M, Daga H, Naito T, Miyazaki M, Okada H, Hayashi H, <u>Tanaka K</u> , Terashima M, Azuma K, Murakami H, Takeda K, Yamamoto N, <u>Nakagawa K</u> .	Phase I and pharmacokinetic study of gefitinib and S-1 combination therapy for advanced adenocarcinoma of the lung.	Cancer Chemother Pharmacol	2013	71(4)	859-865
Kogure Y, Ando M, Saka H, Chiba Y, Yamamoto N, Asami K, Hirashima T, Seto T, Nagase S, Otsuka K, Yanagihara K, Takeda K, Okamoto I, Aoki T, Takayama K, Yamasaki M, Kudo H, Katakami N, Miyazaki M, <u>Nakagawa K</u> .	Histology and Smoking Status Predict Survival of Patients with Advanced Non-Small-Cell Lung Cancer: Results of West Japan Oncology Group (WJOG) Study 3906L.	J Thorac Oncol	2013	8(6)	753-758
Akamatsu H, Inoue A, Mitsudomi T, Kobayashi K, <u>Nakagawa K</u> , Mori K, Nukiwa T, Nakanishi Y, Yamamoto N.	Interstitial Lung Disease Associated with Gefitinib in Japanese Patients with EGFR-mutated Non-small-cell Lung Cancer: Combined Analysis of Two Phase III Trials (NEJ 002 and WJTOG 3405).	Jpn J Clin Oncol	2013	43(6)	664-668
Seto T, Kiura K, Nishio M, <u>Nakagawa K</u> , Maemondo M, Inoue A, Hida T, Yamamoto N, Yoshioka H, Harada M, Ohe Y, Nogami N, Takeuchi K, Shimada T, Tanaka T, Tamura T.	CH5424802 (RO5424802) for patients with ALK-rearranged advanced non-small-cell lung cancer (AF-001JP study): a single-arm, open-label, phase 1-2 study.	Lancet Oncol	2013	14(7)	590-598

Shaw AT, Kim DW, Nakagawa K, Seto T, Crino L, Ahn MJ, De Pas T, Besse B, Solomon BJ, Blackhall F, Wu YL, Thomas M, O'Byrne KJ, Moro-Sibilot D, Camidge DR, Mok T, Hirsh V, Riely GJ, Iyer S, Tassell V, Polli A, Wilner KD, Jänne PA.	Crizotinib versus chemotherapy in advanced ALK-positive lung cancer.	N Engl J Med	2013	368(25)	2385-2394
Yang JC, Wu YL, Chan V, Kurnianda J, Nakagawa K, Saijo N, Fukuoka M, McWalter G, McCormack R, Mok TS.	Epidermal growth factor receptor mutation analysis in previously unanalyzed histology samples and cytology samples from the phase III Iressa Pan-ASia Study (IPASS).	Lung Cancer	2013	83(2)	174-181
Takeda M, Okamoto I, Nakagawa K.	Survival outcome assessed according to tumor response and shrinkage pattern in patients with EGFR mutation-positive non-small-cell lung cancer treated with gefitinib or erlotinib.	J Thorac Oncol	2014	9(2)	200-204
Hayashi H, Arao T, Matsumoto K, Kimura H, Togashi Y, Hirashima Y, Horita Y, Iwasa S, Okita NT, Honma Y, Takashima A, Kato K, Hamaguchi T, Shimada Y, Nakagawa K, Nishio K, Yamada Y.	Biomarkers of reactive resistance and early disease progression during chemotherapy plus bevacizumab treatment for colorectal carcinoma.	Oncotarget	2014	5(9)	2588-2599
Yoshida T, Zhang G, Smith MA, Lopez AS, Bai Y, Li J, Fang B, Koomen J, Rawal B, Fisher KJ, Chen AY, Kitano M, Morita Y, Yamaguchi H, Shibata K, Okabe T, Okamoto I, Nakagawa K, Haura EB.	Tyrosine phosphoproteomics identifies both codrivers and cotargeting strategies for T790M-related EGFR-TKI resistance in non-small cell lung cancer.	Clin Cancer Res	2014	20(15)	4059-4074
Kawakami H, Okamoto I, Okamoto W, Tanizaki J, Nakagawa K, Nishio K.	Targeting MET Amplification as a New Oncogenic Driver.	Cancers (Basel)	2014	6(3)	1540-1552
Hayashi H, Okamoto I, Tanizaki J, Tanaka K, Okuda T, Kato A, Nishimura Y, Nakagawa K.	Cystic Brain Metastasis in Non-Small-Cell Lung Cancer With ALK Rearrangement.	J Clin Oncol	2014	32(36)	e122-124
Seto T, Kato T, Nishio M, Goto K, Atagi S, Hosomi Y, Yamamoto N, Hida T, Maemondo M, Nakagawa K, Nagase S, Okamoto I, Yamanaka T, Tajima K, Harada R, Fukuoka M, Yamamoto N.	Erlotinib alone or with bevacizumab as first-line therapy in patients with advanced non-squamous non-small-cell lung cancer harbouring EGFR mutations (JO25567): an open-label, randomised, multicentre, phase 2 study.	Lancet Oncol	2014	15(11)	1236-1244
Gemma A, Kudoh S, Ando M, Ohe Y, Nakagawa K, Johkoh T, Yamazaki N, Arakawa H, Inoue Y, Ebina M, Kusumoto M, Kuwano K, Sakai F, Taniguchi H, Fukuda Y, Seki A, Ishii T, Fukuoka M.	Final safety and efficacy of erlotinib in the phase 4 POLARSTAR surveillance study of 10 708 Japanese patients with non-small-cell lung cancer.	Cancer Sci	2014	105(12)	1584-1590
Solomon BJ, Mok T, Kim DW, Wu YL, Nakagawa K, Mekhail T, Felip E, Cappuzzo F, Paolini J, Usari T, Iyer S, Reisman A, Wilner KD, Tursi J, Blackhall F; PROFILE 1014 Investigators.	First-line crizotinib versus chemotherapy in ALK-positive lung cancer.	N Engl J Med	2014	371(23)	2167-2177
Kawakami H, Okamoto I, Yonesaka K, Okamoto K, Shibata K, Shinkai Y, Sakamoto H, Kitano M, Tamura T, Nishio K, Nakagawa K.	The anti-HER3 antibody patritumab abrogates cetuximab resistance mediated by heregulin in colorectal cancer cells.	Oncotarget	2014	5(23)	11847-11856

Okamoto I, Miyazaki M, Takeda M, Terashima M, Azuma K, Hayashi H, <u>Kaneda H</u> , Kurata T, <u>Tsurutani J</u> , Seto T, Hirai F, Konishi K, Sarashina A, Yagi N, Kaiser R, Nakagawa K.	Tolerability of Nintedanib (BIBF 1120) in Combination with Docetaxel: A Phase 1 Study in Japanese Patients with Previously Treated Non-Small-Cell Lung Cancer.	J Thorac Oncol	2015	10(2)	346-352
Sakai K, Takeda M, Okamoto I, Nakagawa K, <u>Nishio K</u> .	Multiple regulatory mechanisms of hepatocyte growth factor expression in malignant cells with a short poly(dA) sequence in the HGF gene promoter.	Oncol Lett	2015	9(1)	405-410
Nishio M, Horike A, Murakami H, Yamamoto N, <u>Kaneda H</u> , Nakagawa K, Horinouchi H, Nagashima M, Sekiguchi M, Tamura T.	Phase I study of the HER3-targeted antibody patritumab (U3-1287) combined with erlotinib in Japanese patients with non-small cell lung cancer.	Lung Cancer	2015	88(3)	275-281
<u>Tsurutani J</u> , Kuroi K, <u>Iwasa T</u> , Miyazaki M, Nishina S, Makimura C, Tanizaki J, Okamoto K, Yamashita T, Aruga T, Shigekawa T, Komoike Y, Saeki T, <u>Nakagawa K</u> .	Phase I study of weekly nab-paclitaxel combined with S-1 in patients with human epidermal growth factor receptor type 2-negative metastatic breast cancer.	Cancer Sci	2015		inpress