

. 研究成果の刊行に関する一覧表

研究成果の刊行に関する一覧表

雑誌

| | 発表者氏名 | 論文タイトル名 | 発表誌名 | 巻号 | ページ | 出版年 |
|---|--|--|-----------------------|--------|----------|------|
| 1 | Satoh M, Ohkubo T, Asayama K, Murakami Y, Sakurai M, Nakagawa H, Iso H, Okayama A, Miura K, Imai Y, Ueshima H, Okamura T; Evidence for Cardiovascular Prevention From Observational Cohorts in Japan (EPOCH-JAPAN) Research Group. | Combined Effect of Blood Pressure and Total Cholesterol Levels on Long-Term Risks of Subtypes of Cardiovascular Death: Evidence for Cardiovascular Prevention From Observational Cohorts in Japan. | Hypertension. | 65(3) | 517-24 | 2015 |
| 2 | Asayama K, Satoh M, Murakami Y, Ohkubo T, Nagasawa SY, Tsuji I, Nakayama T, Okayama A, Miura K, Imai Y, Ueshima H, Okamura T; Evidence for Cardiovascular Prevention From Observational Cohorts in Japan (EPOCH-JAPAN) Research Group. | Cardiovascular risk with and without antihypertensive drug treatment in the Japanese general population: participant-level meta-analysis. | Hypertension. | 63(6) | 1189-97 | 2014 |
| 3 | Sugiyama D, Higashiyama A, Wakabayashi I, Kubota Y, Adachi Y, Hayashibe A, Kawamura K, Kuwabara K, Nishimura K, Kadota A, Nishida Y, Hirata T, Imano H, Miyamatsu N, Miyamoto Y, Sawamura T, Okamura T. | The relationship between lectin-like oxidized low-density lipoprotein receptor-1 ligands containing apolipoprotein B and cardio-ankle vascular index in healthy community inhabitants: The KOBE study. | J Atheroscler Thromb. | | In press | 2014 |
| 4 | Kubota Y, Higashiyama A, Imano H, Sugiyama D, Kawamura K, Kadota A, Nishimura K, Miyamatsu N, Miyamoto Y, Okamura T. | Serum polyunsaturated fatty acid composition and serum high-sensitivity C-reactive protein levels in healthy Japanese residents: the KOBE study. | J Nutr Health Aging. | | In press | 2015 |
| 5 | Hitara T, Higashiyama A, Kubota Y, Nishimura K, Sugiyama D, Kadota A, Imano H, Nishikawa T, Miyamatsu N, Miyamoto Y, Okamura T. | HOMA-IR values are associated with glycemic control in Japanese subjects without diabetes or obesity: the KOBE Study. | J Epidemiol. | | In press | 2015 |
| 6 | Xu D, Murakoshi N, Sairenchi T, Irie F, Igarashi M, Nogami A, Tomizawa T, Yamaguchi I, Yamagishi K, Iso H, Ota H, Aonuma K. | Anemia and Reduced Kidney Function as Risk Factors for New Onset of Atrial Fibrillation (from the Ibaraki Prefectural Health Study). | Am J Cardiol. | 115(3) | 328-33 | 2015 |
| 7 | Murakoshi N, Xu D, Sairenchi T, Igarashi M, Irie F, Tomizawa T, Tada H, Sekiguchi Y, Yamagishi K, Iso H, Yamaguchi I, Ota H, Aonuma K. | Prognostic impact of supraventricular premature complexes in community-based health checkups: The Ibaraki Prefectural Health Study. | Eur Heart J. | 36(3) | 170-78 | 2015 |
| 8 | Fujihara K, Sugawara A, Heianza Y, Sairenchi T, Irie F, Iso H, Doi M, Shimano H, Watanabe H, Sone H, Ota H. | Utility of the triglyceride level for predicting incident diabetes mellitus according to the fasting status and body mass index category: the Ibaraki | J Atheroscler Thromb. | 21(11) | 1152-69 | 2014 |

| | | Prefectural Health Study. | | | | |
|----|--|--|-------------------------------|--------|---|------|
| 9 | Tsujimoto T, Sairenchi T, Iso H, Irie F, Yamagishi K, Watanabe H, Tanaka K, Muto T, Ota H. | The dose-response relationship between body mass index and the risk of incident stage ≥ 3 chronic kidney disease in a general Japanese population: the Ibaraki prefectural health study (IPHS). | J Epidemiol. | 24(6) | 444-51 | 2014 |
| 10 | Tsukinoki R, Okamura T, Watanabe M, Kokubo Y, Higashiyama A, Nishimura K, Takegami M, Murakami Y, Okayama A, Miyamoto Y. | Blood pressure, low-density lipoprotein cholesterol, and incidences of coronary artery disease and ischemic stroke in Japanese: the Suita study. | Am J Hypertens. | 27(11) | 1362-69 | 2014 |
| 11 | Nishimura K, Okamura T, Watanabe M, Nakai M, Takegami M, Higashiyama A, Kokubo Y, Okayama A, Miyamoto Y. | Predicting coronary heart disease using risk factor categories for a Japanese urban population, and comparison with the Framingham risk score: the Suita study. | J Atheroscler Thromb. | 21(8) | 784-98 | 2014 |
| 12 | Gotoh S, Hata J, Ninomiya T, Hirakawa Y, Nagata M, Mukai N, Fukuhara M, Ikeda F, Shikata K, Kamouchi M, Kitazono T, Kiyohara Y. | Trends in the incidence and survival of intracerebral hemorrhage by its location in a Japanese community. | Circ J. | 78 | 403-409 | 2014 |
| 13 | Imamura T, Doi Y, Ninomiya T, Hata J, Nagata M, Ikeda F, Mukai N, Hirakawa Y, Yoshida D, Fukuhara M, Kitazono T, Kiyohara Y. | Non-high-density lipoprotein cholesterol and the development of coronary heart disease and stroke subtypes in a general Japanese population: the Hisayama Study. | Atherosclerosis. | 233 | 343-48 | 2014 |
| 14 | Kojima I, Ninomiya T, Hata J, Fukuhara M, Hirakawa Y, Mukai N, Yoshida D, Kitazono T, Kiyohara Y. | A low ankle brachial index is associated with an increased risk of cardiovascular disease: the Hisayama Study. | J Atheroscler Thromb. | 21 | 966-73 | 2014 |
| 15 | Kishimoto H, Hata J, Ninomiya T, Nemeth H, Hirakawa Y, Yoshida D, Kumagai S, Kitazono T, Kiyohara Y. | Midlife and late-life handgrip strength and risk of cause-specific death in a general Japanese population: the Hisayama Study. | J Epidemiol Community Health. | 68 | 663-68 | 2014 |
| 16 | Mukai N, Doi Y, Ninomiya T, Hirakawa Y, Nagata M, Yoshida D, Hata J, Fukuhara M, Nakamura U, Kitazono T, Kiyohara Y. | Trends in the prevalence of type 2 diabetes and prediabetes in community-dwelling Japanese subjects: the Hisayama Study. | J Diabetes Investig. | 5 | 162-69 | 2014 |
| 17 | Mukai N, Yasuda M, Ninomiya T, Hata J, Hirakawa Y, Ikeda F, Fukuhara M, Hotta T, Koga M, Nakamura U, Kang D, Kitazono T, Kiyohara Y. | Thresholds of various glycemic measures for diagnosing diabetes based on prevalence of retinopathy in community-dwelling Japanese subjects: the Hisayama Study. | Cardiovasc Diabetol. | 13(45) | http://www.cardiab.com/content/13/1/45 | 2014 |
| 18 | Izumaru K, Ninomiya T, Nagata M, Usui T, Yoshida D, Yonemoto K, Fukuhara M, Tsuruya K, Kitazono T, Kiyohara Y. | Serum 1,25-dihydroxyvitamin D and the development of kidney dysfunction in a Japanese community: the Hisayama Study. | Circ J. | 78 | 732-37 | 2014 |
| 19 | 立川佳美、Cologne JB、山田美智子、大石和佳、飛田あゆみ、古川恭治、 | 親の放射線被ばくと多因子疾患有病率との関連：被爆二世健康診断調査（第2報） | 広島医学 | 67(4) | 375-78 | 2014 |

| | | | | | | |
|----|---|---|------------------------|---------|---|------|
| | 高橋規郎、中村典、小笠晃太郎、赤星正純、藤原佐枝子、Shore RE. | | | | | |
| 20 | 斎藤重幸 | 高血圧 | 最新肥満症学 日本臨床増刊号 | 72 | 583-85 | 2014 |
| 21 | 斎藤重幸 | JNC8 ガイドライン | 血圧 | 21 | 496-97 | 2014 |
| 22 | 斎藤重幸 | 肥満症・メタボリックシンドローム、高血圧発症・進展の関わり | 臨床高血圧学 日本臨床 | 72 | 269-72 | 2014 |
| 23 | 斎藤重幸 | 日本高血圧学会高血圧治療ガイドライン (JSH) の変遷 | 循環器内科 | 14 | 385-91 | 2014 |
| 24 | 斎藤重幸 | 糖尿病合併高血圧の降圧目標をめぐって | 血圧 | 21 | 926-31 | 2014 |
| 25 | 斎藤重幸 | 高血圧 | Mod Physician. | 35 | 213-15 | 2015 |
| 26 | Yoshida Y, Akasaka H, Saitoh, S, Shimamoto K, Miura T. | Comparative effects of telmisartan and valsartan as add-on agents for hypertensive patients with morning blood pressure insufficiency controlled by amlodipine monotherapy. | Hypertens Res. | 37 | 225-31 | 2014 |
| 27 | Furuhashi M, Moniwa N, Mita T, Fuseya T, Ishimura S, Ohno K, Shibata S, Tanaka M, Watanabe Y, Akasaka H, Ohnishi H, Yoshida H, Takizawa H, Saitoh S, Ura N, Shimamoto K, Miura T. | Urinary angiotensin- converting enzyme 2 in hypertensive patients may be increased by Olmesartan, an angiotensin II Receptor Blocker. | Am J Hypertens. | 28(1) | 15-21 | 2015 |
| 28 | Ihira H, Furuna T, Mizumoto A, Makino K, Saitoh S, Ohnishi H, Shimada H, Makizako H. | Subjective physical and cognitive age among community-dwelling older peoples aged 75 years and Older: differences with chronological age and its associated factors. | Aging Ment Health. | | In press | 2014 |
| 29 | Fuseya T, Furuhashi M, Yuda S, Muranaka A, Kawamukai M, MitaT, Ishimura S, Watanabe Y, HoshinaK, Tanaka M, Ohno K, Akasaka H, OhnishiH, YoshidaH, Saitoh S, Shimamoto K, Miura T. | Elevation of circulating fatty acid-binding protein 4 is independently associated with left ventricular diastolic dysfunction in a general population. | Cardiovasc Diabetol. | 13(126) | http://www.cardiab.com/content/13/1/126 | 2014 |
| 30 | Miki T, Tobisaw T, Sato T, Tanno M, Yano T, Akasaka H, Kuno A, Ogasawara M, MuraseH, Saitoh S, MiuraT. | Does glycemic control reverse dispersion of ventricular repolarization in type 2 diabetes? | Cardiovasc Diabetol. | 13(125) | http://www.cardiab.com/content/13/1/125 | 2014 |
| 31 | Wang C, Yatsuya H, Tamakoshi K, Iso H, Tamakoshi A. | Milk drinking and mortality: findings from the Japan collaborative cohort study. | J Epidemiol. | 25(1) | 66-73 | 2015 |
| 32 | Kumagai Y, Chou WT, Tomata Y, Sugawara Y, Kakizaki M, Nishino Y, Tsuji I. | Dietary Patterns and colorectal cancer risk in Japan: the Ohsaki Cohort Study. | Cancer Causes Control. | 25(6) | 727-36 | 2014 |

| | | | | | | |
|----|---|---|--------------------------|---------|---------|------|
| 33 | Matsumoto A, Satoh M, Kikuya M, Ohkubo T, Hirano M, Inoue R, Hashimoto T, Hara A, Hirose T, Obara T, Metoki H, Asayama K, Hosokawa A, Totsune K, Hoshi H, Hosokawa T, Sato H, Imai Y. | Day-to-Day variability in home blood pressure is associated with cognitive decline: the Ohasama study. | Hypertension. | 63(6) | 1333-38 | 2014 |
| 34 | Imai E, Tsubota-Utsugi M, Kikuya M, Satoh M, Inoue R, Hosaka M, Metoki H, Fukushima N, Kurimoto A, Hirose T, Asayama K, Imai Y, Ohkubo T. | Animal protein intake is associated with higher-level functional capacity in elderly adults: the Ohasama study. | J Am Geriatr Soc. | 62(3) | 426-34 | 2014 |
| 35 | Satoh M, Hosaka M, Asayama K, Kikuya M, Inoue R, Metoki H, Utsugi MT, Hara A, Hirose T, Obara T, Mori T, Totsune K, Hoshi H, Mano N, Imai Y, Ohkubo T. | Aldosterone-to-renin ratio and nocturnal blood pressure decline assessed by self-measurement of blood pressure at home: the Ohasama Study. | Clin Exp Hypertens. | 36(2) | 108-14 | 2014 |
| 36 | Tsubota-Utsugi M, Satoh M, Hosaka M, Inoue R, Asayama K, Hirose T, Metoki H, Kikuya M, Imai Y, Ohkubo T. | Personality traits as predictors of decline in higher-level functional capacity over a 7-year follow-up in older adults: the Ohasama Study. | Tohoku J Exp Med. | 234(3) | 197-207 | 2014 |
| 37 | Satoh M, Matsumoto A, Hara A, Iwamori S, Obara T, Kikuya M, Metoki H, Hosaka M, Asayama K, Takahashi N, Sato H, Mano N, Imai Y, Ohkubo T. | A survey of self-medication practices and related factors in the general population: the Ohasama study. | Yakugaku Zasshi. | 134(12) | 1347-55 | 2014 |
| 38 | Sakurai M, Nakamura K, Miura K, Takamura T, Yoshita K, Nagasawa SY, Morikawa Y, Ishizaki M, Kido T, Naruse Y, Suwazono Y, Sasaki S, Nakagawa H. | Sugar-sweetened beverage and diet soda consumption and the 7-year risk for type 2 diabetes mellitus in middle-aged Japanese men. | Eur J Nutr. | 53(1) | 251-58 | 2014 |
| 39 | Sakurai M, Nakamura K, Miura K, Yoshita K, Takamura T, Nagasawa SY, Morikawa Y, Ishizaki M, Kido T, Naruse Y, Nakashima M, Nogawa N, Suwazono Y, Nakagawa H. | Association between serum thyroid stimulating hormone within the normal range and indices of obesity in Japanese men and women. | Intern Med. | 53 | 669-74 | 2014 |
| 40 | Yoneyama S, Sakurai M, Nakamura K, Morikawa Y, Miura K, Nakashima M, Yoshita K, Ishizaki M, Kido T, Naruse Y, Nogawa K, Suwazono Y, Sasaki S, Nakagawa H. | Associations between rice, noodle, and bread intake and sleep quality in Japanese men and women. | PLoS One. | 9(8) | e105198 | 2014 |
| 41 | Morikawa Y, Nakamura K, Sakurai M, Nagasawa SY, Ishizaki M, Nakashima M, Kido T, Naruse Y, Nakagawa H. | The effect of age on the relationships between work-related factors and heavy drinking. | J Occup Health. | 56 | 141-49 | 2014 |
| 42 | Nakamura K, Sakurai M, Miura K, Morikawa Y, Nagasawa SY, Ishizaki M, Kido T, Naruse Y, | HOMA-IR and the risk of hyperuricemia: A prospective study in non-diabetic Japanese men. | Diabetes Res Clin Pract. | 106 | 154-60 | 2014 |

| | | | | | |
|----|--|---|--------------------------------|-------|----------|
| | Nakashima M, Nogawa K, Suwazono Y, Nakagawa H. | | | | |
| 43 | Nakamura K, Sakurai M, Miura K, Morikawa Y, Nagasawa SY, Ishizaki M, Kido T, Naruse Y, Nogawa K, Nakashikma M, Suwazono Y, Nakagawa H. | Overall sleep status and high sensitivity C-reactive protein: a prospective study in a Japanese population. | J Sleep Res. | 23(6) | 717-27 |
| 44 | Makita S, Onoda T, Ohsawa M, Tanno K, Tanaka F, Omama S, Yoshida Y, Ishibashi Y, Itai K, Sakata K, Ohta M, Kurabayashi T, Ogasawara K, Ogawa A, Okayama A, Nakamura M. | Bradycardia is associated with future cardiovascular diseases and death in men from the general population. | Atherosclerosis. | 236 | 116-20 |
| 45 | Koeda Y, Tanaka F, Segawa T, Ohta M, Ohsawa M, Tanno K, Makita S, Ishibashi Y, Omama S, Onoda T, Nakamura M. | Usefulness of risk grading system using albuminuria for predicting cardiovascular events and all-cause death in chronic kidney disease: a population-based prospective cohort study in Japan. | Int J Cardiol. | 175 | 576-77 |
| 46 | Makita S, Tanaka F, Onoda T, Ohsawa M, Tanno K, Omama S, Yoshida Y, Ishibashi Y, Segawa T, Takahashi T, Satoh K, Itai K, Sakata K, Ohta M, Kurabayashi T, Ogasawara K, Ogawa A, Okayama A, Nakamura M. | Elevated blood pressure at the first measurement predicts cardiovascular disease independently from the subsequent second reading in men, but not in women. | Clin Exp Hypertens. | 37 | 39-44 |
| 47 | Ando A, Tanno K, Ohsawa M, Onoda T, Sakata K, Tanaka F, Makita S, Nakamura M, Omama S, Ogasawara K, Ishibashi Y, Kurabayashi T, Koyama T, Itai K, Ogawa A, Okayama A. | Associations of number of teeth with risks for all-cause mortality and cause-specific mortality in middle-aged and elderly men in the northern part of Japan: the Iwate-KENCO study. | Community Dent Oral Epidemiol. | 42 | 358-65 |
| 48 | Iso H, Imano H, Yamagishi K, Ohira T, Cui R, Noda H, Sato S, Kiyama M, Okada T, Hitsumoto S, Tanigawa T, Kitamura A; CIRCS Investigators. | Fasting and non-fasting triglycerides and risk of ischemic cardiovascular disease in Japanese men and women: the Circulatory Risk in Communities Study (CIRCS). | Atherosclerosis. | 237 | 361-68 |
| 49 | Chei CL, Yamagishi K, Ikeda A, Noda H, Maruyama M, Cui , Imano H, Kiyama M, Kitamura A, Asada T, Iso H; CIRCS Investigators. | C-reactive protein levels and risk of disabling dementia with and without stroke in Japanese: the Circulatory Risk in Communities Study (CIRCS). | Atherosclerosis. | 236 | 438-43 |
| 50 | Cui R, Yamagishi K, Imano H, Ohira T, Tanigawa T, Hitsumoto S, Kiyama M, Okada T, Kitamura A, Iso H; CIRCS investigators. | Relationship between the ankle-brachial index and the risk of coronary heart disease and stroke: the circulatory risk in communities study. | J Atheroscler Thromb. | 21 | 1283-89 |
| 51 | Ishikawa J, Ishikawa S, Kario K. | Relationships between the QTc interval and cardiovascular, stroke, or sudden cardiac mortality in the general Japanese population. | J Cardiol. | | In press |
| | | | | | 2014 |

| | | | | | | |
|----|--|--|---------------------------|-------|----------|------|
| 52 | Ishikawa J, Ishikawa S, Kario K. | Levels of cornell voltage and cornell product for predicting cardiovascular and stroke mortality and morbidity in the general Japanese population. | Circ J. | 78 | 465-75 | 2014 |
| 53 | Kakehi E, Kotani K, Ishikawa S, Gotoh T, Kayaba K, Nakamura Y, Kajii E. | Relationship between screening plasma glucose concentrations and cancer- and all-cause mortality: the Jichi Medical School (JMS) cohort study. | Journal of Public Health. | 22 | 505-11 | 2014 |
| 54 | Yamasaki K, Kayaba K, Ishikawa S. | Soy and Soy Products Intake, All-Cause Mortality, and Cause-Specific Mortality in Japan: The Jichi Medical School Cohort Study. | Asia Pac J Public Health. | | In press | 2014 |
| 55 | Uemura M, Yatsuya H, Hilawe EH, Li Y, Wang C, Chiang C, Otsuka R, Toyoshima, Tamakoshi K, Aoyama A. | Breakfast Skipping is Positively Associated with Incidence of Type 2 Diabetes Mellitus - Evidence from Aichi Workers' Cohort Study. | J Epidemiol. | 25(5) | In press | 2015 |
| 56 | Hilawe EH, Yatsuya H, Li Y, Uemura M, Wang C, Chiang C, Toyoshima H, Tamakoshi K, Zhang Y, Kawazoe N, Aoyama A. | Smoking and Diabetes: Is the Association Mediated by Adiponectin, Leptin, or C-reactive Protein? | J Epidemiol. | 25(2) | 99-109 | 2015 |
| 57 | Toyoshima H, Otsuka R, Hashimoto S, Tamakoshi K, Yatsuya H. | Body mass index-modified relationship of chronic mental stress with resting blood pressure during 5 years in Japanese middle-aged male workers. | Circ J. | 78(6) | 1379-86 | 2014 |
| 58 | Yatsuya H, Nihashi T, Li Y, Hotta Y, Matsushita K, Muramatsu T, Otsuka R, Matsunaga M, Yamashita K, Wang C, Uemura M, Harada A, Fukatsu H, Toyoshima H, Aoyama A, Tamakoshi K. | Independent association of liver fat accumulation with insulin resistance. | Obes Res Clin Pract. | 8(4) | E350-55 | 2014 |

書籍

| 著者氏名 | 論文タイトル名 | 書籍全体の編集者名 | 書籍名 | 出版社名 | 出版地 | 出版年 | ページ |
|------|-------------------|-------------|-------------------------------|--------|-----|------|--------|
| 斎藤重幸 | 血管疾患 | 日本高血圧学会編 | 高血圧専門医ガイドブック | 診断と治療社 | 東京 | 2014 | 144-47 |
| 斎藤重幸 | 糖尿病 | 日本循環器病予防学会編 | 第7版循環器病予防ハンドブック | 保健同人社 | 東京 | 2014 | 273-89 |
| 斎藤重幸 | 疫学からみた高齢者高血圧治療の意義 | 楽木宏美 | 高齢者高血圧の治療と管理 - JSH2014改訂をふまえて | 先端医学社 | 東京 | 2014 | 32-38 |