



Urbanization, Life Style Changes and the Incidence/In-Hospital Mortality of Acute Myocardial Infarction in Japan

– Report From the MIYAGI-AMI Registry Study –

Kiyotaka Hao, MD; Satoshi Yasuda, MD, PhD; Toru Takii, MD, PhD; Yoshitaka Ito, MD, PhD;
Jun Takahashi, MD, PhD; Kenta Ito, MD, PhD; Masaharu Nakayama, MD, PhD;
Nobuyuki Shiba, MD, PhD; Yoshihiro Fukumoto, MD, PhD; Hiroaki Shimokawa, MD, PhD
on behalf of the MIYAGI-AMI Study Investigators

Background: It remains to be examined whether urbanization and lifestyle changes are associated with the incidence and mortality from acute myocardial infarction (AMI) in Japan.

Methods and Results: A total of 19,921 AMI patients (male/female 14,290/5,631) registered by the MIYAGI-AMI Registry Study from 1988 to 2009 were divided into 2 groups according to their residences; inside (urban area, n=7,316) and outside (rural area, n=11,402) of Sendai City. From 1988 to 2009, the incidence of AMI (/100,000 persons/year) increased more rapidly in the rural area (24.2 to 51.4) than in the urban area (31.3 to 40.8) ($P<0.001$), with rapid aging in both areas. Moreover, from 1998 to 2009, the age-adjusted incidence of AMI in young (<44 years) and middle-aged (45–64 years) male patients (both $P<0.05$) in the rural area increased significantly, along with a markedly increased prevalence of dyslipidemia ($P<0.001$). Although in-hospital mortality from AMI decreased in both areas over the last 20 years (both $P<0.001$), it remained relatively higher in female than in male patients and was associated with higher age of the onset, longer elapsing time for admission and lower prevalence of primary coronary intervention in female patients in both areas.

Conclusions: These results demonstrate that urbanization and lifestyle changes have been associated with the incidence and mortality from AMI, although sex differences still remain to be improved. (*Circ J* 2012; **76**: 1136–1144)

Key Words: Acute myocardial infarction; Aging; Life-style; Risk factors; Sex

The incidence and mortality from coronary artery disease (CAD) has been declining in the United States and European countries.^{1–4} These declines have been attributed to the control of risk factors (eg, hypertension, dyslipidemia and smoking) and the improvement in critical care (eg, coronary revascularization therapy).^{5–7} In contrast to the Western countries, in Japan, a highly developed and racially homogeneous country that is rapidly aging, total cholesterol levels and the prevalence of obesity have been increasing as a result of lifestyle Westernization influence since the 1960s.^{8,9} However, the mortality from CAD has been declining and has remained much lower compared with other Western countries from 1960 to 2000.^{9–11} Importantly, there are some differences in lifestyle between people living in rural and urban areas in Japan. Indeed, it was reported that people in urban areas had

greater intakes of fat and cholesterol than those in rural areas in Japan.⁸ However, only a few studies have previously addressed the difference in the incidence and mortality from CAD between the rural and urban areas in Japan.^{8,12}

In order to explore the annual trend for acute myocardial infarction (AMI) in Japan, we have been conducting the MIYAGI-AMI Registry Study for more than 30 years since 1979, where almost all AMI patients in the Miyagi prefecture have been prospectively registered.^{10,13,14} The Miyagi prefecture, which is located in northeastern Japan, includes Sendai City, one of the 19 government-designated cities, and has a typical balance of urban and rural areas in Japan. Sendai City merged with neighboring municipalities in 1987–1988 and the population of Sendai City increased to 1,008,130 in 2000, which accounted for approximately 40% of the population of

Received October 26, 2011; revised manuscript received December 12, 2011; accepted January 5, 2012; released online February 18, 2012
Time for primary review: 14 days


Department of Cardiovascular Medicine, Tohoku University Graduate School of Medicine, Sendai, Japan

The Guest Editor for this article was Masafumi Kitakaze, MD.

Mailing address: Hiroaki Shimokawa, MD, PhD, Department of Cardiovascular Medicine, Tohoku University Graduate School of Medicine, Seiryō-machi, Aoba-ku, Sendai 980-8574, Japan. E-mail: shimo@cardio.med.tohoku.ac.jp

ISSN-1346-9843 doi:10.1253/circj.CJ-11-1233

All rights are reserved to the Japanese Circulation Society. For permissions, please e-mail: cj@j-circ.or.jp



the Miyagi prefecture, which was 2,365,320 in 2000. The population density of Sendai City (1,279/km² in 2000) has been much higher than that of any other parts of the Miyagi prefecture (209/km² in 2000).¹⁵

In the present study, we examined whether urbanization and lifestyle changes were associated with the incidence and mortality from AMI, with special reference to the difference between the urban and rural areas in our MIYAGI-AMI Registry Study.

Methods

The MIYAGI-AMI Registry Study

The MIYAGI-AMI Registry Study is a prospective, multi-center and observational study. As previously reported,^{10,13,14} this registry was established in 1979 and all 43 hospitals with a coronary care unit and/or cardiac catheterization facility in the Miyagi prefecture have been participating (Appendix 1). In the Miyagi prefecture, almost all AMI patients are transferred to one of those participating hospitals via the emergency medical service. This study was approved by the Institutional Review Board of Tohoku University Graduate School of Medicine under the condition that personal data are protected at all times.

In the MIYAGI-AMI Registry Study, the diagnosis of AMI and decision to use reperfusion therapy were made by individual cardiologists in charge. Diagnosis of AMI was made based on the WHO-MONICA criteria.¹⁶ Briefly, it was based on the finding of typical severe chest pain accompanied by abnormal ECG changes and increased serum levels of cardiac enzymes (ie, creatine phosphokinase, aspartate amino transferase and lactate dehydrogenase). Coronary thrombolysis was performed with intravenous administration of urokinase (480–960×10³ IU for 30 min) or alteplase (290–435×10³ IU/kg for 60 min) or with intracoronary administration of urokinase (maximum 960×10³ IU) or alteplase (maximum 6.4×10⁶ IU). Rescue percutaneous coronary intervention (PCI) was performed when thrombolysis was unsuccessful. Primary PCI has been widely performed in the Miyagi prefecture since 1992, as reported previously.^{10,13,14}

The registration form of the MIYAGI-AMI Registry includes the date and time of symptom onset, age, sex, pre-hospital management (eg, use of ambulance, time interval from the onset of symptoms to admission), infarction site, coronary risk factors (hypertension, diabetes mellitus, dyslipidemia and smoking), reperfusion therapies (eg, thrombolysis and/or PCI), and in-hospital outcome (eg, in-hospital mortality). In our MIYAGI-AMI Registry Study, we have revised the registra-

Table. Clinical Characteristics and Outcome of the Study Population								
	Rural area			P value for trend	Urban area			P value for trend
	1998–2001 (n=2,145)	2002–2005 (n=2,699)	2006–2009 (n=2,807)		1998–2001 (n=1,529)	2002–2005 (n=1,508)	2006–2009 (n=1,682)	
Male								
Age (years)	66.2±12.4*	67.0±12.9*	66.7±12.7	0.373	65.0±12.7	65.2±12.9	65.9±12.9	0.046
Age-adjusted incidence of AMI (/10 ⁵ persons/year)								
All	42.3±3.8*	47.2±3.2	47.3±2.5	0.274	55.1±4.7	49.3±10.9	47.9±4.1	0.163
<45 years old	4.9±0.9	5.8±0.7	6.9±1.2	0.018	5.1±0.7	5.7±0.5	6.0±2.7	0.460
45–64 years old	66.6±6.3*	83.2±5.5	88.9±14.9	0.016	91.2±4.9	85.9±21.0	83.7±8.2	0.402
65–74 years old	170.2±32.9	186.3±39.2	179.3±17.8	0.679	228.2±18.1	208.1±56.3	180.1±15.6	0.065
≥75 years old	253.5±47.0*	261.1±62.9	250.8±33.4	0.937	355.0±48.0	277.8±73.4	308.0±19.7	0.207
Hypertension (%)	46.1	59.5*	60.9	<0.001	48.2	54.3	63.0	<0.001
Diabetes mellitus (%)	27.5	32.9	29.5*	0.265	30.6	31.6	34.1	0.070
Dyslipidemia (%)	22.4*	34.1*	41.4	<0.001	32.2	39.0	42.0	<0.001
Smoking (%)	40.6	42.1	40.6	0.956	44.0	41.8	38.6	0.008
In-hospital mortality (%)	7.6	6.8	7.8	0.832	8.8	5.7	8.7	0.997
Female								
Age (years)	74.1±9.7	76.1±11.1	75.3±11.4	0.017	74.4±10.4	74.6±12.0	75.3±11.4	0.224
Age-adjusted incidence of AMI (/10 ⁵ persons/year)								
All	11.5±2.4*	13.6±1.1	13.2±1.0	0.202	15.1±1.2	11.9±2.0	12.4±2.4	0.077
<45 years old	0.2±0.4	0.4±0.2	0.7±0.5	0.114	0.2±0.2	0.5±0.3	0.5±0.7	0.297
45–64 years old	10.5±4.2	13.7±3.1	18.1±4.1	0.102	10.1±1.6	11.0±2.2	16.1±7.1	0.102
65–74 years old	54.5±1.8*	65.0±8.4	56.4±4.4	0.602	84.5±5.8	55.3±6.5	48.9±9.1	<0.001
≥75 years old	100.8±17.4*	135.7±14.9	120.8±7.9	0.076	165.9±13.9	131.4±19.4	129.8±17.2	0.016
Hypertension (%)	55.8	69.3	67.5	<0.001	60.2	63.5	65.0	0.137
Diabetes mellitus (%)	29.3	36.1	35.1	0.032	32.5	33.2	34.5	0.510
Dyslipidemia (%)	25.8	30.9	38.6	<0.001	31.0	37.1	37.7	0.039
Smoking (%)	8.9	6.6*	10.6	0.163	12.1	13.4	14.1	0.383
In-hospital mortality (%)	12.3	11.1	14.5	0.254	14.4	15.3	14.1	0.892

Values are mean ± SD or n (%). *P<0.05 for the difference between rural and urban areas. AMI, acute myocardial infarction. Study population was divided into 2 groups according to the residence: inside (urban area) and outside Sendai City (rural area).

tion form gradually over the last 30 years. Thus, although the incidence of AMI and related data (time of onset, age and sex) are available for the last 30 years, the data on the pre-hospital management, infarction site, coronary risk factors, reperfusion therapies, duration of hospitalization and in-hospital outcome are only available for the last 10–20 years, which were analyzed in the present study.



Data Analysis

In the present study, we have registered a total of 19,921 patients with AMI (male/female 14,290/5,631) over the last 20 years after the municipal merger in 1988. In particular, we have focused on the patients registered between 1998 and 2009 (total, 12,491; male/female, 8,969/3,522), who were divided into 2 groups according to their residences; inside (urban area, n=4,719) and outside Sendai City (rural area, n=7,651), after excluding the patients whose residences were unknown (n=159). We also divided the total observational period of 12 years into the 3 periods: 1998–2001, 2002–2005 and 2006–2009. To calculate the sex- and age-adjusted incidence of AMI (/100,000 person-years), we applied the direct standardization method using the age distribution of the Japanese population from the 2000 census,⁵ as the standard population. In addition, in order to clarify the age-specific trend, we categorized the age at AMI onset into the 4 groups: ≤44 (young), 45–64 (middle-aged), 65–74 (old) and ≥75 years old (high-old).¹⁵

Results are expressed as mean ± SD. Linear trends were examined for continuous variables by using analysis of variance (ANOVA) with repeated measures or the Jonckheere-Terpstra trend test as appropriate, and for categorical variables by using the chi-square test for trend. Differences in mean values were examined with a t-test, Mann-Whitney test or chi-square test as appropriate. Multiple logistic regression analysis was used to examine determinants of risk factor prevalence in AMI patients. Variables used for analysis included: sex, age at onset of AMI (per 10 years), study periods (1998–2001, 2002–2005 and 2006–2009), residence (rural vs. urban), and other risk factors. The odds ratios (ORs) and 95% confidence intervals (95%CI) were calculated. A P-value less than 0.05 were considered to be statistically significant. All statistical analyses were performed using the statistical software SPSS version 18 for Windows.

Results

Over the last 20 years, the incidence of AMI (/100,000 persons/year) significantly increased in both the rural and the urban areas in the Miyagi prefecture (2.1- and 1.3-fold, respectively, both P<0.001) (Figure 1A). Furthermore, the extent of the increase in AMI incidence was greater in the rural area than in the urban area, finally exceeding that in the urban area after 2000. These changes were accompanied with rapid aging






Figure 4. Time interval from the onset of symptoms to hospitalization. The percentage of patients with less than 2 h of elapsing time for hospitalization has significantly increased in rural male patients. The percentage was significantly lower in female patients than in male patients in both areas in 2006–2009. *P<0.05 for the difference between rural and urban areas. †P<0.05 for the difference between the sexes in the same rural or urban areas. ¶P<0.05 for a linear trend.





Figure 5. Prevalence of reperfusion therapy for acute myocardial infarction (AMI). The prevalence of primary percutaneous coronary intervention (PCI) steadily increased in the rural area in both sexes. Importantly, the prevalence of PCI was approximately 10% lower in female patients than in male patients in both rural and urban areas. *P<0.05 for the difference in male patients between rural and urban areas. †P<0.05 for linear trend.




in both areas in the Miyagi prefecture (**Figure 1B**). Following age adjustment (**Figure 1C**), the incidence of AMI in the rural area increased significantly ($P < 0.001$), whereas that in the urban area decreased significantly ($P < 0.001$) in the recent 10-year period (between 1998 and 2009). In contrast, in-hospital mortality significantly decreased in both areas (both $P < 0.001$), but to a greater extent in the rural area (0.5-fold in the rural area and 0.9-fold in the urban area) (**Figure 1D**). In 1998–2001, there was no significant difference in in-hospital mortality between the rural and urban male patients ($P = 0.263$), and in-hospital mortality remained low (~8%) from 1998–2001 to 2006–2009 in both the rural and urban male patients (rural: $P = 0.832$; urban: $P = 0.997$) (**Table**). Importantly, in-hospital mortality of the female patients in both the rural and the urban areas remained doubled compared with the male patients during the study period (**Table**).

The clinical characteristics of the AMI patients in the present study are shown in **Table**. The female patients were approximately 10 years older than the male patients and approximately a half of them were ≥ 75 years-old in 1998–2001 in both areas, with a significant further increase in the rural area (male, $P < 0.001$; female, $P < 0.001$) and such a trend in the urban area (male, $P = 0.054$; female, $P = 0.176$) (**Figure 2**). In 1998–2001, the age-adjusted incidence of AMI was significantly lower in the rural area than in the urban area for both sexes (male, $P = 0.019$; female, $P = 0.035$) (**Table**). However, the difference between the 2 areas became insignificant in 2006–2009 for both sexes (male, $P = 0.824$; female, $P = 0.530$). When investigating the age-specific trend, the significant in-

crease in the age-adjusted incidence of AMI was noted in the young (<44 years-old) and middle age (45–64 years-old) male patients only in the rural area (young, $P = 0.018$; middle age, $P = 0.016$), and the significant decrease was noted in the old (65–74 years-old) and high-old (>75 years-old) female patients in the urban area (old, $P < 0.001$; high-old, $P = 0.016$) (**Table, Figure 3**).

Regarding the time from the onset of AMI to admission, the percentage of the patients with less than 2 h of elapsing time at admission was significantly lower in the rural area than in the urban area for the male patients in 1998–2001 ($P < 0.001$) (**Figure 4**). However, the difference became insignificant in 2006–2009 ($P = 0.051$), accompanied with the significant increase in the percentage in the rural area (rural, $P < 0.001$; urban, $P = 0.082$). Importantly, in the rural female patients, the percentage of patients with less than 2 h of elapsing time at admission remained at a low level (~20%), and the difference between the sexes in the rural area became greater from 1998–2001 ($P = 0.086$) to 2006–2009 ($P < 0.001$). In contrast, the difference between the sexes in the urban area was significant in 2006–2009 ($P = 0.04$). Moreover, the prevalence of primary PCI in the female patients was lower by ~10% compared with the male patients in both areas (**Figure 5**). In the male patients, the prevalence of primary PCI significantly increased only in the rural area from 1998–2001 to 2006–2009 (rural, $P < 0.001$; urban, $P = 0.054$), and a similar trend was also noted in the female patients (rural, $P < 0.001$; urban, $P = 0.176$).

Multivariate analysis of the coronary risk factors in AMI patients showed that the prevalence of hypertension and dys-



lipidemia significantly increased and that of diabetes tended to increase (Figure 6). Hypertension was associated with older age but not with residence, whereas dyslipidemia was associated with younger age and urban residence. Although the prevalence of dyslipidemia in the male patients was significantly lower in the rural area than in the urban area in 1998–2001, it significantly increased in the rural area and the difference between the 2 areas became insignificant in 2006–2009 (Table). Moreover, the progressive increase in the prevalence of dyslipidemia was noted in both areas for both sexes with a more sharp increase in the rural area (Figure 7). Smoking was associated with male sex and younger age, but not with residence (Figure 6), and the prevalence of smoking largely remained unchanged in both areas for both sexes (Figure 7).

Discussion

The novel findings of the present study were that the incidence of AMI increased more rapidly in the rural area than in the urban area, with rapid aging in both areas. Moreover, the incidence of AMI in the rural male patients ≤64 years-old was increased along with the marked increase in the prevalence of dyslipidemia in Japan. Although in-hospital mortality from AMI markedly decreased in both areas over the last 20 years, it remained relatively high in female patients than in male patients in both areas. To the best of our knowledge, this is the first study that demonstrates the association between urbaniza-

tion, life-style changes and the incidence and mortality of AMI in the largest number of patients in Japan.

Comparison of the Incidence of AMI Between Rural and Urban Areas

Although in the United States and European countries, the incidence of CAD has been declining in the last decades,^{1,2,4} the present study demonstrates that the incidence of AMI has been rapidly increasing in both the rural and urban areas over the last 20 years, with a more noted increase in the former than in the latter. However, this tendency has disappeared following age adjustment in recent years only in the urban area, which implied that the increased tendency in the incidence of AMI in the rural area might be not be associated with rapid aging alone in recent years.

There were few studies that addressed the difference in the incidence of CAD between rural and urban areas in Japan. The Akita-Osaka study is the community-based survey, where the residents of the Yao City, Osaka prefecture (an urban community with a total census population of 23,552 in 2000) and those of Ikawa Town, Akita prefecture (a rural community with a total census population of 6,116 in 2000) were compared during the period of 1964–2003.¹² In this study, significant increases in the age-adjusted incidence of AMI and sudden cardiac death were noted in Yao City (in male patients from 1980 to 2003) but not in Ikawa City in both sexes.¹² The present study confirmed the results of the Akita-Osaka study

in the rural and urban areas of the same Miyagi prefecture. The Yamagata AMI Registry study provided more recent data and an age-specific trend in the period of 1993–2007.¹⁷ The population density of the Yamagata prefecture was 133/km² in 2000, which was comparable with that of the rural area in the present study.¹⁵ In this study, the age-adjusted incidence of AMI in the male but not that in the female patients significantly increased. In particular, the male population who were younger than 65 years old showed a marked increase in AMI, a consistent finding with the present results for the rural area. These results indicate that the incidence of AMI has been increasing in the younger male population in the rural areas of Japan. Taken together, unlike the trend in Western countries, it appears that the incidence of AMI has been increasing in Japan to a greater extent in the rural area than in the urban area over the last 20 years and has been associated with rapid aging.

Decreasing In-Hospital Mortality and Improvement in Critical Care

In the present study, the in-hospital mortality from AMI significantly decreased in both the urban and the rural areas over the last 20 years. The present study also demonstrates that primary PCI was performed more frequently in the rural area than in the urban area, along with the shortening in the elapsing time from the onset to hospitalization. The recent progress in critical care might have beneficial effects, overcoming the rapid aging in AMI patients.

In the most recent 10 year period, the in-hospital mortality remained at a low level in male patients, whereas in female patients, the mortality remained doubled compared with the male patients in both the rural and the urban areas. It was previously reported that the poorer outcome of the female AMI patients could be caused by multiple factors, including higher age, higher risk profiles, longer elapsing time from the onset to hospitalization, higher incidence of Killip class ≥ 2 , and less frequent use of primary PCI.^{18–20} Indeed, in the present study, the female patients were approximately 10 years older than the male patients and half of them were older than 75 years and needed a longer time from the onset of AMI to hospitalization in the both areas in 2006–2009. These points might have limited the use of primary PCI with a resultant poor outcome for the female AMI patients in the present study.

Changes in the Prevalence of Coronary Risk Factors in AMI Patients

The WHO-MONICA studies, as well as several Japanese cohort studies, demonstrated that the incidence of cardiovascular diseases increased and were associated with the clustering of risk factors.^{21–23} In the present study, the prevalence of hypertension and dyslipidemia in AMI patients significantly increased in both the rural and urban areas. Importantly, there was a significant difference in the prevalence of dyslipidemia between the rural and urban areas with a marked increase noted in the rural area, especially in those male patients aged ≤ 64 years. Indeed, previous studies demonstrated that dyslipidemia is an independent risk factor in male but not in female patients,^{17,24} and in the Yamagata-AMI Registry study, the increased prevalence of dyslipidemia in the younger male patients with AMI was also associated with an increased incidence of AMI.¹⁷ In the Miyagi prefecture, the intake of animal fat was significantly higher in the rural than in the urban area in 2000 (rural 20.7 g/day vs. urban 23.4 g/day, $P < 0.05$).²⁵ Moreover, in Japan, fat intake and serum levels of total cholesterol were higher in the urban than in the rural areas in

1966; however, the difference in cholesterol levels between the 2 areas became smaller in 1966–1985 along with the influence of Westernization of food habits in the rural area.⁸ Taken together, it might indicate that the increase in the incidence of AMI in younger male patients in the rural area was likely to be associated with the marked increase in the prevalence of dyslipidemia.

The present study also demonstrates the increase in the prevalence of hypertension in AMI patients. In the Tohoku district, including the Miyagi prefecture, the prevalence of hypertension was relatively higher compared with other parts of Japan,^{12, 26} and thus more careful and strict control of risk factors is needed.

The prevalence of smoking remained high not only in the urban areas but also in the rural areas. In particular, in the younger male patients, the prevalence of smoking ($\sim 50\%$) was higher compared with the general Japanese population (36.8% in males and 9.1% in females in 2008).²⁷ Importantly, in the younger urban female patients, it remained more than 30%; 3 times higher than in the general Japanese population.

Study Limitations

Several limitations should be mentioned for the present study. First, although in the Miyagi prefecture, almost all AMI patients are transferred to our participating hospitals via the established emergency medical system, we cannot completely confirm that all patients have been registered in our registry. Second, while the MIYAGI-AMI Registry Study has been conducted over 20 years, the diagnosis of AMI has been changing.²⁸ In the present study, the diagnosis was made on the basis of the WHO-MONICA criteria with creatine kinase (CK).¹⁶ Indeed, troponins are widely used in recent clinical practice and are more sensitive and specific biomarkers of myocyte necrosis than CK,²⁹ which might affect the results. Third, this study is an observational study and cannot reach the cause-effect relationship. Moreover, we did not examine the prevalence of risk factors in control subjects and did not collect the data of medical treatment for prevention, thus we were unable to precisely estimate the influence of risk factors on the incidence of AMI. Finally, in the present study, we did not examine the long-term mortality but only examined in-hospital mortality. The increasing incidence of decreasing in-hospital mortality from AMI in the Japanese population has apparently resulted in the recent increase in the number of patients with ischemic heart failure, as recently demonstrated in our heart failure cohort study, the CHART-1 and the CHART-2 studies.^{30,31} Thus, a more effective strategy to improve the management of post-infarction heart failure needs to be developed.

Conclusions

Our MIYAGI-AMI Registry Study demonstrates that urbanization and life-style changes have been associated with the incidence and mortality of AMI in Japan, although sex differences still remain to be improved.

Acknowledgments

This study was supported, in part, by the grants-in-aid from the Sendai City, the Miyagi Prefecture and the Miyagi Medical Association and the grants-in-aid [H22-Shinkin-004] from the Japanese Ministry of Education, Culture, Sports, Science, and Technology, Tokyo, Japan. We thank all the collaborators in the MIYAGI-AMI Registry Study ([Appendix 1](#)). We also thank Ayako Tsunoda for excellent secretarial assistance.

References

1. Yeh RW, Sidney S, Chandra M, Sorel M, Selby JV, Go AS. Popula-

- tion trends in the incidence and outcomes of acute myocardial infarction. *N Engl J Med* 2010; **362**: 2155–2165.
2. Parikh NI, Gona P, Larson MG, Fox CS, Benjamin EJ, Murabito JM, et al. Long-term trends in myocardial infarction incidence and case fatality in the National Heart, Lung, and Blood Institute's Framingham Heart study. *Circulation* 2009; **119**: 1203–1210.
 3. Unal B, Critchley JA, Capewell S. Explaining the decline in coronary heart disease mortality in England and Wales between 1981 and 2000. *Circulation* 2004; **109**: 1101–1107.
 4. Linnertsjö A, Hammar N, Gustavsson A, Reuterwall C. Recent time trends in acute myocardial infarction in Stockholm, Sweden. *Int J Cardiol* 2000; **76**: 17–21.
 5. Gregg EW, Cheng YJ, Cadwell BL, Imperatore G, Williams DE, Flegal KM, et al. Secular trends in cardiovascular disease risk factors according to body mass index in US adults. *JAMA* 2005; **293**: 1868–1874.
 6. Tamis-Holland JE, Palazzo A, Stebbins AL, Slater JN, Boland J, Ellis SG, et al. Benefits of direct angioplasty for women and men with acute myocardial infarction: Results of the Global Use of Strategies to Open Occluded Arteries in Acute Coronary Syndromes Angioplasty (GUSTO II-B) Angioplasty Substudy. *Am Heart J* 2004; **147**: 133–139.
 7. Hochman JS, Tamis JE, Thompson TD, Weaver WD, White HD, Van de Werf F, et al. Sex, clinical presentation, and outcome in patients with acute coronary syndromes: Global Use of Strategies to Open Occluded Coronary Arteries in Acute Coronary Syndromes IIB Investigators. *N Engl J Med* 1999; **341**: 226–232.
 8. Okayama A, Ueshima H, Marmot M, Elliott P, Choudhury SR, Kita Y. Generational and regional differences in trends of mortality from ischemic heart disease in Japan from 1969 to 1992. *Am J Epidemiol* 2001; **153**: 1191–1198.
 9. Kubo M, Kiyohara Y, Kato I, Tanizaki Y, Arima H, Tanaka K, et al. Trends in the incidence, mortality, and survival rate of cardiovascular disease in a Japanese community: The Hisayama study. *Stroke* 2003; **34**: 2349–2354.
 10. Takii T, Yasuda S, Takahashi J, Ito K, Shiba N, Shirato K, et al. Trends in acute myocardial infarction incidence and mortality over 30 years in Japan: Report from the MIYAGI-AMI Registry Study. *Circ J* 2010; **74**: 93–100.
 11. Ueshima H. Explanation for the Japanese paradox: Prevention of increase in coronary heart disease and reduction in stroke. *J Atheroscler Thromb* 2007; **14**: 278–286.
 12. Kitamura A, Sato S, Kiyama M, Imano H, Iso H, Okada T, et al. Trends in the incidence of coronary heart disease and stroke and their risk factors in Japan, 1964 to 2003: The Akita-Osaka study. *J Am Coll Cardiol* 2008; **52**: 71–79.
 13. Sakurai K, Watanabe J, Iwabuchi K, Koseki Y, Kon-no Y, Fukuchi M, et al. Comparison of the efficacy of reperfusion therapies for early mortality from acute myocardial infarction in Japan: Registry of Miyagi Study Group for AMI (MsAMI). *Circ J* 2003; **67**: 209–214.
 14. Watanabe J, Iwabuchi K, Koseki Y, Fukuchi M, Shinozaki T, Miura M, et al. Declining trend in the in-hospital case-fatality rate from acute myocardial infarction in Miyagi Prefecture from 1980 to 1999. *Jpn Circ J* 2001; **65**: 941–946.
 15. Ministry of Internal Affairs and Communications of Japan. 2000 population census of Japan. Tokyo, Japan: Statistics Bureau, Ministry of Internal Affairs and Communications, Government of Japan, 2001 (in Japanese).
 16. Tunstall-Pedoe H, Kuulasmaa K, Amouyel P, Arveiler D, Rajakangas AM, Pajak A. Myocardial infarction and coronary deaths in the World Health Organization MONICA Project: Registration procedures, event rates, and case-fatality rates in 38 populations from 21 countries in four continents. *Circulation* 1994; **90**: 583–612.
 17. Nishiyama S, Watanabe T, Arimoto T, Takahashi H, Shishido T, Miyashita T, et al. Trends in coronary risk factors among patients with acute myocardial infarction over the last decade: The Yamagata AMI registry. *J Atheroscler Thromb* 2010; **17**: 989–998.
 18. Stone GW, Grines CL, Browne KF, Marco J, Rothbaum D, O'Keefe J, et al. Comparison of in-hospital outcome in men versus women treated by either thrombolytic therapy or primary coronary angioplasty for acute myocardial infarction. *Am J Cardiol* 1995; **75**: 987–992.
 19. Milcent C, Dormont B, Durand-Zaleski I, Steg PG. Gender differences in hospital mortality and use of percutaneous coronary intervention in acute myocardial infarction: Microsimulation analysis of the 1999 nationwide french hospitals database. *Circulation* 2007; **115**: 833–839.
 20. Kosuge M, Kimura K, Kojima S, Sakamoto T, Ishihara M, Asada Y, et al. Sex differences in early mortality of patients undergoing primary stenting for acute myocardial infarction. *Circ J* 2006; **70**: 217–221.
 21. Kuulasmaa K, Tunstall-Pedoe H, Dobson A, Fortmann S, Sans S, Tolonen H, et al. Estimation of contribution of changes in classic risk factors to trends in coronary-event rates across the WHO MONICA Project populations. *Lancet* 2000; **355**: 675–687.
 22. Nakamura T, Tsubono Y, Kameda-Takemura K, Funahashi T, Yamashita S, Hisamichi S, et al. Magnitude of sustained multiple risk factors for ischemic heart disease in Japanese employees: A case-control study. *Jpn Circ J* 2001; **65**: 11–17.
 23. Nakamura Y, Yamamoto T, Okamura T, Kadowaki T, Hayakawa T, Kita Y, et al. Combined cardiovascular risk factors and outcome: NIPPON DATA80, 1980–1994. *Circ J* 2006; **70**: 960–964.
 24. Kawano H, Soejima H, Kojima S, Kitagawa A, Ogawa H, Investigators JACSSJ. Sex differences of risk factors for acute myocardial infarction in Japanese patients. *Circ J* 2006; **70**: 513–517.
 25. Ministry of Health and Welfare of Miyagi prefecture. Report from health and nutrition investigation in Miyagi prefecture, 2000. Miyagi, Japan: Miyagi Prefectural Government, 2002 (in Japanese).
 26. Tomonari T, Fukuda M, Miura T, Mizuno M, Wakamatsu TY, Ichikawa T, et al. Is salt intake an independent risk factor of stroke mortality? Demographic analysis by regions in Japan. *J Am Soc Hypertens* 2011; **5**: 456–462.
 27. Japan Health Promotion and Fitness Foundation. <http://www.healthnet.or.jp/tobacco/product/pd100000.html> (accessed on 26 October 2011).
 28. Yasuda S, Shimokawa H. Acute myocardial infarction: The enduring challenge for cardiac protection and survival. *Circ J* 2009; **73**: 2000–2008.
 29. Jaffe AS, Babuin L, Apple FS. Biomarkers in acute cardiac disease: The present and the future. *J Am Coll Cardiol* 2006; **48**: 1–11.
 30. Shiba N, Shimokawa H. Chronic heart failure in Japan: Implications of the CHART studies. *Vasc Health Risk Manag* 2008; **4**: 103–113.
 31. Shiba N, Nochioka K, Miura M, Kohno H, Shimokawa H, CHART-2 Investigators. Trend of westernization of etiology and clinical characteristics of heart failure patients in Japan: First report from the CHART-2 study. *Circ J* 2011; **75**: 823–833.

Appendix 1. List of Participating Hospitals and Investigators

Fukaya Hospital, Akiho H, MD.
 Hikarigaoka Spellman Hospital, Shimura S, MD.
 Ishinomaki Medical Association; Ishinomaki Municipal Hospital, Akai K, MD.
 Ishinomaki Red-cross Hospital, Sukegawa H, MD.
 JR Sendai Hospital, Honda H, MD.
 Katta General Hospital, Kanno H, MD.
 Kesen-numa Hospital, Ogata K, MD.
 Kurihara Central Hospital, Komatsu S, MD.
 Labour Welfare Corporation Tohoku Rosai Hospital, Komaru T, MD.
 Marumori National Health Insurance Hospital, Otomo M, MD.
 Miyagi Eastern Cardiovascular Institute, Kikuchi Y, MD.
 Miyagi Cancer Center, Owada N, MD.
 Miyagi Cardiovascular and Respiratory Center, Osawa N, MD.
 Mori Hospital, Mori A, MD.
 Nagamachi Hospital, Mitobe H, MD.
 Nishitaga National Hospital, Kitaoka S, MD.
 NTT EAST Tohoku Hospital, Yamada A, MD.
 Oizumi Memorial Hospital, Koiba Y, MD.
 Osaki Citizen Hospital, Hiramoto T, MD.
 Saito Hospital, Otsuka K, MD.
 Saka General Hospital, Watanabe K, MD.
 Sendai Cardiovascular Center, Fujii S, MD.
 Sendai City Hospital, Yagi T, MD.
 Sendai Kosei Hospital, Meguro T, MD.
 Sendai Medical Center, Shinozaki T, MD.
 Sendai Open Hospital Sendai City Medical Center, Kato A, MD.
 Sendai Public Health Insurance Hospital, Oikawa Y, MD.
 Sendai Red-cross Hospital, Wakayama M, MD.
 Sendai Tokushukai Hospital, Fukuchi M, MD.
 Sen-en General Hospital, Hashiguchi R, MD.
 Shichigashuku National Health Insurance Clinic, Nagashima T, MD.
 Shiogama City Hospital, Goto J, MD.
 South Miyagi Medical Center, Inoue K, MD.
 Tohoku Kosai Hospital, Suzuki S, MD.
 Tohoku University Hospital,
 Department of Cardiovascular Medicine, Shimokawa H, MD.
 Department of Cardiovascular Surgery, Saiki Y, MD.
 Department of Gastroenterology, Shimosegawa T, MD.
 Tohoku Welfare and Pension Hospital, Katahira Y, MD.
 Tome Citizen Hospital, Izuma M, MD.
 Toyama Clinic on Tome City, Ishii M, MD.



Original article

Effect of cardiac rehabilitation on muscle mass, muscle strength, and exercise tolerance in diabetic patients after coronary artery bypass grafting

Miho Nishitani (MD)^a, Kazunori Shimada (MD, FJCC)^{a,b,*}, Masayuki Masaki (MD)^a, Satoshi Sunayama (MD)^a, Atsumi Kume (MD)^a, Kosuke Fukao (MD)^a, Eiryu Sai (MD)^a, Tomo Onishi (PhD)^b, Miki Shioya (MS)^b, Hiroyuki Sato (MD)^{b,c}, Taira Yamamoto (MD)^d, Atsushi Amano (MD, FJCC)^d, Hiroyuki Daida (MD, FJCC)^a

^a Department of Cardiovascular Medicine, Juntendo University School of Medicine, Tokyo, Japan

^b Juntendo Sports Clinic, Juntendo University Hospital, Tokyo, Japan

^c Department of General Medicine, Juntendo University School of Medicine, Tokyo, Japan

^d Department of Cardiovascular Surgery, Juntendo University School of Medicine, Tokyo, Japan

ARTICLE INFO

Article history:

Received 7 March 2012

Received in revised form 23 October 2012

Accepted 20 November 2012

Available online 16 January 2013

Keywords:

Cardiac rehabilitation

Coronary artery bypass grafting

Diabetes mellitus

Exercise tolerance

Muscle strength

Muscle mass

ABSTRACT

Background: The effects of cardiac rehabilitation (CR) on muscle mass, muscle strength, and exercise tolerance in patients with diabetes mellitus (DM) who received CR after coronary artery bypass grafting (CABG) have not been fully elucidated.

Methods: We enrolled 78 consecutive patients who completed a supervised CR for 6 months after CABG (DM group, $n = 37$; non-DM group, $n = 41$). We measured mid-upper arm muscle area (MAMA), handgrip power (HGP), muscle strength of the knee extensor (Ext) and flexor (Flex), and exercise tolerance at the beginning and end of CR.

Results: No significant differences in confounding factors, including age, gender, ejection fraction, or number of CR sessions, were observed between the two groups. At the beginning of CR, the levels of Ext muscle strength and peak VO_2 were significantly lower in the DM group than in the non-DM group. At the end of CR, significant improvement in the levels of muscle strength, HGP, and exercise tolerance was observed in both groups. However, the levels of Ext muscle strength, HGP, peak VO_2 , thigh circumference, and MAMA were significantly lower in the DM group than in the non-DM group. In addition, no significant improvement in thigh circumference and MAMA was observed in the DM group. At the end of CR, the levels of thigh circumference and MAMA correlated with Ext and Flex muscle strength as well as with HGP. Percent changes in the levels of Ext muscle strength were significantly correlated with those of MAMA and hemoglobin A1c.

Conclusions: These data suggest that improvement in muscle strength may be influenced by changes in muscle mass and high glucose levels in DM patients undergoing CR after CABG. A CR program, including muscle mass intervention and blood glucose control, may improve deterioration in exercise tolerance in DM patients after CABG.

© 2012 Japanese College of Cardiology. Published by Elsevier Ltd. All rights reserved.

Introduction

Patients with diabetes mellitus (DM) are at increased risk of coronary artery disease (CAD) [1]. Patients with DM are at 2–4 times higher risk of developing CAD and mortality due to CAD compared with non-DM patients [2]. Patients with CAD are treated by lifestyle modification, medical therapy, and coronary revascularization

such as percutaneous coronary intervention and coronary artery bypass grafting (CABG). However, the benefits of revascularization are less and the risks and complications are greater than those in non-DM patients. Previous studies have also reported a high incidence of bypass graft dysfunction and a high mortality even in DM patients who underwent CABG [3].

Cardiac rehabilitation (CR) has numerous benefits such as modification of risk factors and prevention of future cardiovascular events [4]. Improvement in peak VO_2 after CR reduced cardiovascular morbidity and mortality in patients with CAD [5]. However, a previous study demonstrated that the presence of DM was a negative factor for improvement in peak VO_2 [6]. Another report showed a significant inverse relationship between fasting blood glucose

* Corresponding author at: Department of Cardiovascular Medicine, Juntendo University School of Medicine, 2-1-1 Hongo, Bunkyo-ku, Tokyo 113-8421, Japan. Tel.: +81 3 5802 1056; fax: +81 3 5689 0627.

E-mail address: shimakaz@juntendo.ac.jp (K. Shimada).