

- 12) Muraki I et al : Nocturnal intermittent hypoxia and C reactive protein among middle-aged community residents : a cross-sectional survey. *Thorax* 2010 ; **65** : 523-527.
- 13) Muraki I et al : Nocturnal intermittent hypoxia and the development of type 2 diabetes : the Circulatory Risk in Communities Study(CIRCS). *Diabetologia* 2010 ; **53** : 481-488.
- 14) Tanno S et al : Sleep-related intermittent hypoxia and glucose intolerance : a community based study. *Sleep Med* (In press).
- 15) Campos E et al : Relationship between nocturnal intermittent hypoxia, arterial stiffness and cardiovascular risk factors in a community-based population : the Toon Health Study. *J Atheroscler Thromb* (In press).
- 16) Enright PL et al : Prevalence and correlates of snoring and observed apneas in 5,201 older adults. *Sleep* 1996 ; **19** : 531-538.
- 17) Al-Delaimy WK et al : Snoring as a risk factor for type II diabetes mellitus : a prospective study. *Am J Epidemiol* 2002 ; **155** : 387-393.
- 18) Kim J et al : Snoring as an independent risk factor for hypertension in the nonobese population : the Korean Health and Genome Study. *Am J Hypertens* 2007 ; **20** : 819-824.
- 19) Hu FB et al : Prospective study of snoring and risk of hypertension in women. *Am J Epidemiol* 1999 ; **150** : 806-816.
- 20) Shin MH et al : Self-reported snoring and metabolic syndrome : the Korean Multi-Rural Communities Cohort Study. *Sleep Breath* 2014 ; **18** : 423-430.
- 21) Sun L et al : Snoring, inflammatory markers, adipokines and metabolic syndrome in apparently healthy Chinese. *PLoS One* 2011 ; **6** : e27515.
- 22) Li D et al : Self-reported habitual snoring and risk of cardiovascular disease and all-cause mortality. *Atherosclerosis* 2014 ; **235** : 189-195.
- 23) Nagayoshi M et al : Self-reported snoring frequency and incidence of cardiovascular disease : the Circulatory Risk in Communities Study (CIRCS). *J Epidemiol* 2012 ; **22** : 295-301.
- 24) Zhang X et al : Associations of self-reported sleep duration and snoring with colorectal cancer risk in men and women. *Sleep* 2013 ; **36** : 681-688.
- 25) Nieto FJ et al : Sleep-disordered breathing and cancer mortality : results from the Wisconsin Sleep Cohort Study. *Am J Respir Crit Care Med* 2012 ; **186** : 190-194.
- 26) Toffoli S et al : Intermittent hypoxia is a key regulator of cancer cell and endothelial cell interplay in tumours. *FEBS J* 2008 ; **275** : 2991-3002.
- 27) 循環器病の診断と治療に関するガイドライン (2008-2009年度合同研究班報告) : 循環器領域における睡眠呼吸障害の診断・治療に関するガイドライン. *Circulation J* 2010 ; Suppl II : 963-1051.
- 28) Bazzano LA et al : Effect of nocturnal nasal continuous positive airway pressure on blood pressure in obstructive sleep apnea. *Hypertension* 2007 ; **50** : 417-423.
- 29) Campos-Rodriguez F et al : Role of sleep apnea and continuous positive airway pressure therapy in the incidence of stroke or coronary heart disease in women. *Am J Respir Crit Care Med* 2014 ; **189** : 1544-1550.
- 30) Cassar A et al : Treatment of obstructive sleep apnea is associated with decreased cardiac death after percutaneous coronary intervention. *J Am Coll Cardiol* 2007 ; **50** : 1310-1314.

