

## ・研究成果の刊行に関する一覧表

### 書籍

著者氏名	論文タイトル名	書籍全体の編集者名	書籍名	出版社名	出版地	出版年	ページ
榎野博史	糖尿病とCKDの関係	榎野博史	糖尿病×CKD診療ガイド	南山堂	東京	2012	1-4
和田淳	尿検査の意義 - 蛋白尿 -	榎野博史	糖尿病×CKD診療ガイド	南山堂	東京	2012	10-11
江口潤	HbA1cを血糖コントロール指標としてみる場合の注意点	榎野博史	糖尿病×CKD診療ガイド	南山堂	東京	2012	19-22
中司敦子	血糖コントロールの指標としてみる場合の注意点	榎野博史	糖尿病×CKD診療ガイド	南山堂	東京	2012	29-31
松岡孝	糖尿病性足病変とフットケア	中村二郎	Visual 糖尿病臨床のすべて シリーズ 糖尿病性神経障害 - 基礎から臨床のすべて	中山書店	東京	2013	188-195
和田淳	慢性腎臓病はなぜ悪い	伊藤浩	冠動脈疾患のパフォーマンス	南江堂	東京	2013	45-51
松岡孝	糖代謝、脂質代謝を悪化させない 遮断薬の選択	伊藤浩	遮断薬を臨床で生かす！	南江堂	東京	2013	116-118
和田淳	慢性腎臓病患者に使用できる 遮断薬は？	伊藤浩	遮断薬を臨床で生かす！	南江堂	東京	2013	102-104
和田淳	エビデンスに基づくCKD診療ガイドライン2013	福井次矢、北原光夫	今日の治療指針2014	医学書院	東京	2014	1831-1834

### 雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Wada J and Makino H	Inflammation and the pathogenesis of diabetic nephropathy.	<i>Clin Sci</i>	124(3)	139-152	2013
Nakatsuka A, Wada J, Hida K, Hida A, Eguchi J, Teshigawara S, Murakami K, Kanzaki M, Inoue K, Terami T, Katayama A, Ogawa D, Kagechika H, Makino H.	RXR antagonism induces G0/G1 cell cycle arrest and ameliorates obesity by up-regulating the p53-p21 <sup>Cip1</sup> pathway in adipocytes.	<i>J Pathol</i>	226(5)	784-795	2012
Murakami K, Wada J, Ogawa D, Horiguchi CS, Miyoshi T, Sasaki M, Uchida HA, Nakamura Y, Makino H.	The effects of telmisartan treatment on the abdominal fat depot in patients with metabolic syndrome and essential hypertension: Abdominal fat Depot Intervention Program of Okayama (ADIPO).	<i>Diab Vasc Dis Res</i>	10(1)	93-96	2013
Teshigawara S, Wada J, Hida K, Nakatsuka A, Eguchi J, Murakami K, Kanzaki M, Inoe K, Terami T, Katayama A, Iseda I, Matsushita Y, Miyatake N, McDonald JF, Hotta K, Makino H.	Serum vaspin concentrations are closely related to insulin resistance and rs77060950 at SERPINA12 genetically defines distinct group with higher serum levels in Japanese population.	<i>J Clin Endocrinol Metab</i>	97(7)	E1202-1207	2012
Kawai D, Takaki A, Nakatsuka A, Wada J, Tamaki N, Yasunaka T, Koike K, Tsuzaki R, Matsumoto K, Miyake Y, Shiraha H, Morita M, Makino H, Yamamoto K.	Hydrogen-rich water prevents progression of non-alcoholic steatohepatitis and accompanying hepatocarcinogenesis in mice.	<i>Hepatology</i>	56(3)	912-921	2012
Kanzaki M, Wada J, Kikumoto Y, Akagi S, Nakao K, Sugiyama H, Makino H.	The therapeutic potential of synthetic human atrial natriuretic peptide in nephrotic syndrome: a randomized controlled trial.	<i>Int J Nephrol Renovasc Dis</i>	5	91-96	2012

Sugiyama K, Sada KE, Kurosawa M, Wada J, Makino H.	Current status of the treatment of microscopic polyangiitis and granulomatosis with polyangiitis in Japan.	<i>Clin Exp Nephrol</i>	17(1)	51-58	2013
Nakatsuka A, Wada J, Iseda I, Teshigawara S, Higashio K, Murakami K, Kanzaki M, Inoue K, Terami T, Katayama A, Hida K, Eguchi J, Horiguchi CS, Ogawa D, Matsuki Y, Hiramatsu R, Yagita H, Kakuta S, Iwakura Y, Makino H.	Vaspin is an adipokine ameliorating ER stress in obesity as a ligand for cell-surface GRP78/MTJ-1 complex.	<i>Diabetes</i>	61(11)	2823-2832	2012
Tachibana H, Ogawa D, Matsushita Y, Bruemmer D, Wada J, Teshigawara S, Eguchi J, Sato-Horiguchi C, Uchida HA, Shikata K, Makino H.	Activation of liver x receptor inhibits osteopontin and ameliorates diabetic nephropathy.	<i>J Am Soc Nephrol</i>	23(11)	1835-1846	2012
Inoue J, Wada J, Teshigawara S, Hida K, Nakatsuka A, Takatori U, Kojo S, Akagi S, Nakao K, Miyatake N, McDonald JF, Makino H.	The serum vaspin levels are reduced in Japanese chronic hemodialysis patients.	<i>BMC Nephrol</i>	13(1)	163	2012
Sato-Horiguchi C, Ogawa D, Wada J, Tachibana H, Kodera R, Eguchi J, Nakatsuka A, Terami N, Shikata K, Makino H.	Telmisartan attenuates diabetic nephropathy by suppressing oxidative stress in db/db mice.	<i>Nephron Exp Nephrol</i>	121(3-4)	e97-e108	2013

Nakatsuka A, Wada J, Iseda I, Teshigawara S, Higashio K, Murakami K, Kanzaki M, Inoue K, Terami T, Katayama A, Hida K, Eguchi J, Ogawa D, Matsuki Y, Hiramatsu R, Tagita H, Kakuta S, Iwakura Y, Makino H.	Vaspin inhibits apoptosis of endothelial cells as a ligand for cell-surface GRP78/VDAC complex.	<i>Circ Res</i>	112(5)	771-778	2013
Kurose Y, Wada J, Kanzaki M, Teshigawara S, Nakatsuka A, Murakami K, Inoue K, Terami T, Katayama A, Watanabe M, Higushi C, Eguchi J, Miyatake N, Makino H.	Serum galectin-9 levels are elevated in the patients with type 2 diabetes and chronic kidney disease.	<i>BMC Nephrol</i>	14	23	2013
Nakatsuka A, Wada J, Makino H	Cell cycle abnormality in metabolic syndrome and nuclear receptors as an emerging therapeutic target.	<i>Acta Medica Okayama</i>	67(3)	129-134	2013
Inoue K, Wada J, Eguchi J, Nakatsuka A, Teshigawara S, Murakami K, Ogawa D, Terami T, Katayama A, Tone A, Iseda I, Hida K, Yamada M, Ogawa T, Makino H	Urinary fetuin-A is a novel marker for diabetic nephropathy in type 2 diabetes identified by lectin microarray.	<i>PLoS ONE</i>	8(10)	e77118	2013
Terami T, Wada J, Inoue K, Nakatsuka A, Ogawa D, Teshigawara S, Murakami K, Katayama A, Eguchi J, Makino H	Urinary angiotensinogen is a marker for tubular injuries in patients with type 2 diabetes.	<i>Int J Nephrol Renovasc Dis</i>	6	233-240	2013

Fujiwara D, Takahashi K, Suzuki T, Shii M, Nakashima Y, Takekawa S, Yoshida A, Matsuoka T.	Postprandial serum C-peptide value is the optimal index to identify patients with non-obese type 2 diabetes who require multiple daily insulin injection: Analysis of C-peptide values before and after short-term intensive insulin therapy.	<i>Journal of Diabetes Investigation</i>	4(6)	618-625	2013
Ogawa D, Eguchi J, Wada J, Terami N, Hatanaka T, Tachibana H, Nakatsuka A, Horiguchi CS, Nishii N, Makino H.	Nuclear hormone receptor expression in mouse kidney and renal cell lines.	<i>PLoS ONE</i>	9(1)	e85594	2014
Watanabe M, Nakatsuka A, Murakami K, Inoue K, Terami T, Higuchi C, Katayama A, Teshigawara S, Eguchi J, Ogawa D, Watanabe E, Wada J, Makino H.	Pemt deficiency ameliorates endoplasmic reticulum stress in diabetic nephropathy.	<i>PLoS ONE</i>	9(3)	E92647	2014
Miyoshi T, Nakamura K, Yoshida M, Miura D, Oe H, Akagi S, Sugiyama H, Akazawa K, Yonezawa T, Wada J and Ito H.	Effect of vildagliptin, a dipeptidyl peptidase 4 inhibitor, on cardiac hypertrophy induced by chronic beta-adrenergic stimulation in rats.	<i>Cardiovascular Diabetology</i>	13	43	2014
Terami N, Ogawa D, Tachibana H, Hatanaka T, Wada J, Nakatsuka A, Eguchi J, Horiguchi CS, Nishii N, Yamada H, Takei K, Makino H.	Long-term treatment with the sodium glucose cotransporter 2 inhibitor, dapagliflozin, ameliorates glucose homeostasis and diabetic nephropathy in db/db mice.	<i>PLoS ONE</i>	9(6)	e100777	2014
Ono T, Shikata K, Obika M, Miyatake N, Koder R, Hirota D, Wada J, Kataoka H, Ogawa D, Makino H.	Factors associated with remission and/or regression of microalbuminuria in type 2 diabetes mellitus.	<i>Acta Med Okayama</i>	68(4)	235-241	2014

Hishikawa N, Yamashita T, Deguchi K, Wada J, Shikata K, MakinoH, Abe K.	Cognitive and affective functions in diabetic patients associated with diabetes-related factors, white matter abnormality and aging.	<i>Eur J Neurol</i>	Sep 15		2014
---	---	---------------------	--------	--	------