

## Case report

## Clinical features of three cases with pulmonary alveolar proteinosis secondary to myelodysplastic syndrome developed during the course of Behçet's disease



Tomohiro Handa<sup>a</sup>, Takeshi Nakatsue<sup>b</sup>, Motoo Baba<sup>c</sup>, Toshinori Takada<sup>d</sup>,  
Koh Nakata<sup>e,\*</sup>, Haruyuki Ishii<sup>f</sup>

<sup>a</sup>Department of Respiratory Medicine, Graduate School of Medicine, Kyoto University, 54 Kawaharacho Shogoin, Sakyo-ku, Kyoto 606-8507, Japan

<sup>b</sup>Division of Clinical Nephrology and Rheumatology, Niigata University Graduate School of Medical and Dental Sciences, 1-754 Asahimachi-dori, Chuo-ku, Niigata 951-8520, Japan

<sup>c</sup>Department of Respiratory Medicine, Tomishiro Chuo Hospital, 25, Ueda Tomishiro, JP-47, 901-0243, Japan

<sup>d</sup>Division of Respiratory Medicine, Niigata University Graduate School of Medical and Dental Sciences, 1-754 Asahimachi-dori, Chuo-ku, Niigata 951-8520, Japan

<sup>e</sup>Bioscience Medical Research Center, Niigata University, 1-754 Asahimachi Dori, Chuoku, Niigata 951-8520, Japan

<sup>f</sup>Department of Respiratory Medicine, Kyorin University School of Medicine, 6-20-2 Shinkawa, Mitaka, 181-8611, Japan

## ARTICLE INFO

## Article history:

Received 21 December 2012

Received in revised form

16 May 2013

Accepted 20 May 2013

Available online 3 July 2013

## Keywords:

Intestinal ulcer

Myelodysplastic syndrome

Sepsis

Trisomy 8

## ABSTRACT

We have previously reported that myelodysplastic syndrome (MDS) is the most common underlying disease in cases of secondary pulmonary alveolar proteinosis (PAP). Here, we present 3 MDS cases in which PAP developed during the course of Behçet's disease (BD). All patients carried trisomy 8 in the bone marrow. Chest HRCT scans showed variable distribution of ground glass opacities, but none of the scans showed so called "crazy paving appearance". Two patients with intestinal BD who underwent potent immunosuppressive therapy died of sepsis. These findings demonstrate that PAP secondary to MDS may be occasionally associated with BD.

© 2013 The Japanese Respiratory Society. Published by Elsevier B.V. All rights reserved.

Abbreviations: BAL, bronchoalveolar lavage; BD, Behçet's disease; GGO, ground glass opacity; HRCT, high-resolution computed tomography; MAC, Mycobacterium avium complex; MDS, myelodysplastic syndrome; PAP, pulmonary alveolar proteinosis; RA, refractory anemia; RAEB, RA with excess blasts; SLB, surgical lung biopsy; SPAP, secondary PAP; WLL, whole lung lavage

\*Corresponding author. Tel.: +81 25 227 0847; fax: +81 25 227 0377.

E-mail addresses: [hanta@kuhp.kyoto-u.ac.jp](mailto:hanta@kuhp.kyoto-u.ac.jp) (T. Handa), [nakatsue@med.niigata-u.ac.jp](mailto:nakatsue@med.niigata-u.ac.jp) (T. Nakatsue), [mbaba@yuuai.or.jp](mailto:mbaba@yuuai.or.jp) (M. Baba), [ttakada@med.niigata-u.ac.jp](mailto:ttakada@med.niigata-u.ac.jp) (T. Takada), [radical@med.niigata-u.ac.jp](mailto:radical@med.niigata-u.ac.jp) (K. Nakata), [h141@ks.kyorin-u.ac.jp](mailto:h141@ks.kyorin-u.ac.jp) (H. Ishii).

2212-5345/\$ - see front matter © 2013 The Japanese Respiratory Society. Published by Elsevier B.V. All rights reserved.  
<http://dx.doi.org/10.1016/j.resinv.2013.05.005>


## 1. Introduction

Pulmonary alveolar proteinosis (PAP) is a respiratory disease characterized by accumulation of phospholipids and surfactant proteins within the alveolar lumen and terminal bronchioli. PAP is classified into 3 groups on the basis of etiology: autoimmune PAP (APAP), secondary PAP (SPAP), and unclassified PAP. We previously reported that hematological disorders such as myelodysplastic syndrome (MDS) are the main underlying conditions in SPAP, and are observed in more than 70% of the cases of PAP [1]. However, the precise mechanism underlying the pathogenesis of this condition remains unclear.

Behçet's disease (BD) is a chronic, relapsing, inflammatory disease of unknown etiology that presents with oral aphthae, genital ulcers, uveitis, and skin lesions. Intestinal involvement is seen in 3–26% of cases, with a higher frequency in Asian than European countries. The disease is frequently intractable and may present with severe complications such as hemorrhage, perforation, and infection [2].

In a large cohort study of BD, which included 661 cases, PAP was not reported as a complication [3]. However, 2 BD cases with concomitant PAP have been reported in other studies [4,5]. When we studied the incidence of BD in the 40 Japanese cases of SPAP recorded in our database, we found 5 cases, including the 2 mentioned above. In this report, we present the clinical, radiological, and pathological features of the remaining 3 patients with SPAP and BD, and discuss the characteristic disease features common to all 5 cases, including those previously reported.

SPAP was diagnosed as described previously, with confirmation of the absence of serum GM-CSF autoantibodies [1,6]. BD was diagnosed according to the Behçet Disease Research Committee of Japan criteria [7]. Onset of BD was defined by the emergence of at least 2 symptoms attributable to BD [8]. High-resolution computed tomography (HRCT) images of the chest at the time of diagnosis were evaluated by 2 radiologists (M.A and H.I) as previously reported [6]. Consent was obtained from all identified patients by the treating physicians, and the study was approved by The



**Fig. 1** – (a) Chest HRCT in Case 1 showed diffuse apparent interlobular septal thickening. (b) Lung histology on surgical lung biopsy (SLB) (left S8) in Case 1. Elastica van Giesson stain (EVG),  $\times 40$ , of the lung biopsy specimen showed interstitial infiltration of inflammatory cells with many cholesterol clefts in the peribronchovascular areas (arrow heads). Amorphous material was found in the alveolar lumen (arrow), although it was not a predominant finding. (c) Chest HRCT in Case 2 showed predominant bilateral diffuse GGO in the upper and middle lobes. (d) Autopsy findings of the lung (PAS stain,  $\times 100$ ) in Case 2 showed diffuse alveolar septal thickening with mild cellular infiltration and with PAS-positive granular material within the alveolar spaces. (e). Chest HRCT in Case 3. Initial HRCT showed bilateral diffuse GGO, and multiple dense opacities developed later (arrow). The dense opacity had resolved with MAC treatment.

Institutional Review Board of Kyorin University School of Medicine (approval number H23-085, October 19, 2011).

## 2. Case presentation

Case 1 involved a female patient who developed oral and genital ulcers and papulopustular skin lesions at 32 years of age. The patient was diagnosed with intestinal BD 5 years later, and with MDS with refractory anemia (RA) and myelofibrosis at 46 years of age. Intestinal BD was diagnosed and prednisolone treatment was started at a dose of 40 mg/day and tapered to 5 mg/day. At 49 years of age, the patient was diagnosed with PAP on the basis of chest HRCT (Fig. 1a) and typical histologic findings in surgical lung biopsy (SLB) (Fig. 1b). Two years later, she was admitted to the hospital due to an exacerbation of intestinal BD and leukemic transformation of MDS. Despite receiving intensive treatment with steroid pulse, cyclosporine A, and infliximab, the intestinal involvement remained refractory without improvement of

persistent peritonitis. During treatment, the patient developed pneumonia caused by methicillin-resistant *Staphylococcus aureus* combined with *Mycobacterium avium complex* (MAC). This led to sepsis, with a fatal outcome 6 months after admission. The severity of PAP remained unchanged during the entire disease course.

Case 2 involved a male patient who developed oral ulcers and papulopustular skin lesions at 26 years of age. The patient was diagnosed with intestinal BD the following year, and MDS (RA) 5 years later. BD was refractory to long-term treatment with prednisolone combined with sulfasalazine. Thereafter, azathioprine combined with etanercept was administered for several weeks, but did not yield any improvement. At 33 years of age, the patient was admitted for treatment of intractable intestinal BD, when he was diagnosed with PAP on the basis of the findings of chest HRCT (Fig. 1c) and BAL. Whole lung lavage (WLL) could not be performed due to persistent fever, although no pathogen was detected. The patient died of PAP deterioration and septic shock 2 months after admission. Autopsy findings revealed

**Table 1 – Clinical features of secondary pulmonary alveolar proteinosis complicated with Behçet's disease.**

	Case 1	Case 2	Case 3	Literature case 1 [4]	Literature case 2 [5]	Control (n=35)
<i>At diagnosis of PAP</i>						
Age	49	33	50	51	39	53 (24-77)
Gender	F	M	F	F	F	M/F 21/14
Smoking history <sup>a</sup>	NS	S	S	S	S	S/NS/NA 19/11/5
Respiratory symptoms	none	cough	DOE	DOE	cough	16/35 (46%)
Diagnostic procedure	SLB	BS	SLB	BS	BS	BS/SLB 27/8
Serum KL-6 (U/mL)	936	1220	1050	1960	4160	2040 (358-20210)
Serum CEA (ng/mL)	2.0	12.6	NA	3.3	39.5	3.9 (0.5-36.0)
<i>HRCT findings</i>						
GGO pattern <sup>b</sup>	Diffuse	Diffuse	Diffuse	Patchy	Mixed	Diffuse 12/17 (71%)
'Crazy-paving' appearance	-	-	-	+	+	1/17 (6%)
Subpleural sparing	-	-	-	+	-	6/17 (35%)
Thickening of interlobular septa	+	-	-	-	+	4/17 (24%)
<i>Pathological findings</i>						
Distribution	SLB	Autopsy	SLB	N/A	N/A	
Cholesterol clefts	Perilobular	Diffuse	Patchy			
	Remarkable	Rare	Moderate			
<i>Behçet's disease</i>						
HLA-B51	-	-	N/A	N/A	N/A	
Oral ulcer	+	+	+	+	+	
Eye lesion	-	-	+	-	-	
Skin lesion	+	+	+	+	+	
Genital ulcer	+	-	+	+	+	
Intestinal lesion	+	+	-	-	+	
MDS	+	+	+	+	-	22 (63%)
WHO classification	RA	RA	RAEB-2	RAEB-1		
Trisomy 8	+	+	+	+		5/22 (23%)
WPSS	2	1	3	3		
Duration of BD prior to the onset of MDS	14 Years	5 Years	6 Months	14 Years		
Treatment before MDS onset <sup>c</sup>	P, S, C, T	P, S, A, T	Celecoxib	Colchicine		

Data are expressed as median (range). The control cohorts are patients with secondary pulmonary alveolar proteinosis but without Behçet's disease. NS, never smoked; DOE, dyspnea on effort; BS, bronchoscopy; SLB, surgical lung biopsy; RA, refractory anemia; RAEB, RA with excess of blasts; WPSS, WHO classification-based Prognostic Scoring System (1, low; 2, intermediate; 3, high-risk group); HRCT, high-resolution computed tomography; GGO, ground glass opacity and N/A, not available.

<sup>a</sup> S, current or former smoker.

<sup>b</sup> Mixed, mixed patchy geographic and diffuse pattern; patchy, patchy geographic pattern.

<sup>c</sup> P, prednisolone; S, sulfasalazine; C, cyclosporine A; T, TNF  $\alpha$ inhibitors and A, azathioprine.

invasive aspergillosis in the lung, but there was no evidence of leukemic transformation in the bone marrow.

Case 3 involved a female patient who developed uveitis at 38 years of age, which subsequently resolved without treatment. At the age of 48 years, she developed genital ulcers, skin lesions, and oral ulcers, and was diagnosed with complete BD. Six months later, abnormal shadowing was seen on a chest HRCT scan (Fig. 1e), and pancytopenia developed. The patient was diagnosed with PAP and MAC infection by surgical lung biopsy, and MDS (RA with excess blasts [RAEB] type 2) by bone marrow aspiration. Although lung opacity improved with treatment for MAC, the patient showed deterioration of PAP in the subsequent 2 years and required long-term oxygen therapy. The patient underwent WLL and showed improvement of lung opacity and oxygenation.

### 3. Discussion

In this report, we present 3 patients who developed SPAP during the course of BD and MDS. Their clinical, radiological, and pathological data, as well as those of the 2 cases of SPAP and BD already published in the literature [4,5] and the 35 cases of SPAP without BD in the Japanese SPAP registry [1] are summarized in Table 1. The clinical courses of the 3 patients are shown in Fig. 2. Notably, all 3 patients as well as another patient described in the literature [4] carried trisomy 8 in the bone marrow, and the onset of BD consistently preceded the onset of MDS and SPAP in these patients. HRCT features showed diffuse homogenous patterns in the distribution of ground glass opacity with or without thickening of the interlobular septa, which is characteristic for SPAP [6]. Generally, the clinical, serological, and radiological features of SPAP with BD were similar to those seen in the controls (Table 1). Lung pathology findings were variable with respect

to the extent of surfactant accumulation, the amount of cholesterol clefts, and the distribution of lesions. Intestinal BD may be associated with fatal infections due to the use of potent immunosuppressive therapy.

The co-occurrence of BD and MDS has been reported mainly in Japan and Korea [8-16]. These case reports were collected, and the demographic features are summarized in supplementary Table S1. A total of 64 cases of MDS complicated with BD or suspected BD have been reported during the period from 1988 to 2012. The meta-analysis revealed a high frequency of intestinal lesions (66%), trisomy 8 (80%) and a low frequency of ocular lesions (13%). In contrast, a nationwide survey of 3187 BD patients in Japan demonstrated a distinct distribution of intestinal and ocular involvements in 15.5% and 69.1%, respectively [8]. Thus, the disease phenotype of the 5 patients with SPAP and BD (Table 1) is similar to the phenotype of MDS-associated BD described in the literature. This is exemplified by the occurrence of trisomy 8 in the present cases. Since the general frequency of trisomy 8 is only 10-15% in MDS [17], MDS with trisomy 8 is likely to be a risk factor for both BD [9-11] and PAP [4], and they occasionally develop together, as in the 3 cases presented in this report. Immunosuppressive therapy for BD can cause MDS, which may subsequently cause SPAP. However, immunosuppressive drugs had not been administered in both case 3 and the case in the literature [4] until the onset of MDS (Table 1).

Previous large cohort studies have demonstrated a mortality rate of 5-9.8% during a follow up period of 7.7-19 years [18,19]. However, the effect of immunosuppressive treatment or the presence of intestinal lesions on mortality has not been evaluated. Patients with BD complicated with MDS were frequently treated with immunosuppressive drugs (Table S1). Of those, 89% and 94% with and without the intestinal involvements respectively, underwent immunosuppressive




Fig. 2 – Clinical course and prognosis of the patients. Time course of the 3 diseases (BD, Behcet disease; MDS, myelodysplastic syndrome and SPAP, secondary alveolar proteinosis) are shown. Treatments for BD were as follows: prednisolone (5-50 mg/day), sulfasalazine and TNF- $\alpha$ inhibitors in cases 1 and 2, cyclosporine A in case 1, azathioprine in case 2, celecoxib in case 3. Causes of death were as follows: Case 1: leukemic transformation and sepsis due to MRSA and MAC infection; Case 2: sepsis with unknown pathogen and PAP progression.

therapy. It is noteworthy that the mortality rate after such therapy was higher in those with intestinal involvement (41%) than those without involvement (21%). In 5 of 12 deceased cases with intestinal lesions, the cause of death was severe infection. Consistently, 2 fatal cases presented here and another previously published case [5] showed intestinal involvement, and therefore required potent immunosuppressive therapy.

In conclusion, SPAP secondary to MDS is a rare complication during the course of BD. The clinical features other than BD-related findings were not distinguished from those seen in SPAP without BD.

### Funding source

This study was funded by a grant from the Ministry of Health, Labour, and Welfare, Japan (H24-Nanchi-Ippann-Japan-035, and 10103322). The funding source had no role in study design, data collection, or in the decision to submit the paper for publication.

### Conflict of interest

The authors have no conflicts of interest.

### Acknowledgments


We would like to thank Dr. Carmel J. Stock (Interstitial Lung Disease Unit, Royal Brompton Hospital) for English proof-reading. We thank Dr. Aya Nishida (Department of Hematology, Toranomon Hospital), Dr. Michihiro Uchiyama, (Department of Hematology, Suwa Red Cross Hospital), Dr. Yoshikazu Inoue (Department of Diffuse Lung Diseases and Respiratory Failure, NHO Kinki-Chuo Chest Medical Center), Dr. Toshio Ichiwata (Department of Respiratory Medicine, Tokyo Medical University Hachioji Medical Center), Dr. Kohei Ikezoe (Department of Respiratory Medicine, Graduate School of Medicine, Kyoto University), Dr. Emi Hamano (Department of Respiratory Medicine, The University of Tokyo Hospital), Dr. Sonoko Nagai (Kyoto Central Clinic/Clinical Research Center), Dr. Michiaki Mishima (Department of Respiratory Medicine, Graduate School of Medicine, Kyoto University), and Dr. Hajime Goto (Department of Respiratory Medicine, Kyorin University School of Medicine) for their contribution to the clinical assessment. We thank Dr. Masanori Akira (Department of Radiology, NHO Kinki-Chuo Chest Medical Center) for his contribution in the radiological assessment. We also thank Dr. Akira Hebisawa (Department of Pathology, NHO Tokyo Hospital) and Dr. Akihiko Yoshizawa (Department of Laboratory Medicine, Shinshu University Hospital) for their contribution in the histopathological assessment.

### Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.resinv.2013.05.005>.

### REFERENCES

- [1] Ishii H, Tazawa R, Kaneko C, et al. Clinical features of secondary pulmonary alveolar proteinosis: pre-mortem cases in Japan. *Eur Respir J* 2011;37:465-8.
- [2] Iwata S, Saito K, Yamaoka K, et al. Efficacy of combination therapy of anti-TNF- $\alpha$  antibody infliximab and methotrexate in refractory entero-Beçet's disease. *Mod Rheumatol* 2011;21:184-91.
- [3] Alpsoy E, Donmez L, Onder M, et al. Clinical features and natural course of Behçet's disease in 661 cases: a multicentre study. *Br J Dermatol* 2007;157:901-6.
- [4] Nishida A, Miyamoto A, Yamamoto H, et al. Possible association of trisomy 8 with secondary pulmonary alveolar proteinosis in myelodysplastic syndrome. *Am J Respir Crit Care Med* 2011;184:279-80.
- [5] Uchiyama M, Nagao T, Hattori A, et al. Pulmonary alveolar proteinosis in a patient with Behçet's disease. *Respirology* 2009;14:305-8.
- [6] Ishii H, Trapnell BC, Tazawa R, et al. Comparative study of high-resolution CT findings between autoimmune and secondary pulmonary alveolar proteinosis. *Chest* 2009;136:1348-55.
- [7] Mizushima Y. Recent research into Behçet's disease in Japan. *Int J Tissue React* 1988;10:59-65.
- [8] Tada Y, Koarada S, Haruta Y, et al. The association of Behçet's disease with myelodysplastic syndrome in Japan: a review of the literature. *Clin Exp Rheumatol* 2006;24:S115-9 [Erratum in: *Clin Exp Rheumatol* 2007;25:507-8].
- [9] Fujimura T, Yukawa N, Nakashima R, et al. Periodic fever and erythema nodosum associated with MDS with trisomy 8: report of two cases and review of the literature. *Mod Rheumatol* 2010;20:413-9.
- [10] Kawabata H, Sawaki T, Kawanami T, et al. Myelodysplastic syndrome complicated with inflammatory intestinal ulcers: significance of trisomy 8. *Intern Med* 2006;45:1309-14.
- [11] Ahn JK, Cha HS, Koh EM, et al. Behçet's disease associated with bone marrow failure in Korean patients: clinical characteristics and the association of intestinal ulceration and trisomy 8. *Rheumatology (Oxford)* 2008;47:1228-30.
- [12] Lin YC, Liang TH, Chang HN, et al. Behçet's disease associated with myelodysplastic syndrome. *J Clin Rheumatol* 2008;14:169-74.
- [13] Mantzourani MG, Chantziara K, Thanopoulou I, et al. Coexistence of Behçet's disease and chronic myelomonocyte leukemia with trisomy 8: a case report and review of the literature. *Clin Exp Rheumatol* 2009;27:S85-7.
- [14] Kovacs E, Nemeth H, Telek B, et al. Behçet's disease in a patient with myelodysplastic syndrome. *Clin Lymphoma Myeloma* 2009;9:459-61.
- [15] Nakamoto K, Hatano Y, Furumi K, et al. A fatal case of Behçet's disease with every three special types of the complication accompanied with myelodysplastic syndrome and immunodeficiency. *Iryo* 1994;48:1083-7 [in Japanese].
- [16] Cengiz M, Altundag MK, Zorlu AF, et al. Malignancy in Behçet's disease: a report of 13 cases and a review of the literature. *Clin Rheumatol* 2001;20:239-44.
- [17] Lee DS, Kim SH, Seo EJ, et al. Predominance of trisomy 1q in myelodysplastic syndromes in Korea: is there an ethnic difference? A 3-year multi-center study. *Cancer Genet Cytogenet* 2002;132:97-101.
- [18] Saadoun D, Wechsler B, Desseaux K, et al. Mortality in Behçet's disease. *Arthritis Rheum* 2010;62:2806-12.
- [19] Kural-Seyahi E, Fresko I, Seyahi N, et al. The long-term mortality and morbidity of Behçet syndrome: a 2-decade outcome survey of 387 patients followed at a dedicated center. *Medicine (Baltimore)* 2003;82:60-76.



## Standardized serum GM-CSF autoantibody testing for the routine clinical diagnosis of autoimmune pulmonary alveolar proteinosis

Kanji Uchida<sup>a,b</sup>, Koh Nakata<sup>c</sup>, Brenna Carey<sup>a</sup>, Claudia Chalk<sup>a</sup>, Takuji Suzuki<sup>a</sup>, Takuro Sakagami<sup>a</sup>, Diana E. Koch<sup>a</sup>, Carrie Stevens<sup>a</sup>, Yoshikazu Inoue<sup>d</sup>, Yoshitsugu Yamada<sup>b</sup>, Bruce C. Trapnell<sup>a,e,\*</sup>

<sup>a</sup> Cincinnati Children's Hospital Medical Center, 3333 Burnet Avenue, Cincinnati, OH 45229-3039, USA

<sup>b</sup> Graduate School of Medicine, The University of Tokyo, Tokyo, Japan

<sup>c</sup> Niigata University Medical & Dental Hospital, Niigata, Japan

<sup>d</sup> National Hospital Organization, Kinki-Chuo Chest Medical Center, Sakai, Japan

<sup>e</sup> University of Cincinnati College of Medicine, Cincinnati, OH, USA

### ARTICLE INFO

#### Article history:

Received 3 October 2013

Received in revised form 16 November 2013

Accepted 18 November 2013

Available online 23 November 2013

#### Keywords:

Pulmonary alveolar proteinosis

Granulocyte/macrophage-colony stimulating factor

Autoimmune disease

Autoantibodies

Enzyme linked immunosorbent assay

Diagnosis

### ABSTRACT

Autoantibodies against granulocyte/macrophage colony-stimulating factor (GMABs) cause autoimmune pulmonary alveolar proteinosis (PAP) and measurement of the GMAB level in serum is now commonly used to identify this disease, albeit, in a clinical research setting. The present study was undertaken to optimize and standardize serum GMAB concentration testing using a GMAB enzyme-linked immunosorbent assay (GMAB ELISA) to prepare for its introduction into routine clinical use. The GMAB ELISA was evaluated using serum specimens from autoimmune PAP patients, healthy people, and GMAB-spiked serum from healthy people. After optimizing assay components and procedures, its accuracy, precision, reliability, sensitivity, specificity, and ruggedness were evaluated. The coefficient of variation in repeated measurements was acceptable (<15%) for well-to-well, plate-to-plate, day-to-day, and inter-operator variation, and was not affected by repeated freeze-thaw cycles of serum specimens or the reference standards, or by storage of serum samples at  $-80^{\circ}\text{C}$ . The lower limit of quantification (LLOQ) of the PAP patient-derived polyclonal GMAB reference standard (PCRS) was 0.78 ng/ml. Receiver operating characteristic curve analysis identified a serum GMAB level of 5  $\mu\text{g}/\text{ml}$  (based on PCRS) as the optimal cut off value for distinguishing autoimmune PAP serum from normal serum. A pharmaceutical-grade, monoclonal GMAB reference standard (MCRS) was developed as the basis of a new unit of measure for GMAB concentration: one International Unit (IU) of GMAB is equivalent to 1  $\mu\text{g}/\text{ml}$  of MCRS. The median [interquartile range] serum GMAB level was markedly higher in autoimmune PAP patients than in healthy people (21.54 [12.83–36.38] versus 0.08 [0.05–0.14] IU;  $n = 56, 38$ ; respectively;  $P < 0.0001$ ). Results demonstrate that serum GMAB measurement using the GMAB ELISA was accurate, precise, reliable, had an acceptable LLOQ, and could be accurately expressed in standardized units. These findings support the use of this GMAB ELISA for the

**Abbreviations:** PAP, pulmonary alveolar proteinosis; PCRS, polyclonal reference standard; MCRS, monoclonal reference standard; rhGM-CSF, recombinant, human GM-CSF.

\* Corresponding author at: Divisions of Pulmonary Biology and Medicine, Cincinnati Children's Hospital Medical Center, Room 4029, CCRF, 3333 Burnet Avenue, Cincinnati, OH 45229-3039, USA. Tel.: +1 513 636 6361; fax: +1 513 636 3723.

**E-mail addresses:** [uchidak-ane@h.u-tokyo.ac.jp](mailto:uchidak-ane@h.u-tokyo.ac.jp) (K. Uchida), [radical@med.niigata-u.ac.jp](mailto:radical@med.niigata-u.ac.jp) (K. Nakata), [Brenna.Carey@cchmc.org](mailto:Brenna.Carey@cchmc.org) (B. Carey), [Claudia.Chalk@cchmc.org](mailto:Claudia.Chalk@cchmc.org) (C. Chalk), [Takuji.Suzuki@cchmc.org](mailto:Takuji.Suzuki@cchmc.org) (T. Suzuki), [stakuro@med.niigata-u.ac.jp](mailto:stakuro@med.niigata-u.ac.jp) (T. Sakagami), [kochde83@gmail.com](mailto:kochde83@gmail.com) (D.E. Koch), [Carrie.Stevens@cchmc.org](mailto:Carrie.Stevens@cchmc.org) (C. Stevens), [giichi@kch.hosp.go.jp](mailto:giichi@kch.hosp.go.jp) (Y. Inoue), [yamaday-ane@h.u-tokyo.ac.jp](mailto:yamaday-ane@h.u-tokyo.ac.jp) (Y. Yamada), [Bruce.Trapnell@cchmc.org](mailto:Bruce.Trapnell@cchmc.org) (B.C. Trapnell).

0022-1759/\$ – see front matter © 2013 Published by Elsevier B.V.

<http://dx.doi.org/10.1016/j.jim.2013.11.011>

routine clinical diagnosis of autoimmune PAP and introduce a new unit of measure to enable standardized reporting of serum GMAB data from different laboratories.

© 2013 Published by Elsevier B.V.

## 1. Introduction

Autoimmune pulmonary alveolar proteinosis (autoimmune PAP) is a rare disease characterized by alveolar surfactant accumulation, respiratory failure, and an increased risk of opportunistic infections (Trapnell et al., 2003). The disease is strongly associated with granulocyte/macrophage-colony stimulating factor autoantibodies (GMABs) (Kitamura et al., 1999) that neutralize GM-CSF bioactivity (Kitamura et al., 1999; Uchida et al., 2004) and mediate pathogenesis by blocking signaling to alveolar macrophages (Sakagami et al., 2009, 2010) and neutrophils (Uchida et al., 2007). Alveolar macrophages require GM-CSF for terminal differentiation and constitutive regulation of functions including surfactant clearance (Shibata et al., 2001; Bonfield et al., 2003). Without pulmonary GM-CSF signaling, alveolar macrophages have impaired pulmonary surfactant clearance (Ikegami et al., 1996) resulting in the slow, progressive surfactant accumulation and the insidious onset of the clinical manifestations of PAP syndrome. Disruption of GM-CSF signaling by recessive mutations in *CSF2RA* or *CSF2RB*, which encode the GM-CSF receptor  $\alpha$  and  $\beta$  chains, respectively, causes a hereditary form of PAP that is clinically, physiologically, and histologically indistinguishable from autoimmune PAP (Martinez-Moczygemba et al., 2008; Suzuki et al., 2008, 2010, 2011; Tanaka et al., 2011). PAP can also occur in a heterogeneous group of other diseases either as a consequence of an underlying clinical condition presumably affecting alveolar macrophage function (secondary PAP) or mutations in the genes involved surfactant production (e.g., *SFTPB*, *SFTPC*, *ABCA3*, *TTF1*) (congenital PAP, and PAP associated with interstitial lung disease) (Nogee, 2010).

Clinically, the diagnosis of PAP is made based on a compatible history, typical radiologic findings, and characteristic lung biopsy or bronchoalveolar lavage cytology findings. However, while this approach can determine if PAP syndrome is present, it cannot identify the underlying causative disease. The strong association of a high serum GM-CSF autoantibody (GMAB) level with autoimmune PAP (Kitamura et al., 1999; Bonfield et al., 2002; Trapnell et al., 2003; Inoue et al., 2008), development of an ELISA to measure GMABs (Schoch et al., 2002), and demonstration that GMABs actually drive the pathogenesis of autoimmune PAP (Uchida et al., 2007; Sakagami et al., 2009, 2010), support what is now widespread use of serum GMAB measurement for the clinical research diagnosis of autoimmune PAP. The potential clinical use of the GMAB ELISA is further supported by the identification of critical threshold of serum GMAB that is associated with an increased risk of autoimmune PAP (Bendtsen et al., 2007; Uchida et al., 2009; Sakagami et al., 2010). Several GMAB ELISA-based methods have been developed for measurement of serum GMAB including a quantitative method based on use of a neutralizing, polyclonal GMAB reference standard (PCRS) (Kitamura et al., 1999; Schoch et al., 2002) and a non-

neutralizing, monoclonal GMAB reference standard (MCRS) (Inoue et al., 2008) that return values in units of GMAB concentration and a serum titration method that returns values in units of GMAB titer (Kitamura et al., 1999; Bonfield et al., 2002).

The purpose of the present study was to optimize the GMAB ELISA with respect to reagents, experimental protocol, and analysis methods, and then validate it by rigorously establishing its sensitivity, accuracy, precision, and ruggedness to support its clinical use for the diagnosis of autoimmune PAP. Further, we compared the relative performance of a GMAB polyclonal reference standard (PCRS) purified from PAP patient serum against a GMAB monoclonal reference standard (MCRS) prepared under good manufacturing practices for potential use as a reference standard for calibrating the results obtained in different laboratories.

## 2. Materials and methods

### 2.1. Participants

The institutional review board of the Cincinnati Children's Hospital Medical Center approved the study. All participants or their legal guardians gave written informed consent and minors gave assent. Participants included patients with autoimmune PAP ( $n = 96$ ; 45.8% male;  $37.3 \pm 15.7$  years of age at evaluation) referred for evaluation or treatment. The diagnosis of autoimmune PAP was based on clinical and radiographic findings; an open lung biopsy, transbronchial lung biopsy, or cytologic analysis of bronchoalveolar lavage cells and fluid; and a positive GMAB test performed as previously described (Uchida et al., 2004). We also studied healthy people ( $n = 58$ ; 22.4% male;  $30.6 \pm 7.0$  years of age at evaluation) who were nonsmokers with no history of major illness and symptom-free at the time of evaluation.

### 2.2. GMAB polyclonal reference standard

A GMAB polyclonal reference standard (PCRS) was prepared as previously described (Schoch et al., 2002) from a patient with autoimmune PAP (Luisetti et al., 2009). Briefly, GMAB was isolated from plasmapheresis fluid by protein G column chromatography followed by GM-CSF affinity chromatography (Schoch et al., 2002), concentrated by ultrafiltration (Ultra-15; MWCO 30 kDa, Amicon), and re-suspended in phosphate-buffered saline (PBS). The concentration of GMAB was determined previously by comparison to a GMAB reference standard for which the IgG content had been determined by ELISA (Kitamura et al., 2000). The purity of the PCRS was assessed by polyacrylamide gel electrophoresis (ReadyGel®, 5–15% gradient, BioRad Laboratories, Hercules, CA) with 2-mercaptoethanol. A 'PCRS Master Stock' containing

the final purified GMAb at 2 mg/ml in PBS was prepared as 10  $\mu$ l aliquots in 0.5 ml polypropylene tubes (Eppendorf, Hamburg, Germany) and stored at  $-80^{\circ}\text{C}$  with continuous electronic temperature monitoring as previously reported (Uchida et al., 2009). A 'PCRS Working Standard' was prepared by diluting one vial of PCRS Master Stock with ELISA dilution buffer to GMAb concentration of 100 ng/ml and stored as 250  $\mu$ l frozen aliquots as above. GMAb ELISA plate concentration standards were prepared from the PCRS Working Standard by thawing one vial at room temperature and diluting a 120  $\mu$ l aliquot serially (1/2) to create standard concentrations (50, 25, 12.5, 6.25, 3.125, 1.5625 ng/ml). ELISA buffer without PCRS served as a 0 ng/ml control. A serial dilution to 0.78125 ng/ml was used in some experiments to define the lower limit of quantification (LLOQ).

### 2.3. GMAb monoclonal reference standard

A GMAb monoclonal reference standard (MCRS) was previously prepared from a PAP patient-derived, Epstein-Barr virus transformed, immortalized B lymphocyte clone. One of six initial cell clones was used to produce the MCRS by Boehringer Ingelheim pharmaceuticals (Biberach, Germany). Briefly, the B cell clone-derived cDNA was used to stably-transduce Chinese hamster ovary D644 cells (Urlaub et al., 1983). A pool of MCRS-producing cells was subjected to 10 day fed-batch cultures in a 30 and 50 l bio-fermenter and MCRS was purified from culture supernatant using protein A chromatography (MabSelect SuRE™, GE Healthcare, Inc.). Purified MCRS was re-suspended in 25 mM sodium citrate, pH 6.0, 115 mM sodium chloride, 0.01% (w/v) polysorbate 20, pH 6. Its binding affinity for GM-CSF was determined to be 423 pM using KinExA (Sapidyne Instruments, Inc.) equilibrium experiments, which were conducted at room temperature ( $\sim 21^{\circ}\text{C}$ ). The antibody concentration was held constant as the antigen (rGM-CSF, Biomol GmbH, Germany) was titrated in a 2-fold serial dilution. Solutions were allowed to come to equilibrium prior measurement and the free antibodies were captured by rGM-CSF-rabbit Fc coupled to beads. The final MCRS preparation (monoclonal, non-neutralizing, human anti-human GM-CSF IgG, lambda antibody; BI 01049904, Lot 1, 05.08.2009; 500  $\mu\text{g}/500 \mu\text{l}$ ; MCRS Original Stock) was shipped on dry ice with continuous electronic temperature recording to Cincinnati, Ohio. A "MCRS Master Stock" was prepared by thawing the Original Stock tube at room temperature, mixing thoroughly by gentle inversion, dispensing as 25  $\mu$ l aliquots without dilution in 0.5 ml polypropylene tubes (Eppendorf, Hamburg, Germany) and then storing frozen as for the PCRS. A 'MCRS Working Stock' was prepared by diluting one vial (25  $\mu$ l) of MCRS Master Stock with ELISA dilution buffer (4.975 ml; PBS, Tween-20, 1% BSA) to a final concentration of 5  $\mu\text{g}/\text{ml}$ , which was dispensed as 200  $\mu$ l aliquots and stored frozen as above. A 'MCRS Working Standard' was prepared by diluting one vial (200  $\mu$ l) of MCRS Master Stock with ELISA dilution buffer (49.8 ml) to 20 ng/ml and stored frozen in 250  $\mu$ l aliquots as above. GMAb ELISA plate concentration standards were prepared from the MCRS Working Standard as above to create standard concentrations (20, 10, 5, 2.5, 1.25, 0.63, 0.31). ELISA buffer without MCRS serves as the 0 ng/ml control.

### 2.4. GM-CSF capture antigen

Preparations of recombinant, human GM-CSF from several suppliers (Miltenyi Biotech, Auburn, CA, USA; Prospec-Tany Technogene Ltd, Rehovot, Israel, USA; Invitrogen Corp., Camarillo, CA, USA; Genway Biotech, Inc., San Diego, CA, USA; Genzyme, Cambridge, MA, USA) were evaluated as a capture antigen in the GMAb ELISA by measuring the baseline optical absorbance for each preparation in the absence of added serum sample. The rhGM-CSF from Miltenyi Biotech (Fig. 1C, Supplier 2) was chosen for the standard GMAb ELISA and was used throughout this study except where indicated.


### 2.5. Human immunoglobulin G detection antibody

Two anti-human immunoglobulin G (IgG) reagents (goat anti-human IgG-specific F(ab')<sub>2</sub> antibody fragment, #A2290; and intact goat anti-human-IgG-Fc specific antibody, #A6029; both from Sigma-Aldrich, St. Louis, MO, USA) were evaluated as the secondary detection antibody in the GMAb ELISA. The F(ab')<sub>2</sub> antibody was chosen for the standard GMAb ELISA and was used throughout this study except where indicated.

### 2.6. GMAb ELISA

The standard GMAb ELISA used was based on an initial description (Nakata, 1999; Kitamura et al., 2000; Schoch et al., 2002) and subsequent modifications (Uchida et al., 2004) and performed as follows. Microtiter plates (96-well, Maxisorp; Nalge Nunc International, Rochester, NY) were incubated ( $4^{\circ}\text{C}$ , overnight) with 50  $\mu$ l of capture antigen solution (1  $\mu\text{g}/\text{mL}$  rhGM-CSF (Miltenyi except where noted) in phosphate buffered saline (PBS)), washed in PBS containing 0.1% Tween 20 (Sigma-Aldrich, St. Louis, MO, USA), and incubated (room temperature, 1 h) with blocking solution, Stabilcoat® (Surmodics, Eden Prairie, MN, USA). Serum samples (100  $\mu$ l except where indicated) were mixed with sample dilution buffer (PBS, 1% [vol/vol] BSA, 0.1% [wt/vol] Tween 20) to prepare a standard dilution series (1/101, 1/3001, 1/6001 and 1/12,001) for each sample. Aliquots (50  $\mu$ l) of diluted serum or reference standards were pipetted into adjacent microtiter wells, incubated (room temperature, 40 min) to allow binding of GMAb to the capture antigen, and then washed five times with 300  $\mu$ l of wash buffer (PBS, 0.1% [vol/vol] Tween 20). Horse radish peroxidase-conjugated anti-human IgG F(ab')<sub>2</sub> fragment (except where noted) was diluted 1/3000 with sample dilution buffer for use as the detection antibody and 50  $\mu$ l was pipetted into each well. The plates were incubated (room temperature, 30 min) to allow binding of IgG detection antibody to captured GMAb and then rinsed 4 times with wash buffer. Chromogenic substrate solution (50  $\mu$ l; 3,3',5,5'-tetramethylbenzidine; T4444, Sigma-Aldrich) was added to each well and the plates were incubated (room temperature, 15 min) to permit color development, which was stopped by adding 50  $\mu$ l of 1 N H<sub>2</sub>SO<sub>4</sub>. Optical absorbance at wavelength of 450 nm was measured using a Benchmark® ELISA plate reader using Microplate Manager software, version 5.21 (Bio-Rad Laboratories, Hercules, CA, USA) as described by the manufacturer.





**Fig. 1.** Effects of the capture antigen on GMAb ELISA performance. A. Non-glycosylated versus glycosylated GM-CSF capture antigen. Serum GMAb concentration measured with the GMAb ELISA using GM-CSF produced in *E. coli* (non-glycosylated form) or in yeast (glycosylated form). Each symbol represents the mean of 2 determinations on serum samples from each autoimmune PAP patient or healthy person ( $n = 20$  and  $10$ , respectively). The GMAb concentration measured for each sample with each capture antigen was not different ( $P = 0.796$ ,  $n = 30$ ; Mann-Whitney Rank Sum Test). B. Bland and Altman analysis. All data shown in panel A were evaluated by Bland and Altman analysis as described in the Materials and methods section. C. Source of *E. coli*-derived GM-CSF capture antigen. ELISA plates were coated with GM-CSF ( $1 \mu\text{g/ml}$ ) produced in *E. coli* obtained from several suppliers (1–5) and then used in the GMAb ELISA with sham samples (i.e., buffer instead of serum) to determine the effect on background optical absorbance. Each bar represents the mean ( $\pm$ SD) of 4 separate determinations. The background absorbance was different for all comparisons except between suppliers 1, 4, and 5 ( $P < 0.001$ ,  $n = 4$  each; ANOVA with pairwise multiple comparison procedures by the Holm-Sidak method).

GMAb concentration was determined as follows. The mean optical absorbance of the PCRS in replicate wells was plotted (x-axis) against the known GMAb concentration (y-axis) and quadratic regression analysis (except where noted) was performed to determine the equation relating

optical absorbance to PCRS concentration (ng/ml on plate) using Microsoft Excel (Microsoft Corp, Seattle, WA, USA). The mean optical absorbance of replicate wells for each sample was substituted for  $x$  in the regression equation to determine the concentration of GMAb in the well in ng/ml, which was multiplied by the dilution factor to determine the GMAb concentration in serum in  $\mu\text{g/ml}$ .

Quality control standards consisting of duplicate serum samples of known GMAb concentration were run on every plate. All results from any plate for which the quality control replicates differed by more than 15% were rejected, which occurred in one of eighty plates in this study.

### 2.7. Statistical analysis


Numerical data were evaluated for a normal distribution using the Shapiro-Wilk test and for equal variance using the Levene median test. Parametric data are presented as means ( $\pm$ SD) and nonparametric data are presented as medians (interquartile range [IQR]). Statistical comparisons of parametric data were made with Student's  $t$ -test for two-group comparisons and with one-way analysis of variance with post hoc analysis by the Holm-Sidak method for comparisons among three or more groups. Nonparametric data were compared with the use of the Mann-Whitney Rank-Sum test. Agreement between assay methods was made using Bland-Altman difference analysis. Receiver operating characteristic (ROC) curve analysis and other statistical analyses were done using SigmaPlot software (Version 12.3, Systat Software, San Jose, CA, USA).  $P$  values less than 0.05 were considered to indicate statistical significance. All experiments were repeated at least twice, with similar results.

## 3. Results

### 3.1. Optimization of GMAb ELISA components and procedure

GMAb concentration measured with the GMAb ELISA using *E. coli*- or yeast-derived GM-CSF as the capture antigen was similar indicating that glycosylation had little effect on capture antigen function (Fig. 1A, B). Because prior studies indicated that some rhGM-CSF preparations had a high background absorption, GM-CSF from multiple sources was evaluated. Background absorbance in the GMAb ELISA varied significantly for rhGM-CSF from different suppliers (Fig. 1C). Thus every batch of rhGM-CSF considered for use as the capture antigen must be checked for acceptability to ensure a low background absorbance since higher values will affect the cutoff between normal and disease (see below). Non-glycosylated rhGM-CSF produced in *E. coli* was chosen for use in the standard GMAb ELISA to minimize potential non-specific binding to the carbohydrate moiety.

Use of an  $F(ab')_2$  fragment (#A2290) or intact IgG (#A6029) as the anti-human IgG secondary (detection) antibody in the GMAb ELISA gave similar results except at higher GMAb concentrations where the intact IgG gave slightly higher values (Fig. 2A, B). Since  $F(ab')_2$  gave good results and was used in our prior reports (Schoch et al., 2002; Trapnell et al., 2003; Uchida et al., 2004, 2007, 2009), it was chosen for routine use in the standard GMAb ELISA to maximize comparability to previously reported results.




**Fig. 2.** Effects of the secondary detection antibody on GMAb ELISA performance. A. Use of intact goat-anti-human IgG or an F(ab')<sub>2</sub> fragment as the detection antibody. Serum GMAb concentration measured with the GMAb ELISA in parallel using either an F(ab')<sub>2</sub> fragment (A2290) or intact anti-human IgG (A6029) as the detection antibody. Each symbol represents the mean of 3 determinations on the serum sample from a different autoimmune PAP patient. The median GMAb concentration (not shown) determined with the intact IgG secondary antibody was slightly higher than with the F(ab')<sub>2</sub> secondary antibody (128 [51–234] vs 118 [43–187] µg/mL;  $P < 0.001$ ;  $n = 29$  serum samples; Wilcoxon Signed Rank test). B. Bland and Altman analysis. All data shown in panel A were evaluated by Bland and Altman analysis as described in the Materials and methods section.

Patient-derived GMAb PCRS contained three molecular species similar in molecular mass to IgG1, IgG2, (predominant species of roughly equal proportion) and IgG3, and IgG4

**Fig. 3.** Characterization and performance of a PAP patient-derived GMAb polyclonal reference standard (PCRS). A. Purity and composition of the PCRS. PCRS, prepared as described in the Materials and methods section, commercially available IgG heavy chain isotype standards (IgGκ 1, 2, 3, or 4), or molecular weight markers (MWM) were subjected to polyacrylamide gel electrophoresis under reducing conditions, Coomassie blue staining, and photography as described in the Materials and methods section. B. Optical absorbance of the PCRS as a function of concentration. The PCRS was serially diluted and evaluated as the standard in the GMAb ELISA as described in the Materials and methods section. Optical absorbance increased smoothly in proportion with PCRS concentration. Regression analysis using a quadratic equation yielded a correlation coefficient ( $R^2$ ) of 0.9998. Each point represents the mean ( $\pm$ SD) of 5 separate measurements. C. Effect of regression method used on percent error of the PCRS curve fit. Results from 5 independent, simultaneously conducted experiments determining the optical absorbance of serial dilutions of the PCRS were subjected individually to linear, quadratic, or logarithmic regression analysis and the mean ( $\pm$ SD) percent deviation at each concentration was determined. The percent error of GMAb concentration was determined as the measured value minus the expected value divided by the expected value and multiplied by 100. The mean ( $\pm$ SD) correlation coefficients for regression analysis of 5 separate experiments (not shown) were  $0.9999 \pm 0.0001$  (quadratic),  $0.9969 \pm 0.0029$  (log<sub>10</sub>), and  $0.9819 \pm 0.0043$  (linear).


(minor species; 0.2–3.5%) (Fig. 3A) consistent with our prior report that GMABs from autoimmune PAP patients and healthy people are composed primarily of IgG1 and IgG2 with negligible amounts of IgG3 and IgG4 (Uchida et al.,



2009). The  $IC_{50}$  of the PCRS was previously shown to be 10.3 molecules of GMAb per molecule of GM-CSF (Sakagami et al., 2010), similar to values reported for GMAb from a single patient (10.6), GMAb from serum pooled from 11 patients (10.3) (Uchida et al., 2007), or the mean value for GMAb from 11 patients determined individually ( $7.05 \pm 3.81$ ) (Uchida et al., 2004). In the GMAb ELISA, optical absorbance increased as a smooth, slightly curved function of PCRS concentration over a range from 0.781 to 50 ng/ml (Fig. 3B). Quadratic regression analysis of optical absorbance versus PCRS concentration yielded an outstanding correlation coefficient and gave a lower percent deviation for repeated measurements over the entire PCRS concentration range than did either linear or log regression analysis (Fig. 3C).

Evaluation of the initial serum aliquot volume used to prepare the standard serum dilutions for measurement in the GMAb ELISA revealed an important source of assay variability. The mean serum GMAb concentration was slightly higher with 10  $\mu$ l initial serum aliquots than with 100  $\mu$ l aliquots but did not reach statistical significance (Fig. 4A). However, the percent deviation was significantly greater for smaller aliquots (Fig. 4B). The total amount of serum transferred to the first dilution tube includes that intended for measurement (i.e., serum inside the pipet tip) plus unintentionally included serum (i.e., serum adherent to the outside and end of the pipet tip), these findings suggest that unintentionally included serum comprised a greater fraction of the total amount transferred when the initial serum aliquot was smaller. The net effect was a tendency to overestimate GMAb concentration and markedly increased variability of repeated measurements.

Since dilution is required to reduce the optical absorbance of PAP patient serum samples on the ELISA plate into a readable range and experience has shown that the serum GMAb concentration determined can vary with the dilution used to make the measurement, we evaluated the effect of the serum dilution used for determining GMAb concentration on variability of the results for both PAP patients and healthy people. As expected, the percent deviation of GMAb concentration was higher for diluted samples with optical absorbance values falling outside the range of reference standard values ('out-of-range') (Fig. 5A) at all dilutions (Fig. 5B) in both PAP patients and healthy controls. In contrast, the percent deviation was lowest and similar at all dilutions for PAP patient serum when the absorbance was 'in-range' (Fig. 5B). Most PAP patients had an in-range absorbance value at a dilution of 1/3000 or 1/6000, with slightly fewer at





**Fig. 4.** Effect of serum aliquot volume on GMAb ELISA performance. A. Effect on serum GMAb concentration. An initial volume of serum either 10 or 100  $\mu$ l was used to measure serum GMAb concentration with the GMAb ELISA. Bars represent the mean ( $\pm$ SD) of 20 independent determinations using the same autoimmune patient serum. The gray line represents the mean for the data obtained using the 100  $\mu$ l aliquots of serum. The difference in median (IQR) GMAb concentration for the assay performed using 10 vs 100  $\mu$ l initial serum aliquots did not reach statistical significance (64.9 [52.6–73.4] vs 59.3 [56.0–64.8]  $\mu$ g/mL;  $P = 0.525$ ,  $n = 20$  samples; Mann-Whitney Rank Sum Test). B. Effect on percent deviation of the serum GMAb concentration. Data from the experiment shown in panel A were used to calculate the percent deviation of the 20 determinations. The mean ( $\pm$ SD) percent deviation for the assay performed when using 10  $\mu$ l initial serum aliquots was markedly higher than with 100  $\mu$ l aliquots ( $17.1 \pm 8.9$  vs  $7.1 \pm 5.2$ ;  $P < 0.001$ ;  $n = 20$  samples; Student's  $t$  test).

1/12,000 and none at 1/100 (Fig. 5B–C). In healthy people, all dilutions greater than 1/100 had an out-of-range absorbance with the percent deviation increasing in proportion to the dilution (Fig. 5B) and all had an in-range or undetectable value at a dilution of 1/100 (Fig. 5C). These results indicate that any dilution can be used to determine serum GMAb concentration as long as the optical absorbance of the diluted sample is within the range of the reference standard values, or below the level of detection as in some healthy people.

The specificity of GMAb detection by the GMAb ELISA was evaluated using PAP patient serum before and after GMAb

**Fig. 5.** Effect of the choice of serum dilution on determination of GMAb concentration. A. Effect of the sample absorbance to that of the reference standard range. Separate series of standard dilutions (1/100, 1/3000, 1/6000, 1/12,000) were prepared from serum from 46 autoimmune PAP patients and 32 healthy people and GMAb concentration were measured for each dilution (184 for autoimmune PAP and 128 for healthy controls) using the GMAb ELISA. Individual measurements were defined as 'in-range' if the optical absorbance for the dilution was within the range of the optical absorbance values for the reference standards, 'low' (and out of range) if the optical absorbance was less than that of the lowest reference standard, and 'high' (and out of range) if the optical absorbance was greater than that of the highest reference standard. Only 'in-range' values were used to calculate the mean GMAb concentration. In healthy people, the percent deviation for 'in range' values was small (\*) and no measurements were greater than the highest reference standard (\*\*). Bars represent the mean ( $\pm$ SD) and the number of determinations is shown above the error bars. B. Effect of serum dilution factor on percent deviation. Data are derived from the experiments shown in panel A but shown here as function of the specimen dilution. In healthy people, the percent deviation for 'in-range' values was small at the 1/101 dilution (\*) and no determinations were in-range at higher dilutions (\*\*). Bars represent the mean ( $\pm$ SD) and the number of determinations is shown above the error bars. C. Effect of serum dilution on the frequency of obtaining 'in range' measurements. Data are derived from the experiments shown in panels A and B. The percentage of determinations that are 'in-range' for each of the standard dilutions (1/100, 1/3000, 1/6000, and 1/12,000) is shown (bars). Gray bar represent measurements that were 'in-range' and the clear bar represents measurements in which GMABs were undetectable. The cumulative percent of determinations that are 'in-range' at each successively greater dilution is shown (open symbols). In healthy people, GMAB values below the LLOQ were accepted as normal and counted as being in-range.





**Fig. 6.** Specificity of the GMAB ELISA. Serum from autoimmune PAP patients ( $n = 3$ ) was subjected to GM-CSF affinity chromatography and the unbound fraction (flow-through) and bound fraction (after elution from the column) from each was collected as previously described (Uchida et al., 2004). The GMAB concentration was measured in serum aliquots taken before chromatography and the unbound and bound fractions. All data was normalized to the serum GMAB concentration for each patient, which was set to 100%. Bars represent the mean ( $\pm$ SD) amount of GMAB present in serum (hatched bar), or recovered in the Unbound (open bar) and Bound (solid bar) fractions.

depletion by GM-CSF affinity chromatography performed as previously described (Uchida et al., 2004). GMAB was readily detected in serum before depletion, essentially undetectable in the unbound fraction (i.e., column flow-through), and approximately 60% of the serum value was detected in the bound fraction (i.e., bound to the column) recovered after elution with acidic buffer and neutralization (Fig. 6). These results indicate that the GMAB ELISA is highly specific for detection of GMAB in serum.

### 3.2. GMAB ELISA performance

The accuracy of the GMAB ELISA was evaluated using the PCRS with 'spiked' samples containing known concentrations of exogenously added, purified GMABs. The standard error for measurements of GMAB concentration in serial dilutions

**Table 1**  
Concentration response of the calibration curve and accuracy of measuring GMAB concentration in human serum using the GMAB ELISA.

Expected concentration, ng/ml	Measured concentration, ng/ml (Mean $\pm$ SD)	Coefficient of variation (100 $\times$ SD / Mean)	Standard error (%) (Mean $\pm$ SD)
<b>Calibration curve<sup>a</sup></b>			
50	51.9 $\pm$ 2.3	4.39	4.3 $\pm$ 3.9
25	24.4 $\pm$ 0.7	2.79	2.5 $\pm$ 2.7
12.5	12.2 $\pm$ 0.4	5.07	4.0 $\pm$ 3.5
6.25	5.8 $\pm$ 0.4	6.09	7.7 $\pm$ 5.2
3.125	3.1 $\pm$ 0.2	6.35	5.3 $\pm$ 2.5
1.5625	1.3 $\pm$ 0.1	10.58	14.8 $\pm$ 7.0
0.78125	0.8 $\pm$ 0.2	19.41	16.7 $\pm$ 12.1
<b>Spiked samples<sup>b</sup></b>			
10	10.1 $\pm$ 0.6	6.32	5.4 $\pm$ 2.9
20	19.0 $\pm$ 1.1	5.89	6.2 $\pm$ 4.0
40	41.8 $\pm$ 1.7	4.18	4.5 $\pm$ 4.2

<sup>a</sup> Prepared using the autoimmune PAP-patient derived PCRS by serial dilution in sample dilution buffer.

<sup>b</sup> Spiked samples were dilution buffers spiked with known concentration of purified GMAB.

of the PCRS ranging from 50 ng/ml to 1.563 ng/ml (plate concentration) was less than 15% (Table 1) in accordance with FDA guidance regarding diagnostic assay performance (Anonymous, 2001). The standard error of the measurement at a GMAB PCRS concentration of 0.78125 ng/ml in the well was  $16.7 \pm 12.1\%$  (Table 1), which is less than the limit of 20% specified by FDA guidance to define the LLOQ for the GMAB ELISA. The standard error of measurement of GMAB concentration in 'spiked' human serum samples containing 10, 20, 40 ng/ml GMAB revealed a standard error of less than 7% (Table 1) in accordance with FDA guidance criteria for assay accuracy (Anonymous, 2001).

The precision of GMAB ELISA measurements with respect to well-to-well variability, plate-to-plate variability, day-to-day variability was evaluated using the three samples from PAP patients. The coefficient of variation for measurement of GMAB concentration in patient sera was less than 15% in accordance with FDA guidance criteria for assay precision (Anonymous, 2001). The coefficient of variation for measurement of GMAB concentration in spiked samples was also less than 15% (not shown). The GMAB concentrations in PAP patient serum measured repeatedly using the GMAB ELISA by two different operators were similar (Table 2).

Repeated cycles of freezing and thawing of serum had no effect on measurement of GMAB concentration at low or high GMAB concentrations (Table 3). Storage of serum samples at room temperature for 6 h or at  $-80^\circ\text{C}$  for three months had no detectable effect on measurement of GMAB concentration (Table 3). Storage of the PCRS at room temperature for 6 h had no detectable effect on measurement of GMAB concentration (Table 3).

The optimized GMAB ELISA was used to measure serum GMAB concentration in people previously diagnosed with autoimmune PAP and in healthy, asymptomatic people. The serum GMAB concentration in patients with autoimmune PAP (94.13 [34.64–158.80]  $\mu\text{g/ml}$ ;  $n = 44$ ) was markedly higher than in healthy people (0.28 [0.20–0.51]  $\mu\text{g/ml}$ ;  $n = 38$ ) (Fig. 7A). Serum GMAB concentration was skewed towards higher concentrations in PAP patients but there was a clear separation between the two groups (Fig. 7B). ROC curve analysis in this limited group of patients and controls showed a good sensitivity and specificity for a diagnosis of autoimmune PAP and an optimal cut off value for serum GMAB of 5.0  $\mu\text{g/ml}$  (Fig. 7C, D).

### 3.3. A new GMAB reference standard as the basis for a standardized unit of measure

A new immunoglobulin G<sub>1</sub> subclass, monoclonal GMAB reference standard (MCRS) was prepared as described in the Materials and methods section and characterized as for the PCRS (Fig. 3). The MCRS had a molecular mass similar to IgG1 (Fig. 8A). Functional evaluation in the GMAB ELISA demonstrated a smooth, slightly curved increase in optical absorbance with increasing MCRS concentration over a range from 0.3125 to 20 ng/ml (Fig. 8B). Quadratic regression analysis of the optical absorbance versus MCRS concentration yielded an outstanding correlation coefficient and gave a lower percent error over the entire MCRS concentration range than did linear or logarithmic regression analysis (Fig. 8C).

**Table 2**  
Precision of measuring GMAb concentration in human serum.<sup>a</sup>

Serum samples <sup>c</sup>	Coefficient of variation <sup>a</sup>			Operator to operator <sup>b</sup> (Mean ± SD) (mcg/ml)		
	Well-to-well	Plate-to-plate	Day-to-day	Operator 1	Operator 2	P value
PAP patient 1	10.3	11.7	11.6	100.4 ± 4.5	97.8 ± 7.0	0.27
PAP patient 2	9.1	8.3	5.4	142.1 ± 12.9	138.4 ± 11.1	0.61
PAP patient 3	4.5	10.0	7.2	24.0 ± 2.5	25.5 ± 2.1	0.47

<sup>a</sup> Coefficient of variation was calculated as defined in the methods. Standard deviation × 100 / mean.

<sup>b</sup> GMAb concentration of three serum samples were evaluated by two different operators. Data are expressed as mean.

<sup>c</sup> GMAb concentration of three serum samples were evaluated to validate precision of the assay.

To determine the relationship between GMAb concentration determined using the PCRS ([GMAb]<sub>PCRS</sub>) with that determined using the MCRS ([GMAb]<sub>MCRS</sub>), standard dilutions of sera from PAP patients and healthy people were measured with both reference standards included on the same plate. [GMAb]<sub>MCRS</sub> values were ~5-fold less than corresponding [GMAb]<sub>PCRS</sub> values; the ratio varied slightly with concentration resulting in a nonlinear relationship (Fig. 9A). Regression analysis using linear, quadratic, and cubic fit equations was done to define the relationship between [GMAb]<sub>PCRS</sub> and [GMAb]<sub>MCRS</sub>. Cubic regression gave the best fit (Fig. 9A). To confirm this, [GMAb]<sub>PCRS</sub> values were converted to values equivalent to [GMAb]<sub>MCRS</sub> using each set of regression parameters just described and the percent error was determined for each measurement. Conversion using cubic equation parameters gave the lowest for percent error, which was

negligible at all GMAb concentrations >0.2 ng/ml in diluted serum samples (Fig. 9B). Thus, the relationship between [GMAb]<sub>PCRS</sub> and [GMAb]<sub>MCRS</sub> was best described as cubic polynomial. To establish the parameters needed to routinely convert [GMAb]<sub>PCRS</sub> values into values equivalent of [GMAb]<sub>MCRS</sub>, six independent experiments similar to and including the one shown above (Panel A) were evaluated simultaneously, which yielded an excellent correlation coefficient (Fig. 9C). Thus, the relationship between the GMAb concentration determined with these two reference standards is given by the following equation:

$$F(x) = 0.00002 \cdot x^3 - 0.0028 \cdot x^2 + 0.3026 \cdot x - 0.0575;$$

where x is the GMAb concentration in diluted serum in units of ng/ml determined with the PCRS (i.e., [GMAb]<sub>PCRS</sub>) and F(x) is the GMAb concentration in ng/ml determined with the MCRS (i.e., [GMAb]<sub>MCRS</sub>). To standardize reporting, one international unit (IU) of GMAb is hereby defined as a GMAb concentration equivalent to 1 µg/ml measured using the GMAb ELISA as described in the Materials and methods section with the MCRS as the reference standard.

To validate the mathematical approach for conversion of [GMAb]<sub>PCRS</sub> data to international units equivalent to values determined with the MCRS, serum GMAb was measured in 12 additional, separate autoimmune PAP patients using the PCRS (described above) and simultaneously using the MCRS. A plot of [GMAb]<sub>PCRS</sub> after conversion to IU against [GMAb]<sub>MCRS</sub> revealed a linear relationship with an outstanding correlation (Fig. 10A). Bland and Altman analysis confirmed this agreement for the two values with a difference of less than 5 IU at all concentrations below 60 IU (Fig. 10B).

To determine the range of normal and abnormal values for GMAb concentration in international units, serum GMAb in 56 autoimmune PAP patients and 38 healthy individuals were measured using the GMAb ELISA with the PCRS and converted to international units using the equation parameters identified (Fig. 11).

#### 4. Discussion

In this study, we optimized a GMAb ELISA (Schoch et al., 2002) and evaluated its accuracy, precision, and reliability for measuring serum GMAb concentration. The assay performed very well in distinguishing patients with autoimmune PAP from healthy people. A monoclonal GMAb reference standard (MCRS) was developed as the basis for a new proposed international unit of measure for GMAb concentration and a method for

**Table 3**  
Stability of samples and reagents used in GM-CSF autoantibody assay.

Condition/sample <sup>a</sup>	Concentration		P value <sup>b</sup>
	(Mean ± SD) µg/ml		
	Before	After	
Freeze–thaw (×3) <sup>c</sup>			
Low	9.398 ± 0.9	10.46 ± 0.67	0.17
High	37.20 ± 2.44	37.66 ± 1.61	0.77
Short term sample storage <sup>d</sup>			
Low	9.398 ± 0.9	9.82 ± 0.69	0.56
High	37.20 ± 2.44	38.64 ± 2.18	0.49
Long term sample storage <sup>e</sup>			
Low	9.398 ± 0.9	9.75 ± 0.05	0.54
High	37.20 ± 2.44	38.19 ± 1.10	0.55
Short term reagent storage <sup>f</sup>			
Diluted PCRS, low concentration	8.45 ± 0.56	7.82 ± 0.75	0.31
Diluted PCRS, high concentration	27.85 ± 1.21	27.79 ± 1.09	0.95

<sup>a</sup> GMAb concentration was measured in separate serum aliquots (n = 3) from two patients.


<sup>b</sup> P values were determined by comparison of before and after values using Student's *t*-test.

<sup>c</sup> GMAb concentration was measured in serum samples subjected to freezing to –80 °C and thawing at room temperature three times.

<sup>d</sup> GMAb concentration was measured in serum samples before and after keeping at temperature for 6 h.

<sup>e</sup> GMAb concentration was measured in serum samples before and after keeping in –80 °C freezer for 3 months.

<sup>f</sup> GMAb concentration was measured in aliquots of PCRS diluted to low or high concentration (indicated) before or 6 h after being kept on the counter at room temperature for 6 h.



**Fig. 7.** Measurement and ROC analysis of serum GMAb concentration in autoimmune PAP patients and healthy controls. A. Serum GMAb levels in autoimmune PAP patients and healthy people. Serum GMAb concentration was measured using the GMAb ELISA with the PCR in 44 autoimmune PAP patients and 38 healthy controls as described in the text. B. Histogram of the distribution of serum GMAb concentrations in autoimmune PAP patients and healthy people. Data represented as a frequency distribution of serum GMAb concentrations in autoimmune PAP patients (filled bars) and healthy people (hatched bars). C–D. Receiver operating characteristic (ROC) curve analysis of serum GMAb ELISA test results for 44 autoimmune PAP patients and 38 healthy people. Standard ROC characteristic analysis was performed to determine the sensitivity and specificity for the data shown in panel A. The area under the curve was 1.0 (C), and, at a cut off value for GMAb of 5.0  $\mu\text{g/ml}$  determined by the software, the sensitivity and specificity of the GMAb ELISA were both 100% (D).

converting GMAb data obtained with the PCR into international units was developed. These results facilitate the comparison of serum GMAb concentration testing obtained in different laboratories thereby facilitating research on this rare disease.

Several modifications to the original GMAb ELISA were important in improving assay performance. First, identification

of a capture antigen preparation with low baseline optical absorbance minimized non-specific background. Second, use of an increased initial serum aliquot volume to prepare specimen dilutions improved precision. Third, regression of reference standard data with a quadratic equation resulted in a better fit than with linear or logarithmic equations, which have been used previously. Fourth, restriction of absorbance


**Fig. 8.** Characterization and performance of pharmaceutical-grade, GMAb monoclonal reference standard (MCRS). A. Purity of the MCRS. The MCRS was prepared as described in the Materials and methods section. MCRS, commercially available IgG heavy chain isotype standards (IgG<sub>1</sub>, 2, 3, or 4), or molecular weight markers (MWM) were subjected to polyacrylamide gel electrophoresis under reducing conditions, Coomassie blue staining, and photography as described in the Materials and methods section. B. Optical absorbance of the MCRS as a function of concentration. The MCRS was serially diluted and evaluated as the standard in the GMAb ELISA as described in the Materials and methods section. Optical absorbance increased smoothly in proportion with MCRS concentration. Regression analysis using a quadratic equation yielded a correlation coefficient ( $R^2$ ) of 0.999. C. Effect of regression method used on percent error of the MCRS curve fit. Results from 6 independent, simultaneously conducted experiments determining the optical absorbance of serial dilutions of the MCRS were subjected to linear, quadratic, or logarithmic regression analysis and the percent deviation at each concentration was determined. The percent error of the  $[\text{GMAb}]_{\text{PCR}}$  measurement was calculated as  $[\text{GMAb}]_{\text{PCR}}$  minus  $[\text{GMAb}]_{\text{MCRS}}$  divided by  $[\text{GMAb}]_{\text{MCRS}}$  multiplied by 100; where  $[\text{GMAb}]_{\text{PCR}}$  are the unconverted values (None) or values after conversion using the linear, quadratic, or cubic regression equation parameters (indicated) and  $[\text{GMAb}]_{\text{MCRS}}$  is the value actually determined using the MCRS (assumed to be the true value). The mean ( $\pm$ SD) correlation coefficients for regression analysis of 6 separate experiments (not shown) were  $0.999 \pm 0.0004$  (quadratic),  $0.990 \pm 0.008$  ( $\log_{10}$ ), and  $0.982 \pm 0.0168$  (linear).

values used for calculating serum GMAb concentration to those within range of the reference standards improved accuracy. Further, any sample dilution with an 'in-range' absorbance value was acceptable for determining the GMAB


concentration. In contrast, some variables had little influence on GMAB ELISA performance, e.g., glycosylation of the capture antigen and use of an F(ab')<sub>2</sub> fragment or intact IgG as the anti-human IgG detection antibody.

These results help establish a basis for the routine clinical use of the GMAB ELISA for diagnosis of autoimmune PAP. First, the accuracy, precision, and reliability of the GMAB ELISA were within performance parameter guidelines established by the United States Food and Drug Administration (Anonymous, 2001). Second, the ranges of serum GMAB concentrations in autoimmune PAP patients and healthy people were similar to prior reports utilizing the GMAB ELISA with a PCRS (Trapnell et al., 2003; Uchida et al., 2004, 2007, 2009). Third, the optimal cut off value of 5 µg/ml for the upper limit of normal determined by ROC curve analysis in this study is consistent with passive immunization studies in non-human primates identifying a serum GMAB concentration of 5 µg/ml as the critical threshold above which GM-CSF bioactivity was completely neutralized and the risk of autoimmune PAP in passively immunized animals is increased (Sakagami et al., 2009, 2010). At this cut off value, the sensitivity and specificity of the assay were both 100%, which is improved compared to a prior report utilizing the assay prior to optimization (Presneill et al., 2004). This is important given the low prevalence of autoimmune PAP (Inoue et al., 2008). Notwithstanding, the present study was not designed to establish the range of normal and abnormal serum GMAB levels or the sensitivity and specificity of the GMAB ELISA for a diagnosis of autoimmune PAP. It is also necessary to establish and validate the cutoff values used to identify autoimmune PAP and to validate the use of GMAB ELISA testing and these parameters for the diagnosis of autoimmune PAP. To this end, a study designed for this purpose involving subjects from Germany, Italy, Japan, and the United States (the MICEPAP study) has been undertaken and will be subsequently reported elsewhere.

One limitation of the GMAB ELISA is that it measures both neutralizing and non-neutralizing GMABs. This should not be a problem when the total serum GMAB levels is high as in most autoimmune PAP patients but could be for values near the cutoff as can occur in some healthy people (Uchida et al., 2009). In such cases, the measurement of serum GM-CSF neutralizing capacity using a functional assay may help determine if such increases are functionally important, i.e., disrupt GM-CSF signaling. Several cell-based assays are useful for this including inhibition of GM-CSF-dependent TF1 cell growth (Uchida et al., 2004, 2009) or the GM-CSF-stimulated increase in whole blood leukocyte surface CD11b levels (Uchida et al., 2007) or GM-CSF stimulated increase in signal inducer of transcription 5 (STAT5) phosphorylation in blood leukocytes (Suzuki et al., 2008). Another limitation relates to small increases in serum GMAB observed in diseases not associated with development of PAP (Bendtzen et al., 2007; Han et al., 2009). For example, in 272 pediatric and 88 adult patients with Crohn's disease who did not have PAP, the median serum GMAB concentrations were 2.4 and 11.7 µg/ml, respectively (Han et al., 2009). Functional testing was helpful and indicated that GM-CSF signaling was reduced but not abolished in these patients. Since the clinical symptoms of autoimmune PAP do not occur in patients without significant radiographic findings, combining GMAB testing with routine








chest computed tomography will likely resolve any discrepancy potentially arising from intermediate GMAB values close to the cutoff. Further, in a typical clinical setting, GMAB testing would likely be considered after radiographic evaluation had suggested a diagnosis of PAP. Another limitation is the GMAB ELISA only detects free GMAB and not GMAB bound to GM-CSF, which could underestimate the GMAB concentration at the very low levels in individuals without PAP. This is illustrated by considering the following: 1) up to 7.8 GMAB molecules can bind to each GM-CSF molecule (Uchida et al., 2004); 2) more than 99% of serum GM-CSF is bound to GMAB in healthy people and PAP patients (Uchida et al., 2009); 3) total serum GM-CSF, i.e., unbound and GMAB-bound GM-CSF, is ~3048 pg/ml in healthy people and ~2360 in aPAP patients (Uchida et al., 2009); 4) the serum GMAB level in healthy people was 280 ng/ml (this report). Assuming 7.8 GMAB molecules bind to each GM-CSF molecule and the molecular mass for GMAB is ten times that of GM-CSF, the amount of GMAB bound to GM-CSF would be 7.8 multiplied by 3048 multiplied 10, or ~237 ng/ml. Thus, in healthy people, the percentage of GMAB detected by the ELISA would equal unbound GMAB (280 ng/ml) divided by total GMAB (unbound and GM-CSF-bound; 280 + 237 ng/ml) multiplied by 100 (conversion to percent) or approximately 54% of total GMAB. In autoimmune PAP patients, by similar calculations, ~99.8% of GMAB would be unbound and therefore detected by the GMAB ELISA. From above calculations, it is anticipated that GM-CSF bound to GMAB minimally interfere with the diagnostic threshold. Further, the lower limit of detection (0.78 ng/ml) is well below the cutoff for a positive result (5 µg/ml).


The observation that optical absorbance was greater for the MCRS than for the PCRS can be interpreted in terms of

**Fig. 9.** Conversion of GMAB concentration data into International units. **A.** Comparison of Serum GMAB concentration determined with the PCRS and MCRS. Separate sets of standard dilutions (1/100, 1/3000, 1/6000, 1/12,000) were prepared from serum from 14 autoimmune PAP patients and 1 healthy person. All standard dilutions (60 total) were used to measure GMAB concentration with the PCRS ( $[GMAB]_{PCRS}$ ) and MCRS ( $[GMAB]_{MCRS}$ ) as described in the legends in Figs. 3 and 7, respectively, in parallel on the same plate. All in-range values ( $n = 41$ ; defined in the legend to Fig. 5) are shown and were used for analysis. The relationship between GMAB concentration determined using each of the two reference standards was evaluated by linear, quadratic (not shown to improve readability), or cubic regression analysis. The correlation coefficients for regression analysis were 0.980 (linear), 0.998 (quadratic), and 1.00 (cubic). **B.** Effect of regression method and GMAB concentration range on the accuracy of conversion of  $[GMAB]_{PCRS}$  to international units. Using the data and regression equation parameters determined above (panel A), the values for GMAB concentration determined with the PCRS ( $[GMAB]_{PCRS}$ ) were converted mathematically to values that would have been obtained if the MCRS had been used (i.e., equivalent to  $[GMAB]_{MCRS}$ ). The percent error of the  $[GMAB]_{PCRS}$  measurement was calculated as  $[GMAB]_{PCRS}$  minus  $[GMAB]_{MCRS}$  divided by  $[GMAB]_{MCRS}$  multiplied by 100; where  $[GMAB]_{PCRS}$  are the unconverted values (None) or values after conversion using the linear, quadratic, or cubic regression equation parameters (indicated) and  $[GMAB]_{MCRS}$  is the value actually determined using the MCRS (assumed to be the true value). The dashed line represents a percent error of zero. **C.** Determining the equation to convert  $[GMAB]_{PCRS}$  data to international units. Data from six independent experiments similar to and including the one shown in panel A were combined and cubic regression analysis was done to determine the equation needed to convert data for GMAB concentration determined using the PCRS into units equivalent to that determined using the MCRS. In total, serum samples from 55 autoimmune PAP patients and 12 healthy controls were used for determining the equation. The correlation coefficient ( $R^2$ ) was 0.98343.



**Fig. 10.** Validation of the equation to convert  $[GMAb]_{PCRS}$  data to international units. A. Comparison of  $[GMAb]_{PCRS}$  data before and after conversion to international units with data determined directly with the MCRS. The converted PCRS-based data described above was plotted against the MCRS-based data and evaluated by linear regression analysis. The correlation coefficient was 0.984. See text for further details. B. Bland and Altman analysis. The data shown in panel A were used to evaluate the error for the comparison between  $[GMAb]_{PCRS}$  converted to international units and  $[GMAb]_{MCRS}$  by Bland and Altman analysis as described in the Materials and methods section.

epitope differences: the PCRS has polyclonal epitopes distributed throughout the entire GM-CSF molecule while the MCRS targets a single epitope. In the GMAb ELISA, a lower optical absorbance indicates reduced secondary (detection) antibody binding, which could be caused by either 1) reduced binding of GMAb to capture antigen or 2) reduced binding of the detection antibody to captured GMAb (i.e., fewer 'targets for detection or reduced efficiency of target detection). Assuming that GMAb capture has reached equilibrium, theoretically, reduced binding of PCRS in the GMAb ELISA (compared to MCRS) could be explained by 'epitope hiding', perhaps of less-avid GMAb species, or by reduced access to GM-CSF



**Fig. 11.** Normal and abnormal ranges for GMAb concentration expressed in units of the PCRS, MCRS, and international units. Serum GMAb concentration in 56 autoimmune PAP patients and 38 healthy people was measured using the GMAb ELISA with the PCRS and converted into international units using the equation parameters determined by the analysis shown in Fig. 9C and was also simultaneously measured with the MCRS on the same plate. Box plots show the median (line) and IQR (box top and bottom), and 5th, 95 CI (circles) for patients with autoimmune 56 PAP patients (aPAP), and 38 healthy people (HP).

adjacent to bulkier complexes of polyvalent PCRS. Neither of these mechanisms seems likely since capture antigen is in molar excess of GMAb. Alternatively, since the binding ratio can be as high as 7.8 molecules of GMAb per GM-CSF molecule for the PCRS and only one for MCRS, reduced binding of the detection antibody could be explained by increased steric hindrance for the polyvalent PCRS–antigen complexes than for monovalent MCRS–antigen complexes. While we believe the latter is most likely, these mechanisms are not mutually exclusive and additional studies would be needed to elucidate the mechanism.

The availability of a high-quality, monoclonal GMAb reference standard (e.g., MCRS) should facilitate research on autoimmune PAP, which is challenged by low prevalence like many rare diseases. Specifically, the use of a common international unit of measure for GMAb concentration based on the MCRS would improve comparability of test results from different laboratories. To this end, efforts are underway to establish the GMAb ELISA in multiple reference laboratories in Japan, Europe, Asia, and the United States. The MCRS and a common set of serum specimens will be used to calibrate serum GMAb test results between laboratories. Conceivably, these reagents could also allow GMAb data obtained by other methods, e.g., serial dilution GMAb titer data (Bonfield et al., 2002), to be expressed in a common unit of measure. Finally, the availability of a validated method for converting data obtained with other GMAb reference standards to international units may facilitate meta-analysis of published data from small cases series, which are common in rare diseases.

#### Acknowledgments

We thank Barbara Enenkel, Boehringer Ingelheim for producing the MCRS (BI1049904), Barbara Kistler, PhD for helpful discussions during the course of this work and

writing of the manuscript, and Carrie Stevens for help with collection of human blood specimens. This work was supported by grants from the National Heart Lung and Blood Institute (R01 HL085453; B.C.T.), National Center for Research Resources (U54 RR0198498, B.C.T.), the National Institute for Health and Human Development, Japan Society for the Promotion of Science (Category B International, 19390403, Y.I. and K.U.) and from the Japanese Ministry of Health, Labor, and Welfare (H24-Nanchitou-ippan-035, Y.I. and K.N.).

## References

- Anonymous, 2001. Guidance for industry bioanalytical method validation. Guidance Documents, Vol. 2012. U.S. Department of Health and Human Services, Food and Drug Administration, Center for Drug Evaluation and Research (CDER), Center for Veterinary Medicine (CVM), Bethesda.
- Bendtsen, K., Svenson, M., Hansen, M.B., Busch, T., Bercker, S., Kaisers, U., Uchida, K., Beck, D.C., Trapnell, B.C., 2007. GM-CSF autoantibodies in pulmonary alveolar proteinosis. *N. Engl. J. Med.* 356, 2001.
- Bonfield, T.L., Russell, D., Burgess, S., Malur, A., Kavuru, M.S., Thomassen, M.J., 2002. Autoantibodies against granulocyte macrophage colony-stimulating factor are diagnostic for pulmonary alveolar proteinosis. *Am. J. Respir. Cell Mol. Biol.* 27, 481.
- Bonfield, T.L., Raychaudhuri, B., Malur, A., Abraham, S., Trapnell, B.C., Kavuru, M.S., Thomassen, M.J., 2003. PU.1 regulation of human alveolar macrophage differentiation requires granulocyte-macrophage colony-stimulating factor. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 285, L1132.
- Han, X., Uchida, K., Jurickova, I., Koch, D., Willson, T., Samson, C., Bonkowski, E., Trauernicht, A., Kim, M.O., Tomer, G., Dubinsky, M., Plevy, S., Kugathasan, S., Trapnell, B.C., Denson, L.A., 2009. Granulocyte-macrophage colony-stimulating factor autoantibodies in murine ileitis and progressive ileal Crohn's disease. *Gastroenterology* 136 (1261–71), e1.
- Ikegami, M., Ueda, T., Hull, W., Whitsett, J.A., Mulligan, R.C., Dranoff, G., Jobe, A.H., 1996. Surfactant metabolism in transgenic mice after granulocyte macrophage-colony stimulating factor ablation. *Am. J. Physiol.* 270, L650.
- Inoue, Y., Trapnell, B.C., Tazawa, R., Arai, T., Takada, T., Hizawa, N., Kasahara, Y., Tatsumi, K., Hojo, M., Ichihata, T., Tanaka, N., Yamaguchi, E., Eda, R., Oishi, K., Tsuchihashi, Y., Kaneko, C., Nukiwa, T., Sakatani, M., Krischer, J.P., Nakata, K., 2008. Characteristics of a large cohort of patients with autoimmune pulmonary alveolar proteinosis in Japan. *Am. J. Respir. Crit. Care Med.* 177, 752.
- Kitamura, T., Tanaka, N., Watanabe, J., Uchida, K., Kanegasaki, S., Yamada, Y., Nakata, K., 1999. Idiopathic pulmonary alveolar proteinosis as an autoimmune disease with neutralizing antibody against granulocyte/macrophage colony-stimulating factor. *J. Exp. Med.* 190, 875.
- Kitamura, T., Uchida, K., Tanaka, N., Tsuchiya, T., Watanabe, J., Yamada, Y., Hanaoka, K., Seymour, J.F., Schoch, O.D., Doyle, I., Inoue, Y., Sakatani, M., Kudoh, S., Azuma, A., Nukiwa, T., Tomita, T., Katagiri, M., Fujita, A., Kurashima, A., Kanegasaki, S., Nakata, K., 2000. Serological diagnosis of idiopathic pulmonary alveolar proteinosis. *Am. J. Respir. Crit. Care Med.* 162, 658.
- Luisetti, M., Rodi, G., Perotti, C., Campo, J., Mariani, F., Pozzi, E., Trapnell, B.C., 2009. Plasmapheresis for treatment of pulmonary alveolar proteinosis. *Eur. Respir. J.* 33, 1220.
- Martinez-Moczygemba, M., Doan, M.L., Elidemir, O., Fan, L.L., Cheung, S.W., Lei, J.T., Moore, J.P., Tavana, G., Lewis, L.R., Zhu, Y., Muzny, D.M., Gibbs, R.A., Huston, D.P., 2008. Pulmonary alveolar proteinosis caused by deletion of the GM-CSFRalpha gene in the X chromosome pseudoautosomal region 1. *J. Exp. Med.* 205, 2711.
- Nakata, K., 1999. Anti-GM-CSF Autoantibodies and Reagents Required for Its Evaluation. In: J.P. Office (Ed.) Japan Patent Office, Number 4372904, Japan.
- Nogee, L.M., 2010. Genetic basis of children's interstitial lung disease. *Pediatr. Allergy Immunol. Pulmonol.* 23, 15.
- Presneill, J.J., Nakata, K., Inoue, Y., Seymour, J.F., 2004. Pulmonary alveolar proteinosis. *Clin. Chest Med.* 25, 593 (viii).
- Sakagami, T., Uchida, K., Suzuki, T., Carey, B.C., Wood, R.E., Wert, S.E., Whitsett, J.A., Trapnell, B.C., Luisetti, M., 2009. Human GM-CSF autoantibodies and reproduction of pulmonary alveolar proteinosis. *N. Engl. J. Med.* 361, 2679.
- Sakagami, T., Beck, D., Uchida, K., Suzuki, T., Carey, B.C., Nakata, K., Keller, G., Wood, R.E., Wert, S.E., Ikegami, M., Whitsett, J.A., Luisetti, M., Davies, S., Krischer, J.P., Brody, A., Ryckman, F., Trapnell, B.C., 2010. Patient-derived GM-CSF autoantibodies reproduce pulmonary alveolar proteinosis in non-human primates. *Am. J. Respir. Crit. Care Med.* 182, 49.
- Schoch, O.D., Schanz, U., Koller, M., Nakata, K., Seymour, J.F., Russi, E.W., Boehler, A., 2002. BAL findings in a patient with pulmonary alveolar proteinosis successfully treated with GM-CSF. *Thorax* 57, 277.
- Shibata, Y., Berclaz, P.Y., Chroneos, Z.C., Yoshida, M., Whitsett, J.A., Trapnell, B.C., 2001. GM-CSF regulates alveolar macrophage differentiation and innate immunity in the lung through PU.1. *Immunity* 15, 557.
- Suzuki, T., Sakagami, T., Rubin, B.K., Nogee, L.M., Wood, R.E., Zimmerman, S.L., Smolarek, T., Dishop, M.K., Wert, S.E., Whitsett, J.A., Grabowski, G., Carey, B.C., Stevens, C., van der Loo, J.C., Trapnell, B.C., 2008. Familial pulmonary alveolar proteinosis caused by mutations in CSF2RA. *J. Exp. Med.* 205, 2703.
- Suzuki, T., Sakagami, T., Young, L.R., Carey, B.C., Wood, R.E., Luisetti, M., Wert, S.E., Rubin, B.K., Kevill, K., Chalk, C., Whitsett, J.A., Stevens, C., Nogee, L.M., Campo, J., Trapnell, B.C., 2010. Hereditary pulmonary alveolar proteinosis: pathogenesis, presentation, diagnosis, and therapy. *Am. J. Respir. Crit. Care Med.* 182, 1292.
- Suzuki, T., Maranda, B., Sakagami, T., Catellier, P., Couture, C., Carey, B., Chalk, C., Trapnell, B., 2011. Hereditary pulmonary alveolar proteinosis caused by CSF2RB mutations. *Eur. Respir. J.* 37, 201.
- Tanaka, T., Motoi, N., Tsuchihashi, Y., Tazawa, R., Kaneko, C., Nei, T., Yamamoto, T., Hayashi, T., Tagawa, T., Nagayasu, T., Kuribayashi, F., Ariyoshi, K., Nakata, K., Morimoto, K., 2011. Adult-onset hereditary pulmonary alveolar proteinosis caused by a single-base deletion in CSF2RB. *J. Med. Genet.* 48, 205.
- Trapnell, B.C., Whitsett, J.A., Nakata, K., 2003. Pulmonary alveolar proteinosis. *N. Engl. J. Med.* 349, 2527.
- Uchida, K., Nakata, K., Trapnell, B.C., Terakawa, T., Hamano, E., Mikami, A., Matsushita, I., Seymour, J.F., Oh-Eda, M., Ishige, I., Eishi, Y., Kitamura, T., Yamada, Y., Hanaoka, K., Keicho, N., 2004. High-affinity autoantibodies specifically eliminate granulocyte-macrophage colony-stimulating factor activity in the lungs of patients with idiopathic pulmonary alveolar proteinosis. *Blood* 103, 1089.
- Uchida, K., Beck, D.C., Yamamoto, T., Berclaz, P.Y., Abe, S., Staudt, M.K., Carey, B.C., Filippi, M.D., Wert, S.E., Denson, L.A., Puchalski, J.T., Hauck, D.M., Trapnell, B.C., 2007. GM-CSF autoantibodies and neutrophil dysfunction in pulmonary alveolar proteinosis. *N. Engl. J. Med.* 356, 567.
- Uchida, K., Nakata, K., Suzuki, T., Luisetti, M., Watanabe, M., Koch, D.E., Stevens, C.A., Beck, D.C., Denson, L.A., Carey, B.C., Keicho, N., Krischer, J.P., Yamada, Y., Trapnell, B.C., 2009. Granulocyte/macrophage-colony-stimulating factor autoantibodies and myeloid cell immune functions in healthy subjects. *Blood* 113, 2547.
- Urlaub, G., Kas, E., Carothers, A.M., Chasin, L.A., 1983. Deletion of the diploid dihydrofolate reductase locus from cultured mammalian cells. *Cell* 33, 405.

## ORIGINAL ARTICLE

## CYFRA 21-1 as a disease severity marker for autoimmune pulmonary alveolar proteinosis

TORU ARAI,<sup>1,2</sup> YOSHIKAZU INOUE,<sup>2</sup> CHIKATOSHI SUGIMOTO,<sup>3</sup> YASUSHI INOUE,<sup>4</sup> KEIKO NAKAO,<sup>4</sup> NAOKO TAKEUCHI,<sup>4</sup> AKIKO MATSUMURO,<sup>2</sup> MASAKI HIROSE,<sup>2</sup> KOH NAKATA<sup>5</sup> AND SEIJI HAYASHI<sup>4</sup>

<sup>1</sup>Department of Respiratory Medicine, <sup>2</sup>Department of Diffuse Lung Diseases and Respiratory Failure, <sup>3</sup>Division of Clinical Trial, <sup>4</sup>Department of Internal Medicine, National Hospital Organization, Kinki-Chuo Chest Medical Center, Osaka and <sup>5</sup>Bioscience Medical Research Center, Niigata University Medical and Dental Hospital, Niigata, Japan

### ABSTRACT

**Background and objective:** Serum markers, including Krebs von den Lungen (KL-6), surfactant protein (SP)-D, SP-A and carcinoembryonic antigen (CEA), are reported to reflect autoimmune pulmonary alveolar proteinosis (APAP) disease severity. We evaluated serum CYFRA21-1 levels as a marker of APAP.

**Methods:** In addition to KL-6, SP-D and CEA, we prospectively measured serum CYFRA 21-1 levels in 48 patients with APAP, consecutively diagnosed between 2002 and 2010. Diagnostic usefulness of CYFRA 21-1 was determined from 68 patients with interstitial lung diseases by receiver operator characteristic curve analysis. We evaluated the association between these serum markers and other disease severity markers, including pulmonary function parameters, alveolar-arterial oxygen gradient, British Medical Research Council score reflecting shortness of breath, and disease severity score. CYFRA 21-1 localization in the lung was examined by immunohistochemistry.

**Results:** Receiver operator characteristic curve demonstrated that CYFRA 21-1 effectively identified APAP. Serum CYFRA 21-1 levels at diagnosis were significantly associated with the measured disease severity parameters. Following whole lung lavage ( $n = 10$ ) and granulocyte-macrophage colony-stimulating factor (GM-CSF) inhalation ( $n = 20$ ), serum CYFRA 21-1 levels were significantly decreased. Responders ( $n = 11$ ) to GM-CSF inhalation revealed significantly higher serum CYFRA 21-1 levels than non-responders ( $n = 9$ ). Serum CYFRA 21-1 appeared to be a significant predictor of effectiveness of GM-CSF based on regression analysis. Immunohistochemistry showed that CYFRA 21-1 was localized on hyperplastic alveolar type II cells and lipoproteinaceous substances in alveoli.

**Conclusions:** Serum CYFRA 21-1 is a sensitive and useful serum marker for diagnosis and evaluation of disease severity of APAP, and may predict the response to GM-CSF inhalation.

### SUMMARY AT A GLANCE

Serum CYFRA 21-1 is useful for the diagnosis of autoimmune pulmonary alveolar proteinosis (APAP) by receiver operating characteristic curve analysis and significantly correlated with other disease severity markers of APAP. Serum CYFRA 21-1 might be a predictor of responsiveness to GM-CSF inhalation therapy in APAP.

**Key words:** alveolar epithelial hyperplasia, granulocyte-macrophage colony-stimulating factor, inhalation therapy, rare lung disease, whole lung lavage.

**Abbreviations:** APAP, autoimmune pulmonary alveolar proteinosis; CEA, carcinoembryonic antigen; GM-CSF, granulocyte-macrophage colony-stimulating factor; KL-6, Krebs von den Lungen; PaO<sub>2</sub>, partial pressure of oxygen; SP, surfactant protein; WLL, whole lung lavage.

### INTRODUCTION

Pulmonary alveolar proteinosis is characterized by the accumulation of phospholipids and surfactant protein (SP) in alveolar spaces.<sup>1</sup> Impaired alveolar macrophage function underlies the pathogenesis of idiopathic pulmonary alveolar proteinosis, which is caused by a deficiency of the granulocyte-macrophage colony-stimulating factor (GM-CSF) in the presence of anti-GM-CSF autoantibodies.<sup>2,3</sup> Hence, in recent years, idiopathic pulmonary alveolar proteinosis has been referred to as autoimmune pulmonary alveolar proteinosis (APAP) when anti-GM-CSF autoantibodies are detected in the blood.<sup>4</sup> The levels of SP-A,<sup>5</sup> SP-B,<sup>6</sup> SP-D,<sup>7</sup> carcinoembryonic antigen (CEA),<sup>8</sup> Krebs von den Lungen (KL-6)<sup>9</sup> and lactate dehydrogenase<sup>10</sup> are elevated in the sera of idiopathic pulmonary alveolar proteinosis patients. Previously, Seymour and Presneill showed that SP-A and lactate dehydrogenase levels correlated with idiopathic pulmonary alveolar proteinosis disease severity.<sup>10</sup> Moreover, the levels of CEA<sup>11</sup> and KL-6<sup>12</sup> in

Correspondence: Yoshikazu Inoue, National Hospital Organization Kinki-Chuo Chest Medical Center, 1180 Nagasone-Cho, Kita-Ku, Sakai, Osaka 591-8555, Japan. Email: giichi@kch.hosp.go.jp

Received 27 November 2012; invited to revise 17 May and 1 August 2012; revised 3 June and 9 August 2013; accepted 28 August 2013 (Associate Editor: Yuben Moodley).