

Pfizer, Eli Lilly, and JCR Pharmaceuticals and has received speaker's fees from Chugai Pharmaceutical. All other authors state that they have no conflicts of interest.

Acknowledgments

We thank the patients and their family members who participated in this study. We also thank Minako Takaki and Reiko Onai for technical support, and Mitsuko Itoh for kind assistance in English usage. This study was supported by a Grant-in-Aid from the Ministry of Education, Science, Sports, and Culture of Japan.

Authors' roles: Study design: TI and SK. Patients' sample and clinical data collection: YO, ET, AY, TY, and RH. Study conduct and data collection: TI, JM, and HI. Exome sequencing data analysis: JM, KD, JY, HI, SM, and ST. Data interpretation: TI, JM, KD, SM, ST, and SK. Study support and intellectual input: SM, ST, and SK. Drafting manuscript: TI. Revising manuscript content: SM, ST, and SK. Approving final version of manuscript: all authors. TI, SM, and SK take responsibility for the integrity of the data analysis.

References

1. Kenny FM, Linarelli L. Dwarfism and cortical thickening of tubular bones. Transient hypocalcemia in a mother and son. *Am J Dis Child.* 1966;111(2):201–7.
2. Caffey J. Congenital stenosis of medullary spaces in tubular bones and calvaria in two proportionate dwarfs—mother and son; coupled with transitory hypocalcemic tetany. *Am J Roentgenol Radium Ther Nucl Med.* 1967;100(1):1–11.
3. Fanconi S, Fischer JA, Wieland P, et al. Kenny syndrome: evidence for idiopathic hypoparathyroidism in two patients and for abnormal parathyroid hormone in one. *J Pediatr.* 1986;109(3):469–75.
4. Lee WK, Vargas A, Barnes J, Root AW. The Kenny-Caffey syndrome: growth retardation and hypocalcemia in a young boy. *Am J Med Genet.* 1983;14(4):773–82.
5. Franceschini P, Testa A, Bogetti G, et al. Kenny-Caffey syndrome in two sibs born to consanguineous parents: evidence for an autosomal recessive variant. *Am J Med Genet.* 1992;42(1):112–6.
6. Sabry MA, Zaki M, Abul Hassan SJ, et al. Kenny-Caffey syndrome is part of the CATCH 22 haploinsufficiency cluster. *J Med Genet.* 1998;35(1):31–6.
7. Sabry MA, Farag TI, Shaltout AA, et al. Kenny-Caffey syndrome: an Arab variant? *Clin Genet.* 1999;55(1):44–9.
8. Parvari R, Hershkovitz E, Grossman N, et al. Mutation of TBCE causes hypoparathyroidism-retardation-dysmorphism and autosomal recessive Kenny-Caffey syndrome. *Nat Genet.* 2002;32(3):448–52.
9. Tadaki H, Tokuhiko E, Shiga K, Kikuchi N, Mukai N, Fujieda K. A case of a 2-year-old girl with Kenny-Caffey Syndrome Type 2. *Clin Pediatr Endocrinol.* 2005;14(2 Suppl):22.
10. Oda Y, Ono R, Hiwatari M, Iwasaki H, Namai Y, Iimori Y. A case report: three-year-old boy of Kenny-Caffey Syndrome Type 2. *Clin Pediatr Endocrinol.* 2000;9(2):140.
11. Yorifuji T, Muroi J, Uematsu A. Kenny-Caffey syndrome without the CATCH 22 deletion. *J Med Genet.* 1998;35(12):1054.
12. Izumi Y, Tanae A, Kuratsuji T, et al. A case of 12-year-old boy with Kenny syndrome associated with hypomagnesemia and humoral immunodeficiency [in Japanese]. *Shoninaika.* 1987;19(10):1503–2.
13. Unger S, Gořna MW, Le Bécéc A, et al. FAM111A mutations result in hypoparathyroidism and impaired skeletal development. *Am J Hum Genet.* 2013 May 14. [Epub ahead of print].
14. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics.* 2009;25(14):1754–60.
15. Li H, Handsaker B, Wysoker A, et al. The sequence alignment/map format and SAMtools. *Bioinformatics.* 2009;25(16):2078–9.
16. Kong A, Frigge ML, Masson G, et al. Rate of de novo mutations and the importance of father's age to disease risk. *Nature.* 2012;488(7412):471–5.
17. Fine DA, Rozenblatt-Rosen O, Padi M, et al. Identification of FAM111A as an SV40 host range restriction and adenovirus helper factor. *PLoS Pathog.* 2012;8(10):e1002949.
18. Akamatsu S, Takata R, Haiman CA, et al. Common variants at 11q12, 10q26 and 3p11.2 are associated with prostate cancer susceptibility in Japanese. *Nat Genet.* 2012;44(4):426–9, S421.
19. Shiang R, Thompson LM, Zhu YZ, et al. Mutations in the transmembrane domain of FGFR3 cause the most common genetic form of dwarfism, achondroplasia. *Cell.* 1994;78(2):335–42.
20. Weinstein LS, Shenker A, Gejman PV, Merino MJ, Friedman E, Spiegel AM. Activating mutations of the stimulatory G protein in the McCune-Albright syndrome. *N Engl J Med.* 1991;325(24):1688–95.
21. Gensure RC, Mäkitie O, Barclay C, Chan C, Depalma SR, Bastepe M, Abuzahra H, Couper R, Mundlos S, Sillence D, Ala Kokko L, Seidman JG, Cole WG, Jüppner H. A novel COL1A1 mutation in infantile cortical hyperostosis (Caffey disease) expands the spectrum of collagen-related disorders. *J Clin Invest.* 2005;115(5):1250–7.
22. Courtens W, Wuys W, Poot M, Szuhai K, Wauters J, Reyniers E, Eleveld M, Diaz G, Nöthen MM, Parvari R. Hypoparathyroidism-retardation-dysmorphism syndrome in a girl: A new variant not caused by a TBCE mutation—clinical report and review. *Am J Med Genet A.* 2006;140(6):611–7.
23. Shoback D. Clinical practice. Hypoparathyroidism. *N Engl J Med.* 2008;359(4):391–403.
24. Zajac JD, Danks JA. The development of the parathyroid gland: from fish to human. *Curr Opin Nephrol Hypertens.* 2008;17(4):353–6.
25. Boynton JR, Pheasant TR, Johnson BL, Levin DB, Streeten BW. Ocular findings in Kenny's syndrome. *Arch Ophthalmol.* 1979;97(5):896–00.
26. Parvari R, Diaz GA, Hershkovitz E. Parathyroid development and the role of tubulin chaperone E. *Horm Res.* 2007;67(1):12–21.
27. Wilson MG, Maronde RF, Mikity VG, Shinno NW. Dwarfism and congenital medullary stenosis (Kenny syndrome). *Birth Defects Orig Artic Ser.* 1974;10(12):128–32.

