

uses limited resources effectively to control NTDs and other infectious diseases in Africa.

## Supporting Information

**Figure S1 Sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) analysis of purified antigens.**

(PDF)

**Figure S2 Assay stability after coupling microspheres with antigens.**

(PDF)

**Table S1 BIC results of mixture models.**

(PDF)

**Table S2 Sensitivities and specificities in positive and negative controls using cut-off values calculated by the finite mixture model-based MFI value distributions in population-based serological survey.**

(PDF)

**Table S3 Seroprevalence by site, sex, and age group for pathogens measured simultaneously using a micro-**

**sphere-based multi-serological assay on samples from the Kwale site.**

(PDF)

**Table S4 Seroprevalence by site, sex, and age group for pathogens measured simultaneously using a microsphere-based multi-serological assay on samples from the Mbita site.**

(PDF)

## Acknowledgments

We thank all residents in the Kwale and Mbita HDSS sites and the health workers who participated in the multiplex project. We also are grateful to Mr. Tetsuo Yanagi for support and suggestions during the development of antigen preparation. This paper is published with the permission of the Director of KEMRI.

## Author Contributions

Conceived and designed the experiments: YF SK SMNz MMw MK KH. Performed the experiments: YF SMNz MMw CT JK AWM MMi. Analyzed the data: YF SK SMNz TH KK. Contributed reagents/materials/analysis tools: YF SMNz MMw SMNj CT JK IK SM MN MOO YI MIn MIt HT KI TT LMY DM RH SH MC MK MMi. Wrote the paper: YF SK SMNz.

## References

- Hotez PJ, Kamath A (2009) Neglected tropical diseases in sub-Saharan Africa: review of their prevalence, distribution, and disease burden. *PLoS Negl Trop Dis* 3: e412.
- Department of Control of Neglected Tropical Diseases (2006) Neglected Tropical Diseases. Geneva: World Health Organization.
- Hotez PJ, Mistry N, Rubinstein J, Sachs JD (2011) Integrating neglected tropical diseases into AIDS, tuberculosis, and malaria control. *N Engl J Med* 364: 2086–2089.
- Solomon AW, Engels D, Bailey RL, Blake IM, Brooker S, et al. (2012) A diagnostics platform for the integrated mapping, monitoring, and surveillance of neglected tropical diseases: rationale and target product profiles. *PLoS Negl Trop Dis* 6: e1746.
- Vignali DA (2000) Multiplexed particle-based flow cytometric assays. *J Immunol Methods* 243: 243–255.
- Khan IH, Ravindran R, Yee J, Ziman M, Lewinsohn DM, et al. (2008) Profiling antibodies to *Mycobacterium tuberculosis* by multiplex microbead suspension arrays for serodiagnosis of tuberculosis. *Clin Vaccine Immunol* 15: 433–438.
- Go YY, Wong SJ, Branscum AJ, Demarest VL, Shuck KM, et al. (2008) Development of a fluorescent-microsphere immunoassay for detection of antibodies specific to equine arteritis virus and comparison with the virus neutralization test. *Clin Vaccine Immunol* 15: 76–87.
- Clavijo A, Hole K, Li M, Collignon B (2006) Simultaneous detection of antibodies to foot-and-mouth disease non-structural proteins 3ABC, 3D, 3A and 3B by a multiplexed Luminex assay to differentiate infected from vaccinated cattle. *Vaccine* 24: 1693–1704.
- Watson DS, Reddy SM, Brahmakshatriya V, Lupiani B (2009) A multiplexed immunoassay for detection of antibodies against avian influenza virus. *J Immunol Methods* 340: 123–131.
- Moss DM, Priest JW, Boyd A, Weinkopff T, Kucrova Z, et al. (2011) Multiplex bead assay for serum samples from children in Haiti enrolled in a drug study for the treatment of lymphatic filariasis. *Am J Trop Med Hyg* 85: 229–237.
- Griffin SM, Chen IM, Fout GS, Wade TJ, Egorov AI (2011) Development of a multiplex microsphere immunoassay for the quantitation of salivary antibody responses to selected waterborne pathogens. *J Immunol Methods* 364: 83–93.
- Tachibana H, Cheng XJ, Masuda G, Horiki N, Takeuchi T (2004) Evaluation of recombinant fragments of *Entamoeba histolytica* Gal/GalNAc lectin intermediate subunit for serodiagnosis of amoebiasis. *J Clin Microbiol* 42: 1069–1074.
- Takagi H, Islam MZ, Itoh M, Islam AU, Saifuddin Ekram AR, et al. (2007) Short report: production of recombinant kinesin-related protein of *Leishmania donovani* and its application in the serodiagnosis of visceral leishmaniasis. *Am J Trop Med Hyg* 76: 902–905.
- Ishii K, Hisaeda H, Duan X, Imai T, Sakai T, et al. (2006) The involvement of immunoproteasomes in induction of MHC class I-restricted immunity targeting *T* oxoplasma SAG1. *Microbes and Infection/Institut Pasteur* 8: 1045–1053.
- Dissanayake S, Zheng H, Dreyer G, Xu M, Watawana L, et al. (1994) Evaluation of a recombinant parasite antigen for the diagnosis of lymphatic filariasis. *Am J Trop Med Hyg* 50: 727–734.
- Garnett JW, Jr., Schwirumbeck PL, Nelson JA, Truax AB, Oldstone MB (1987) Diagnosis of AIDS by using a 12-amino acid peptide representing an immunodominant epitope of the human immunodeficiency virus. *J Infect Dis* 156: 261–267.
- Kulshrestha A, Gupta A, Verma N, Sharma SK, Tyagi AK, et al. (2005) Expression and purification of recombinant antigens of *Mycobacterium tuberculosis* for application in serodiagnosis. *Protein Expr Purif* 44: 75–85.
- Waterboer T, Schr P, Pawlita M (2006) Suppression of non-specific binding in serological Luminex assays. *J Immunol Methods* 309: 200–204.
- Pickering JW, Larson MT, Martins TB, Copple SS, Hill HR (2010) Elimination of false-positive results in a luminex assay for pneumococcal antibodies. *Clin Vaccine Immunol* 17: 185–189.
- Infectious Disease Surveillance Center, National Institute of Infectious Diseases, Japan (2012) Annual Surveillance Data: Notifiable Diseases. Available: <http://idsc.nih.go.jp/idwr/ydata/report-Ea.html>. Accessed 17 April 2014.
- Kimura E, Itoh M (2011) Filariasis in Japan some 25 years after its eradication. *Trop Med Health* 39: 57–63.
- Nakamura-Uchiyama F, Hiromatsu K, Ishiwata K, Sakamoto Y, Nawa Y (2003) The current status of parasitic diseases in Japan. *Intern Med* 42: 222–236.
- Otsubo H, Yamaguchi K (2008) Current risks in blood transfusion in Japan. *Jpn J Infect Dis* 61: 427–433.
- Sakikawa M, Noda S, Hanaoka M, Nakayama H, Hojo S, et al. (2012) Anti-Toxoplasma antibody prevalence, primary infection rate, and risk factors in a study of toxoplasmosis in 4,466 pregnant women in Japan. *Clin Vaccine Immunol* 19: 365–367.
- Kaneko S, K'Opioji J, Kiche I, Wanyua S, Goto K, et al. (2012) Health and Demographic Surveillance System in the Western and Coastal Areas of Kenya: An Infrastructure for Epidemiologic Studies in Africa. *J Epidemiol* 22: 276–285.
- Bentwich Z, Kalinkovich A, Weisman Z (1995) Immune activation is a dominant factor in the pathogenesis of African AIDS. *Immunol Today* 16: 187–191.
- Yazdanbakhsh M, Kremsner PG, van Ree R (2002) Allergy, parasites, and the hygiene hypothesis. *Science* 296: 490–494.
- Everitt BS (1996) An introduction to finite mixture distributions. *Stat Methods Med Res* 5: 107–127.
- Rota MC, Massari M, Gabutti G, Guido M, De Donno A, et al. (2008) Measles serological survey in the Italian population: interpretation of results using mixture model. *Vaccine* 26: 4403–4409.
- Baughman AL, Bisgard KM, Edwards KM, Guris D, Decker MD, et al. (2004) Establishment of diagnostic cutoff points for levels of serum antibodies to pertussis toxin, filamentous hemagglutinin, and fimbriae in adolescents and adults in the United States. *Clin Diagn Lab Immunol* 11: 1045–1053.
- Deb P (2007) FMM: Stata module to estimate finite mixture models. Available: <http://ideas.repec.org/c/boc/bocode/s456895.html>. Accessed 2 January 2014.
- Youden WJ (1950) Index for rating diagnostic tests. *Cancer* 3: 32–35.
- Ngugi AK, Bottomley C, Kleinschmidt I, Wagner RG, Kakooza-Mwesige A, et al. (2013) Prevalence of active convulsive epilepsy in sub-Saharan Africa and associated risk factors: cross-sectional and case-control studies. *Lancet Neurol* 12: 253–263.
- Marlet MV, Sang DK, Ritmeijer K, Muga RO, Onsongo J, et al. (2003) Emergence or re-emergence of visceral leishmaniasis in areas of Somalia, north-eastern Kenya, and south-eastern Ethiopia in 2000–01. *Trans R Soc Trop Med Hyg* 97: 515–518.

35. Romero HD, Silva Ldc A, Silva-Vergara ML, Rodrigues V, Costa RT, et al. (2009) Comparative study of serologic tests for the diagnosis of asymptomatic visceral leishmaniasis in an endemic area. *Am J Trop Med Hyg* 81: 27–33.
36. Yang B, Chen Y, Wu L, Xu L, Tachibana H, et al. (2012) Seroprevalence of *Entamoeba histolytica* infection in China. *Am J Trop Med Hyg* 87: 97–103.
37. Njenga SM, Mwandawiro CS, Wamae CN, Mukoko DA, Omar AA, et al. (2011) Sustained reduction in prevalence of lymphatic filariasis infection in spite of missed rounds of mass drug administration in an area under mosquito nets for malaria control. *Parasites & vectors* 4: 90.
38. Njomo DW, Amuyenzu-Nyamongo M, Magambo JK, Njenga SM (2012) The role of personal opinions and experiences in compliance with mass drug administration for lymphatic filariasis elimination in Kenya. *PLoS One* 7: e48395.
39. Rao KV, Eswaran M, Ravi V, Gnanasekhar B, Narayanan RB, et al. (2000) The *Wuchereria bancrofti* orthologue of *Brugia malayi* SXPI and the diagnosis of bancroftian filariasis. *Mol Biochem Parasitol* 107: 71–80.
40. Levine MM, Young CR, Black RE, Takeda Y, Finkelstein RA (1985) Enzyme-linked immunosorbent assay to measure antibodies to purified heat-labile enterotoxins from human and porcine strains of *Escherichia coli* and to cholera toxin: application in serodiagnosis and seroepidemiology. *J Clin Microbiol* 21: 174–179.
41. Kotloff KL, Nataro JP, Blackwelder WC, Nasrin D, Farag TH, et al. (2013) Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multicenter Study, GEMS): a prospective, case-control study. *Lancet* 382: 209–222.
42. Qadri F, Svenssonholm AM, Faruque AS, Sack RB (2005) Enterotoxigenic *Escherichia coli* in developing countries: epidemiology, microbiology, clinical features, treatment, and prevention. *Clin Microbiol Rev* 18: 465–483.
43. Ministry of Public Health and Sanitation; Republic of Kenya (2011) National Multi-year Strategic Plan for control of Neglected Tropical diseases 2011–2015.
44. Lammie PJ, Moss DM, Brook Goodhew E, Hamlin K, Krolewiecki A, et al. (2012) Development of a new platform for neglected tropical disease surveillance. *Int J Parasitol* 42: 797–800.

