

Impact of HIV Infection on Colorectal Tumors: A Prospective Colonoscopic Study of Asian Patients

Yohei Hamada, MD,* Naoyoshi Nagata, MD,† Takeshi Nishijima, MD,*‡ Takuro Shimbo, MD, PhD,§
Naoki Asayama, MD,† Yoshihiro Kishida, MD,† Katsunori Sekine, MD,† Shohei Tanaka, MD,†
Tomonori Aoki, MD,† Kazuhiro Watanabe, MD,† Junichi Akiyama, MD,† Toru Igari, MD,||
Masashi Mizokami, MD, PhD,¶ Naomi Uemura, MD, PhD,# and Shinichi Oka, MD, PhD*‡

Background: Non-AIDS defining cancer has recently become a major problem in HIV-infected patients. Little has been reported on whether HIV infection is a risk factor for colorectal adenoma, especially in Asians.

Methods: The study was conducted under a prospective cross-sectional design and included all adults who underwent colonoscopy. Subjects were matched by age and sex to compare the prevalence of colorectal adenoma, adenocarcinoma, polyps, and other tumors. Detailed risk factors were assessed, including lifestyle habits, medications, comorbidities, gastrointestinal symptom rating scale, HIV-associated factors, and human papillomavirus infection. To evaluate the effects of HIV infection on adenoma, the odds ratio (OR) was estimated by multivariate logistic regression.

Results: A total of 177 HIV-infected patients and 177 controls were selected for analysis. No significant difference was noted in the prevalence of adenoma ($n = 29$ vs. 40 , $P = 0.14$). Multivariate analysis adjusted by baseline demographics and risk factors showed that HIV is not associated with increased risk of adenoma (adjusted OR = 0.66 , $P = 0.16$). Kaposi's sarcoma was more common in HIV-infected patients ($n = 6$ vs. 0 , $P = 0.03$). Among HIV-infected patients, advanced age was an independent and significant risk factor for adenoma (adjusted OR = 2.28 , $P < 0.01$). CD4 count, HIV-

RNA, history of antiretroviral treatment, and oncogenic human papillomavirus infection were not risk factors for adenoma.

Conclusions: HIV infection was not identified as risk for adenoma in Asian patients. However, advanced age was independently associated with increased risk of adenoma. HIV-infected patients should not miss screening opportunity for colorectal adenoma and other gastrointestinal malignancies.

Key Words: colorectal cancer, colorectal adenoma, oncogenic HPV infection, Japan, gastrointestinal malignancy

(*J Acquir Immune Defic Syndr* 2014;65:312–317)

INTRODUCTION

The introduction of highly active antiretroviral therapy (HAART) has significantly improved the morbidity and mortality of HIV-infected patients.^{1,2} However, the incidence of non-AIDS defining cancer has increased with prolongation of life expectancy of HIV-infected patients.^{3–6}

Colorectal cancer is the third most commonly diagnosed cancer in male patients and the second in female patients, and a major cause of death worldwide.⁷ To prevent the development of colorectal cancer and death, removal of premalignant lesion, adenoma, is effective, and screening is recommended in patients aged 50 years and older.^{8–11} Recent studies from western countries have suggested that higher incidence of colonic adenoma in patients with HIV infection compared with the general population.^{12–14} Furthermore, HIV-infected patients are at high risk of oncogenic human papillomavirus (HPV) infection, and the potential role of HPV infection in the development of colorectal cancer has been suggested.^{15–17} However, little is known about the risk of adenoma in HIV-infected patients compared with the general population.

In the past, the incidence of colorectal cancer was lower in Asia compared with Western countries.¹⁸ However, the incidence has increased lately in Asian countries, including Japan, and is currently comparable with that in western countries.^{18,19} Nevertheless, all previous studies on colorectal adenoma in HIV-infected patients were only from the United States, and there are no available data in Asia.^{12–14} This study reports the findings of a prospective cross-sectional colonoscopic study that compared the prevalence of colorectal

Received for publication July 9, 2013; accepted September 20, 2013.

From the *AIDS Clinical Center and †Department of Gastroenterology and Hepatology, National Center for Global Health and Medicine, Tokyo, Japan; ‡Center for AIDS Research, Kumamoto University, Kumamoto, Japan; §Research and Informatics, International Clinical Research Center Research Institute, and ||Pathology Division of Clinical Laboratory, National Center for Global Health and Medicine, Tokyo, Japan; and ¶The Research Center for Hepatitis and Immunology, Department of Gastroenterology and Hepatology, and #Department of Gastroenterology and Hepatology, National Center for Global Health and Medicine, Kohnodai Hospital, Chiba, Japan.

The authors have no conflicts of interest to disclose.

Supported by Grants-in Aid for AIDS research from the Japanese Ministry of Health, Labor, and Welfare (H23-AIDS-001) and the Global Center of Excellence Program (Global Education and Research Center Aiming at the Control of AIDS) from the Japanese Ministry of Education, Science, Sports, and Culture.

Correspondence to: Naoyoshi Nagata, MD, Department of Gastroenterology and Hepatology, National Center for Global Health and Medicine, 1-21-1, Toyama, Shinjuku-ku, Tokyo 162-8655, Japan (e-mail: nnagata_ncgm@yahoo.co.jp).

Copyright © 2013 by Lippincott Williams & Wilkins

adenoma in HIV-infected patients with HIV-negative patients in Japan.

METHODS

Study Design, Setting, and Participants

We conducted a prospective cross-sectional single-center study in adults who underwent colonoscopy between September 2009 and July 2012 at the endoscopy unit of the National Center for Global Health and Medicine (NCGM). NCGM has one of the largest clinics for patients with HIV infection in Japan with more than 3500 registered patients as of May 2013. The institutional review board at NCGM approved this study. The study was conducted according to the principles expressed in the Declaration of Helsinki.

The following inclusion criteria were used in this study: (1) aged 18 years or older, (2) independent in activities of daily living, (3) able to understand written documents and to write, and (4) asymptomatic but desired screening for colorectal cancer, or presented with continuous or intermittent lower gastrointestinal (GI) symptoms. The following exclusion criteria were used in this study: (1) contraindication or patient refusal of total colonoscopy, (2) colonoscopy for follow-up evaluation during the study period, and (3) previous diagnosis of either adenoma or adenocarcinoma. All inclusion and exclusion criteria were fulfilled before patients were enrolled in the study. Each HIV-infected patient was matched with 1 HIV-negative patient based on age in 5-year age-bands and sex.

Clinical Factors

A detailed questionnaire was completed at the endoscopy unit on the same day of colonoscopy. Patients were asked about their (1) lifestyle habits (smoking history and alcohol consumption), (2) medications [nonsteroidal anti-inflammatory drugs (NSAIDs) and low-dose aspirin], and (3) comorbidities (hypertension, diabetes mellitus, and coronary heart disease) in a face-to-face interview with the medical staff. With regard to medication history, prescriptions and medical records were reviewed in addition to information provided by the patients to avoid omissions. The survey form included photographs of all these oral drugs, which are approved in Japan. Regular use of medication was defined as oral administration starting at least 1 year before the interview. The smoking index was evaluated among ever and daily smokers and was defined as the number of cigarettes per day multiplied by the number of smoking years. Then, smoking index was categorized into nil, <400, 400–799, and >800. Alcohol consumption was calculated and categorized into nondrinker, light (1–180 g/wk), moderate (181–360 g/wk), and heavy drinker (>360 g/wk). To evaluate lower GI symptoms, the GI symptom rating scale rating on a 7-graded Likert scale was used.^{20,21} The GI symptom rating scale consists of 15 questions covering lower GI symptoms: increased flatus, decreased passage of stools, increased passage of stools, loose stools, hard stools, urgent

need for defecation, and feeling of incomplete evacuation. Positive symptoms were defined as score ≥ 3 .

For HIV-infected patients, CD4 cell count, HIV viral load, history of HAART, and sexual behavior were also obtained. CD4 cell counts within 1 week and HIV-RNA viral load within 1 month were used in the analysis, and positive result for real-time HIV-RNA was defined as ≥ 40 copies per milliliter. Sexual behavior was defined as men who have sex with men or heterosexuality. Furthermore, immediately following colonoscopy, rectal swabs (DNAPAP cervical sampler; Qiagen, Gaithersburg, MD) were obtained. Rectal samples were analyzed for HPV-DNA and genotyping by means of polymerase chain reaction-invasion assay as described previously.²² HPV types 16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, and 59 were defined as oncogenic HPV.²³

Diagnosis of Colorectal Adenoma, Adenocarcinoma, and Non-Neoplastic Polyps

After intestinal lavage with 2 L of solution containing polyethylene glycol, colonoscopy was performed by experienced gastroenterologists by using an electronic high-resolution video endoscope (model CFH260; Olympus Optical, Tokyo, Japan). The location of all lesions was recorded in electronic endoscopic database (Olympus Medical Systems; Solemio Endo). All visualized lesions were biopsied and histologically assessed by experienced pathologists.

Statistical Analysis

Baseline characteristics were compared using the unpaired Student *t* test or χ^2 test (Fisher exact test) for quantitative or qualitative variables, respectively. To estimate the effect of HIV infection on adenoma, multivariate logistic regression analysis was performed adjusted for age, sex, and possible risk factors for adenoma (these included smoking and alcohol consumption, diabetes mellitus, coronary artery diseases, and NSAIDs and aspirin use). In addition, we conducted uni- and multivariate logistic regression analysis in HIV-infected patients to elucidate the impact of other factors on adenoma related to HIV-infected patients (CD4 count, HIV-RNA, history of HAART, sexual behavior, and oncogenic HPV infection).

Statistical significance was defined at 2-sided *P* values < 0.05. We estimated the odds ratios (ORs) and 95% confidence intervals (CIs). All statistical analyses were performed using the Statistical Package for Social Sciences version 17.0 (SPSS, Chicago, IL).

RESULTS

Participants

A total of 177 HIV-infected patients and 177-HIV-negative controls were selected for analysis after the application of the aforementioned exclusion criteria and age matching (Fig. 1). The baseline characteristics are listed in Table 1. The study subjects were mostly men, Asians, and comparatively young. HIV-infected patients were more likely to be smokers




FIGURE 1. Flow diagram of patient selection.

and on treatment with NSAIDs. In contrast, aspirin was mostly used by the control subjects. All other major background parameters were similar in the 2 groups. With regard to the clinical symptoms, there was no difference in GI symptom scores other than increased passage of stools but there was no difference in the proportion of asymptomatic patients between the 2 groups. In patients with HIV infection, the median CD4 count was 371/ μ L (interquartile range, 121–579), 29.4% of the patients were treatment naive, 75.4% had HPV infection, and 71.5% were infected with oncogenic HPV. The most frequently identified HPV types were type 16 (41%), followed by type 58 (35%), 59 (33%), 52 (27%), 31 (25%), 33 (25%), 51 (19%), 18 (18%), 35 (14%), 39 (13%), 56 (11%), and type 45 (7%).

Prevalence of Colorectal Adenoma, Adenocarcinoma, Non-Neoplastic Polyps, and Other Tumors

Adenomas were identified in 29 (16.4%) patients with HIV infection and in 40 (22.6%) control subjects, and the incidence was not significantly different between the 2 groups (Table 2). Classification of the adenoma according to size (<5, 5–9, and ≥ 10 mm) showed that HIV-negative subjects tended to have mainly adenomas measuring <5 mm ($P = 0.08$) although this difference did not reach statistical significance. The incidences of adenocarcinoma and hyperplastic polyps were higher in patients without HIV infection, although the differences in the rates were not statistically significant ($P > 0.05$). In contrast, Kaposi's sarcoma was diagnosed only in HIV-infected patients ($P = 0.03$).

Uni- and multivariate analyses showed that HIV infection did not correlate with higher prevalence of adenoma (Table 3, adjusted OR = 0.66; 95% CI: 0.37 to 1.18; $P = 0.16$). Multivariate analysis identified age as an independent and significant factor associated with increased risk of

adenoma (adjusted OR = 1.72; 95% CI: 1.29 to 2.29; $P < 0.01$). All other factors did not correlate with adenoma by multivariate analysis.

Factors Associated With Colorectal Adenoma in Patients With HIV Infection

Age was an independent factor associated with increased risk of adenoma by uni- and multivariate analysis (adjusted OR = 2.28; 95% CI: 1.37 to 3.80; $P < 0.01$; Table 4). High CD4 count, low HIV-RNA, and history of HAART were associated with prevalence of adenoma by univariate analysis, although these factors were not significant on multivariate analysis. Oncogenic HPV infection was not associated with adenoma.

DISCUSSION

This study demonstrated that HIV infection was not an independent risk for colorectal adenoma after adjustment for variables known to be related to adenoma. In HIV-infected patients, only age was associated with increased risk of colorectal adenoma, whereas CD4 count, HIV-RNA, and HPV infection were not associated with adenoma by multivariate analysis. To our knowledge, this is the first study that compared the prevalence of colorectal adenoma between patients with and without HIV infection in Asia.

Previous reports suggested possible relation between HIV infection and increased risk of colorectal adenoma.^{12–14} Bini et al¹² investigated the prevalence of adenoma in 2382 patients (165 HIV-infected patients and 2217 controls) who underwent screening sigmoidoscopy. Their study identified a high incidence of adenoma in HIV-infected patients and that the risk of such lesion was higher in patients with low CD4 count and long-term HIV infection. The same group also conducted a prospective study of 408 patients who underwent total colonoscopy in the United States.¹³ They included only

TABLE 1. Clinical Characteristics of Patients With and Without HIV Infection

	HIV-Positive Patients (n = 177)	HIV-Negative Patients (n = 177)	P
Age, yr (IQR)	42 (37–50)	42 (37–50)	0.99
Male gender (%)	167 (94.4)	167 (94.4)	1.00
Asian (%)	171 (96.6)	176 (99.4)	0.12
Cigarette smoking (%)			
Never smoker	58 (32.8)	78 (44.1)	
Smoking index			
<400	89 (50.3)	60 (33.9)	
400–799	22 (12.4)	25 (14.1)	
>800	8 (4.5)	14 (7.9)	0.02*
Alcohol consumption (%)			
Nondrinker	77 (43.5)	59 (33.3)	
Light drinker	82 (46.3)	86 (48.6)	
Moderate drinker	13 (7.9)	24 (13.6)	
Heavy drinker	5 (2.8)	8 (4.5)	0.09
Current NSAIDs use (%)†	27 (15.3)	13 (7.3)	0.02*
Current aspirin use (%)	3 (1.7)	11 (6.2)	0.03*
Diabetes mellitus (%)	9 (5.1)	17 (9.6)	0.10
Coronary vascular disease (%)	6 (3.4)	5 (2.8)	0.76
Asymptomatic, %	33.5‡	36.6‡	0.55
GI symptoms score			
Increased flatus (SD)	1.9 (1.1)	2.0 (1.4)	0.80
Decreased passage of stools (SD)	1.8 (1.3)	1.8 (1.3)	0.76
Increased passage of stools (SD)	2.7 (2.0)	2.2 (1.6)	0.03*
Loose stools (SD)	2.4 (1.6)	2.1 (1.4)	0.13
Hard stools (SD)	1.7 (1.2)	1.6 (1.0)	0.84
Urgent need for defecation (SD)	2.3 (1.7)	2.1 (1.6)	0.14
Feeling of incomplete evacuation (SD)	2.2 (1.3)	2.2 (1.4)	0.61
CD4 count (IQR)	371 (121–579)	NA	NA
HIV-RNA log ₁₀ /mL (IQR)	1.6 (1.6–3.8)	NA	NA
Treatment naive (%)	52 (29.4)	NA	NA
MSM (%)	135 (76.3)	NA	NA
HPV infection (%)	98/130 (75.4)	NA	NA
Oncogenic HPV (%)	93/130 (71.5)	NA	NA

**P* < 0.05.
†None of the patients was on selective cox-2 inhibitor.
‡There were 1 missing data in HIV-positive group and 2 in HIV-negative group, thus comparisons were made between 59/176 (33.5%) of HIV-positive and 64/175 (36.6%) of HIV-negative patients.
IQR, interquartile range; SD, standard deviation; MSM, men who have sex with men; NA, not applicable.

asymptomatic patients aged 50 years or older and found a high rate of colonic neoplasm, including adenoma, in HIV-infected patients. They also reported that patients with HIV infection who were not on treatment with HAART and those with a positive family history of colorectal cancer were at higher risk for colonic neoplasm. In contrast, the study of Kothari et al,¹⁴ which included 130 HIV-infected patients and 779 controls who underwent screening colonoscopy, did not find

TABLE 2. Prevalence of Colorectal Adenoma, Adenocarcinoma, Non-Neoplastic Polyps, and Other Tumors

	HIV-Positive Patients (n = 177)	HIV-Negative Patients (n = 177)	P
Any adenoma	29 (16.4%)	40 (22.6%)	0.14
Adenoma, <5 mm	21 (11.9%)	33 (18.6%)	0.08
Adenoma, 5–9 mm	12 (6.8%)	10 (5.6%)	0.66
Adenoma, ≥10 mm	0	4 (2.3%)	0.12
Adenocarcinoma	0	5 (2.8%)	0.06
Hyperplastic polyp	17 (9.6%)	28 (15.8%)	0.08
Other tumors	6 (33.9%)	3 (17.0%)	0.502
Kaposi's sarcoma	6 (33.9%)	0	0.03*
Malignant lymphoma	0	0	1.00
Carcinoid tumor	0	1 (0.6%)	1.000
Lipoma	0	2 (1.1%)	0.499

**P* < 0.05.

significant difference in the prevalence of adenoma between the 2 groups. Similarly, our study showed similar prevalence of adenoma in patients with and without HIV infection. These differences may be explained by differences in sample size, populations, and different inclusion criteria. The abovementioned previous studies included only asymptomatic patients whereas this study included many patients with GI symptoms. Taken together, these results suggest lack of consensus on this issue. Thus, it is still unclear whether HIV infection is truly associated with increased risk of colorectal adenoma. Bini et al¹² suggested that the low immune status associated with HIV infection may enhance the development of adenoma; however, CD4 count did not correlate with adenoma in our study. Furthermore, HIV itself is also suggested to play a role in oncogenesis.²⁴ There is limited information on this issue, and further studies are needed to clarify the association between HIV infection and colorectal adenoma.

In this study, advanced age correlated with increased risk of adenoma in HIV-infected patients. Excision of adenoma prevents colon cancer and screening colonoscopy is recommended for individuals aged 50 years or older.^{8,10,11} However, it has been suggested that colorectal cancer screening is underused in HIV-infected patients.²⁵ In addition, patients with HIV infection are at higher risk for other GI malignancies such as Kaposi's sarcoma, anal cancer, and GI lymphoma than general population,^{26–28} and these patients are sometimes asymptomatic.^{28–31} Therefore, we believe that screening colonoscopy is important in HIV-infected patients, especially those aged 50 years or older.

The association between HPV infection and colorectal cancer is controversial.³² Although 2 recent studies argued against such association, a recent meta-analysis study demonstrated increased risk of colorectal cancer with HPV infection.^{17,33,34} Because previous reports suggested increased prevalence of colorectal adenoma in HIV-infected patients, in whom the prevalence of HPV infection is known to be higher than that in the general population,¹⁵ we hypothesized

TABLE 3. Uni- and Multivariate Analysis to Estimate the Risk for Adenoma

	Unadjusted OR (95% CI)	P	Adjusted OR (95% CI)	P
HIV infection	0.67 (0.39 to 1.14)	0.14	0.66 (0.37 to 1.18)	0.16
Age per 10 yrs	1.96 (1.53 to 2.53)	<0.01*	1.72 (1.29 to 2.29)	<0.01*
Male gender	0.71 (0.25 to 2.03)	0.52	0.92 (0.28 to 3.05)	0.89
Smoking	1.60 (1.19 to 2.13)	<0.01*	1.35 (0.98 to 1.86)	0.06
Alcohol consumption	0.89 (0.63 to 1.26)	0.51	0.83 (0.56 to 1.22)	0.34
Current NSAIDs use	0.70 (0.28 to 1.75)	0.45	0.94 (0.35 to 2.53)	0.91
Current aspirin use	4.48 (1.52 to 13.3)	<0.01*	11.8 (0.52 to 6.44)	0.35
Diabetes mellitus	2.37 (1.00 to 5.56)	0.05	1.39 (0.54 to 3.60)	0.49
Coronary heart disease	3.63 (1.08 to 12.3)	0.04	1.30 (0.30 to 5.54)	0.72

**P* < 0.05.

that oncogenic HPV infection may be a risk factor for adenoma in patients with HIV. However, our results did not find such association.

Fecal occult blood test is a useful screening tool for the detection of colorectal cancers.¹⁰ However, fecal blood test is also positive in various GI diseases such as asymptomatic colitis and Kaposi's sarcoma.^{35,36} Thus, the diagnostic accuracy of fecal occult blood test may be less than ideal in HIV-infected patients and accordingly was not used in all subjects in this study. Instead, we assessed the clinical symptoms because we hypothesized that differences in GI symptoms might affect the prevalence of colorectal adenoma. Nevertheless, the proportion of asymptomatic patients was not different between the 2 groups.

Important strengths of this study includes its prospective study design, detailed assessment of GI symptoms and other GI tumors, first study in Asia, and conducting total colonoscopy in all subjects. However, there are several limitations to our study. First, because our study population was younger than those in previous studies, the prevalence might have been underestimated compared with other studies. It is well known that the risk of colorectal cancer increases with age.³⁷ Thus, the young age of our study subjects and the small sample size of our study could have masked any association between HIV infection and colorectal adenoma. Similar to the study by Bini et al,¹³ which

examined the relation between HIV infection and colorectal adenoma, larger studies on patients aged 50 years or older will be needed in Asia. Second, because we included both symptomatic and asymptomatic patients who underwent diagnostic colonoscopy, a selection bias could not be ruled out in our study. As a result, it is possible that the control group could have included patients suspected to have colon cancer, whereas HIV-infected patients tended to include those who were referred for colonoscopy based on the suspicion of opportunistic infections, which might have led to the higher prevalence of adenoma in the control group. However, the background characteristics and proportion of asymptomatic patients were similar between the 2 groups. Third, although we collected detailed information on risk factors of adenoma, we could not collect data on factors such as obesity and family history of colon cancer as reported previously,^{38,39} and these might have influenced the results.

In conclusion, the incidence of adenoma was not significantly different between patients with and without HIV infection. However, it should be noted that 16.4% HIV-infected patients had adenoma and its risk increased with age. As the issue of aging in patients with HIV infection is growing, the results of this study carry certain significance. Thus, HIV-infected patients should not miss screening opportunities for colorectal adenoma and other HIV-related malignancies.

TABLE 4. Uni- and Multivariate Analyses to Estimate the Risk for Adenoma in HIV-Infected Patients

	Unadjusted OR (95% CI)	P	Adjusted OR (95% CI)	P
Age	2.49 (1.66 to 3.79)	<0.01*	2.28 (1.37 to 3.80)	<0.01*
CD4 count per 10/ μ L	1.02 (1.00 to 1.03)	0.02*	1.01 (0.99 to 1.03)	0.54
HIV-RNA log10/mL	0.40 (0.21 to 0.76)	<0.01*	0.50 (0.18 to 1.37)	0.18
Treatment naive	0.15 (0.03 to 0.64)	0.01*	1.31 (0.12 to 14.49)	0.83
MSM	0.52 (0.22 to 1.24)	0.14	0.66 (0.19 to 2.26)	0.51
Oncogenic HPV	0.25 (0.10 to 0.65)	<0.01*	0.50 (0.17 to 1.47)	0.21

**P* < 0.05.

MSM, men who have sex with men.

ACKNOWLEDGMENTS

The authors are grateful to Hisae Kawashiro (Clinical Research Coordinator) for help with data collection. The authors thank all other clinical staff at the AIDS Clinical Center and all the staff of the endoscopy unit.

REFERENCES

1. Palella FJ Jr, Delaney KM, Moorman AC, et al. Declining morbidity and mortality among patients with advanced human immunodeficiency virus infection. HIV Outpatient Study Investigators. *N Engl J Med*. 1998;338:853–860.
2. Murphy EL, Collier AC, Kalish LA, et al. Highly active antiretroviral therapy decreases mortality and morbidity in patients with advanced HIV disease. *Ann Intern Med*. 2001;135:17–26.
3. Crum-Cianflone N, Hullsiek KH, Marconi V, et al. Trends in the incidence of cancers among HIV-infected persons and the impact of antiretroviral therapy: a 20-year cohort study. *AIDS*. 2009;23:41–50.
4. Shiels MS, Cole SR, Kirk GD, et al. A meta-analysis of the incidence of non-AIDS cancers in HIV-infected individuals. *J Acquir Immune Defic Syndr*. 2009;52:611–622.
5. Sackoff JE, Hanna DB, Pfeiffer MR, et al. Causes of death among persons with AIDS in the era of highly active antiretroviral therapy: New York City. *Ann Intern Med*. 2006;145:397–406.
6. Palella FJ Jr, Baker RK, Moorman AC, et al. Mortality in the highly active antiretroviral therapy era: changing causes of death and disease in the HIV outpatient study. *J Acquir Immune Defic Syndr*. 2006;43:27–34.
7. Jemal A, Bray F, Center MM, et al. Global cancer statistics. *CA Cancer J Clin*. 2011;61:69–90.
8. Winawer SJ, Zauber AG, Ho MN, et al. Prevention of colorectal cancer by colonoscopic polypectomy. The National Polyp Study Workgroup. *N Engl J Med*. 1993;329:1977–1981.
9. Zauber AG, Winawer SJ, O'Brien MJ, et al. Colonoscopic polypectomy and long-term prevention of colorectal-cancer deaths. *N Engl J Med*. 2012;366:687–696.
10. Levin B, Lieberman DA, McFarland B, et al. Screening and surveillance for the early detection of colorectal cancer and adenomatous polyps, 2008: a joint guideline from the American Cancer Society, the US Multi-Society Task Force on Colorectal Cancer, and the American College of Radiology. *CA Cancer J Clin*. 2008;58:130–160.
11. Calonge N, Petitti DB, DeWitt TG, et al. Screening for colorectal cancer: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2008;149:627–637.
12. Bini EJ, Park J, Francois F. Use of flexible sigmoidoscopy to screen for colorectal cancer in HIV-infected patients 50 years of age and older. *Arch Intern Med*. 2006;166:1626–1631.
13. Bini EJ, Green B, Poles MA. Screening colonoscopy for the detection of neoplastic lesions in asymptomatic HIV-infected subjects. *Gut*. 2009;58:1129–1134.
14. Kothari ND, Engelson ES, Drake V, et al. Effect of HIV infection on the prevalence of colorectal adenomas during screening colonoscopy. *J Clin Gastroenterol*. 2010;44:77–78.
15. Mbulawa ZZ, Marais DJ, Johnson LF, et al. Impact of human immunodeficiency virus on the natural history of human papillomavirus genital infection in South African men and women. *J Infect Dis*. 2012;206:15–27.
16. Burnett-Hartman AN, Newcomb PA, Potter JD. Infectious agents and colorectal cancer: a review of *Helicobacter pylori*, *Streptococcus bovis*, JC virus, and human papillomavirus. *Cancer Epidemiol Biomarkers Prev*. 2008;17:2970–2979.
17. Damin DC, Ziegelmann PK, Damin AP. Human papillomavirus infection and colorectal cancer risk: a meta-analysis. *Colorectal Dis*. 2013;15:e420–e428.
18. Center MM, Jemal A, Ward E. International trends in colorectal cancer incidence rates. *Cancer Epidemiol Biomarkers Prev*. 2009;18:1688–1694.
19. Sung JJ, Lau JY, Goh KL, et al. Increasing incidence of colorectal cancer in Asia: implications for screening. *Lancet Oncol*. 2005;6:871–876.
20. Svedlund J, Sjodin I, Dotevall G. GSR— a clinical rating scale for gastrointestinal symptoms in patients with irritable bowel syndrome and peptic ulcer disease. *Dig Dis Sci*. 1988;33:129–134.
21. Revicki DA, Wood M, Wiklund I, et al. Reliability and validity of the Gastrointestinal Symptom Rating Scale in patients with gastroesophageal reflux disease. *Qual Life Res*. 1998;7:75–83.
22. Tadokoro K, Akutsu Y, Tanaka K, et al. Comparative quantitative analysis of 14 types of human papillomavirus by real-time polymerase chain reaction monitoring Invader reaction (Q-Invader assay). *Diagn Microbiol Infect Dis*. 2010;66:58–64.
23. Doorbar J, Quint W, Banks L, et al. The biology and life-cycle of human papillomaviruses. *Vaccine*. 2012;30(suppl 5):F55–F70.
24. Deeken JF, Tjen-A-Looi A, Rudek MA, et al. The rising challenge of non-AIDS-defining cancers in HIV-infected patients. *Clin Infect Dis*. 2012;55:1228–1235.
25. Reinhold JP, Moon M, Tenner CT, et al. Colorectal cancer screening in HIV-infected patients 50 years of age and older: missed opportunities for prevention. *Am J Gastroenterol*. 2005;100:1805–1812.
26. Franceschi S, Lise M, Clifford GM, et al. Changing patterns of cancer incidence in the early- and late-HAART periods: the Swiss HIV Cohort Study. *Br J Cancer*. 2010;103:416–422.
27. Machalek DA, Poynten M, Jin F, et al. Anal human papillomavirus infection and associated neoplastic lesions in men who have sex with men: a systematic review and meta-analysis. *Lancet Oncol*. 2012;13:487–500.
28. Beck PL, Gill MJ, Sutherland LR. HIV-associated non-Hodgkin's lymphoma of the gastrointestinal tract. *Am J Gastroenterol*. 1996;91:2377–2381.
29. Nagata N, Shimbo T, Yazaki H, et al. Predictive clinical factors in the diagnosis of gastrointestinal Kaposi's sarcoma and its endoscopic severity. *PLoS One*. 2012;7:e46967.
30. Abbas A, Yang G, Fakih M. Management of anal cancer in 2010. Part 1: overview, screening, and diagnosis. *Oncology (Williston Park)*. 2010;24:364–369.
31. Heise W, Arasteh K, Mostertz P, et al. Malignant gastrointestinal lymphomas in patients with AIDS. *Digestion*. 1997;58:218–224.
32. Lorenzon L, Ferri M, Pillozzi E, et al. Human papillomavirus and colorectal cancer: evidences and pitfalls of published literature. *Int J Colorectal Dis*. 2011;26:135–142.
33. Khoury JD, Tannir NM, Williams MD, et al. Landscape of DNA virus associations across human malignant cancers: analysis of 3,775 cases using RNA-Seq. *J Virol*. 2013;87:8916–8926.
34. Burnett-Hartman AN, Feng Q, Popov V, et al. Human papillomavirus DNA is rarely detected in colorectal carcinomas and not associated with microsatellite instability: the Seattle colon cancer family registry. *Cancer Epidemiol Biomarkers Prev*. 2013;22:317–319.
35. Okamoto M, Kawabe T, Ohata K, et al. Amebic colitis in asymptomatic subjects with positive fecal occult blood test results: clinical features different from symptomatic cases. *Am J Trop Med Hyg*. 2005;73:934–935.
36. Nagata N, Nakashima R, Nishiumura S, et al. Gastrointestinal Kaposi's sarcoma: diagnosis and clinical features. In: Butler EJ, ed. *Sarcoma*. New York, NY: Nova Science Publishers; 2012:153–165.
37. Eddy DM. Screening for colorectal cancer. *Ann Intern Med*. 1990;113:373–384.
38. Imperiale TF, Ransohoff DF. Risk for colorectal cancer in persons with a family history of adenomatous polyps: a systematic review. *Ann Intern Med*. 2012;156:703–709.
39. Renehan AG, Tyson M, Egger M, et al. Body-mass index and incidence of cancer: a systematic review and meta-analysis of prospective observational studies. *Lancet*. 2008;371:569–578.

DNA methylation profiling can classify HIV-associated lymphomas

Akihiro Matsunaga^a, Tsunekazu Hishima^b, Noriko Tanaka^c,
Maria Yamasaki^{c,d}, Lui Yoshida^{a,e}, Makoto Mochizuki^{f,i},
Junko Tanuma^g, Shinichi Oka^g, Yukihiro Ishizaka^a,
Mari Shimura^a and Shotaro Hagiwara^h

Background: HIV-positive patients have a 60-fold to 200-fold increased incidence of non-Hodgkin lymphomas, including Burkitt lymphoma, diffuse large B-cell lymphoma, and primary central nervous system lymphoma. HIV-associated lymphomas frequently have features such as extranodal involvement, decreased responses to standard chemotherapy, and high relapse rates, which indicate a poor prognosis. General pathological features do not clearly differentiate HIV-associated lymphomas from non-HIV lymphomas.

Methods: To investigate the features of HIV-associated lymphomas, we performed genome-wide DNA methylation profiling of HIV and non-HIV lymphomas using Illumina GoldenGate Methylation Cancer Panel I and Illumina Infinium HumanMethylation450 BeadChip microarrays. DNA methylation profiles in HIV-associated and non-HIV lymphomas were characterized using unsupervised hierarchical clustering analyses.

Results: The analyses of promoter regions revealed unique DNA methylation profiles in HIV-associated lymphomas, suggesting profile differences compared with non-HIV lymphomas, which implies specific gene regulation in HIV-associated lymphoma involving DNA methylation. Based on HumanMethylation450 BeadChip data, 2541 target sites were selected as differing significantly in comparisons between HIV-associated and non-HIV-associated lymphomas using Wilcoxon's rank-sum test ($P < 0.05$) and $\Delta\beta$ values more than 0.30. Recurrent cases of HIV-associated lymphoma had different profiles compared with nonrecurrent HIV lymphomas.

Conclusion: DNA methylation profiling indicated that 2541 target sites differed significantly in HIV-associated lymphoma, which may partly explain the poor prognosis. Our data indicate that the methylation profiles of target genes have potential in elucidating HIV-associated lymphomagenesis and can serve as new prognostic markers.

© 2014 Wolters Kluwer Health | Lippincott Williams & Wilkins

AIDS 2014, **28**:503–510

Keywords: CpG islands, DNA methylation microarray, HIV, HIV-associated lymphomas, poor prognosticators

^aDepartment of Intractable Diseases, Research Institute, National Center for Global Health and Medicine, Shinjuku, ^bDepartment of Pathology, Tokyo Metropolitan Cancer and Infectious Diseases Center, Komagome Hospital, Honkomagome, Bunkyo, ^cBiostatistics Section, Department of Clinical Research and Informatics, Clinical Research Center, National Center for Global Health and Medicine, Shinjuku, ^dDepartment of Human Genetic, School of Medicine, The University of Tokyo, Bunkyo, Tokyo, ^eGraduate School of Frontier Sciences, The University of Tokyo, Kashiwa, Chiba, ^fDepartment of Pathology, ^gAIDS Clinical Center, ^hDivision of Hematology, Internal Medicine, National Center for Global Health and Medicine Hospital, Shinjuku, and ⁱDepartment of Pathology, School of Medicine, Kyorin University, Mitaka, Tokyo, Japan.

Correspondence to Dr Mari Shimura, Department of Intractable Diseases, Research Institute, National Center for Global Health and Medicine, 1-21-1 Toyama, Shinjuku, Tokyo 162-8655, Japan.

Tel: +81 3 32027181; fax: +81 3 32027364; e-mail: mshimura@ri.ncgm.go.jp

Received: 9 July 2013; revised: 18 October 2013; accepted: 18 October 2013.

DOI:10.1097/QAD.0000000000000120

Introduction

The incidence of non-Hodgkin's lymphoma is 60-fold to 200-fold higher in patients with HIV infection [1,2]. Most HIV-associated lymphomas are high-grade B-cell lymphomas such as diffuse large B-cell lymphoma, Burkitt lymphoma, and primary central nervous system lymphoma. The clinical course is often aggressive, with a poor prognosis [2]. Since the introduction of highly active antiretroviral therapy, the risk for opportunistic infections and the incidence of AIDS-defining malignancies, including HIV-associated lymphomas, have declined, and prognoses have improved. Nevertheless, lymphomas remain a major cause of death for HIV-infected patients [3]. It is important to identify differences between HIV-associated lymphomas and non-HIV lymphomas, as their clinical and general pathological features do not clearly distinguish them [2]. Recent studies have revealed that the DNA methylation patterns can differentiate among disease subtypes, suggesting that epigenetic DNA alterations are related to carcinogenesis [4,5]. Epigenetic silencing of functionally important genes may contribute to the development of lymphomas [5,6], and promoter hypermethylation of CpG islands (CGIs) in some genes has been reported in aggressive-phenotype lymphoma with a poor prognosis [7]. In this study, we examined DNA methylation of CGIs in a promoter region clustered with HIV-associated lymphomas and non-HIV lymphomas, and investigated the prognostic significance of DNA methylation. Our findings contribute to an understanding of the lymphomagenesis of HIV-associated lymphomas and suggest specific DNA methylation as a useful prognostic biomarker.

Methods

Patients

HIV-associated lymphoma is a pathologically diagnosed malignant lymphoma in HIV patients. Two cohorts were studied. Cohort I consisted of 11 HIV-associated and 18 non-HIV lymphoma patients who visited Tokyo Metropolitan Cancer and Infectious Diseases Center, Komagome Hospital (CICK), and two non-HIV lymphoma patients who visited the National Center for Global Health and Medicine Hospital (NCGM). Cohort II included nine HIV-associated and 12 non-HIV lymphoma patients who visited NCGM. Formalin-fixed, paraffin-embedded tissues and fresh-frozen tissues were collected from NCGM and CICK, following approval by the ethics committees of both hospitals and in accordance with the Declaration of Helsinki. All patients gave written informed consent for their tissue to be used and for review of their clinical records. Diagnosis was made using the 2008 WHO classification [2]. Hematologists reviewed the tumor specimens and classified them histologically as diffuse large B-cell lymphoma, Burkitt

lymphoma, primary central nervous system lymphoma, follicular lymphoma, or Hodgkin's lymphoma. Non-HIV lymphoma samples were randomly selected from among the Burkitt lymphomas, diffuse large B-cell lymphomas, follicular lymphoma, and Hodgkin's lymphoma. Epstein-Barr virus (EBV) status was determined by Epstein-Barr encoded RNA (EBER) *in situ* hybridization and Southern blotting. BCL2 expression was examined by immunostaining.

HumanMethylation450 microarray analysis

Cohort I was analyzed using an Infinium HumanMethylation450 BeadChip microarray [8], which covered 485 577 methylation sites. Genomic DNA was isolated using a DNeasy mini kit (QIAGEN, Valencia, California, USA) according to the manufacturer's protocol. After 1 µg of DNA was ligated at 24°C for 30 min, the reaction was stopped by 5 min at 95°C (REPLI-g FFPE kit; QIAGEN) [9]. The DNA was subjected to genome-wide DNA methylation profiling using an Infinium HumanMethylation450 BeadChip (Illumina, San Diego, California, USA) [8], according to the manufacturer's instructions. The methylation status of specific cytosines is indicated by the β value, with 1 indicating complete methylation and 0 indicating no methylation. We first filtered the probes and samples using the Bioconductor IMA package to load files created by Illumina GenomeStudio software, using the IMA.methy450R function. With this package, we performed filtering steps using the IMA.methy450PP function. The inclusion criteria were as follows: sample call rate, more than 99.5%; detection *P* value, <0.05; site call rate, more than 90%; probes with no SNPs based on snpsite.txt provided in the IMA package [10]; and probes outside the XY chromosomes. We converted the initial file created by Illumina GenomeStudio to a new file to reflect the filtering results. The data were normalized by entering the filtered data into the Bioconductor lumi package [11]. Using the lumi package, methylation data were first analyzed by the color balance check and then scaled based on the mean of all probes, using methylation simple scaling normalization (SSN) implemented in the lumi package. The Infinium array methylation data are available in the Gene Expression Omnibus database under the accession number GSE42372.

Cancer Panel I microarray analysis

Cohort II was analyzed using the Illumina GoldenGate Methylation Cancer Panel I microarray, a cancer-focused methylation analysis covering 1505 CpG loci from 807 genes (Illumina) [12]. Genomic DNA was isolated (Agencourt FormaPure kit; Beckman Coulter, Brea, California, USA), subjected to sodium bisulfite conversion, labeled with fluorescent dyes, and hybridized to the microarrays according to the manufacturer's protocol. The methylation status of specific cytosines was indicated by the β value (1, complete methylation; 0, no methylation). Only probes with detection *P* value at

<0.01 were used for the analyses. The X chromosome loci were removed from the analysis, leaving 1421 CpG loci. Raw average β values were not normalized and were used for analyses as per the manufacturer's recommendations. The GoldenGate array methylation data are available in the Gene Expression Omnibus database under the accession number GSE42626.

For the statistical analysis, enrichment analysis of target genes, validation by combined bisulfite restriction analysis (COBRA), and bisulfite DNA sequences, see the Supplementary Methods, <http://links.lww.com/QAD/A441>.

Results

To identify differences between HIV-associated and non-HIV lymphomas, genome-wide DNA methylation array analyses were performed using Infinium HumanMethylation450 BeadChip technology. DNA from formalin-fixed and paraffin-embedded or fresh-frozen lymphoma tissues collected from the 11 HIV-positive and 20 HIV-negative Asian patients in Cohort I was analyzed (Table 1). DNA methylation throughout the genome was examined using probes targeting six gene regions (Fig. 1a): within 1500 bps of a transcription start site (TSS1500), within 200 bps of a transcription start site (TSS200), and the 5' untranslated region (5'UTR), first exon (1stExon), body, and 3' untranslated region (3'UTR) and intergenic regions. Three HIV-negative lymphomas were excluded from the analyses in the filtering steps (see Methods for details). The differences in methylation status between HIV-associated and non-HIV lymphomas were significantly greater for CGIs in the

various target regions, compared with non-CGI methylation (Supplementary Fig. 1, <http://links.lww.com/QAD/A441>). Hierarchical clustering analysis of CGI methylation markers of TSS1500, TSS200, 5'UTR, and 1stExon (Fig. 1b) produced roughly two groups that distinguished HIV-associated lymphomas from non-HIV lymphomas (Groups 1 and 2; Fig. 1b, upper left), with a few exceptions. By contrast, the analysis of non-CGI methylation and CGI methylation in the body and 3'UTR and intergenic gene targets did not give clear groupings (Fig. 1b, upper right and lower images, Supplementary Fig. 2, <http://links.lww.com/QAD/A441>). As all HIV patients in this study were men (Table 1), we next analyzed male patients only. The CGI results for TSS1500, TSS200, 5'UTR, and 1stExon again clustered into two groups (Supplementary Fig. 3, <http://links.lww.com/QAD/A441>), suggesting that gender does not affect the results. Generally, patients with HIV-associated lymphomas were younger than patients with non-HIV lymphomas (Table 1) [13]. When we excluded age-related target sites, as previously suggested [14], the analysis of CGI methylation in TSS1500, TSS200, 5'UTR, and 1stExon again produced two groups that distinguished between HIV-associated and non-HIV lymphomas (Supplementary Fig. 4, <http://links.lww.com/QAD/A441>). These results suggest that DNA methylation of CGIs in promoter regions (TSS1500, TSS200, 5'UTR, and 1stExon) probably distinguishes HIV-associated from non-HIV lymphomas. Among the targets measured, those with a significant absolute difference between HIV-associated and non-HIV lymphomas were used for further analyses (Supplementary Methods, <http://links.lww.com/QAD/A441>). Compared with non-HIV lymphoma DNA, HIV-associated lymphoma DNA tended to be hypomethylated (Fig. 1c). Representative genes were used to validate the array analyses. Using COBRA, three of the five non-HIV lymphomas cases were methylated as positive controls, whereas none of the HIV-associated lymphomas was detected as methylated at either *RARRES1* or *FGF5* (Fig. 1d, upper). Bisulfite DNA sequencing gave consistent results (Fig. 1d, lower), confirming this tendency toward hypomethylation in Group 1 (Fig. 1d). These findings encouraged us to examine previously analyzed cases in Cohort II.

Data from nine HIV-associated lymphoma samples derived from the first visit of Cohort II, which had been previously analyzed using Illumina GoldenGate Methylation Cancer Panel I (see Methods), were used for hierarchical clustering analyses. The results showed two apparent methylation profiles for HIV-associated lymphomas (Groups 3 and 4, Fig. 2a). The genes with a significant absolute difference between two clusters were used for further analyses (Supplementary Method, <http://links.lww.com/QAD/A441>). Group 3 tended to be hypermethylated compared with Group 4 (Fig. 2b). COBRA indicated that all of the Group 3 cases were

Table 1. Patient characteristics of lymphoma samples for Human Methylation450 (450K) microarray analysis in Cohort I.

Items examined		HIV	Non-HIV	P value (HIV vs. non-HIV)
Sex	Female	0	10	0.0049*
	Male	11	10	
Age	Mean	45.27	64.35	0.018*
	SD	16.92	10.60	
Histology	BL	2	3	0.57
	DLBCL	8	17	
	HD	1	0	
Stage	I & II	3	5	0.63
	III & IV	8	12	
	ND	0	3	
EBV	+	3	7	0.22
	-	8	9	
	ND	0	4	

The statistical significance of differences in the categorical variables was calculated by Fisher's exact test or Wilcoxon's rank-sum test. BL, Burkitt lymphoma; DLBCL, diffuse large B-cell lymphoma; EBV, Epstein-Barr virus; HD, Hodgkin's disease; ND, not determined; SD, standard deviation.

* $P < 0.05$




Fig. 1. Methylation profile analysis of HIV-associated and non-HIV lymphoma DNA in Cohort I, using Infinium HumanMethylation450 BeadChip technology. (a) Schematic of the gene regions examined for methylation. (b) Hierarchical clustering analysis of CpG island (CGI) and non-CGI methylation of lymphoma DNA in Cohort I. The analysis of CGI methylation in the promoter regions (TSS1500, TSS200, 5'UTR, and 1stExon) produced two groups that distinguished between HIV-associated lymphomas (Group 1, Gr. 1) and non-HIV lymphomas (Group 2, Gr. 2). TSS, transcription start site; AU p value, approximately unbiased P value computed using multiscale bootstrap resampling. (c) Numbers of hypermethylation or hypomethylation targets in HIV-associated lymphomas compared with non-HIV-lymphomas. (d) Validation by combined bisulfite restriction analysis (COBRA) and bisulfite DNA sequences. Retinoic acid receptor responder 1 (*RARRES1*) and fibroblast growth factor 5 (*FGF5*) are representative targets in the array analysis. Me, methylated allele or methylated control; UMe, unmethylated allele or unmethylated control; open circle, unmethylated CpG site; solid circle, methylated CpG site; HIV, HIV-associated lymphoma; non-HIV, non-HIV lymphoma.




Fig. 2. Methylation profile clustering of HIV-associated lymphoma DNA in Cohort II, using Cancer Panel I. Cancer Panel I microarray analysis was performed for nine HIV-associated lymphomas in Cohort II. The color bar indicates hypermethylation and hypomethylation. Hierarchical clustering analysis of methylation gave two groups: Group 3 (Gr. 3) and Group 4 (Gr. 4). (b) Numbers of hypermethylation or hypomethylation targets in Group 3 compared with Group 4. (c) Validation by combined bisulfite restriction analysis (COBRA) and bisulfite DNA sequences. *BMP2* (bone morphogenetic protein 2), *SKI* (oncogene), and *PTCH2* (patched 2) are representative targets in the array analysis. Me, methylated allele or methylated control; UMe, unmethylated allele or unmethylated control; open circle, unmethylated CpG site; solid circle, methylated CpG site.

methylated, whereas fewer in Group 4 were methylated among those tested (Fig. 2c, upper). Bisulfite DNA sequencing clearly showed that Group 3 was highly methylated (Fig. 2c, lower), confirming the tendency toward hypermethylation in Group 3. Two cases in Group 3 subsequently showed recurrence, representing a significant patient characteristic ($P=0.083$), if 0.1 was considered a significant level (Table 2). In another case in Group 3, a tumor mass appeared in the cervical spinal cord about 17 months later, although recurrence was not confirmed pathologically. Notably, the methylation profile of nonrecurrent HIV-associated lymphomas (Group 4) did not differ significantly from that of non-HIV lymphomas (non-Group 3, Supplementary Fig. 5 and Supplementary Table 1, <http://links.lww.com/QAD/A441>). These data suggest that recurrent HIV-associated lymphomas have a specific methylation profile.

Discussion

The prognosis of HIV-associated lymphoma has improved with the development of HIV and cancer therapies [15]. Nevertheless, it is important to identify the mechanism responsible for the aggressiveness of HIV-associated lymphomas. Our data suggest that the DNA methylation profile is a molecular indicator of prognosis.

In the methylation analyses, we examined nine or 11 HIV-associated lymphomas. This number was relatively small because of the small HIV-positive population in Japan [13]. Even so, our data clearly suggest that DNA

methylation profiles, especially CGI methylation in promoter regions, differ between HIV-associated and non-HIV lymphomas. As the tumor location varies in HIV-associated lymphoma [2], it is essential to know whether tumor location influenced our analyses. Lymph nodes were the most frequent tumor location and were broadly similar in Groups 1 and 2 ($P=0.45$; Supplementary Fig. 6a, <http://links.lww.com/QAD/A441>), although Group 1 had more extra-node variation, probably due to the high proportion of HIV-associated lymphoma. It is noteworthy that Group 1 had narrower correlation distances than Group 2, indicating that the DNA methylation profiles in Group 1 were quite similar, although Group 1 included various tumor locations (Supplementary Fig. 6b, <http://links.lww.com/QAD/A441>). Additionally, the lymph node cases in Group 1 were very dissimilar from the lymph node cases in Group 2. The data suggested that the clustered results were not due to tumor location. The differences between the profiles may not be related to antiretroviral therapy either, as only two HIV-positive lymphomas in Cohort I were treated with antiretroviral therapy. Coinfections such as EBV with HIV may influence DNA methylation profiles, but we found no significant difference between HIV-associated and non-HIV lymphomas in terms of EBV infection status in our study. However, we cannot exclude the influence of HIV infection on methylation profiles. One of our validation genes, *RARRES1*, is a cancer methylation target [16] that is differentially expressed in various tumors [17,18], although its clinical relevance to lymphomas remains unknown. *FGF5* is reported to be a bone metastasis-related gene related to angiogenesis [19]. As angiogenic growth factors have been implicated in a

Table 2. Patient characteristics of lymphoma samples in Cohort II for Cancer Panel I.

Items examined		HIV-associated lymphomas		P value (Group 3 vs. Group 4)
		Group 3	Group 4	
Sex	Female	1	0	0.33
	Male	2	6	
Age	Mean	36.66	35.00	1.00
	SD	5.77	13.78	
Histology	BL	2	3	1.00
	DLBCL	1	1	
	HD	0	2	
Bcl-2	+	0	1	1.00
	–	3	5	
Stage	I & II	0	2	0.50
	III & IV	3	4	
EBV	+	1	4	0.52
	–	2	2	
Recurrence	+	2	0	0.083
	–	1 ^a	6	
IPI score ^b	0 or 1	0	1	1.00
	2 or 3	1	1	
	4 or 5	2	2	

The statistical significance of differences in the categorical variables was calculated by Fisher’s exact test or Wilcoxon’s rank-sum test. BL, Burkitt lymphoma; DLBCL, diffuse large B-cell lymphoma; EBV, Epstein–Barr virus; HD, Hodgkin’s disease.

^aA tumor mass appeared in the cervical spinal cord about 17 months later, although recurrence was not confirmed pathologically.

^bIPI, International Prognostic Index for non-HD [stage, lactate dehydrogenase (LDH), performance status, age]. SD, standard deviation.

poor prognosis in non-Hodgkin lymphomas [20], hypomethylated *FGF5* may similarly influence the prognosis in HIV-associated lymphomas. Note that several significant pathways related to cell adhesion were found (Supplementary Table 2, <http://links.lww.com/QAD/A441>). Of these, those involving laminins, collagens, N-cadherin, and caveolin2 were significantly hypomethylated in HIV-associated lymphomas, suggesting that their increased expression initiates and promotes tumors and results in a poor prognosis [21–23]. These data partly support the poor prognosis seen in HIV-associated lymphomas.

Clustering analysis of the Cohort II data obtained using Cancer Panel I placed recurrent or suspicious and nonrecurrent HIV-associated lymphomas into separate groups, suggesting that recurrence of HIV-associated lymphomas is attributable to specific gene regulation involving DNA methylation. *PTCH2*, which was used for validation, was a significant component of the Hedgehog signaling pathway (Supplementary Table 3, <http://links.lww.com/QAD/A441>), which is related to relapse rate in carcinomas [24]. The data imply that the DNA methylation profile is a good indicator of prognosis. Recently, specific methylation targets have been reported as candidates for new biomarkers of prognosis or metastasis [25,26]. Careful determinations in more cases will identify biomarkers for recurrence in HIV-associated lymphomas.

To our knowledge, this is the first report using molecular technology to distinguish HIV-associated lymphomas from non-HIV lymphomas. Our findings contribute to the understanding of HIV-associated lymphomagenesis and suggest new prognostic biomarkers.

Acknowledgements

This study was supported by a Grant-in-Aid for Research on Advanced Medical Technology from the Ministry of Health, Labor, and Welfare of Japan (Grant number: H22-AIDS-I-002, and H25-AIDS-I-002), a Grant for International Health Research (104-A), Center of Excellence Research from the Ministry of Education, Sports, Culture, Science, and Technology of Japan, and Grants-in-Aid from the Japan Science and Technology Agency.

Author contributions: S.H. is responsible for the study design and the clinical data. S.H., M.S. and Y.I. planned the experiments; J.T. and S.O. analyzed the clinical information; T.H. and M.M. prepared the pathology samples; A.M. prepared gDNA from clinical samples; A.M. and M.S. analyzed the microarray data; N.T., M.Y., and L.Y. were in charge of the statistical analyses and bioinformatics; and M.S. and S.H. wrote the article. All authors discussed the results of the article.

Conflicts of interest

There are no conflicts of interest.

References

1. Cote TR, Biggar RJ, Rosenberg PS, Devesa SS, Percy C, Yellin FJ, et al. **Non-Hodgkin's lymphoma among people with AIDS: incidence, presentation and public health burden.** *Int J Cancer* 1997; **73**:645–650.
2. Raphael M, Said J, Borisch B, Cesarman E, Harris NL. **Lymphomas associated with HIV infection.** In: Swerdlow SH, Campo E, Harris NL, Jaffe E, Pillar SA, Stein H, et al. editors. *WHO Classification of tumours of haematopoietic and lymphoid tissues*. 4th ed. Lyon: IARC; 2008. pp. 340–342.
3. Simard EP, Engels EA. **Cancer as a cause of death among people with AIDS in the United States.** *Clin Infect Dis* 2010; **51**:957–962.
4. Borinstein SC, Conerly M, Dzieciatkowski S, Biswas S, Washington MK, Trobridge P, et al. **Aberrant DNA methylation occurs in colon neoplasms arising in the azoxymethane colon cancer model.** *Mol Carcinog* 2010; **49**:94–103.
5. Martin-Subero JL, Kreuz M, Bibikova M, Bentink S, Ammerpohl O, Wickham-Garcia E, et al. **New insights into the biology and origin of mature aggressive B-cell lymphomas by combined epigenomic, genomic, and transcriptional profiling.** *Blood* 2009; **113**:2488–2497.
6. Richter J, Ammerpohl O, Martin-Subero JL, Montesinos-Rongen M, Bibikova M, Wickham-Garcia E, et al. **Array-based DNA methylation profiling of primary lymphomas of the central nervous system.** *BMC Cancer* 2009; **9**:455.
7. Amara K, Trimeche M, Ziadi S, Laatiri A, Hachana M, Korbi S. **Prognostic significance of aberrant promoter hypermethylation of CpG islands in patients with diffuse large B-cell lymphomas.** *Ann Oncol* 2008; **19**:1774–1786.
8. Bibikova M, Barnes B, Tsan C, Ho V, Klotzle B, Le JM, et al. **High density DNA methylation array with single CpG site resolution.** *Genomics* 2011; **98**:288–295.
9. Thirlwell C, Eymard M, Feber A, Teschendorff A, Pearce K, Lechner M, et al. **Genome-wide DNA methylation analysis of archival formalin-fixed paraffin-embedded tissue using the Illumina Infinium HumanMethylation27 BeadChip.** *Methods* 2010; **52**:248–254.
10. Wang D, Yan L, Hu Q, Sucheston LE, Higgins MJ, Ambrosone CB, et al. **IMA: an R package for high-throughput analysis of Illumina's 450K Infinium methylation data.** *Bioinformatics* 2012; **28**:729–730.
11. Davis S, Bilke S. methylumi: Handle Illumina methylation data. Bioconductor R package version 132 2010. <http://www.bioconductor.org/packages/release/bioc/html/methylumi.html> [Accessed 24 September 2012].
12. Bibikova M, Lin Z, Zhou L, Chudin E, Garcia EW, Wu B, et al. **High-throughput DNA methylation profiling using universal bead arrays.** *Genome Res* 2006; **16**:383–393.
13. Hagiwara S, Yotsumoto M, Odawara T, Ajisawa A, Uehira T, Nagai H, et al. **Non-AIDS-defining hematological malignancies in HIV-infected patients: an epidemiological study in Japan.** *AIDS* 2013; **27**:279–283.
14. Teschendorff AE, Menon U, Gentry-Maharaj A, Ramus SJ, Weisenberger DJ, Shen H, et al. **Age-dependent DNA methylation of genes that are suppressed in stem cells is a hallmark of cancer.** *Genome Res* 2010; **20**:440–446.
15. Shiels MS, Pfeiffer RM, Gail MH, Hall HI, Li J, Chaturvedi AK, et al. **Cancer burden in the HIV-infected population in the United States.** *J Natl Cancer Inst* 2011; **103**:753–762.
16. Youssef EM, Chen XQ, Higuchi E, Kondo Y, Garcia-Manero G, Lotan R, et al. **Hypermethylation and silencing of the putative tumor suppressor Tazarotene-induced gene 1 in human cancers.** *Cancer Res* 2004; **64**:2411–2417.
17. Davidson B, Stavnes HT, Risberg B, Nesland JM, Wohlschlaeger J, Yang Y, et al. **Gene expression signatures differentiate adenocarcinoma of lung and breast origin in effusions.** *Hum Pathol* 2012; **43**:684–694.
18. Kloth M, Goering W, Ribarska T, Arsov C, Sorensen KD, Schulz WA. **The SNP rs6441224 influences transcriptional activity and prognostically relevant hypermethylation of RARRES1 in prostate cancer.** *Int J Cancer* 2012; **131**:E897–E904.

19. Casimiro S, Luis I, Fernandes A, Pires R, Pinto A, Gouveia AG, et al. Analysis of a bone metastasis gene expression signature in patients with bone metastasis from solid tumors. *Clin Exp Metastasis* 2012; **29**:155–164.
20. Salven P, Orpana A, Teerenhovi L, Joensuu H. Simultaneous elevation in the serum concentrations of the angiogenic growth factors VEGF and bFGF is an independent predictor of poor prognosis in non-Hodgkin lymphoma: a single-institution study of 200 patients. *Blood* 2000; **96**:3712–3718.
21. Saha S, Lo PK, Duan X, Chen H, Wang Q. Breast tumour initiating cell fate is regulated by microenvironmental cues from an extracellular matrix. *Integr Biol (Camb)* 2012; **4**:897–904.
22. Montenegro RC, de Vasconcellos MC, Barbosa GD, Burbano RM, Souza LG, Lemos TL, et al. A novel o-naphthoquinone inhibits N-cadherin expression and blocks melanoma cell invasion via AKT signaling. *Toxicol In Vitro* 2013; **27**:2076–2083.
23. Ando T, Ishiguro H, Kimura M, Mitsui A, Mori Y, Sugito N, et al. The overexpression of caveolin-1 and caveolin-2 correlates with a poor prognosis and tumor progression in esophageal squamous cell carcinoma. *Oncol Rep* 2007; **18**:601–609.
24. Chaudary N, Pintilie M, Hedley D, Fyles AW, Milosevic M, Clarke B, et al. Hedgehog pathway signaling in cervical carcinoma and outcome after chemoradiation. *Cancer* 2012; **118**:3105–3115.
25. Bougel S, Lhermitte B, Gallagher G, de Flaugergues JC, Janzer RC, Benhattar J. Methylation of the hTERT promoter: a novel cancer biomarker for leptomeningeal metastasis detection in cerebrospinal fluids. *Clin Cancer Res* 2013; **19**:2216–2223.
26. Huang RL, Gu F, Kirma NB, Ruan J, Chen CL, Wang HC, et al. Comprehensive methylome analysis of ovarian tumors reveals hedgehog signaling pathway regulators as prognostic DNA methylation biomarkers. *Epigenetics* 2013; **8**:624–634.

Slow Turnover of HIV-1 Receptors on Quiescent CD4⁺ T Cells Causes Prolonged Surface Retention of gp120 Immune Complexes *In Vivo*

Yasuhiro Suzuki^{1,2*}, Hiroyuki Gatanaga², Natsuo Tachikawa^{2,3}, Shinichi Oka²

1 The Department of Infectious Diseases, Graduate School of Medicine, Tohoku University, Sendai, Japan, **2** AIDS Clinical Center, National Center for Global Health and Medicine, Tokyo, Japan, **3** The Department of Infectious Diseases, Yokohama Municipal Citizen's Hospital, Yokohama, Japan

Abstract

Peripheral blood CD4⁺ T cells in HIV-1⁺ patients are coated with Ig. However, the causes and consequences of the presence of Ig⁺ CD4⁺ T cells remain unknown. Previous studies have demonstrated the rapid turnover of viral receptors (VRs) on lymphoma and tumor cells. The present study investigates the turnover of VRs on peripheral quiescent CD4⁺ T cells (qCD4s), which are the most abundant peripheral blood CD4⁺ T cells. Utilizing pharmacological and immunological approaches, we found that the turnover of VRs on qCD4s is extremely slow. As a result, exposure to gp120 or HIV-1 virions *in vitro* causes gp120 to remain on the surface for a long period of time. It requires approximately three days for cell-bound gp120 on the surface to be reduced by 50%. In the presence of patient serum, gp120 forms surface immune complexes (ICs) that are also retained for a long time. Indeed, when examining the percentages of Ig⁺ CD4⁺ T cells at different stages of HIV-1 infection, approximately 70% of peripheral resting CD4⁺ T cells (rCD4s) were coated with surface VRs bound to slow-turnover gp120-Ig. The levels of circulating ICs in patient serum were insufficient to form surface ICs on qCD4s, suggesting that surface ICs on qCD4s require much higher concentrations of HIV-1 exposure such as might be found in lymph nodes. In the presence of macrophages, Ig⁺ CD4⁺ T cells generated *in vitro* or directly isolated from HIV-1⁺ patients were ultimately phagocytosed. Similarly, the frequencies and percentages of Ig⁺ rCD4s were significantly increased in an HIV-1⁺ patient after splenectomy, indicating that Ig⁺ rCD4s might be removed from circulation and that non-neutralizing anti-envelope antibodies could play a detrimental role in HIV-1 pathogenesis. These findings provide novel insights for vaccine development and a rationale for using Ig⁺ rCD4 levels as an independent clinical marker.

Citation: Suzuki Y, Gatanaga H, Tachikawa N, Oka S (2014) Slow Turnover of HIV-1 Receptors on Quiescent CD4⁺ T Cells Causes Prolonged Surface Retention of gp120 Immune Complexes *In Vivo*. PLoS ONE 9(2): e86479. doi:10.1371/journal.pone.0086479

Editor: Derya Unutmaz, New York University, United States of America

Received: July 10, 2013; **Accepted:** December 9, 2013; **Published:** February 6, 2014

Copyright: © 2014 Suzuki et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by a grant-in-aid for AIDS research from the Ministry of Health, Labor, and Welfare of Japan and by the Japanese Foundation of AIDS Prevention. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: suzuki39@med.tohoku.ac.jp

Introduction

The most immunogenic HIV-1 molecules for the elicitation of an antibody (Ab) response appear to be envelope (env) glycoproteins, and high titers of anti-gp120 and anti-gp41 Abs are observed in HIV-1-infected patients (HIV-1⁺ Pts) [1–3]. However, it is apparent that the neutralizing Ab response in infected patients is weak compared with non-neutralizing HIV Abs [4]. Therefore, non-neutralizing Abs are dominant in the circulation of HIV-1⁺ Pts. Nevertheless, the role of non-neutralizing anti-env Abs in HIV-1 infection remains unclear. More than 95% of the body's CD4⁺ T cells reside in lymphoid tissues, which are the major sites for HIV-1 replication, CD4⁺ T cell depletion [5], and development of anti-env Ab-secreting B cells [3,6]. CD4⁺ T cells continuously travel between the blood, the lymphatic system, and lymph nodes (LNs) and re-circulate into the blood over a period of approximately 1 d [7–9]; therefore, most peripheral blood CD4⁺ T cells are recent emigrants from the LNs. Because a large proportion of HIV-1 is produced in the LNs (10¹⁰–10¹¹ virions/d) [7,10–14], it is assumed that target CD4⁺ T cells in LNs are continuously exposed to high concentrations of HIV-1 as well as anti-env Abs. In the presence of HIV-1⁺ Pt serum, gp120 forms

surface immune complexes (sICs) on HIV-1-infected cells or uninfected cells coated with gp120 *in vitro* [15]. Natural killer (NK) cells have been shown to be able to eliminate gp120/HIV-1-coated or HIV-infected target cells by Ab-dependent cell-mediated cytotoxicity (ADCC) [1,15–20]. However, compared with the distribution in non-lymphoid organs, a relatively small number of NK cells are present in the LNs [21]; therefore, the organs where sICs appear to form on target cells and the effector cells that can eliminate sIC⁺ cells seem to be segregated *in vivo*.

For practical reasons, the dynamics of viral receptors (VRs) and cell-bound gp120/HIV-1 have been extensively studied in both lymphoma and VR-transfected cancer cells. The cell-surface CXCR4 receptors on lymphoma [22,23] and HeLa cells [22–24] are rapidly internalized, and approximately 100% of the cell-surface CXCR4 pools are exchanged every 5 h (in lymphoma cell lines) and 40 min (in HeLa cells). Moreover, cell-bound gp120 has been shown to be internalized in 2 h in Jurkat cells [25], 1 h in CD4-transfected HeLa cells [23,24], and 1–2 h in U937 cells [24]; therefore, the gp120-VR complex is believed to be rapidly removed from the surface of target cells. Consequently, even if gp120/HIV-1-VR complexes form on CD4⁺ T cells *in vivo*, it has

been thought that the complex would disappear from the cell surface before encountering ADCC effector cells. Collectively, it is believed that cell-bound gp120 or HIV-1 on VRs on CD4⁺ T cells have a limited effect on the destruction of HIV-1-exposed cells *in vivo*. In contrast, substantial percentages of CD4⁺ T cells in HIV-1⁺ Pts are shown to be coated with Ig [26,27]. Because the gp120-VR complex was thought to be rapidly removed from the cell surface, it was also believed that sICs on CD4⁺ T cells in HIV-1⁺ Pts mainly reflect the non-specific attachment of Ig-virion complexes (known as circulating immune complexes; cICs) in serum to the cellular surface [28].

The most abundant HIV-1 target cells *in vivo* are quiescent CD4⁺ T cells (qCD4s) because they comprise more than 90% of both peripheral and lymphoid T cells [14,29]. However, the dynamics of cell-surface molecules on quiescent cells are generally shown to be slower than on cancer or activated cells [30]. Furthermore, qCD4s have been shown to have unique biological characteristics, particularly the possession of static cortical actin barriers [31,32] and abundant expression of SAMHD1, a deoxynucleoside triphosphate triphosphohydrolase, to prevent reverse transcription of HIV-1 RNA [33].

Here, we first reevaluated the turnover dynamics of VRs in qCD4s compared with lymphoma cells. We then examined the dynamics of cell-bound gp120 in qCD4s. gp120/HIV-1-exposed qCD4s were further exposed to anti-env Abs to form sICs and to examine their pathological effects. We also investigated the characteristics of sICs on CD4⁺ T cells purified from HIV-1-infected Pts and conducted a longitudinal analysis of the changing levels of sIC⁺ CD4⁺ T cells in peripheral blood from HIV-1⁺ Pts under various conditions.

Results

Slow Turnover of VRs in Dense Resting CD4⁺ T Cells

We first thoroughly reevaluated the turnover dynamics of VRs and cell-bound gp120 or HIV-1 on qCD4s by employing highly purified dense resting CD4⁺ T cells (drCD4s) from healthy donors. drCD4s are purified from resting CD4⁺ T cells as a dense fraction using discontinuous density gradients of Percoll (see **Materials and Methods**) [34]. We have previously shown that these drCD4s are largely in the G₀ phase of the cell cycle, do not produce detectable cytokines, and are highly resistant to spontaneous cell death; therefore, drCD4s are a useful tool for observing biological responses over a long period while avoiding a decrease in viability and spontaneous cell activation in cell culture [34].

To investigate how the dynamics of the receptor are influenced by cellular state, we first examined the effect of cellular activation on VR surface expression. In agreement with previous studies [34–36], CXCR4 was rapidly internalized following anti-CD3 Ab-induced activation (**Fig. 1A left and 1B**). In contrast, CD4 expression remained virtually unaffected by anti-CD3 Ab treatment (**Fig. 1A right**). The addition of IL-2 or anti-CD28 Ab exposure along with anti-CD3 Ab treatment only had a marginal effect on initial CXCR4 internalization; however, these additional stimuli slightly enhanced the restoration of surface CXCR4 expression after 72 h (**Fig. 1A left**). In contrast, surface CD4 expression remained unaffected (**Fig. 1A right**). Collectively, we conclude that anti-CD3 Ab-triggered CD4⁺ T cell activation significantly alters CXCR4 dynamics but only has a marginal effect on CD4 dynamics.

We next evaluated VR turnover kinetics in qCD4s compared with lymphoma cells or activated cells. In these experiments, in addition to cycloheximide (a protein synthesis inhibitor), retrograde trafficking of internalized molecules and anterograde

transport from the endoplasmic reticulum to the Golgi complex was blocked using Brefeldin-A (BFA) (see the schematic description of the inhibitors of protein turnover in **Fig. S1A**). Previous studies have shown that the effect of BFA on protein transport is greatest soon after treatment; therefore, the rate of reduction of VRs was determined after the first 2–3 h. CXCR4 expression was modestly reduced (approximately 25% in a 3-h assay) in T cell lymphoma A3.01 cells by BFA (**Fig. S1B left**). Some of the BFA-induced CXCR4 reduction (approximately 10% after 3 h) was caused by blockage of the transport of newly synthesized molecules, as shown using treatment with cycloheximide (**Fig. S1B left**). Because the transport of newly synthesized CXCR4 appeared to be suppressed by BFA, cycloheximide plus BFA did not produce any additive effects on CXCR4 reduction. Therefore, in agreement with a previous report [24], CXCR4 expression levels in A3.01 cells appear to be maintained by both recycling and replacement at relatively rapid rates. Next, we utilized qCD4s that were activated by 72 h of anti-CD3 Ab plus anti-CD28 Ab exposure but still had low CXCR4 expression (**Fig. 1A left**). In contrast with A3.01 cells, when we examined the pharmacological effects on VRs after 72 h of anti-CD3 Ab plus anti-CD28 Ab activation in qCD4s, CXCR4 expression was significantly reduced by both BFA (approximately 70% after 2.5 h) and cycloheximide treatments (approximately 60% after 2.5 h) (**Fig. S1B right; see also Fig. S1A**). Again, because the transport of newly synthesized CXCR4 appeared to be suppressed by BFA, cycloheximide plus BFA did not show any additive effects on CXCR4 reduction, suggesting that the reduced CXCR4 surface expression on activated qCD4s after 72 h of exposure was linked to rapid turnover due to greater degradation of CXCR4 than replacement by both newly synthesized and recycled molecules. In agreement with these results, RT-PCR analysis revealed that CXCR4 mRNA transcripts increased approximately 3.5-fold in activated qCD4s relative to qCD4s (**data not shown**). Utilizing confocal microscopy, we found that a significant portion of intracellular CXCR4 colocalized with the late endosomal/lysosomal marker LAMP-1 and the early endosomal marker Rab5 [37] in activated qCD4s, whereas, such colocalization was not observed in qCD4s (**data not shown**), suggesting the degradation of the CXCR4 proteins that are enhanced in activated qCD4s. Collectively, these results suggest that the CXCR4 turnover rate was enhanced because protein degradation predominated over replacement by both newly synthesized and recycled molecules; consequently, CXCR4 expression remains low in activated qCD4s.

In contrast, exposure of qCD4s to BFA minimally reduced CXCR4 expression levels following 16 h of incubation (approximately 3% and 20% after 2.5 h and 16 h, respectively), and exposure to both cycloheximide and Actinomycin-D (ActD), a DNA transcription suppressor, did not affect CXCR4 expression levels (**Fig. 1C left**). Again, cycloheximide plus BFA did not show any additive effects on CXCR4 expression. These results suggest that CXCR4 expression in qCD4s is stable and that a small fraction (approximately 3% over 3 h) of surface CXCR4 is continually internalized and recycled back to the surface. In contrast, CD4 expression in qCD4s was unaffected by exposure to BFA, cycloheximide, and ActD after 24 h (**Fig. 1C right**), indicating that CD4 turnover in qCD4s is more stable than CXCR4. Given that the inhibitors' effects on protein transport/synthesis are not complete, it seems reasonable to propose that the actual turnover rate of VRs may be faster.

To further confirm the CXCR4 turnover results described above, we monitored CXCR4 turnover by employing T22, a peptide that binds to CXCR4 and blocks the binding of anti-CXCR4 mAb 12G5 [38]. The binding of 12G5 to CXCR4 was




Figure 1. HIV-1/gp120 remains on the surface of qCD4s for a long period of time due to slow VR turnover. (a, b) Time course of surface VR expression (a) and representative FACS of CXCR4/CCR5 expression on qCD4s (b) following a variety of activation stimuli. (c) The effect of BFA (10 μ M), cycloheximide (50 μ M) and ActD (20 μ M) on the surface expression of CXCR4 (left) and CD4 (right) on qCD4s. (d, e) Confocal micrographs of CD4, CXCR4, and gp120 in qCD4s that were exposed or not exposed to the indicated strain of gp120 or HIV-1 before (d) or after (e)

16 h of anti-CD3 Ab exposure. qCD4s with (e) or without (d) permeabilization were stained with anti-CD4 goat polyclonal Abs (Cy3, red), anti-CXCR4 mouse mAbs (Qdot 655, blue), and anti-gp120 rabbit antiserum (Cy2, green). (f) Time course of cell-bound gp120, sICs (left panel), or surface CD4 expression (right panel) on gp120_{IIIIB}-pulsed or untreated qCD4s. The gp120_{IIIIB}-pulsed qCD4s were further incubated with HIV-1⁺ Pt serum (Pt-serum) to form sICs or untreated and cultured in the absence or presence of anti-CD3 Abs. The effect of T22 pre-exposure on cell-bound gp120_{IIIIB} in anti-CD3 Ab stimulation (α CD3+T22) was also examined. (g) Time course of cell-bound gp120, sICs, or surface CD4 expression on HIV-1_{Lai} (Lai)- (left), HIV-1_{BaL} (BaL)-, or clinical isolate (Clinical)- (right) pulsed qCD4s. HIV-1-pulsed qCD4s were further incubated with HIV-1⁺ Pt serum (Pt-serum) to form sICs or untreated and cultured in the absence or presence of anti-CD3 Abs. The effect of enfuvirtide (Enf) exposure was also examined. (h) The amount and location of cell-bound gp120 in gp120_{IIIIB}-pulsed qCD4s that were cultured in the absence or presence of anti-CD3 Abs were assessed by confocal microscopy (upper) or by western blotting (bottom). The lower numbers indicate the value by densitometry. (i) Time course of chemotaxis inhibition on gp120_{IIIIB}-, gp120_{BaL}- (upper), or HIV-1_{Lai}- (bottom) pulsed qCD4s. Chemotaxis of gp120- or HIV-1-pulsed or non-pulsed qCD4s toward the indicated chemokines was evaluated using a transwell assay. Bars, SD. The data here are representative of at least three independent experiments.

doi:10.1371/journal.pone.0086479.g001

initially completely blocked by T22 exposure but gradually recovered (Fig. S1C). The duration and level of T22 occupancy on CXCR4 molecules over time is mainly influenced by four factors, namely CXCR4 internalization and degradation, CXCR4 recycling, *de novo* CXCR4 synthesis, and a steady level of T22 detachment from CXCR4; however, of these four factors, the rate of T22 detachment from CXCR4 is less likely to be affected by the level of CXCR4 turnover. Therefore, the duration of T22-CXCR4 occupancy should represent the level of intracellular replacement and degradation of CXCR4. The calculated times required for T22 occupation to fall to 50% of CXCR4 molecules were approximately 30 h, 8 h, and 8.5 h in qCD4s, activated qCD4s, and A3.01 cells, respectively (Fig. S1C). Although there are concerns that partial inhibition and/or cytotoxicity of the inhibitors may interfere with an accurate determination of VR dynamics, an estimation performed with either inhibitors or T22 showed similar trends in A3.01 cells, qCD4s, and activated qCD4s. Therefore, we conclude that CXCR4 turnover in qCD4s is truly stable, with a small fraction of CXCR4 slowly recycled, whereas CXCR4 turnover is significantly more rapid in both lymphoma and activated qCD4s. In this respect, it has been shown that a rigid layer of cortical actin exists in qCD4s [31], which may be partially linked to the slow turnover of VRs in qCD4s.

Slow Turnover of Cell-bound HIV-1/gp120 on Dense Resting CD4⁺ T Cells

We next studied the dynamics of cell-bound gp120 in qCD4s before and after activation. Cells were exposed to the indicated subtypes of gp120 or HIV-1 for 30 min, thoroughly washed, and cultured at 37°C. As expected, confocal microscopy revealed that gp120, CD4, and CXCR4 colocalized on the surface in X4-gp120 (gp120_{IIIIB})- or X4-HIV-1 (HIV-1_{Lai})-exposed qCD4s, whereas CXCR4 was not recruited to gp120-CD4 complexes on R5-gp120 (gp120_{BaL})-exposed qCD4s (Fig. 1D). The rates of surface gp120 reduction in both X4-gp120- and X4-HIV-1-exposed qCD4s were extremely slow, and we calculated that cell-bound gp120 was reduced to 50% at the surface approximately 3 d after exposure (Fig. 1F left, open circles, 1G left, open circles, and Fig. S2A). However, the rate of reduction of cell-bound X4-gp120 was rather rapid during the initial 20 h in both X4-gp120- and X4-HIV-1-exposed qCD4s (Fig. 1F left, open circles and 1G left, open circles). This observation may reflect that the rate of VR turnover is slightly enhanced by gp120 inducing VR-mediated signaling. In contrast, anti-CD3 Ab stimulation of X4-gp120- or X4-HIV-1-exposed qCD4s led to the rapid internalization of gp120-CD4-CXCR4 ternary complexes (see induction of gp120 internalization in Fig. 1F left and G left, open squares; see that CD4 and CXCR4 co-internalization only occurred in X4-gp120- or X4-HIV-1-exposed cells in Fig. 1E). Additionally, anti-CD3 Ab stimulation only induced CD4 down-regulation in X4-gp120- or X4-HIV-1-treated cells, which indicates that CD4 co-mobilizes

with CXCR4 through gp120 (Fig. 1E, 1F right and 1G left; see also Fig. S2A). However, T22 pre-exposure, which inhibits the association of gp120 with CXCR4, abrogated anti-CD3 Ab-induced gp120 internalization (Fig. 1F, closed triangles), suggesting that association with CXCR4 is essential for gp120 internalization. Therefore, the rapid internalization of gp120 in anti-CD3 Ab-stimulated qCD4s was mainly directed by internalized CXCR4.

When cell-bound gp120 stability was assessed by western blotting, approximately 70% of the gp120 was detected on the surface of qCD4s after 36 h of cell culture (Fig. 1H, lower panel; see also gp120 colocalizes with CD4 and CXCR4 after 36 h of cell culture in Fig. 1H, upper panel), and the results were comparable to those from FACS (see Fig. 1F left, open circles). In contrast, approximately 95% of the gp120 that was initially bound was degraded within 36 h of anti-CD3 Ab treatment (Fig. 1H, lower panel; see also Fig. 1F left, open squares). Because CCR5 expression was limited to approximately 10% of peripheral qCD4s (see Fig. 1B), most of the qCD4-bound R5-gp120 could bind to CD4 alone, and the dynamics of cell-bound R5-gp120 followed the dynamics of CD4. As anticipated, R5-HIV-1 (both the experimental strain (BaL) and the clinical isolate (Clinical)) on the qCD4 cell surface was retained for slightly longer than X4-HIV-1 (Fig. 1G right, open circles and open diamonds). Collectively, these results clearly demonstrate that irrespective of HIV-1 subtype, gp120 bound to qCD4s remains on the surface for a long time.

Because gp120 can be rapidly dissociated from virions by soluble CD4 [39], we hypothesized that gp120 dissociates from virions after HIV-1 becomes attached to surface CD4 and persists on VRs. To investigate this possibility, we used enfuvirtide (Enf) to inhibit virus and target membrane fusion [40], and we examined the dynamics of gp120 and p24, an HIV-1 capsid antigen, in HIV-1-exposed qCD4s. The dynamics of surface gp120 within 24 h of HIV-1 exposure in qCD4s were comparable between Enf-treated and untreated cells (Fig. 1G left, compare open circles with open diamonds). Similarly, western blotting analysis revealed that approximately 70% of the initially attached p24 disappeared from both Enf-treated and untreated cells after 24 h (Fig. S2B). However, early HIV-1 DNA products were only detected in untreated qCD4s (data not shown). Therefore, irrespective of HIV-1 cell entry or uncoating, binding of HIV-1 to VRs appears to lead to the dissociation of gp120 from HIV-1, and the dissociated gp120 remains on VRs.

We then inquired whether gp120 directly associates with VRs for a prolonged period or associates with another molecule on the cell surface when cells are exposed to gp120 or HIV-1. Because the spatial resolution of confocal microscopy is not sufficient to determine gp120 directly associates with VRs accurately, we employed a transwell chemotaxis assay to examine the effect of X4-HIV-1 or gp120 exposure on CXCL12-induced chemotaxis.

Given that X4-gp120 blocks the binding of CXCL12 to CXCR4, the initial exposure of qCD4s to X4-gp120 or X4-HIV-1 abrogated CXCL12-induced chemotaxis (**Fig. 1I, see open triangles (upper panel) and open squares (lower panel)**). However, CXCL12-induced chemotaxis was not suppressed by R5-gp120 exposure (**Fig. 1I upper panel, see open squares**). Consistent with the kinetics of cell-bound X4-gp120 or X4-HIV-1 in qCD4s, inhibition of CXCL12-induced chemotaxis was sustained for more than 3 d. In contrast, the migration of qCD4s toward CCL19 was not abrogated in X4-gp120- or X4-HIV-1-exposed qCD4s, serving as a control for the functional integrity of the cells to respond to other chemokines. These results cannot rule out the possibility that gp120 binds to other cell surface molecules but do clearly show that cell-bound X4-gp120 or X4-HIV-1 forms gp120-CD4-CXCR4 ternary surface complexes on qCD4s for prolonged periods.

Slow Turnover of Ig-gp120 sICs on Dense Resting CD4⁺ T Cells

Several studies have shown that in the presence of serum from HIV-1⁺ Pts, sICs can form on HIV-1-infected cells or gp120-exposed uninfected cells (e.g., [15]). We tested whether patient serum contains sufficient anti-env Abs to allow the formation of sICs on gp120-pre-exposed qCD4s. Although the amount of sICs on qCD4s was proportional to the concentration of exposed gp120, the levels of sICs varied among patients, reflecting different levels of anti-env Abs in the serum of HIV-1⁺ Pts (**Fig. S3 right panel**; see also the **left panel**, which demonstrates the relationship between the concentration of exposed gp120 and the level of CD4 occupancy by gp120 by utilizing the gp120-blocking anti-CD4 mAb Leu3a and the gp120-non-blocking anti-CD4 mAb CD4V4). We then examined the turnover of cell-bound gp120. The turnover of cell-bound gp120 was not significantly affected, even in the presence of patient serum (**compare open circles vs. open triangles (Fig. 1F and 1G) or open diamonds vs. open squares (Fig. 1G right)**). Collectively, serum from HIV-1⁺ Pts always contained sufficient levels of anti-env Abs to form sICs, and the kinetics of surface gp120 were extremely slow in qCD4s regardless of whether cell-bound gp120 formed sICs.

Resting CD4⁺ T Cells from Acutely and Chronically HIV-1-infected Subjects are Coated with IgG and IgM

If gp120 turnover on qCD4s *in vivo* is similar to that observed *in vitro*, we should detect sICs on qCD4s from the peripheral blood of HIV-1⁺ Pts. For technical convenience, to easily detect qCD4s by FACS, we examined the presence of sICs in peripheral blood CD25⁻ CD69⁻ CD4⁺ CD3⁺ cells (designated here as resting CD4⁺ T cells; rCD4s). We utilized biotinylated anti-IgG F(ab')₂ and/or anti-IgM F(ab')₂ Abs to prevent non-specific surface binding through the Fc portion. Sixteen individuals with asymptomatic chronic HIV-1 infection, four individuals with acute HIV-1 infection, and ten healthy individuals were examined.

In agreement with previous studies[26–28], means of 78.18±11.77% (± SD) and 42.18±19.73% (± SD) of peripheral blood rCD4s from sixteen chronic HIV-1⁺ Pts stained positive with anti-IgG and anti-IgM, respectively, whereas no sIC⁺ rCD4s were detected in healthy donors (**Fig. 2A and B**). In contrast, means of 48.22±22.69% (± SD) and 72.10±9.20% (± SD) of peripheral blood rCD4s from four acute HIV-1⁺ Pts were positive for anti-IgG and anti-IgM, respectively (**Fig. 2A**). To more clearly demonstrate that peripheral blood rCD4s from HIV-1⁺ Pts were coated with Ig, rCD4s were purified from HIV-1⁺ Pts, lysed, and

immunoblotted with anti-IgG Ab. As shown in **Fig. 2C**, IgG was only detected in rCD4 lysates from HIV-1⁺ Pts but not from healthy donors. The level of IgG detected by immunoblotting correlated with the mean fluorescence intensities (MFIs) of surface IgG on rCD4s as detected by FACS (**Fig. 2C**, the numbers above the panel indicate the MFIs of IgG in rCD4s). Furthermore, utilizing confocal microscopy, we found that Igs colocalized with surface CD4 on rCD4s purified from HIV-1⁺ Pts (**Fig. 2D**, see three-dimensional reconstruction confocal micrograph). These results collectively confirm that Igs are attached to CD4 on peripheral blood rCD4s in HIV-1⁺ Pts.

cICs in the Serum of Viremic HIV-1⁺ Pts are Sufficient to form sICs on B Cells but not Resting CD4⁺ T Cells

It has been reported that B cells and T cells from HIV-1⁺ Pts are covered with complement-opsonized cICs [41] or auto-Abs [27]. Therefore, we tested whether serum from HIV-1⁺ Pts contains sufficient levels of cICs or auto-Abs to form sICs on rCD4s. Before proceeding with the experiments, we first sought to determine whether complement receptors or the Fc receptor were expressed in B cells and rCD4s. As shown in **Table S1**, complement receptors (CRs) 1, 2, and 3 and FcγRII were expressed on B cells but not rCD4s from both healthy donors and HIV-1⁺ Pts; these findings suggest that cICs with or without complement opsonization could theoretically bind to B cells through the Fc region of IgG to Fc receptors and/or complement opsonization to CRs but not to rCD4s.

Because B cells intrinsically express IgG and/or IgM on the cell surface, it is difficult to directly detect surface cICs using anti-IgG or anti-IgM Abs without interfering with their expression. Therefore, to clearly identify cell-bound cICs, we utilized a purified CD20⁺ IgG^{dull} IgM^{dull} population from the blood of healthy donors (see **Fig. S4B, upper panel**). When this purified subset of B cells (CD20⁺ IgG^{dull} IgM^{dull}) was incubated with the patients' serum, the percentages of B cells coated with cICs as detected by anti-IgG or anti-IgM Abs approximately paralleled viral loads (VLs) in the serum samples (**Fig. S4A and B**). In contrast, no cIC binding was detected on B cells incubated with serum from healthy donors or aviremic HIV-1⁺ Pts. Similarly, an *in situ* hybridization assay demonstrated that HIV-1 RNA was detected on all B cells incubated with serum from viremic HIV-1⁺ Pts but not from healthy donors or aviremic HIV-1⁺ Pts (**Fig. S4C and D**). In contrast, when rCD4s were incubated with serum from HIV-1⁺ Pts, virtually no cICs bound to the rCD4 cell surface (**Fig. S4E left column**). However, when gp120 pre-exposed cells were utilized, sICs were easily detected on all rCD4s incubated with serum from HIV-1⁺ Pts but not from healthy donors (**Fig. S4E middle column**). When rCD4s were exposed to 10 mg/ml of purified IgG from serum from HIV-1⁺ Pts, no Ig was detected on the rCD4s (**Fig. S4E right column**). These results suggest that CR or FcγRII expression is critical for efficient cIC binding to the cell surface. Collectively, we can conclude that cIC levels in serum from viremic HIV-1⁺ Pts are insufficient to form sICs on rCD4s and that auto-Abs to rCD4s are either non-existent or below the limit of detection in serum from HIV-1⁺ Pts.

The Dynamics of sICs on Resting CD4⁺ T Cells from HIV-1-infected Subjects show Similar Kinetics to gp120-Igs

We attempted to clarify whether sIC⁺ rCD4s in the peripheral blood of HIV⁺ Pts were also caused by cell-bound gp120. To this end, we first studied the dynamics of sICs in rCD4s purified from HIV⁺ Pts and sought to determine whether the dynamics are similar to Ig-gp120-VRs. The estimated mean duration of a 50%




Figure 2. sICs of IgG or IgM on purified rCD4s from HIV-1⁺ Pts is molecularly linked to surface CD4 and shows slow turnover. (a) Summary of the percentages of IgG⁺ rCD4s or IgM⁺ rCD4s in healthy individuals (H), acute HIV-1⁺ individuals (A), and chronic pre-symptomatic HIV-1⁺ individuals (C) before, after 3 yrs of complete suppression of VL (<50 copies/ml) with ART (Treated), or untreated for 3 yrs (Untreated). Bars, SD. (b) Representative FACS of IgG expression on rCD4s from HIV-1⁺ Pts. (c) Anti-IgG Ab immunoblotting of purified HIV-1⁺ Pt rCD4 lysates. For the