

**雑誌：発表者氏名，論文タイトル名，発表誌名，巻号，ページ，出版年**

【全国油症班が厚生労働科学研究費の補助を受けて刊行】

責任編集者 古江増隆，

福岡医学雑誌特集号「油症と PCB 及びダイオキシン関連化合物」研究報告第 24 集 .

福岡医学雑誌，104(4) : 71-169

古江増隆（厚生労働省全国油症治療研究班 班長）.序言，

福岡医学雑誌 104(4) : 71-72 , 2013

辻 博

油症における特異抗核抗体の検討.

福岡医学雑誌104(4):73-77, 2013

松本伸哉、赤羽学、神奈川芳行、梶原淳睦、内博史、古江増隆、今村知明 .

油症患者におけるダイオキシン類異性体ごとの症状ならびに異性体間の濃度と半減期の関係、

福岡医学雑誌 104(4):78-84, 2013

内 博史、古江増隆 .

油症患者における血清アディポカイン濃度に関する研究.

福岡医学雑誌104(4):85-87, 2013

鍬塚 大、小池雄太、清水和宏、宇谷厚志 .

油症患者におけるIL-21、TGF- の検討 .

福岡医学雑誌104(4):88-90, 2013

小池雄太、鍬塚 大、清水和宏、宇谷厚志 .

油症認定患者における制御性 T 細胞産生サイトカイン IL-10、IL-35 の検討 .

福岡医学雑誌104(4):91-94, 2013

吉村俊朗、川崎涼子、吉村俊祐、宮崎禎一郎、向井晃弘、米澤武人 .

長崎県力ネミ油症検診者の血清尿酸値.

福岡医学雑誌104(4):95-99, 2013

川崎五郎、吉富 泉、梅田正博 .

油症患者における頸関節症に関する臨床的検討 .

福岡医学雑誌104(4):100-103, 2013

**雑誌：発表者氏名，論文タイトル名，発表誌名，巻号，ページ，出版年**

梶原淳睦、戸高尊、平川博仙、堀就英、安武大輔、宮脇崇、飛石和大、高尾佳子、平田輝昭、内博史、古江増隆。

血液中ダイオキシン，PCB 類濃度測定のクロスチェック。

福岡医学雑誌104(4):104-109, 2013

戸高尊、内博史、平川博仙、梶原淳睦、古江増隆。

油症患者血液中ダイオキシン類分析における新しい大量注入法の検討。

福岡医学雑誌104(4):110-117, 2013

戸高尊、内博史、平川博仙、高尾佳子、梶原淳睦、古江増隆。

2004 年から2010 年の期間における油症患者血液中ダイオキシン類濃度の推移。

福岡医学雑誌104(4):118-127, 2013

飛石和大、鈴木茂、戸高尊、平川博仙、堀就英、梶原淳睦、平田輝昭、飯田隆雄、内博史、古江増隆。

LC/MS/MS を用いた血液試料中の水酸化ポリ塩化ビフェニル(OH-PCBs)測定法の改良。

福岡医学雑誌104:128-134, 2013

飛石和大、鈴木茂、戸高尊、平川博仙、堀就英、梶原淳睦、平田輝昭、飯田隆雄、内博史、古江増隆。

2010 年度油症認定患者血液中の水酸化ポリ塩化ビフェニル(OH-PCBs)濃度。

福岡医学雑誌104(4):135-142, 2013

武田知己、服部友紀子、藤井美彩子、田浦順樹、石井祐次、山田英之。

ダイオキシン母体曝露による胎児副腎ステロイド合成系への影響と性差。

福岡医学雑誌104(4):143-151, 2013

堀就英、山本貴光、石黒靖尚、飯田隆雄、梶原淳睦、平田輝昭、内博史、古江増隆。

血液中PCB 異性体分離分析におけるアルカリ分解温度の検討。

福岡医学雑誌104(4):152-160, 2013

太田 千穂、原口浩一、加藤善久、遠藤哲也、古賀信幸。

2,2,3,4,5,5-六塩素化ビフェニル(CB146)のin vitro代謝の動物種差。

福岡医学雑誌104(4):161-169, 2013

**雑誌：発表者氏名，論文タイトル名，発表誌名，巻号，ページ，出版年**

**【古江 増隆】**

Mitoma C, Mine Y, Utani A, Imafuku S, Muto M, Akimoto T, Furue M, Uchi H. Current skin symptoms of Yusho patients exposed to high levels of 2,3,4,7,8-pentachlorodibenzofuran and polychlorinated biphenyls in 1968. *Chemosphere* in press

Liu G, Asanoma K, Takao T, Tsukimori K, Uchi H, Furue M, Kato K, Wake N. Aryl hydrocarbon receptor SNP-130 C/T associates with dioxins susceptibility through regulating its receptor activity and downstream effectors including interleukin 24. *Toxicol lett* 2015 Jan; 232(2): 384-392.

Kenjiro Takei, Chikage Mitoma, Akiko Hashimoto-Hachiya, Hiroshi Uchi, Masakazu Takahara, Gaku Tsuji, Makiko Kido-Nakahara, Takeshi Nakahara, Masutaka Furue. Antioxidant soybean tar Glyteer® rescues Th2-mediated downregulation of filaggrin expression via aryl hydrocarbon receptor.

*J Dermatol* in press

Kenjiro Takei, Chikage Mitoma, Akiko Hashimoto-Hachiya, Masakazu Takahara, Gaku Tsuji, Takeshi Nakahara, Masutaka Furue. Galactomyces ferment filtrate prevents Th2-mediated reduction of filaggrin in an aryl hydrocarbon receptor-dependent manner. *Clin Exp Dermatol* in press

Taura J, Takeda T, Fujii M, Hattori Y, Ishii Y, Kuroki H, Tsukimori K, Uchi H, Furue M, Yamada H. 2,3,4,7,8-Pentachlorodibenzofuran is far less potent than 2,3,7,8-tetrachlorodibenzo-p-dioxin in disrupting the pituitary-gonad axis of the rat fetus. *Toxicol Appl Pharmacol*. 2014 Sep 16. pii: S0041-008X(14)00326-3. doi: 10.1016/j.taap.2014.09.001.

Onozuka D, Hirata T, Furue M. Net survival after exposure to polychlorinated biphenyls and dioxins: The Yusho study. *Environ Int*. 2014 Dec;73:28-32.

Furue M, Takahara M, Nakahara T, Uchi H. Role of AhR/ARNT system in skin homeostasis. *Arch Dermatol Res*. 2014 Nov;306(9):769-79.

Wu Z, Uchi H, Morino-Koga S, Shi W, Furue M. Resveratrol inhibition of human keratinocyte proliferation via SIRT1/ARNT/ERK dependent downregulation of aquaporin 3. *J Dermatol Sci*. 2014 Jul;75(1):16-23.

Wu Z, Uchi H, Morino-Koga S, Nakamura-Satomura A, Kita K, Shi W, Furue M. Z-Ligustilide inhibits benzo(a)pyrene-induced CYP1A1 upregulation in cultured human keratinocytes via ROS-dependent Nrf2 activation. *Exp Dermatol*. 2014 Apr;23(4):260-5.

**雑誌：発表者氏名，論文タイトル名，発表誌名，巻号，ページ，出版年**

Kuwatsuka Y, Shimizu K, Akiyama Y, Koike Y, Ogawa F, Furue M, Utani A. Yusho patients show increased serum IL-17, IL-23, IL-1 $\beta$ , and TNF $\alpha$  levels more than 40 years after accidental polychlorinated biphenyl poisoning. *J Immunotoxicol.* 2014 Jul-Sep;11(3):246-9.

三苦千景、内 博史、古江増隆. 油症の診断と最新の知見. *食品衛生研究* 64(11): 15-21, 2014

Kohda F, Takahara M, Hachiya A, Takei K, Tsuji G, Yamamura K, Furue M. Decrease of reactive oxygen species and reciprocal increase of nitric oxide in human dermal endothelial cells by *Bidens pilosa* extract: a possible explanation of its beneficial effect on livedo vasculopathy. *J Dermatol Sci.* 2013 Oct;72(1):75-7.

Morino-Koga S, Uchi H, Tsuji G, Takahara M, Kajiwara J, Hirata T, Furue M. Reduction of CC-chemokine ligand 5 by aryl hydrocarbon receptor ligands. *J Dermatol Sci.* 2013 Oct;72(1):9-15.

Matsumoto S, Akahane M, Kanagawa Y, Kajiwara J, Todaka T, Yasukawa F, Uchi H, Furue M, Imamura T. Individuals' half-lives for 2,3,4,7,8-penta-chlorodibenzofuran (PeCDF) in blood: Correlation with clinical manifestations and laboratory results in subjects with Yusho. *Chemosphere.* 2013 Aug;92(7):772-7.

Tsujimoto S, Ishida T, Takeda T, Ishii Y, Onomura Y, Tsukimori K, Takechi S, Yamaguchi T, Uchi H, Suzuki SO, Yamamoto M, Himeno M, Furue M, Yamada H. Selenium-binding protein 1: Its physiological function, dependence on aryl hydrocarbon receptors, and role in wasting syndrome by 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Biochim Biophys Acta.* 2013 Jun;1830(6):3616-24

Li D, Takao T, Tsunematsu R, Morokuma S, Fukushima K, Kobayashi H, Saito T, Furue M, Wake N, Asanoma K. Inhibition of AHR transcription by NF1C is affected by a single-nucleotide polymorphism, and is involved in suppression of human uterine endometrial cancer. 2013 Oct 10;32(41):4950-9.

Tsukimori K, Uchi H, Tokunaga S, Yasukawa F, Chiba T, Kajiwara J, Hirata T, Furue M. Blood levels of PCDDs, PCDFs, and coplanar PCBs in Yusho mothers and their descendants: association with fetal Yusho disease. *Chemosphere.* 2013 Feb;90(5):1581-8.

松本伸哉、赤羽学、神奈川芳行、梶原淳睦、内博史、古江増隆、今村知明、油症患者におけるダイオキシン類異性体ごとの症状ならびに異性体間の濃度と半減期の関係、福岡医学雑誌104:78-84, 2013

梶原淳睦、戸高尊、平川博仙、堀就英、安武大輔、宮脇崇、飛石和大、高尾佳子、平田輝昭、内博史、古江増隆、血液中ダイオキシン、PCB類濃度測定のクロスチェック、福岡医学雑誌104:104-109, 2013

**雑誌：発表者氏名，論文タイトル名，発表誌名，巻号，ページ，出版年**

戸高尊、内博史、平川博仙、梶原淳睦、古江増隆、油症患者血液中ダイオキシン類分析における新しい大量注入法の検討、福岡医学雑誌104:110-117, 2013

戸高尊、内博史、平川博仙、高尾佳子、梶原淳睦、古江増隆、2004 年から2010 年の期間における油症患者血液中ダイオキシン類濃度の推移、戸高尊・内博史・平川博仙、福岡医学雑誌104:118-127, 2013

飛石和大、鈴木茂、戸高尊、平川博仙、堀就英、梶原淳睦、平田輝昭、飯田隆雄、内博史、古江増隆、LC/MS/MS を用いた血液試料中の水酸化ポリ塩化ビフェニル (OH-PCBs) 測定法の改良、福岡医学雑誌104:128-134, 2013

飛石和大、鈴木茂、戸高尊、平川博仙、堀就英、梶原淳睦、平田輝昭、飯田隆雄、内博史、古江増隆、2010 年度油症認定患者血液中の水酸化ポリ塩化ビフェニル (OH-PCBs) 濃度、福岡医学雑誌104:135-142, 2013

堀就英、山本貴光、石黒靖尚、飯田隆雄、梶原淳睦、平田輝昭、内博史、古江増隆、血液中PCB 異性体分離分析におけるアルカリ分解温度の検討、福岡医学雑誌104:152-160, 2013

Koga T, Ishida T, Takeda T, Ishii Y, Uchi H, Tsukimori K, Yamamoto M, Himeno M, Furue M, Yamada H. Restoration of dioxin-induced damage to fetal steroidogenesis and gonadotropin formation by maternal co-treatment with -lipoic acid. PLoS One. 2012;7(7):e40322..

Chiba T, Chihara J, Furue M. Role of the Arylhydrocarbon Receptor (AhR) in the Pathology of Asthma and COPD. J Allergy (Cairo). 2012;2012:372384.

Tsukimori K, Yasukawa F, Uchi H, Furue M, Morokuma S. Sex ratio in two generations of the Yusho cohort. Epidemiology. 2012 Mar;23(2):349-50.

Todaka T, Hirakawa H, Kajiwara J, Onozuka D, Sasaki S, Miyashita C, Yoshioka E, Yuasa M, Kishi R, Iida T, Uchi H, Furue M. Concentrations of polychlorinated dibenzo-p-dioxins, polychlorinated dibenzofurans, and polychlorinated biphenyls in blood and breast milk collected from pregnant women in Sapporo City, Japan. Chemosphere. 2011 Dec;85(11):1694-700.

Tsukimori K, Uchi H, Mitoma C, Yasukawa F, Chiba T, Todaka T, Kajiwara J, Yoshimura T, Hirata T, Fukushima K, Wake N, Furue M. Maternal exposure to high levels of dioxins in relation to birth weight in women affected by Yusho disease. Environ Int. 2012 Jan;38(1):79-86.

Tsuji G, Takahara M, Uchi H, Matsuda T, Chiba T, Takeuchi S, Yasukawa F, Moroi Y, Furue M. Identification of ketoconazole as an AhR-Nrf2 activator in cultured human keratinocytes: the basis of its anti-inflammatory effect. J Invest Dermatol. 2012 Jan;132(1):59-68.

## 雑誌：発表者氏名，論文タイトル名，発表誌名，巻号，ページ，出版年

Uchi H, Yasukawa F, Kiryu H, Hashimoto K, Furue M. Infundibular cyst with seborrheic verruca-like cyst walls in a patient with Yusho disease exposed to dioxins. Eur J Dermatol. 2012 Oct; 122(5):687-8.

### 【赤羽 学】

Matsumoto S, Akahane M, Kanagawa Y, Kajiwara J, Todaka T, Yasukawa F, Uchi H, Furue M, Imamura T. Individuals' half-lives for 2,3,4,7,8-penta-chlorodibenzofuran (PeCDF) in blood: Correlation with clinical manifestations and laboratory results in subjects with Yusho. Chemosphere. 2013 Aug;92(7):772-7. doi: 10.1016/j.chemosphere.2013.04.005. Epub 2013 May 4.

Shinya Matsumoto, Manabu Akahane, Yoshiyuki Kanagawa, Jumboku Kajiwara, Hiroshi Uchi, Masataka Furue, Tomoaki Imamura :DISTRIBUTION OF PENTA-CHLORODIBENZOFURAN (PECDF) HALF LIFES IN YUSHO PATIENTS. The 33th International Symposium on Halogenated Persistent Organic Pollutants and POPs (Dioxin 2013) August 25-30, 2013 Daegu, Korea

松本伸哉、赤羽学、神奈川芳行、梶原淳睦、内博史、古江増隆、今村知明. 油症患者におけるダイオキシン類異性体ごとの症状ならびに異性体間の濃度と半減期の関係 福岡医学雑誌. 第104巻、4号、78-84、2013.

### 【内 博史】

Takei K, Mitoma C, Hashimoto-Hachiya A, Uchi H, Takahara M, Tsuji G, Kido-Nakahara M, Nakahara T, Furue M. Antioxidant soybean tar Glyteer® rescues Th2-mediated downregulation of filaggrin expression via aryl hydrocarbon receptor.

J Dermatol in press

Wu Z, Uchi H, Morino-Koga S, Shi W, Furue M. Resveratrol inhibition of human keratinocyte proliferation via SIRT1/ARNT/ERK dependent downregulation of aquaporin 3. J Dermatol Sci. 2014 Jul; 75(1):16-23.

Wu Z, Uchi H, Morino-Koga S, Nakamura-Satomura A, Kita K, Shi W, Furue M. Z-Ligustilide inhibits benzo(a)pyrene-induced CYP1A1 upregulation in cultured human keratinocytes via ROS-dependent Nrf2 activation. Exp Dermatol. 2014 Apr;23(4):260-5.

Morino-Koga S, Uchi H, Tsuji G, Takahara M, Kajiwara J, Hirata T, Furue M. Reduction of CC-chemokine ligand 5 by aryl hydrocarbon receptor ligands. J Dermatol Sci. 2013 Oct;72(1):9-15.

Tsukimori K, Uchi H, Tokunaga S, Yasukawa F, Chiba T, Kajiwara J, Hirata T, Furue M. Blood levels of PCDDs, PCDFs, and coplanar PCBs in Yusho mothers and their descendants: association with fetal Yusho disease. Chemosphere. 2013 Feb;90(5):1581-8.

## 雑誌：発表者氏名，論文タイトル名，発表誌名，巻号，ページ，出版年

Tsukimori K, Yasukawa F, Uchi H, Furue M, Morokuma S. Sex ratio in two generations of the Yusho cohort. *Epidemiology.* 2012 Mar;23(2):349-50.

Tsukimori K, Uchi H, Mitoma C, Yasukawa F, Chiba T, Todaka T, Kajiwara J, Yoshimura T, Hirata T, Fukushima K, Wake N, Furue M. Maternal exposure to high levels of dioxins in relation to birth weight in women affected by Yusho disease. *Environ Int.* 2012 Jan;38(1):79-86.

Tsuji G, Takahara M, Uchi H, Matsuda T, Chiba T, Takeuchi S, Yasukawa F, Moroi Y, Furue M. Identification of ketoconazole as an AhR-Nrf2 activator in cultured human keratinocytes: the basis of its anti-inflammatory effect. *J Invest Dermatol.* 2012 Jan;132(1):59-68.

Uchi H, Yasukawa F, Kiryu H, Hashimoto K, Furue M. Infundibular cyst with seborrheic verruca-like cyst walls in a patient with Yusho disease exposed to dioxins. *Eur J Dermatol.* 2012 Oct 1;22(5):687-8.

内 博史 油症の現況と慢性ダイオキシン類中毒の治療. 西日本皮膚科 76(6): 535-539, 2014

### 【平田 輝明（福岡県保健環境研究所）】

Onozuka Daisuke. Hirata Teruaki. Furue Masataka. Net survival after exposure to polychlorinated biphenyls and dioxins: the Yusho study. *Environ Int.* 73. 28-32. 2014.

Miyawaki T, Todaka T, Hirakawa H, Hori T, Kajiwara J, Hirata T, Uchi H, Furue M. Polychlorinated biphenyls concentrations in blood of Yusho patients during medical check-ups performed in 2012. *Organohalogen Compounds.* 76. 1457-1459. 2014.

Jumboku Kajiwara, Takashi Todaka, Hironori Hirakawa, Takashi Miyawaki, Tamiko Ikeno, Cihiro Miyasita, Sachiko Itoh, Seiko Sasaki, Reiko Kishi, Masataka Furue. Dioxins and PCBs concentrations in umbilical cord blood collected in Hokkaidou, Japan. *Organohalogen Compounds.* 76. 1481-1484. 2014.

Hori Tsuguhide, Takashi Miyawaki, Koji Takahashi, Daisuke Yasutake, Takamitsu Yamamoto, Jumboku Kajiwara, Takahiro Watanabe. Concentration of Dechlorane Plus in fish samples collected in Kyushu district, western Japan. *Organohalogen Compounds.* 76. 900-903. 2014.

Koji Takahashi, Tsuguhide Hori, Jumboku Kajiwara, Takahiro Watanabe. Determination of hexabromocyclododecane in fish samples collected from Japanese markets. *Organohalogen Compounds.* 76. 930-933. 2014.

**雑誌：発表者氏名，論文タイトル名，発表誌名，巻号，ページ，出版年**

Kiyomi Tsukimori, Hirohi Uchi, Shouji Tokunaga, Fumiko Yasukawa, Takahiro Chiba, Jumboku Kajiwara, Teruakai Hirata, Masutake Furue, et al. Blood levels of PCDDs, PCDFs, and coplanar PCBs in Yusho mothers and their descendants: Association with fetal Yusho disease. *Chemosphere*. 90(5). 1581-1588. 2013.

Morino-Koga S, Uchi H, Tsuji G, Takahara M, Kajiwara J, Hirata T, Furue M. Reduction of CC-chemokine ligand 5 by aryl hydrocarbon receptor ligands. *J Dermatol Sci*. 72(1). 9-15. 2013.

梶原淳睦・戸高 尊・平川博仙・堀 就英・安武大輔・宮脇 崇・飛石和大・高尾佳子・平田輝昭・内 博史・古江増隆・血液中ダイオキシン、PCB濃度測定のクロスチェック. 福岡医学雑誌. 104(4). 104-109. 2013.

Tobiishi K, Suzuki S, Todaka T, Hirakawa H, Hori T, Kajiwara J, Hirata T, Iida T, Uchi H, Furue M. Improvement of measurement method for hydroxylated polychlorinated biphenyls (OH-PCBs) in blood samples using LC/MS/MS. *Fukuoka Igaku Zasshi*. 104(4). 128-135. 2013.

Tobiishi K, Suzuki S, Todaka T, Hirakawa H, Hori T, Kajiwara J, Hirata T, Iida T, Uchi H, Furue M. Concentration of hydroxylated polychlorinated biphenyls (OH-PCBs) in the blood of Yusho patients in 2010. *Fukuoka Igaku Zasshi*. 104(4). 136-142. 2013.

堀 就英・山本貴光・石黒靖尚・飯田隆雄・梶原淳睦・平田輝昭・内 博史・古江増隆・血液中PCB異性体分離分析におけるアルカリ分解温度の検討. 福岡医学雑誌. 104(4). 152-160. 2013.

Hori T, Takahashi K, Kajiwara J, Morokuma S, Otera Y, Yamamoto T, Kawamoto T, Hirata T, Furue M, Tsukimori K, et al. CHARACTERIZATION OF DIOXINS IN SETS OF PLACENTAL TISSUE, MATERNAL BLOOD AND UMBILICAL CORD BLOOD SAMPLES COLLECTED FROM JAPANESE PREGNANT WOMEN. *Organohalogen Compounds*. 74, 763-766. 2012.

Tobiishi K, Todaka T, Hirakawa H, Hori T, Kajiwara J, Hirata T, Sasaki S, Miyashita C, Yoshioka E, Yuasa M, Kishi R, Iida T, Uchi H, Furue M, et al. Determination of hydroxylated polychlorinated biphenyls (OH-PCBs) in the blood of pregnant women by LC/MS/MS. *Organohalogen Compounds*. 74. 43-45. 2012.

Miyawaki T, Hori T, Todaka T, Hirakawa H, Kajiwara J, Hirata T, Uchi H, Furue M : CONCENTRATIONS OF POLYCHLORINATED BIPHENYLS IN BLOOD OF YUSHO PATIENTS COLLECTED FROM MEDICAL CHECK-UPS IN 2010. *Organohalogen Compounds*. 74. 885-887. 2012.

Tsukimori K, Uchi H, Mitoma C, Yasukawa F, Chiba T, Todaka T, Kajiwara J, Yoshimura T, Hirata T, Fukushima K, Wake N, Furue M, et al. Maternal exposure to high levels of dioxins in relation to birth weight in women affected by Yusho disease. *Environ Int*. 38(1). 79-86. 2012.

## 雑誌：発表者氏名，論文タイトル名，発表誌名，巻号，ページ，出版年

### 【吉村 健清】

二塚信，吉村健清.

食品汚染による中毒の認定をめぐる最近の動向 - 水俣病・力ネミ油症.

医学のあゆみ, 244 (10): 941-943, 2013.

Takesumi Yoshimura.

Yusho: 43 years later.

Kaohsiung Journal of Medical Sciences. 28 (2), S49-S52, 2012.

Pham TM, Ozasa K, Kubo T, Fujino Y, Sakata R, Grant EJ, Matsuda S, Yoshimura T.

Age-Period-Cohort Analysis of Chronic Obstructive Pulmonary Disease Mortality in Japan, 1950-2004.

J Epidemiol. 22(4):302-7, 2012.

吉村健清.

EBM と臨床研究 - 痘学の役割 - .

JOHNS, Feb;28(2):156-158, 2012.

Tsukimori,T.,Uchi,H.,Mitoma,C.,Yasukawa,F.,Chiba,T.,Todaka,T.,Kajiwara,J.,Yoshimura ,T.,Hirata,T.,Fukushima,K.,Wake,N.,Furue,M.

Maternal exposure to high levels of dioxins in relation to birth weight in women affected by Yusho disease.

Environmental International, 38;79-86, 2012.

### 【古賀 信幸】

C. Ohta, K. Haraguchi, Y. Kato, T. Endo, O. Kimura, and N. Koga, Distribution and excretion of 2,2',3,4',5,5',6-heptachlorobiphenyl (CB187) and its metabolites in rats and guinea pigs. Chemosphere, 118, 5-11 (2015).

O. Kimura,C. Ohta, N. Koga, K. Haraguchi, Y. Kato, and T. Endo, Carrier-mediated uptake of nobiletin, a citrus polymethoxyflavonoid, in human intestinal Caco-2 cells. Food Chem., 154, 145-150 (2014).

C. Ohta, K. Haraguchi, Y. Kato, T. Endo, N. Koga, The participation of rat CYP3A enzymes in the metabolism of 2,2',4,5,5'-pentachlorobiphenyl (CB101). Organohalogen Compounds, 76, 466-469(2014).

Y. Kato, K. Haraguchi, M. Onishi, S. Ikushiro, T. Endo, C. Ohta, N. Koga, S. Yamada and M. Degawa, 3,3',4,4'-Tetrachlorobiphenyl-mediated decrease of serum thyroxine level in C57BL/6 and DBA/2 mice occurs mainly through enhanced accumulation of thyroxine in the liver. Biol. Pharm. Bull., 37, 504-509 (2014).

Y. Kato, M. Onishi, K. Haraguchi, S. Ikushiro, C. Ohta, N. Koga, T. Endo, S. Yamada and M. Degawa, A possible mechanism for 2,3',4,4',5-pentachlorobiphenyl-mediated decrease in serum thyroxine level in mice. Biol. Pharm. Bull., 36(10), 1594-1601 (2013).

### 雑誌：発表者氏名，論文タイトル名，発表誌名，巻号，ページ，出版年

C. Ohta, K. Haraguchi, Y. Kato, T. Endo, O. Kimura, N. Koga, In vitro metabolism of 2,2',4,4',5-pentachlorobiphenyl (CB99) by rat and guinea pig liver microsomes. *Organohalogen Compounds*, 75, 587-590(2013).

太田千穂，原口浩一，加藤善久，遠藤哲也，古賀信幸，2,2',3,4',5,5'-六塩素化ビフェニル(CB146)の in vitro 代謝の動物種差. 福岡医誌, 104(4), 161-169 (2013).

C. Ohta, K. Haraguchi, Y. Kato, T. Endo and N. Koga, Involvement of rat CYP3A enzymes in the metabolism of 2,2',3,4',5',6-hexachlorobiphenyl (CB149). *Organohalogen Compounds*, 74, 1475-1478(2012).

Y. Kato, S. Tamaki, K. Haraguchi, S. Ikushiro, M. Sekimoto, C. Ohta, T. Endo, N. Koga, S. Yamada and M. Degawa, 2,2',4,5,5'-Pentachlorobiphenyl-mediated inhibition of a serum T<sub>4</sub>-transthyretin complex formation is one of causes for the PCB-induced changes in the serum and hepatic T<sub>4</sub> levels in mice. *Organohalogen Compounds*, 74, 1377-1380(2012).

### 【月森 清巳】

Tsukimori K, Uchi H, Tokunaga S, Yasukawa F, Chiba T, Kajiwara J, Hirata T, Furue M. Blood levels of PCDDs, PCDFs, and coplanar PCBs in Yusho mothers and their descendants: association with fetal Yusho disease. *Chemosphere*. 90(5):1581-8, 2013.

Tsukimori K, Morokuma S, Hori T, Takahashi K, Hirata T, Otera Y, Fukushima K, Kawamoto T, Wake N. Characterization of placental transfer of polychlorinated dibenzo-p-dioxins, dibenzofurans and polychlorinated biphenyls in normal pregnancy. *J Obstet Gynaecol Res*. 39(1):83-90, 2013.

Tsukimori K, Yasukawa F, Uchi H, Furue M, Morokuma S. Sex ratio in two generations of the Yusho cohort. *Epidemiology*. 23(2):349-50, 2012.

Tsukimori K, Uchi H, Mitoma C, Yasukawa F, Chiba T, Todaka T, Kajiwara J, Yoshimura T, Hirata T, Fukushima K, Wake N, Furue M. Maternal exposure to high levels of dioxins in relation to birth weight in women affected by Yusho disease. *Environ Int*. 38(1):79-86, 2012.

### 【辻 博】

辻 博：油症における特異抗核抗体の検討. 福岡医学雑誌 104 : 73-77, 2013 .

### 【山田 英之】

Hattori Y., Takeda T., Fujii M., Taura J., Ishii Y., Yamada H. Dioxin-induced fetal growth retardation: the role of a preceding attenuation in the circulating level of glucocorticoid. *Endocrine*, 47: 572-580, 2014.

**雑誌：発表者氏名，論文タイトル名，発表誌名，巻号，ページ，出版年**

Taura J., Takeda T., Fujii M., Hattori Y., Ishii Y., Kuroki H., Tsukimori K., Uchi H., Furue M., Yamada H. 2,3,4,7,8-Pentachlorodibenzofuran is far less potent than 2,3,7,8-tetrachlorodibenzo-*p*-dioxin in disrupting the pituitary-gonad axis of the rat fetus. *Toxicol. Appl. Pharmacol.*, 281: 48-57, 2014.

Takeda T., Taura J., Hattori Y., Ishii Y., Yamada H. Dioxin-induced retardation of development through a reduction in the expression of pituitary hormones and possible involvement of an aryl hydrocarbon receptor in this defect: a comparative study using two strains of mice with different sensitivities to dioxin. *Toxicol. Appl. Pharmacol.*, 278: 220-229, 2014.

Takeda T., Fujii M., Hattori Y., Yamamoto M., Shimazoe T., Ishii Y., Himeno M., Yamada H. Maternal exposure to dioxin imprints sexual immaturity of the pups through fixing the status of the reduced expression of expression of hypothalamic gonadotropin-releasing hormone. *Mol. Pharmacol.*, 85: 74-82, 2014.

武田知起，服部友紀子，藤井美彩紀，田浦順樹，石井祐次，山田英之。

ダイオキシン母体曝露による胎児副腎ステロイド合成系への影響と性差。福岡医学雑誌，104 (4): 143-151 , 2013.

Tsujimoto S, Ishida T, Takeda T, Ishii Y, Onomura Y, Tsukimori K, Takechi S, Yamaguchi T, Uchi H, Suzuki SO, Yamamoto M, Himeno M, Furue M, Yamada H. Selenium-binding protein 1: its physiological function, dependence on aryl hydrocarbon receptors, and role in wasting syndrome by 2,3,7,8-tetrachlorodibenzo-*p*-dioxin.

*Biochim. Biophys. Acta*, 1830: 3616-3624 , 2013.

Koga T., Ishida T., Takeda T., Ishii Y., Uchi H., Tsukimori K., Yamamoto M., Himeno M., Furue M., Yamada H. Restoration of dioxin-induced damage to fetal steroidogenesis and gonadotropin formation by maternal co-treatment with -lipoic acid. *PLoS ONE*, 7: e40322, 2012.

Takeda T., Fujii M., Taura J., Ishii Y., Yamada H. Dioxin silences gonadotropin expression in perinatal pups by inducing histone deacetylases: a new insight into the mechanism for the imprinting of sexual immaturity by dioxin. *J. Biol. Chem.*, 287: 18440-18450, 2012.

**【宇谷 厚志】**

Kuwatsuka Y, Shimizu K, Akiyama Y, Koike Y, Ogawa F, Furue M, Utani A: Yusho patients show increased serum IL-17, IL-23, IL-1beta, and TNFalpha levels more than 40 years after accidental polychlorinated biphenyl poisoning. *J Immunotoxicol* 11(3): 246-249, 2014.

