

Table 3. Multivariate Analysis of Pre- and Intraoperative Factors Predicting Postoperative Pancreatic Fistula (International Study Group of Pancreatic Fistula Grade B or Higher)

Factors	Multivariate analysis		
	Odds ratio	95 % CI	p Value
Soft pancreas (SWV < 1.54 m/s)	38.3	5.82–445	0.001
Male sex	8.87	1.71–67.8	0.018
Age ≤ 70 y	7.69	1.44–62.3	0.030
BMI ≤ 25.0 kg/m ²	8.26	1.08–94.4	0.059
MPD diameter ≤ 3 mm	2.32	0.50–11.6	0.285
Blood loss > 1,000 mL	4.78	0.41–79.2	0.233
Pathology, pancreatic cancer	0.40	0.07–1.93	0.259
Presence of diabetes mellitus	1.43	0.24–9.63	0.699

BMI, body mass index; MPD, main pancreatic duct; SWV, shear wave velocity.

the surface of the pancreas and/or patients had difficulty holding their breath during the examination. Preoperative ARFI imaging cannot be used to evaluate severely obese patients because to measure SWV, this method requires visualization of the pancreatic parenchyma within 8 cm from the surface of the body, with minimum compression to the abdominal wall. However, intraoperative ARFI imaging, done by setting a transducer directly onto the pancreatic surface, may enable measurement of pancreatic SWV even in obese patients. Another limitation of ARFI imaging is that it correlates with the degree of pathologic fibrosis but not with fatty infiltration, which may also be associated with pancreatic function and risk of postoperative PF.^{10,11} Additional studies are needed to enhance the ability of ARFI imaging to predict postoperative PF by, for example, including the pancreatic echogenicity reflecting fatty infiltration and intraoperative pancreatic SWVs, into estimations and optimizing SWV cut-off values in a larger series of patients.

CONCLUSIONS

In conclusion, preoperative assessment of pancreatic elasticity by ARFI imaging enabled estimation of the degree of pathologic fibrosis in pancreatic tissue. Preoperative determination of pancreatic exocrine function and risk of postoperative PF may be useful in designing appropriate surgical procedures and postoperative management for individual patients undergoing pancreatic resection.

Author Contributions

Study conception and design: Harada, Ishizawa, Inoue

Acquisition of data: Harada, Tanaka

Analysis and interpretation of data: Harada, Ishizawa, Aoki, Sakamoto, Hasegawa, Sugawara

Drafting of manuscript: Harada, Ishizawa, Inoue
Critical revision: Fukayama, Kokudo

Acknowledgment: The authors acknowledge the significant contributions made by Drs A Nakazawa, Y Okinaga, and J Kaneko, members of this study group.

REFERENCES

- Yeo CJ, Cameron JL, Sohn TA, et al. Six hundred fifty consecutive pancreaticoduodenectomies in the 1990s: pathology, complications, and outcomes. *Ann Surg* 1997;226:248–257.
- Fuks D, Piessen G, Huet E, et al. Life-threatening postoperative pancreatic fistula (grade C) after pancreaticoduodenectomy: incidence, prognosis, and risk factors. *Am J Surg* 2009;197:702–709.
- Mckay A, Mackenzie S, Sutherland FR, et al. Meta-analysis of pancreaticojunostomy versus pancreaticogastrostomy reconstruction after pancreaticoduodenectomy. *Br J Surg* 2006;93:929–936.
- Topal B, Fieuws S, Aerts R, et al. Pancreaticojunostomy versus pancreaticogastrostomy reconstruction after pancreaticoduodenectomy for pancreatic or periampullary tumours: a multicenter randomized trial. *Lancet Oncol* 2013;14:655–662.
- Pratt WB, Maithel SK, Vanounou T, et al. Clinical and economic validation of the International Study Group of Pancreatic Fistula (ISGPF) classification scheme. *Ann Surg* 2007;245:443–451.
- Kawai M, Kondo S, Yamaue H, et al. Predictive risk factors for clinically relevant pancreatic fistula analyzed in 1,239 patients with pancreaticoduodenectomy: multicenter data collection as a project study of pancreatic surgery by the Japanese Society of Hepato-Biliary-Pancreatic Surgery. *J Hepatobiliary Pancreat Sci* 2011;18:601–608.
- Yeo CJ, Cameron JL, Maher MM, et al. A prospective randomized trial of pancreaticogastrostomy versus pancreaticojunostomy after pancreaticoduodenectomy. *Ann Surg* 1995;222:588–592.
- Yang YM, Tian XD, Zhuang Y, et al. Risk factors of pancreatic leakage after pancreaticoduodenectomy. *World J Gastroenterol* 2005;28:2456–2461.
- Kah Heng CA, Salleh I, San TS, et al. Pancreatic fistula after distal pancreatectomy: incidence, risk factors and management. *ANZ J Surg* 2010;80:619–623.
- Gaujoux S, Cortes A, Couvelard A, et al. Fatty pancreas and increased body mass index are risk factors of pancreatic fistula after pancreaticoduodenectomy. *Surgery* 2010;148:15–23.
- Mathur A, Pitt HA, Marine M, et al. Fatty pancreas: a factor in postoperative pancreatic fistula. *Ann Surg* 2007;246:1058–1064.
- Akamatsu N, Sugawara Y, Komagome M, et al. Risk factors for postoperative pancreatic fistula after pancreaticoduodenectomy: the significance of the ratio of the main pancreatic duct to the pancreas body as a predictor of leakage. *J Hepatobiliary Pancreat Sci* 2010;17:322–328.
- Wada K, Traverso LW. Pancreatic anastomotic leak after the Whipple procedure is reduced using the surgical microscope. *Surgery* 2006;139:735–742.
- Muscaria F, Suc B, Kirzin S, et al. Risk factors for mortality and intra-abdominal complications after pancreaticoduodenectomy:

- multivariate analysis in 300 patients. *Surgery* 2006;139:591–598.
15. Pratt WB, Callery MP, Vollmer CM Jr. Risk prediction for development of pancreatic fistula using the ISGPF classification scheme. *World J Surg* 2008;32:419–428.
 16. Dong X, Zhang B, Kang MX, et al. Analysis of pancreatic fistula according to the International Study Group on Pancreatic Fistula classification scheme for 294 patients who underwent pancreaticoduodenectomy in a single center. *Pancreas* 2011;40:222–228.
 17. Yeh TS, Jan YY, Jeng LB, et al. Pancreaticojejunal anastomotic leak after pancreaticoduodenectomy—multivariate analysis of perioperative risk factors. *J Surg Res* 1997;67:119–125.
 18. Piscaqlia F, Marinelli S, Botta S, et al. The role of ultrasound elastographic techniques in clinic liver disease: Current status and future perspectives. *Eur J Radiol* 2014;83:450–455.
 19. Friedrich-Rust M, Wunder K, Kriener S, et al. Liver fibrosis in viral hepatitis: noninvasive assessment with acoustic radiation force impulse imaging versus transient elastography. *Radiology* 2009;252:595–604.
 20. Bamber J, Cosgrove D, Dietrich CF, et al. EFSUMB guidelines and recommendations on the clinical use of ultrasound elastography. Part 1: Basic principles and technology. *Ultraschall Med* 2013;34:169–184.
 21. Fahey BJ, Nightingale KR, Nelson RC, et al. Acoustic radiation force impulse imaging of the abdomen: demonstration of feasibility and utility. *Ultrasound Med Biol* 2005;31:1185–1198.
 22. Guzmán-Aroca F, Frutos-Bernal MD, Bas A, Luján-Mompeán JA, et al. Detection of non-alcoholic steatohepatitis in patients with morbid obesity before bariatric surgery: preliminary evaluation with acoustic radiation force impulse imaging. *Eur Radiol* 2012;22:2525–2532.
 23. Vermehren J, Polta A, Zimmermann O, et al. Comparison of acoustic radiation force impulse imaging with transient elastography for the detection of complications in patients with cirrhosis. *Liver Int* 2012;32:852–858.
 24. Son CY, Kim SU, Han WK, et al. Normal liver elasticity values using acoustic radiation force impulse imaging: a prospective study in healthy living liver and kidney donors. *J Gastroenterol Hepatol* 2012;27:130–136.
 25. Zhang P, Zhou P, Tian SM, et al. Application of acoustic radiation force impulse imaging for the evaluation of focal liver lesion elasticity. *Hepatobiliary Pancreat Dis Int* 2013;12:165–170.
 26. Meng W, Zhang G, Wu C, et al. Preliminary results of acoustic radiation force impulse (ARFI) ultrasound imaging of breast lesions. *Ultrasound Med Biol* 2011;37:1436–1443.
 27. Friedrich-Rust M, Romenski O, Meyer G, et al. Acoustic radiation force impulse-imaging for the evaluation of the thyroid gland: a limited patient feasibility study. *Ultrasonics* 2012;52:69–74.
 28. D'Onofrio M, Crosara S, Canestrini S, et al. Virtual analysis of pancreatic cystic lesion fluid content by ultrasound acoustic radiation force impulse quantification. *J Ultrasound Med* 2013;32:647–651.
 29. Lee TK, Kang CM, Park MS, et al. Prediction of postoperative pancreatic fistulas after pancreatectomy: assessment with acoustic radiation force impulse elastography. *J Ultrasound Med* 2014;33:781–786.
 30. Hasegawa K, Kokudo N, Sano K, et al. Two-stage pancreaticojejunostomy in pancreaticoduodenectomy: a retrospective analysis of short-term results. *Am J Surg* 2008;196:3–10.
 31. Yamashita S, Sakabe M, Ishizawa T, et al. Visualization of the leakage of pancreatic juice using a chymotrypsin-activated fluorescent probe. *Br J Surg* 2013;100:1220–1228.
 32. Konishi T, Hiraishi M, Kubota K, et al. Segmental occlusion of the pancreatic duct with prolamine to prevent fistula formation after distal pancreatectomy. *Ann Surg* 1995;221:165–170.
 33. Yashima Y, Sasahira N, Isayama H, et al. Acoustic radiation force impulse elastography for noninvasive assessment of chronic pancreatitis. *J Gastroenterol* 2012;47:427–432.
 34. Klöppel G, Maillet B. Pseudocysts in chronic pancreatitis: a morphological analysis of 57 resection specimens and 9 autopsy pancreata. *Pancreas* 1991;6:266–274.
 35. Youden WJ. Index for rating diagnostic tests. *Cancer* 1950;3:32–35.
 36. Yoshioka R, Saiura A, Koga R, et al. Risk factors for clinical pancreatic fistula after pancreatectomy: analysis of consecutive 100 patients. *World J Surg* 2010;34:121–125.
 37. Sledzianowski JF, Duffas JP, Muscari F, et al. Risk factors for mortality and intra-abdominal morbidity after distal pancreatectomy. *Surgery* 2005;137:180–185.
 38. Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 2004;240:205–213.
 39. Toshima T, Shirabe K, Takeishi K, et al. New method for assessing liver fibrosis based on acoustic radiation force impulse: a special reference to the difference between right and left liver. *J Gastroenterol* 2011;46:705–711.
 40. Mishra PK, Saluja SS, Gupta M, et al. Blumgart's technique of pancreaticojejunostomy: an appraisal. *Dig Surg* 2011;28:281–287.
 41. Kennedy EP, Yeo CJ. Dunking pancreaticojejunostomy versus duct-to-mucosa anastomosis. *J Hepatobiliary Pancreat Surg* 2011;18:769–774.
 42. Hamilton NA, Porembka MR, Johnston FM, et al. Mesh reinforcement of pancreatic transection decreases incidence of pancreatic occlusion failure for left pancreatectomy: a single-blinded, randomized controlled trial. *Ann Surg* 2012;255:1037–1042.
 43. Diener MK, Seiler CM, Rossion I, et al. Efficacy of stapler versus hand-sewn closure after distal pancreatectomy (DIS-PACT): a randomized, controlled multicenter trial. *Lancet* 2011;377:1514–1522.
 44. Peng SY, Wang JW, Lau WY, et al. Conventional versus binding pancreaticojejunostomy after pancreaticoduodenectomy: a prospective randomized trial. *Ann Surg* 2007;245:692–698.
 45. Ke S, Ding XM, Gao J, et al. A prospective, randomized trial of Roux-en-Y reconstruction with isolated pancreatic drainage versus conventional loop reconstruction after pancreaticoduodenectomy. *Surgery* 2013;153:743–752.
 46. Poon RT, Fan ST, Lo CM, et al. External drainage of pancreatic duct with a stent to reduce leakage rate of pancreaticojejunostomy after pancreaticoduodenectomy: a prospective randomized trial. *Ann Surg* 2007;246:425–435.

Supplementary Table 1. Reproducibility of Shear Wave Velocity Measurement of the Spleen in Healthy Volunteers

	Spleen		
HV1 Examiner	①	②	③
Measured value	2.50	2.04	2.30
	2.44	2.30	1.89
	2.42	2.16	2.06
	2.09	2.04	2.15
	2.00	2.60	2.16
	2.32	2.23	2.06
	2.01	2.15	2.06
	2.44	2.26	2.23
	1.80	2.20	2.07
	2.50	2.02	2.43
Median value	2.37 (1.80–2.50)	2.18 (2.02–2.60)	2.11 (1.89–2.43)
Average value	2.25 ± 0.25	2.20 ± 0.17	2.14 ± 0.15
p value			0.549
HV2 Examiner			
Measured value	2.02	2.39	2.35
	2.05	2.13	2.56
	1.98	2.51	2.44
	2.36	1.89	1.85
	2.14	2.21	2.08
	2.40	2.21	2.15
	2.41	2.11	2.57
	2.01	2.20	2.54
	2.24	2.10	2.22
	2.54	2.19	1.89
Median value	2.19 (1.98–2.54)	2.20 (1.89–2.51)	2.29 (1.85–2.57)
Average value	2.22 ± 0.20	2.22 ± 0.17	2.27 ± 0.27
p value			0.697
HV3 Examiner			
Measured value	2.11	2.34	2.87
	1.90	2.51	2.57
	2.47	2.51	2.30
	2.83	2.21	2.30
	2.38	2.54	2.71
	2.89	2.53	2.56
	2.66	2.87	2.32
	2.83	2.37	2.17
	2.96	2.25	2.36
	2.34	2.53	2.24
Median value	2.57 (1.90–2.96)	2.51 (2.21–2.87)	2.34 (2.17–2.87)
Average value	2.54 ± 0.36	2.47 ± 0.19	2.44 ± 0.23
p value			0.633

Supplementary Table 2. Reproducibility of Shear Wave Velocity Measurement of the Right Liver in Healthy Volunteers

	Right Liver		
HV1 Examiner	①	②	③
Measured value	1.15	1.20	1.25
	1.31	1.27	1.31
	1.19	1.34	1.30
	1.37	1.34	1.29
	1.20	1.32	1.31
	1.34	1.28	1.30
	1.20	1.32	1.28
	1.23	1.23	1.34
	1.25	1.29	1.27
	1.30	1.27	1.37
Median value	1.24 (1.15–1.37)	1.29 (1.20–1.34)	1.3 (1.25–1.37)
Average value	1.25 ± 0.07	1.29 ± 0.05	1.30 ± 0.03
p value			0.281
HV2 Examiner			
Measured value	1.01	1.08	1.05
	0.99	1.00	1.01
	1.13	1.15	1.02
	1.04	1.11	1.03
	1.05	1.08	1.02
	1.11	1.10	0.93
	1.09	1.07	1.03
	1.08	1.11	1.13
	1.12	0.92	1.09
	1.13	0.95	0.99
Median value	1.09 (0.99–1.13)	1.08 (0.92–1.15)	1.03 (0.93–1.13)
Average value	1.08 ± 0.05	1.06 ± 0.08	1.03 ± 0.05
p value			0.243
HV3 Examiner			
Measured value	1.25	1.26	1.16
	1.20	1.22	1.24
	1.24	1.16	1.22
	1.18	1.22	1.32
	1.19	1.20	1.15
	1.16	1.27	1.40
	1.20	1.18	1.26
	1.26	1.19	1.20
	1.25	1.16	1.34
	1.26	1.23	1.40
Median value	1.22 (1.16–1.26)	1.21 (1.16–1.27)	1.25 (1.15–1.40)
Average value	1.22 ± 0.04	1.21 ± 0.04	1.27 ± 0.09
p value			0.317

Supplementary Table 3. Reproducibility of Shear Wave Velocity Measurement of the Left Liver in Healthy Volunteers

	Left Liver		
HV1 Examiner	①	②	③
Measured value	1.17	1.27	1.10
	1.04	1.27	1.05
	1.22	1.09	1.03
	0.95	1.28	1.27
	1.02	1.16	1.17
	1.21	1.26	1.21
	1.06	0.96	1.20
	1.32	1.22	1.08
	1.33	0.97	1.18
	1.33	1.13	1.21
Median value	1.19 (0.95–1.33)	1.19 (0.96–1.28)	1.18 (1.03–1.27)
Average value	1.17 ± 0.14	1.16 ± 0.12	1.15 ± 0.08
p value			0.859
HV2 Examiner			
Measured value	0.93	1.02	1.03
	1.21	1.01	0.98
	0.99	1.14	1.07
	1.03	1.1	1.12
	0.97	1.19	1.04
	1.3	0.96	1.14
	1.24	1.02	1.08
	0.76	1.07	1.09
	1.04	1.05	1.03
	1.16	0.93	1
Median value	1.04 (0.76–1.30)	1.04 (0.93–1.19)	1.06 (0.98–1.14)
Average value	1.06 ± 0.16	1.05 ± 0.08	1.06 ± 0.05
p value			0.791
HV3 Examiner			
Measured value	1.46	1.35	1.61
	1.49	1.24	1.16
	1.27	1.49	1.28
	1.56	1.32	1.13
	1.06	1.43	1.36
	1.70	1.26	1.47
	1.40	1.47	1.25
	1.39	1.58	1.33
	1.48	1.58	1.60
	1.64	1.53	1.27
Median value	1.47 (1.06–1.70)	1.45 (1.24–1.58)	1.31 (1.13–1.61)
Average value	1.45 ± 0.18	1.43 ± 0.13	1.35 ± 0.17
p value			0.359

Supplementary Table 4. Reproducibility of Shear Wave Velocity Measurement of the Pancreas in Healthy Volunteers

	Pancreas		
HV1 Examiner	①	②	③
Measured value	0.83	0.96	1.00
	1.30	0.96	0.93
	1.10	1.23	1.07
	0.94	0.93	1.21
	1.03	1.05	1.07
	1.07	1.10	1.02
	1.06	1.02	1.12
	0.84	1.13	0.95
	0.80	1.11	0.99
	0.88	1.00	1.00
Median value	0.99 (0.80–1.30)	1.04 (0.93–1.23)	1.01 (0.93–1.21)
Average value	0.99 ± 0.16	1.05 ± 0.09	1.04 ± 0.08
p value			0.437
HV2 Examiner			
Measured value	1.06	1.25	1.38
	0.93	1.35	1.28
	1.20	1.27	1.40
	1.48	1.31	1.17
	1.17	1.26	1.24
	1.47	1.27	1.38
	1.51	1.21	1.47
	1.51	1.38	1.20
	1.21	1.15	1.29
	1.05	1.21	1.20
Median value	1.21 (0.93–1.51)	1.27 (1.15–1.38)	1.29 (1.17–1.47)
Average value	1.26 ± 0.22	1.27 ± 0.07	1.30 ± 0.10
p value			0.785
HV3 Examiner			
Measured value	1.19	0.98	0.98
	0.89	0.94	1.05
	0.98	1.10	0.98
	0.96	0.88	1.02
	1.15	1.13	1.10
	0.95	0.93	1.03
	0.97	1.03	0.91
	1.02	1.11	1.11
	1.24	1.14	1.09
	0.93	1.19	1.10
Median value	0.98 (0.89–1.24)	1.07 (0.88–1.19)	1.04 (0.91–1.11)
Average value	1.03 ± 0.12	1.04 ± 0.11	1.04 ± 0.07
p value			0.945

Supplementary Table 5. Reproducibility of Shear Wave Velocity Measurement of the Pancreas in Patients Undergoing Pancreatoduodenectomy

	IPMN		
Patient 1 Examiner	①	②	③
Measured value	1.51	1.47	1.69
	1.72	1.72	1.27
	1.34	1.55	1.53
	1.70	1.51	1.75
	1.66	1.52	1.37
	1.39	1.43	1.68
	1.62	1.62	1.45
	1.32	1.40	1.49
	1.66	1.63	1.66
	1.38	1.61	1.67
Median value	1.57 (1.32–1.72)	1.54 (1.40–1.72)	1.60 (1.27–1.75)
Average value	1.53 ± 0.16	1.55 ± 0.10	1.56 ± 0.16
p value			0.919
Bile Duct Cancer			
Patient 2 Examiner	①	②	③
Measured value	1.30	1.40	1.38
	1.39	1.47	1.30
	1.19	1.28	1.69
	1.33	1.37	1.41
	1.27	1.48	1.16
	1.54	1.32	1.13
	1.33	1.30	1.26
	1.57	1.46	1.55
	1.39	1.55	1.27
	1.38	1.29	1.46
Median value	1.36 (1.19–1.57)	1.39 (1.28–1.55)	1.34 (1.13–1.69)
Average value	1.37 ± 0.12	1.39 ± 0.09	1.36 ± 0.17
p value			0.863
Pancreatic Cancer			
Patient 3 Examiner	①	②	③
Measured value	0.84	1.17	0.96
	0.91	0.99	1.12
	1.07	0.92	0.89
	0.94	1.09	1.21
	0.90	1.21	0.98
	1.25	0.99	0.89
	1.21	1.00	1.25
	1.30	1.18	0.95
	0.92	1.22	1.34
	1.16	0.95	0.89
Median value	1.01 (0.84–1.30)	1.05 (0.92–1.22)	0.97 (0.89–1.34)
Average value	1.05 ± 0.17	1.07 ± 0.12	1.05 ± 0.17
p value			0.927

