

## Author Contributions

Conceived and designed the experiments: DO. Performed the experiments: NT DO HT TH CSH. Analyzed the data: JW AN JE NN HY. Wrote the paper: DO KT HM.

## References

- Gilbertson DT, Liu J, Xue JL, Louis TA, Solid CA, et al. (2005) Projecting the number of patients with end-stage renal disease in the United States to the year 2015. *J Am Soc Nephrol* 16: 3736–3741.
- Kanwar YS, Sun L, Xie P, Liu FY, Chen S (2011) A glimpse of various pathogenetic mechanisms of diabetic nephropathy. *Annu Rev Pathol* 6: 395–423.
- Navarro-Gonzalez JF, Mora-Fernandez C, Muros de Fuentes M, Garcia-Perez J (2011) Inflammatory molecules and pathways in the pathogenesis of diabetic nephropathy. *Nat Rev Nephrol* 7: 327–340.
- Giacco F, Brownlee M (2010) Oxidative stress and diabetic complications. *Circ Res* 107: 1058–1070.
- Hediger MA, Rhoads DB (1994) Molecular physiology of sodium-glucose cotransporters. *Physiol Rev* 74: 993–1026.
- Wright EM (2001) Renal Na(+)-glucose cotransporters. *Am J Physiol Renal Physiol* 280: F10–18.
- Chao EC, Henry RR (2010) SGLT2 inhibition—a novel strategy for diabetes treatment. *Nat Rev Drug Discov* 9: 551–559.
- Chen LH, Leung PS (2013) Inhibition of the sodium glucose co-transporter-2: its beneficial action and potential combination therapy for type 2 diabetes mellitus. *Diabetes Obes Metab* 15: 392–402.
- Han S, Hagan DL, Taylor JR, Xin L, Meng W, et al. (2008) Dapagliflozin, a selective SGLT2 inhibitor, improves glucose homeostasis in normal and diabetic rats. *Diabetes* 57: 1723–1729.
- Bouchie A (2013) SGLT2 inhibitors enter crowded diabetes space. *Nat Biotechnol* 31: 469–470.
- Plosker GL (2012) Dapagliflozin: a review of its use in type 2 diabetes mellitus. *Drugs* 72: 2289–2312.
- DeFronzo RA, Hompesch M, Kasichayanula S, Liu X, Hong Y, et al. (2013) Characterization of Renal Glucose Reabsorption in Response to Dapagliflozin in Healthy Subjects and Subjects With Type 2 Diabetes. *Diabetes Care*.
- Fujimori Y, Katsuno K, Nakashima I, Ishikawa-Takemura Y, Fujikura H, et al. (2008) Remogliflozin etabonate, in a novel category of selective low-affinity sodium glucose cotransporter (SGLT2) inhibitors, exhibits antidiabetic efficacy in rodent models. *J Pharmacol Exp Ther* 327: 268–276.
- Macdonald FR, Peel JE, Jones HB, Mayers RM, Westgate L, et al. (2010) The novel sodium glucose transporter 2 inhibitor dapagliflozin sustains pancreatic function and preserves islet morphology in obese, diabetic rats. *Diabetes Obes Metab* 12: 1004–1012.
- Chen L, Klein T, Leung PS (2012) Effects of combining linagliptin treatment with BI-38335, a novel SGLT2 inhibitor, on pancreatic islet function and inflammation in *db/db* mice. *Curr Mol Med* 12: 995–1004.
- Matsushita Y, Ogawa D, Wada J, Yamamoto N, Shikata K, et al. (2011) Activation of peroxisome proliferator-activated receptor delta inhibits streptozotocin-induced diabetic nephropathy through anti-inflammatory mechanisms in mice. *Diabetes* 60: 960–968.
- Okada S, Shikata K, Matsuda M, Ogawa D, Usui H, et al. (2003) Intercellular adhesion molecule-1-deficient mice are resistant against renal injury after induction of diabetes. *Diabetes* 52: 2586–2593.
- Ogawa D, Asanuma M, Miyazaki I, Tachibana H, Wada J, et al. (2011) High glucose increases metallothionein expression in renal proximal tubular epithelial cells. *Exp Diabetes Res* 2011: 534872.
- Kodera R, Shikata K, Kataoka HU, Takatsuka T, Miyamoto S, et al. (2011) Glucagon-like peptide-1 receptor agonist ameliorates renal injury through its anti-inflammatory action without lowering blood glucose level in a rat model of type 1 diabetes. *Diabetologia* 54: 965–978.
- Whaley JM, Tirmenstein M, Reilly TP, Poucher SM, Saye J, et al. (2012) Targeting the kidney and glucose excretion with dapagliflozin: preclinical and clinical evidence for SGLT2 inhibition as a new option for treatment of type 2 diabetes mellitus. *Diabetes Metab Syndr Obes* 5: 135–148.
- Jurczak MJ, Lee HY, Birkenfeld AL, Jornayvaz FR, Frederick DW, et al. (2011) SGLT2 deletion improves glucose homeostasis and preserves pancreatic beta-cell function. *Diabetes* 60: 890–898.
- Nagata T, Fukuzawa T, Takeda M, Fukazawa M, Mori T, et al. (2013) Tofogliflozin, a novel sodium-glucose co-transporter 2 inhibitor, improves renal and pancreatic function in *db/db* mice. *Br J Pharmacol* 170: 519–531.
- Vlassara H, Cai W, Chen X, Serrano EJ, Shobha MS, et al. (2012) Managing chronic inflammation in the aging diabetic patient with CKD by diet or sevelamer carbonate: a modern paradigm shift. *J Gerontol A Biol Sci Med Sci* 67: 1410–1416.
- Ohga S, Shikata K, Yozai K, Okada S, Ogawa D, et al. (2007) Thiazolidinedione ameliorates renal injury in experimental diabetic rats through anti-inflammatory effects mediated by inhibition of NF-kappaB activation. *Am J Physiol Renal Physiol* 292: F1141–1150.
- Tachibana H, Ogawa D, Matsushita Y, Bruemmer D, Wada J, et al. (2012) Activation of liver X receptor inhibits osteopontin and ameliorates diabetic nephropathy. *J Am Soc Nephrol* 23: 1835–1846.
- Forbes JM, Cooper ME (2013) Mechanisms of diabetic complications. *Physiol Rev* 93: 137–188.
- Kojima N, Williams JM, Takahashi T, Miyata N, Roman RJ (2013) Effects of a new SGLT2 inhibitor, luseogliflozin, on diabetic nephropathy in T2DN rats. *J Pharmacol Exp Ther* 345: 464–472.
- Vallon V, Rose M, Gerasimova M, Satriano J, Platt KA, et al. (2013) Knockout of Na-glucose transporter SGLT2 attenuates hyperglycemia and glomerular hyperfiltration but not kidney growth or injury in diabetes mellitus. *Am J Physiol Renal Physiol* 304: F156–167.
- Cherney DZ, Perkins BA, Soleymanlou N, Maione M, Lai V, et al. (2014) Renal hemodynamic effect of sodium-glucose cotransporter 2 inhibition in patients with type 1 diabetes mellitus. *Circulation* 129: 587–597.
- Tahara A, Kurosaki E, Yokono M, Yamajuku D, Kihara R, et al. (2013) Effects of SGLT2 selective inhibitor ipragliflozin on hyperglycemia, hyperlipidemia, hepatic steatosis, oxidative stress, inflammation, and obesity in type 2 diabetic mice. *Eur J Pharmacol* 715: 246–255.

## ORIGINAL ARTICLE

# Fluorescence virus-guided capturing system of human colorectal circulating tumour cells for non-invasive companion diagnostics

Kunitoshi Shigeyasu,<sup>1</sup> Hiroshi Tazawa,<sup>1,2</sup> Yuuri Hashimoto,<sup>1</sup> Yoshiko Mori,<sup>1</sup> Masahiko Nishizaki,<sup>1</sup> Hiroyuki Kishimoto,<sup>1</sup> Takeshi Nagasaka,<sup>1</sup> Shinji Kuroda,<sup>1</sup> Yasuo Urata,<sup>3</sup> Ajay Goel,<sup>4</sup> Shunsuke Kagawa,<sup>1</sup> Toshiyoshi Fujiwara<sup>1</sup>

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/gutjnl-2014-306957>).

<sup>1</sup>Department of Gastroenterological Surgery, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama, Japan

<sup>2</sup>Center for Innovative Clinical Medicine, Okayama University Hospital, Okayama, Japan

<sup>3</sup>Oncolys BioPharma, Inc., Tokyo, Japan

<sup>4</sup>Division of Gastroenterology, Department of Internal Medicine, Charles A. Sammons Cancer Center and Baylor Research Institute, Baylor University Medical Center, Dallas, Texas, USA

## Correspondence to

Professor Toshiyoshi Fujiwara, Department of Gastroenterological Surgery, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, 2-5-1 Shikata-cho, Kita-ku, Okayama 700-8558, Japan; [toshi\\_f@md.okayama-u.ac.jp](mailto:toshi_f@md.okayama-u.ac.jp)

Received 10 February 2014

Revised 20 April 2014

Accepted 8 May 2014

Published Online First

28 May 2014

## ABSTRACT

**Background** Molecular-based companion diagnostic tests are being used with increasing frequency to predict their clinical response to various drugs, particularly for molecularly targeted drugs. However, invasive procedures are typically required to obtain tissues for this analysis. Circulating tumour cells (CTCs) are novel biomarkers that can be used for the prediction of disease progression and are also important surrogate sources of cancer cells. Because current CTC detection strategies mainly depend on epithelial cell-surface markers, the presence of heterogeneous populations of CTCs with epithelial and/or mesenchymal characteristics may pose obstacles to the detection of CTCs.

**Methods** We developed a new approach to capture live CTCs among millions of peripheral blood leukocytes using a green fluorescent protein (GFP)-expressing attenuated adenovirus, in which the telomerase promoter regulates viral replication (OBP-401, TelomeScan).

**Results** Our biological capturing system can image epithelial and mesenchymal tumour cells with telomerase activities as GFP-positive cells. After sorting, direct sequencing or mutation-specific PCR can precisely detect different mutations in *KRAS*, *BRAF* and *KIT* genes in epithelial, mesenchymal or epithelial–mesenchymal transition-induced CTCs, and in clinical blood samples from patients with colorectal cancer.

**Conclusions** This fluorescence virus-guided viable CTC capturing method provides a non-invasive alternative to tissue biopsy or surgical resection of primary tumours for companion diagnostics.

## INTRODUCTION

The rapid evolution of genetic and genomic technologies in regards to predictive pharmacogenetic biomarkers for molecularly targeted therapies (eg, monoclonal antibodies and small-molecule tyrosine kinase inhibitors) have resulted in tremendous advances in personalised oncologic treatment.<sup>1</sup> The current commonly used biomarkers include human epidermal growth factor receptor 2 (HER2) for the use of trastuzumab in breast and gastric cancer,<sup>2,3</sup> *KRAS* for the use of cetuximab and panitumumab in colorectal cancer,<sup>4</sup> echinoderm microtubule associated protein like 4-anaplastic lymphoma kinase (EML4-ALK) for the use of crizotinib, and epidermal growth factor receptor (EGFR) for the use of

## Significance of this study

### What is already known about this subject?

- The molecular characterisation of circulating tumour cells (CTCs) based on genetic alterations facilitates the administration of molecular targeted drugs for preventing metastatic progression in individual patients with cancer.
- Heterogeneous populations of CTCs with epithelial and/or mesenchymal characteristics make it difficult to detect the entire CTC population because CTC detection mainly depends on epithelial cell surface markers.

### What are the new findings?

- Our fluorescence virus OBP-401 selectively labelled human CTCs with green fluorescent protein (GFP) among millions of peripheral blood leukocytes.
- Our biological capturing system can image both epithelial and mesenchymal tumour cells with telomerase activities as GFP-positive cells.

### How might it impact on clinical practice in the foreseeable future?

- Because current CTC detection strategies mainly depend on epithelial cell surface markers, the presence of heterogeneous populations of CTCs with epithelial and/or mesenchymal characteristics may pose obstacles to the detection of CTCs.
- The fluorescence virus-based biological capture system is a promising tool for monitoring genetic alterations in epithelial and mesenchymal types of CTCs.

erlotinib and gefitinib, in non-small cell lung cancer,<sup>5,6</sup> and BCR–ABL for the use of tyrosine kinase inhibitors in chronic myeloid leukaemia.<sup>7</sup> Companion diagnostic assays are designed to accompany specific therapies and help guide selection of patients according to expected drug responses. While the use of these assays has led to a shift in paradigms from disease-based therapeutic regimens to molecular target-based protocols,<sup>8,9</sup>



► <http://dx.doi.org/10.1136/gutjnl-2014-307647>



**To cite:** Shigeyasu K, Tazawa H, Hashimoto Y, et al. *Gut* 2015;**64**:627–635.

## GI cancer

many of these molecular diagnostic modalities have onerous requirements to obtain specimen, such as needle core biopsies or surgical sampling of tumour tissues that can be invasive.

Circulating tumour cells (CTCs), first described in 1869 by Ashworth,<sup>10</sup> are often present in the peripheral blood of patients with advanced cancers. However, as CTCs are very rare within the bloodstream, detection of CTCs can be difficult. The most commonly used CTC detection method is the CellSearch system,<sup>11 12</sup> which can enrich CTCs using magnetised antibodies that target the major epithelial cell surface marker, epithelial cell adhesion molecule (EpCAM). More recently, genetic analysis of the *EGFR* gene using the EpCAM-dependent CTC-chip detection system has been described for the surveillance of CTCs in patients with lung cancers.<sup>13</sup> CTCs are thought to contain the metastasis-initiating tumour cells that form metastatic colonies at distant organs,<sup>14 15</sup> but recent studies have suggested that there are heterogeneous populations that include CTCs with both epithelial and mesenchymal characteristics,<sup>16</sup> which are associated with the epithelial–mesenchymal transition (EMT).<sup>17</sup> Recently, EpCAM-positive and EpCAM-negative CTCs from patients with breast cancer have been shown to exhibit high potential to metastasise to the lung and brain, respectively, in nude mice.<sup>18 19</sup> In patients with colorectal cancer, not only captured cytokeratin (CK)-positive CTCs, but also co-captured CK-negative cells have been shown to possess complex aneuploidy.<sup>20</sup> Moreover, it has been reported that *plstin3* (*PLS3*), which is a novel marker for EMT, was detected in EpCAM-positive and EpCAM-negative CTCs in patients with colorectal cancer with distant metastasis.<sup>21</sup> These findings indicate the presence of CTCs without epithelial markers in patients with colorectal cancer. Therefore, development of a CTC capture system that functions independent of the epithelial cell marker is required to precisely assess the sensitivity of highly metastatic tumour cells to molecularly targeted drugs.

Epithelial and mesenchymal types of malignant tumour cells possess high telomerase activity to maintain the length of telomere during aberrant cell proliferation, suggesting the potential of telomerase activity as a general tumour marker<sup>22</sup> and therapeutic target.<sup>23</sup> We previously developed a green fluorescent protein (GFP)-expressing telomerase-specific replication-competent adenovirus (OBP-401, TelomeScan) that drives the adenoviral *E1A* and *E1B* genes under the *hTERT* gene promoter for telomerase-dependent virus replication. OBP-401 enables the visualisation of viable epithelial and mesenchymal types of human tumour cells with telomerase activity as GFP-positive cells.<sup>24 25</sup> OBP-401-mediated GFP labelling is a useful method to detect viable CTCs in patients with gastrointestinal cancers<sup>26 27</sup> and ovarian cancers.<sup>28</sup> The present study extends our previous work by exploring the potential of an OBP-401-based biological CTC capture system for the surveillance of genetic mutations in viable CTCs as a novel non-invasive companion diagnostic strategy.

## MATERIALS AND METHODS

### Cell lines

The human colorectal cancer cell lines, SW480, HCT116 and HT29; the human pancreatic cancer cell line, Panc1; the human lung cancer cell line, A549 and H1299; the human gastrointestinal stromal tumour (GIST) cell line, GIST882; and the human normal oesophageal fibroblasts, FEF3, were purchased from the American Type Culture Collection. All cell lines were cultured according to the manufacturer's specifications. There are four types of *KRAS* gene mutations (G12D, G12 V, G12S, G13D) in Panc1, SW480, A549 and HCT116 cells, respectively. HT29 cells have one mutation (V600E) in the *BRAF* gene, whereas

GIST882 cells harbour one mutation (K642E) in the *KIT* gene. Normal FEF3 cells have no mutations in the *KRAS*, *BRAF* or *KIT* genes.

To obtain the EMT-induced human cancer cells, A549 cells were treated with transforming growth factor  $\beta$  (TGF- $\beta$ ) (10 ng/mL) for 72 h. EMT induction was defined as a morphological change to spindle type and a change in the EMT-related marker expression, including downregulation of epithelial markers (EpCAM and E-cadherin) and upregulation of the mesenchymal marker (N-cadherin).

### Recombinant adenovirus

OBP-401 is a telomerase-specific replication-competent adenovirus variant, in which the *hTERT* gene promoter drives the expression of *E1A* and *E1B* genes that are linked to an internal ribosome entry site and in which the *GFP* gene is inserted into the E3 region under a cytomegalovirus (CMV) promoter (figure 1A).<sup>24–26</sup> OBP-401 was purified by ultracentrifugation using CsCl step gradients. Viral titres were determined by a plaque-forming assay using 293 cells, and the virus was stored at  $-80^{\circ}\text{C}$ .

### Immunocytochemical staining

The cells seeded on tissue culture chamber slides were fixed in 4% paraformaldehyde for 15 min on ice. The slides were subsequently incubated with the PE-conjugated mouse anti-EpCAM antibody (BioLegend, San Diego, California, USA) for 1 h on ice. Then the slides were analysed using an inverted fluorescence microscope (Olympus; Tokyo, Japan).

### Flow cytometry


The cells ( $1 \times 10^5$  cells) were labelled with primary mouse antibodies for EpCAM, E-cadherin, N-cadherin (BioLegend) and coxsackievirus and adenovirus receptor (CAR) for 30 min on ice and were analysed using flow cytometry (FACS Array; Becton Dickinson, Mountain View, California, USA).

### CTC model

CTC models were established by incubation with tumour cell lines (SW480, HCT116, HT29, Panc1, EMT-induced A549 and GIST882 cells) in 5 mL of blood (containing approximately  $3.5 \times 10^7$  white blood cells) from a healthy volunteer.

### DNA extraction from CTC models and clinical samples

The protocol for DNA extraction from the CTC models or clinical samples is shown in figure 2A and online supplementary figure S1. Approximately 5 mL of blood was incubated with lysis buffer containing ammonium chloride to remove the red blood cells (RBCs). These cancer cells were then infected with OBP-401 at  $1 \times 10^6$  plaque-forming units (PFU) and incubated for 24 h. Thereafter, the cell pellets were labelled with anti-CD45 antibody conjugated with PE and sequentially sorted by FACS Aria (Becton Dickinson, San Jose, California, USA). We set the P1 gate to obtain viable cells, the P2 gate to detect GFP-positive cells without intrinsic fluorescence and the P3 gate to detect only GFP-positive tumour cells without the haematopoietic CD45 marker. Cells in the P2 or P3 gates were collected and stored temporarily at  $-30^{\circ}\text{C}$ . DNA was extracted from captured cells using a QiaAMP DNA Mini kit (Qiagen, Valencia, California, USA). The DNA solution was mixed with DNA polymerase, and each primer was subjected to PCR analysis. Five-millilitre blood samples were collected with consent from patients with colorectal cancer, according to a protocol



**Figure 1** OBP-401-mediated green fluorescent protein (GFP) expression in human cancer cells with different levels of EpCAM expression. (A) Schematic DNA structure of OBP-401 (TelomeScan). OBP-401 is a telomerase-specific replication-competent adenovirus variant in which the hTERT promoter element drives expression of the *E1A* and *E1B* genes linked with internal ribosome entry sites (IRES). The *GFP* gene is inserted under the cytomegalovirus (CMV) promoter into the E3 region. (B) Immunofluorescence staining of epithelial cell adhesion molecule (EpCAM) in four human cancer cell lines (Panc1, SW480, HCT116, and HT29 cells). EpCAM expression under fluorescence microscopy (top panels) and phase-contrast microscopy (bottom panels). Original magnification:  $\times 100$ . (C) Flow cytometric analysis of EpCAM expression in four human cancer cell lines. Cells are incubated with anti-EpCAM antibody. An isotype-matched normal mouse IgG1 is used as a control. (D) Cells re-infected with OBP-401 at multiplicity of infection (MOI) of 10, 100 or 1000 plaque-forming units per cell and assessed for GFP expression under fluorescence microscopy 24 h after infection. (E) Expression of CAR is analysed using flow cytometry in four human cancer cell lines.

approved by the institutional review board at Okayama University Graduate School (Receipt No. 1537).

#### Gene mutation analysis by direct sequencing

Taq polymerase, forward primer and reverse primer were mixed with eluted DNA solution, and DNA was amplified using the PCR Thermal Cycler. Primer sequences and PCR settings are shown in online supplementary table S1. Using the PCR products, the sequence of each gene was analysed with an ABI PRISM 3100 Genetic Analyzer (Life Technologies, Carlsbad, California, USA).

#### Gene mutation analysis by ASB-PCR

ASB-PCR for the KRAS and BRAF genes was performed with a primer set of a TaqMan Mutation Detection Assays (Applied Biosystems, Foster City, California, USA), as described in a previous report.<sup>29</sup> This assay amplifies only mutant alleles with mutant-specific primers and prevents the amplification of wild-type alleles using blocking oligonucleotides. Genetic mutations of target genes were analysed with the StepOnePlus real-time PCR system (Applied Biosystems). Genotyping Master mix and Mutation Detection Assay were mixed with two sets of eluted DNA solution, and this mixture was applied to real-time PCR analysis. The mutation detection method was customised as follows. The PCR cycle number was set to 70 for the efficient amplification of small copy numbers of target genes. A total cell count was restricted to less than 50 cells/well to prevent non-specific amplification of wild-type alleles. The sensitivity and

specificity were analysed using a mixture of KRAS/BRAF wild-type and mutant cells. Genetic mutation was recognised as positive when the amplification for mutant alleles using a specific primer was detected.

## RESULTS


### Fluorescence imaging of human cancer cells with differential EpCAM expression

OBP-401 (TelomeScan) was previously constructed by inserting the *GFP* gene under the control of the CMV promoter at the deleted E3 region of the telomerase-specific replication-selective type 5 adenovirus OBP-301 (Telomelysin) (figure 1A). To assess the potential of OBP-401-mediated biological imaging, we used four epithelial types of human cancer cell lines (Panc1, SW480, HCT116 and HT29) that differentially express EpCAM in immunocytochemistry (figure 1B) and fluorescence-activated cell sorting (FACS) analysis (figure 1C). All cell lines could be visualised by OBP-401-induced GFP expression in a dose-dependent manner, independently of EpCAM expression (figure 1D). The expression level of CAR, which is associated with adenovirus infectivity, was similar among all cell lines (figure 1E). These results suggest that OBP-401-mediated biological imaging is a useful method to detect human cancer cells regardless of high or low EpCAM expression.

### Fluorescence-guided isolation of CTCs with multi-laser FACS

We used OBP-401 to establish a simple ex vivo method to capture viable human CTCs in the peripheral blood for genetic

## GI cancer



**Figure 2** A simple fluorescence virus-guided capturing system of circulating tumour cells (CTCs). (A) Cell isolation steps in the OBP-401-based CTC capturing system. CTC models containing spiked human cancer cells in 5 mL of blood sample or clinical blood samples obtained from patients with cancer are incubated with red blood cell (RBC) lysis buffer for 6 min. After centrifugation, cell pellets are then infected with OBP-401 at  $1 \times 10^6$  plaque-forming units and incubated for 24 h. Thereafter, cells are incubated with anti-CD45 antibody, and the cell pellet is sorted by fluorescence-activated cell sorting (FACS). DNA extracted from FACS-sorted green fluorescent protein (GFP)-positive cells is subjected to direct sequencing or allele-specific blocker PCR (ASB-PCR) analysis. (B) Each gate is set to capture the GFP-positive CTCs by FACS analysis. After isolating only viable cells at the P1 gate, the P2 and P3 gates are set to exclude the intrinsic fluorescence-positive cells and CD45-positive normal blood cells, respectively. (C) Representative image of GFP-positive CTCs in blood sample containing SW480 cells after infection with OBP-401. Original magnification:  $\times 200$ .

analysis. By spiking a certain number of human cancer cells that have different types of genetic mutations in the *KRAS* or *BRAF* gene in 5 mL of blood from healthy volunteers, we made CTC models with different types of genetic mutations. As illustrated in figure 2A and online supplementary figure S1, following the lysis of RBCs in 5 mL aliquots of CTC models or whole blood samples obtained from patients, the cell pellets were subsequently incubated with OBP-401 at  $1 \times 10^6$  PFU for 24 h, labelled with anti-CD45 antibody conjugated with phycoerythrin (PE), and sequentially sorted by FACS. In preliminary experiments using CTC models, we found suitable conditions for sorting only GFP-positive CTCs by excluding auto-fluorescent allophycocyanin-positive cells at the P2 gate and haematopoietic CD45-positive cells at the P3 gate (figure 2B). The GFP-positive cells could be detected in the CTC model under a fluorescence microscope (figure 2C).


#### Genetic analysis of OBP-401-labelled GFP-positive cells using direct sequencing

FACS-isolated GFP-positive CTCs at the P3 gate were analysed genetically by direct sequencing (see online supplementary table 1). The expected genetic mutations in the *KRAS* or *BRAF* gene were precisely detected in all CTC models comprising four human cancer cell lines by direct sequencing (figure 3A and see online supplementary figure S2), indicating that the OBP-401-based biological capture system is effective for the

collection of CTCs expressing various levels of the EpCAM marker. Recent studies have demonstrated that a heterogeneous population of CTCs is present within individual patients with cancer and that these CTCs have epithelial and mesenchymal markers, suggesting the diverse genetic variations with wild-type and mutant-type genes in the populations of CTCs. To evaluate the minimum purity limitation of mutant-type CTCs for genetic analysis using direct sequencing, SW480 cells (*KRAS* G12V mutant) were mixed with H1299 cells (*KRAS* wild-type) at a 50%, 40%, 30%, 20% or 10% purity ratio. *KRAS* gene mutations could be detected by direct sequencing only in samples containing more than a 30% purity ratio of SW480 cells (figure 3B). Thus, high purity of mutant-type CTCs in heterogeneous populations is necessary for detection of genetic alterations by direct sequencing.

#### Genetic analysis of OBP-401-labelled GFP-positive cells using ASB-PCR

To further increase the sensitivity to detect genetic alterations in heterogeneous populations of CTCs, we next evaluated the potential of the Allele-Specific Blocker (ASB)-PCR method using four types of mutation-specific primers for the *KRAS* or *BRAF* genes. Before analysing the human cancer cells, we confirmed that there was no amplification of PCR products in human normal fibroblasts with wild-type *KRAS* and *BRAF* genes or in blood obtained from normal healthy volunteers by ASB-PCR



**Figure 3** Genetic mutation analysis of human circulating tumour cells (CTCs) by direct sequencing and mutation-specific PCR. (A) Detection of *KRAS* or *BRAF* gene mutation in the CTC models containing 10 human cancer cells by direct sequencing of green fluorescent protein (GFP)-positive cells at the P3 gate. The number of cells in the P3 gate and the mutation pattern in each model are indicated. (B) The minimal purity of tumour cells for direct sequencing to detect the expected gene mutations is evaluated. SW480 (*KRAS* G12V) cells were mixed with H1299 (*KRAS* wild-type) cells at 50%, 40%, 30%, 20% and 10% of purity ratios. DNA is extracted from cell mixtures, and the *KRAS* gene mutation is analysed by direct sequencing. (C) Allele-specific blocker (ASB)-PCR-mediated detection of *KRAS* and *BRAF* gene mutations in GFP-positive cells at the P2 or P3 gate in the CTC models containing as few as 10 SW480 cells and HT29 cells. When *KRAS* and *BRAF* genes contain targeted mutations, mutation-specific curves cross their threshold of detection. (D) Detection of *KRAS* gene mutation by direct sequencing of GFP-positive cells at the P2 gate without CD45 depletion requires at least 50 SW480 cells in the CTC model.

with mutation-specific primers (see online supplementary figure S3).

When we analysed five human cancer cells mixed with 100 human normal fibroblasts at a purity ratio of approximately 5%, ASB-PCR, using all types of primers, detected the expected mutations in the GFP-positive cells (see online supplementary table 2). In the CTC models containing 10 human cancer cells

with different types of *KRAS* and *BRAF* gene mutations, ASB-PCR analysis detected the expected genetic mutations in the GFP-positive cells at the P3 gate (table 1). Moreover, ASB-PCR analysis could detect the genetic alterations in the GFP-positive cells at the P2 gate without exclusion of CD45-positive normal blood cells (figure 3C), whereas at least 50 tumour cells were required for direct sequencing in the

**Table 1** Data for mutation-specific PCR for the genetic analysis of CTC models

CTC model		FACS analysis				Genetic analysis				
Cancer cells	Cell type	Gene status	Number of cancer cells	Gate	Number of GFP-positive cells	Purity of cancer cells (%)	Primer	Amplification	Ct values	
									1st PCR	2nd PCR
Panc1	Epithelial	KRAS G12D	10	P2	29	34.5	KRAS G12D	+	37.1	36.1
				P3	6	100.0	KRAS G12D	+	35.3	38.2
SW480	Epithelial	KRAS G12V	10	P2	105	9.5	KRAS G12V	+	45.0	56.5
				P3	13	76.9	KRAS G12V	+	41.6	52.0
HCT116	Epithelial	KRAS G13D	10	P2	23	43.5	KRAS G13D	+	47.5	37.2
				P3	18	55.6	KRAS G13D	+	37.0	44.0
HT29	Epithelial	BRAF V600E	10	P2	34	29.4	BRAF V600E	+	34.0	NA
				P3	9	100.0	BRAF V600E	+	40.0	NA
EMT-induced A549	Mesenchymal	KRAS G12S	10	P2	77	13.0	KRAS G12S	+	41.9	43.5
				P3	17	58.8	KRAS G12S	+	51.1	64.7

CTC, circulating tumour cell; EMT, epithelial-mesenchymal transition; FACS, fluorescence-activated cell sorting; GFP, green fluorescent protein; NA, not amplified.

## GI cancer

presence of CD45-positive cells at the P2 gate (figure 3D). These results suggest that the ASB-PCR method is more simple and sensitive than direct sequencing for detection of genetic alterations in heterogeneous populations of CTCs.

### Fluorescence-guided capture of EMT-induced and mesenchymal CTCs


Induction of EMT in CTCs has recently been demonstrated in patients with advanced breast cancers.<sup>17</sup> EMT-induced CTCs frequently formed metastatic colonies in the brain and lung of nude mice,<sup>19</sup> suggesting that highly malignant EMT-induced CTCs can be detected to predict metastatic progression in patients with cancer. We used A549 human lung cancer cells with *KRAS* gene mutation (G12S) and EpCAM-negative GIST882 mesenchymal human tumour cells with the *KIT* gene mutation (K642E), which is frequently mutated in more than 70% of GISTs.<sup>30</sup> OBP-401 infection efficiently induced GFP expression in both cell lines in a dose-dependent manner (figure 4A).

When treated with the EMT inducer, TGF- $\beta$ , A549 cells showed spindle-shape morphological changes (figure 4B) and altered EMT-related biomarker expression, such as EpCAM and E-cadherin downregulation and N-cadherin upregulation

(figure 4C). In contrast, CAR expression was not affected after TGF- $\beta$  treatment (figure 4C). Therefore, OBP-401 efficiently induced GFP expression in the TGF- $\beta$ -treated A549 cells (figure 4D). In addition, GIST882 cells were confirmed to be EpCAM negative (figure 4E). When 10 EMT-induced A549 cells were spiked in blood samples, the expected genetic mutation (G12S) in the *KRAS* gene was detected by direct sequencing and by ASB-PCR analysis (figure 4F–G and table 1). In contrast, the expected *KIT* gene mutation could be detected at the P3 gate by direct sequencing in the CTC model containing 100 GIST882 cells (figure 4F) but not in that with 10 cells, presumably due to low expression of CAR. These results suggest that the targeted genetic mutations in EMT-induced and mesenchymal CTCs are also detectable by the OBP-401-based CTC capture system, although the sensitivity is dependent on the CAR expression.

### Detection of genetic mutations in CTCs in patients with colorectal cancer

Finally, the blood samples obtained from eight patients with *KRAS*- or *BRAF*-mutated colorectal cancers were analysed by the OBP-401-based CTC capture system and by ASB-PCR



**Figure 4** Fluorescence virus-guided capture and genetic mutation analysis of human mesenchymal or epithelial-mesenchymal transition (EMT)-induced tumour cells. (A) A549 human lung cancer cells and GIST882 human gastrointestinal stromal tumour cells are infected with OBP-401 at a multiplicity of infection (MOI) of 10, 100 or 1000 plaque-forming units (PFU) per cell. Green fluorescent protein (GFP) expression is assessed under fluorescence microscope 24 h after virus infection. (B) Morphological change of A549 cells treated with transforming growth factor  $\beta$  (TGF- $\beta$ ). A549 cells are treated with TGF- $\beta$  (10 ng/mL) for 72 h and stained with crystal violet. Original magnification:  $\times 200$ . (C) Flow cytometric analysis of epithelial (epithelial cell adhesion molecule (EpCAM) and E-cadherin) and mesenchymal (N-cadherin) cell surface marker and CAR expression in A549 cells treated with or without TGF- $\beta$ . (D) GFP expression in TGF- $\beta$ -treated A549 cells after infection with OBP-401 at an MOI of 100 PFU per cell for 24 h. Original magnification:  $\times 200$ . (E) Flow cytometric analysis of epithelial (EpCAM and E-cadherin) and mesenchymal (N-cadherin) cell-surface markers and coxsackievirus and adenovirus receptor (CAR) expression in GIST882 cells. (F) Detection of *KRAS* and *KIT* gene mutations by direct sequencing of GFP-positive cells at the P3 gate requires 10 TGF- $\beta$ -treated EMT-induced A549 cells and 100 GIST882 cells in the CTC models, respectively. (G) Detection of *KRAS* gene mutations in GFP-positive cells at the P2 or P3 gate in the CTC models containing as few as 10 of TGF- $\beta$ -treated A549 cells by allele-specific blocker (ASB)-PCR. Mutation-specific curves for the *KRAS* gene cross their threshold of detection. (H) Representative CT images of patients with colon cancer and lung, spleen and ovary metastases. The primary tumours and CTCs show the *BRAF* V600E mutation.

technology. In preliminary experiments, the number of GFP-positive cells at the P3 gate was less than 10 cells in some clinical blood samples and, therefore, we performed ASB-PCR analysis using GFP-positive cells at the P2 gate. Among the eight blood samples from patients with various stages of colorectal cancer, the same *KRAS* and *BRAF* gene mutations as in the primary tumours were detected in the CTCs of two patients with advanced colorectal cancer (figure 4H and table 2). The other six patients showed no detectable genetic abnormalities in blood samples, although *KRAS* gene mutations were observed in their primary tumours. Three patients without metastatic lesions did not have a large CTC count, and chemotherapeutic treatment in the other three patients with metastatic disease may have resulted in a reduced number of CTCs. Although further large-scale clinical trials are required, our results suggest that the OBP-401-based telomerase-dependent biological CTC capture system is useful for genetic analysis of CTCs in blood samples from patients with cancer.

## DISCUSSION

The co-development of a targeted therapy together with its companion diagnostic test, which guides selection of patients and provides surrogate markers to monitor responses, is a key part of personalised medicine. The selection of targeted therapies for individual patients is currently made by analysing the primary tumours, although there are very few cells within the primary tumours that are responsible for metastasis or recurrence, and these cells may have additional genetic abnormalities. The present study demonstrated that CTCs obtained non-invasively are a promising alternative to surgically resected or biopsied tumour tissues for real-time molecular characterisation. A telomerase-dependent biological CTC capture system was clinically useful for the detection of mutations in different target genes, such as *KRAS*, *BRAF* and *KIT*, even in EpCAM-negative cells among highly heterogeneous CTC populations.

We applied telomerase-specific OBP-401 to selectively label human neoplastic cells with GFP signals and confirmed its broad infectivity independent of EpCAM expression, which was consistent with observations from our previous reports that OBP-401 induced GFP expression in epithelial and mesenchymal types of tumour cells.<sup>24 25</sup> Recent studies demonstrated that highly metastatic tumour cells are involved in EpCAM-positive and EpCAM-negative subpopulations of CTCs in the blood of patients with breast cancer.<sup>18 19</sup> During anticancer treatment, the characteristics of CTCs dynamically alternate between epithelial and mesenchymal types of CTCs within individual

patients with cancer.<sup>17</sup> Further, platelet-derived TGF- $\beta$  secretion induces EMT with metastatic potential in CTCs.<sup>31</sup> These findings indicate that single CTCs frequently turn the EMT switch on or off in the microenvironment of the bloodstream. In contrast, high telomerase activity is a general functional biomarker for stabilisation of the telomere in epithelial and mesenchymal malignant tumour cells during aberrant proliferation. In fact, high *hTERT* mRNA levels have been detected in the blood samples of patients with cancer.<sup>32–34</sup> Moreover, hTERT overexpression has been shown to be positively associated with EMT induction in human cancer cells.<sup>35</sup> When the telomerase activity in the CTCs is suppressed in circulating cells, these CTCs undergo programmed cell death (ie, apoptosis or senescence). Thus, the telomerase activity may be superior to unstable epithelial cell marker as a general biomarker for the detection of viable CTCs in the blood. Moreover, GFP-labelled CTCs by OBP-401 infection are considered to be useful for direct determination of drug sensitivity and metastatic potential, and determination of tumour heterogeneity.<sup>36–39</sup>

A number of approaches based on the physical and biological properties of CTCs have been studied to distinguish CTCs from surrounding normal haematopoietic cells and to capture them for further analysis. The CellSearch system, which is the only test approved by the US Food and Drug Administration to detect CTCs, uses magnetised antibodies against EpCAM for positive selection and uses CD45 for leukocyte depletion. Another popular technology for CTC enrichment is a microfluidic-based device called the CTC-chip; this device can isolate and analyse CTCs using EpCAM-coated microposts. Our OBP-401-based CTC detection has been previously compared with the CellSearch assay in patients with metastatic breast cancer.<sup>40</sup> Although both assays exhibited comparable detection rates, the number of CTC-positive cells between both assays was not significantly correlated. Nine out of 50 (18%) cases were positive by both methods, while 12 (24%) and 18 (36%) patients showed positive cells with the OBP-401 assay and the CellSearch assays individually, respectively. We speculate that CTCs detected by OBP-401 primarily detect EpCAM-negative tumour cells while the CellSearch method detects epithelial non-tumour cells as well, including circulating fibroblasts.

Our strategy involves conventional FACS to capture OBP-401-labelled GFP-positive CTCs. OBP-401 infection increases the signal-to-background ratio as a tumour-specific probe, because the fluorescence signal can be amplified only in viable human tumour cells by viral replication. We excluded the autofluorescence-positive cells at the P2 gate and the

**Table 2** Data for mutation-specific PCR for the genetic analysis of patient samples

Patients		FACS analysis				Genetic analysis			
Tumour site	Stage	Gene status of primary tumour	Metastasis	Gate	Number of GFP-positive cells	Primer	Amplification	Ct values	
								1st PCR	2nd PCR
Colon	I	<i>KRAS</i> G13D	None	P2	6	<i>KRAS</i> G13D	–	NA	NA
Colon	II	<i>KRAS</i> G13D	None	P2	20	<i>KRAS</i> G13D	–	NA	NA
Colon	II	<i>KRAS</i> G12D	Liver	P2	95	<i>KRAS</i> G12D	+	55.1	61.0
Colon	III	<i>KRAS</i> G13D	None	P2	913	<i>KRAS</i> G13D	–	NA	NA
Colon	III	<i>BRAF</i> V600E	Lung, spleen, ovary	P2	138	<i>BRAF</i> V600E	+	63.0	NA
Colon	IV	<i>KRAS</i> G12D	Liver	P2	14	<i>KRAS</i> G12D	–	NA	NA
Colon	IV	<i>KRAS</i> G12V	Liver	P2	74	<i>KRAS</i> G12V	–	NA	NA
Colon	IV	<i>KRAS</i> G12V	Lung	P2	53	<i>KRAS</i> G12V	–	NA	NA

CTC, circulating tumour cell; FACS, fluorescence-activated cell sorting; GFP, green fluorescent protein; NA, not amplified.



haematopoietic CD45-positive cells at the P3 gate. When at least 10 human cancer cells were spiked in 5 mL of blood from a healthy volunteer, the number of GFP-positive cells detected at the P3 gate was almost the same as the number of spiked tumour cells, suggesting that the P3 gate contains pure CTCs. However, the P2 gate may be contaminated with non-CTC cells. Indeed, ASB-PCR analysis detected the expected gene mutations in the *KRAS* and *BRAF* genes at the P2 gate, whereas the P3 gate was necessary when direct sequencing was applied. Recently, a combination of the CellSearch system and genetic analysis was also performed to detect genetic mutations in rare CTCs from patients with cancer. Mostert *et al*<sup>41</sup> compared the three types of PCR-based genetic analysis of CTCs, and ASB-PCR, used in our study, was the most sensitive method for detecting *KRAS* and *BRAF* gene mutations in the CTCs from patients with metastatic colorectal cancers. In addition, as our data demonstrated that direct sequencing was limited if CTC-derived DNA had more than 30% purity, we conclude that, together with FACS-isolated OBP-401-infected GFP-expressing CTCs, ASB-PCR is a suitable assay for non-invasive companion diagnostics in patients with cancer. The specificity of the ASB-PCR assay allowed us to use the P2 gate for clinical samples even in the presence of non-CTC cells.

Mutation in *KRAS* and *BRAF* genes is highly associated with resistance to the anti-EGFR antibody, cetuximab, in patients with colorectal cancer.<sup>42–43</sup> In fact, the appearance of *KRAS* gene mutant DNA is associated with resistance to cetuximab in patients with *KRAS* wild-type colorectal cancers.<sup>44</sup> In patients with colorectal cancer, the frequency of the *KRAS* and *BRAF* gene mutations is significantly higher in liver metastasis than in primary tumours,<sup>45</sup> and *KRAS* and *BRAF* gene mutant status is significantly associated with poor outcomes.<sup>46</sup> These findings suggest that genetic analysis for the *KRAS* and *BRAF* gene mutation in CTCs can be used as a 'liquid biopsy' to monitor resistance to cetuximab and to predict metastatic potential in patients with *KRAS* wild-type colorectal cancers.

It is also worth noting that the OBP-401-based biological CTC capture system is applicable to the genetic analysis of CTCs with mesenchymal characteristics, including GISTs and osteosarcomas,<sup>25</sup> although the CellSearch system is also useful for detection of epithelial CTCs. Approximately 80% of GIST cells harbour a mutation in the *KIT* gene,<sup>30</sup> which is significantly associated with disease recurrence and poor outcomes.<sup>47</sup> Recently, the small-molecule tyrosine-kinase inhibitor imatinib has been shown to be effective against *KIT*-mutated GIST that is refractory to conventional chemotherapy.<sup>48</sup> In contrast, bone and soft tissue sarcoma cells, which make up one of the most notorious types of malignant mesenchymal tumours, are also detectable as GFP-positive cells by OBP-401 infection.<sup>25</sup> Frequent lung metastasis has been shown to be a poor prognostic factor in patients with osteosarcoma, but the potential of CTC enumeration in patients with osteosarcoma remains to be elucidated. Thus, the characterisation of CTCs, using the OBP-401-based biological CTC capture system, may be a useful strategy for monitoring metastatic progression in patients with GIST or osteosarcomas, as well as those with epithelial malignant tumours.

The combination of the OBP-401-based CTC capture system and genetic analysis using ASB-PCR detected *KRAS* and *BRAF* mutations in blood samples obtained from patients with colorectal cancer, and these mutations were identical to those seen in the primary tumours. This novel 'liquid biopsy' via a simple blood test could be carried out in real time and enables optimised and timely decisions for therapeutic intervention. However, the technology has to be further validated in large

clinical studies with defined endpoints. In addition, one limitation of our study was that it was difficult for ASB-PCR to detect uncommon genetic abnormalities. Regardless, when frequently occurring genetic mutations are targeted for the surveillance of CTCs, the ASB-PCR method would be a useful and highly sensitive method for detecting small numbers of CTCs with genetic mutations. In contrast, if the identification of genetic traits in highly metastatic CTCs is the main goal, a genome-wide approach should be considered for the genetic analysis of CTCs. For example, genome-wide transcriptome analysis has been performed to identify a wide range of copy number alterations in entire CTC populations, using array-comprehensive genomic hybridisation (aCGH).<sup>49</sup> Moreover, genetic analysis in a single CTC has been recently used to clarify global gene alterations using aCGH and next-generation sequencing.<sup>50–51</sup> Thus, the comprehensive analysis of genetic alterations in individual CTCs from patients with cancer would provide novel insight into the identification of genetic signatures associated with metastatic progression.

In summary, we established a telomerase-dependent biological CTC capture system for genotyping of epithelial, mesenchymal, and EMT-induced types of CTCs using OBP-401 and FACS analysis. This technology facilitates the surveillance of genetic alterations in viable CTCs in patients with cancer. Large-scale clinical studies of this strategy are warranted.

**Acknowledgements** We thank Yukinari Isomoto and Tomoko Sueishi for their technical support.

**Contributors** Conception and design: HT, SK, TF; development of methodology: KS, HT, YH, TN; acquisition of data: KS, YH, YM; analysis and interpretation of data: KS, HT, HK, SK, TF; writing, review and/or revision of the manuscript: KS, HT, AG, TF; administrative, technical, or material support: MN, SK, YU; study supervision: TF.

**Funding** This study was supported by grants-in-aid from the Ministry of Education Culture, Sports, Science and Technology, Japan (T. Fujiwara, H. Tazawa, S. Kagawa) and grants from the Ministry of Health, Labour and Welfare, Japan (T. Fujiwara).

**Competing interests** Yasuo Urata is the president and CEO of Oncolys BioPharma, Inc., the manufacturer of OBP-401 (TelomeScan). Hiroshi Tazawa and Toshiyoshi Fujiwara are consultants for Oncolys BioPharma, Inc. The other authors have no real or potential conflicts of interest to declare.

**Ethics approval** Okayama University.

**Provenance and peer review** Not commissioned; externally peer reviewed.

## REFERENCES

- Ong FS, Das K, Wang J, *et al*. Personalized medicine and pharmacogenetic biomarkers: progress in molecular oncology testing. *Expert Rev Mol Diagn* 2012;12:593–602.
- Hurvitz SA, Hu Y, O'Brien N, *et al*. Current approaches and future directions in the treatment of HER2-positive breast cancer. *Cancer Treat Rev* 2013;39:219–29.
- Pietrantonio F, De Braud F, Da Prat V, *et al*. A review on biomarkers for prediction of treatment outcome in gastric cancer. *Anticancer Res* 2013;33:1257–66.
- Anderson SM. Laboratory methods for *KRAS* mutation analysis. *Expert Rev Mol Diagn* 2011;11:635–42.
- Kwak EL, Bang YJ, Camidge DR, *et al*. Anaplastic lymphoma kinase inhibition in non-small-cell lung cancer. *N Engl J Med* 2010;363:1693–703.
- Maemondo M, Inoue A, Kobayashi K, *et al*. Gefitinib or chemotherapy for non-small-cell lung cancer with mutated EGFR. *N Engl J Med* 2010;362:2380–8.
- Cross NC, White HE, Muller MC, *et al*. Standardized definitions of molecular response in chronic myeloid leukemia. *Leukemia* 2012;26:2172–5.
- Zieba A, Grannas K, Soderberg O, *et al*. Molecular tools for companion diagnostics. *N Biotechnol* 2012;29:634–40.
- Fridlyand J, Simon RM, Walrath JC, *et al*. Considerations for the successful co-development of targeted cancer therapies and companion diagnostics. *Nat Rev Drug Discov* 2013;12:743–55.
- Ashworth T. A case of cancer in which cells similar to those in the tumours were seen in blood after death. *Aust Med J* 1869;14:146–9.
- Cristofanilli M, Budd GT, Ellis MJ, *et al*. Circulating tumor cells, disease progression, and survival in metastatic breast cancer. *N Engl J Med* 2004;351:781–91.
- Miller MC, Doyle GV, Terstappen LW. Significance of circulating tumor cells detected by the CellSearch system in patients with metastatic breast colorectal and prostate cancer. *J Oncol* 2010;2010:617421.

- 13 Maheswaran S, Sequist LV, Nagrath S, *et al.* Detection of mutations in EGFR in circulating lung-cancer cells. *N Engl J Med* 2008;359:366–77.
- 14 Chaffer CL, Weinberg RA. A perspective on cancer cell metastasis. *Science* 2011;331:1559–64.
- 15 Joosse SA, Pantel K. Biologic challenges in the detection of circulating tumor cells. *Cancer Res* 2013;73:8–11.
- 16 Konigsberg R, Obermayr E, Bises G, *et al.* Detection of EpCAM positive and negative circulating tumor cells in metastatic breast cancer patients. *Acta Oncol* 2011;50:700–10.
- 17 Yu M, Bardia A, Wittner BS, *et al.* Circulating breast tumor cells exhibit dynamic changes in epithelial and mesenchymal composition. *Science* 2013;339:580–4.
- 18 Baccelli I, Schneeweiss A, Riethdorf S, *et al.* Identification of a population of blood circulating tumor cells from breast cancer patients that initiates metastasis in a xenograft assay. *Nat Biotechnol* 2013;31:539–44.
- 19 Zhang L, Ridgway LD, Wetzel MD, *et al.* The identification and characterization of breast cancer CTCs competent for brain metastasis. *Sci Transl Med* 2013;5:180ra48.
- 20 Pecot CV, Bischoff FZ, Mayer JA, *et al.* A novel platform for detection of CK+ and CK– CTCs. *Cancer Discov* 2011;1:580–6.
- 21 Yokobori T, Iinuma H, Shimamura T, *et al.* Platin3 is a novel marker for circulating tumor cells undergoing the epithelial–mesenchymal transition and is associated with colorectal cancer prognosis. *Cancer Res* 2013;73:2059–69.
- 22 Hiyama E, Hiyama K. Telomerase as tumor marker. *Cancer Lett* 2003;194:221–33.
- 23 Kyo S, Inoue M. Complex regulatory mechanisms of telomerase activity in normal and cancer cells: how can we apply them for cancer therapy? *Oncogene* 2002;21:688–97.
- 24 Kishimoto H, Kojima T, Watanabe Y, *et al.* In vivo imaging of lymph node metastasis with telomerase-specific replication-selective adenovirus. *Nat Med* 2006;12:1213–19.
- 25 Sasaki T, Tazawa H, Hasei J, *et al.* A simple detection system for adenovirus receptor expression using a telomerase-specific replication-competent adenovirus. *Gene Therapy* 2013;20:112–18.
- 26 Kojima T, Hashimoto Y, Watanabe Y, *et al.* A simple biological imaging system for detecting viable human circulating tumor cells. *J Clin Invest* 2009;119:3172–81.
- 27 Ito H, Inoue H, Sando N, *et al.* Prognostic impact of detecting viable circulating tumour cells in gastric cancer patients using a telomerase-specific viral agent: a prospective study. *BMC Cancer* 2012;12:346.
- 28 Takakura M, Kyo S, Nakamura M, *et al.* Circulating tumour cells detected by a novel adenovirus-mediated system may be a potent therapeutic marker in gynaecological cancers. *Br J Cancer* 2012;107:448–54.
- 29 Morlan J, Baker J, Sinicropi D. Mutation detection by real-time PCR: a simple, robust and highly selective method. *PLoS One* 2009;4:e4584.
- 30 Heinrich MC, Blanke CD, Druker BJ, *et al.* Inhibition of KIT tyrosine kinase activity: a novel molecular approach to the treatment of KIT-positive malignancies. *J Clin Oncol* 2002;20:1692–703.
- 31 Labelle M, Begum S, Hynes RO. Direct signaling between platelets and cancer cells induces an epithelial-mesenchymal-like transition and promotes metastasis. *Cancer Cell* 2011;20:576–90.
- 32 Wang JY, Lin SR, Wu DC, *et al.* Multiple molecular markers as predictors of colorectal cancer in patients with normal perioperative serum carcinoembryonic antigen levels. *Clin Cancer Res* 2007;13:2406–13.
- 33 Uen YH, Lin SR, Wu DC, *et al.* Prognostic significance of multiple molecular markers for patients with stage II colorectal cancer undergoing curative resection. *Ann Surg* 2007;246:1040–6.
- 34 Bertorelle R, Briarava M, Rampazzo E, *et al.* Telomerase is an independent prognostic marker of overall survival in patients with colorectal cancer. *Br J Cancer* 2013;108:278–84.
- 35 Liu Z, Li Q, Li K, *et al.* Telomerase reverse transcriptase promotes epithelial–mesenchymal transition and stem cell-like traits in cancer cells. *Oncogene* 2013;32:4203–13.
- 36 Mitsiades CS, Mitsiades NS, Bronson RT, *et al.* Fluorescence imaging of multiple myeloma cells in a clinically relevant SCID/NOD in vivo model: biologic and clinical implications. *Cancer Res* 2003;63:6689–96.
- 37 Kolostova K, Pinterova D, Hoffman RM, *et al.* Circulating human prostate cancer cells from an orthotopic mouse model rapidly captured by immunomagnetic beads and imaged by GFP expression. *Anticancer Res* 2011;31:1535–9.
- 38 Menen RS, Pinney E, Kolostova K, *et al.* A rapid imageable in vivo metastasis assay for circulating tumor cells. *Anticancer Res* 2011;31:3125–8.
- 39 Menen R, Zhao M, Zhang L, *et al.* Comparative chemosensitivity of circulating human prostate cancer cells and primary cancer cells. *Anticancer Res* 2012;32:2881–4.
- 40 Kim SJ, Masago A, Tamaki Y, *et al.* A novel approach using telomerase-specific replication-selective adenovirus for detection of circulating tumor cells in breast cancer patients. *Breast Cancer Res Treat* 2011;128:765–73.
- 41 Mostert B, Jiang Y, Sieuwerts AM, *et al.* KRAS and BRAF mutation status in circulating colorectal tumor cells and their correlation with primary and metastatic tumor tissue. *Int J Cancer* 2013;133:130–41.
- 42 Lievre A, Bachet JB, Le Corre D, *et al.* KRAS mutation status is predictive of response to cetuximab therapy in colorectal cancer. *Cancer Res* 2006;66:3992–5.
- 43 Laurent-Puig P, Cayre A, Manceau G, *et al.* Analysis of PTEN, BRAF, and EGFR status in determining benefit from cetuximab therapy in wild-type KRAS metastatic colon cancer. *J Clin Oncol* 2009;27:5924–30.
- 44 Misale S, Yaeger R, Hobor S, *et al.* Emergence of KRAS mutations and acquired resistance to anti-EGFR therapy in colorectal cancer. *Nature* 2012;486:532–6.
- 45 Oliveira C, Velho S, Moutinho C, *et al.* KRAS and BRAF oncogenic mutations in MSS colorectal carcinoma progression. *Oncogene* 2007;26:158–63.
- 46 Umeda Y, Nagasaka T, Mori Y, *et al.* Poor prognosis of KRAS or BRAF mutant colorectal liver metastasis without microsatellite instability. *J Hepatobiliary Pancreat Sci* 2013;20:223–33.
- 47 Taniguchi M, Nishida T, Hirota S, *et al.* Effect of c-kit mutation on prognosis of gastrointestinal stromal tumors. *Cancer Res* 1999;59:4297–300.
- 48 Tuveson DA, Willis NA, Jacks T, *et al.* STI571 inactivation of the gastrointestinal stromal tumor c-KIT oncoprotein: biological and clinical implications. *Oncogene* 2001;20:5054–8.
- 49 Magbanua MJ, Sosa EV, Roy R, *et al.* Genomic profiling of isolated circulating tumor cells from metastatic breast cancer patients. *Cancer Res* 2013;73:30–40.
- 50 Ramskold D, Luo S, Wang YC, *et al.* Full-length mRNA-Seq from single-cell levels of RNA and individual circulating tumor cells. *Nat Biotechnol* 2012;30:777–82.
- 51 Heitzer E, Auer M, Gasch C, *et al.* Complex tumor genomes inferred from single circulating tumor cells by array-CGH and next-generation sequencing. *Cancer Res* 2013;73:2965–75.