

- [12] Chen Y-W, Hwang KC, Yen C-C, Lai Y-L. Fullerene derivatives protect against oxidative stress in RAW 264.7 cells and ischemia-reperfusion lungs. *Am J Physiol* 2004;287: R21–R6.
- [13] Mirkov SM, Djordjevic AN, Andric NI, Andric SA, Kostic TS, Bogdanovic GM, et al. Nitric oxide-scavenging activity of polyhydroxylated fullereneol,  $C_{60}(OH)_{24}$ . *Nitric Oxide* 2004;11:201–7.
- [14] Injac R, Perse M, Cerne M, Potocnik N, Radic N, Govedarica B, et al. Protective effects of fullereneol  $C_{60}(OH)_{24}$  against doxorubicin-induced cardiotoxicity and hepatotoxicity in rats with colorectal cancer. *Biomaterials* 2009;30:1184–96.
- [15] Nakagawa Y, Suzuki T, Ishii H, Nakae D, Ogata A. Cytotoxic effects of hydroxylated fullerenes on isolated rat hepatocytes via mitochondrial dysfunction. *Arch Toxicol* 2011;85:1429–40.
- [16] Nakagawa Y, Suzuki T, Nakajima K, Inomata A, Ogata A, Nakae D. Effects of N-acetyl-L-cysteine on target sites of hydroxylated fullerene-induced cytotoxicity in isolated rat hepatocytes. *Arch Toxicol* 2014;88:115–26.
- [17] Li C, Martin Bradley C. Trends in emergency department visits attributable to acetaminophen overdoses in the United States: 1993–2007. *Pharmacoepidemiol Drug Saf* 2011;20:810–8.
- [18] Craig DGN, Bates CM, Davidson JS, Martin KG, Hayes PC, Simpson KJ. Overdose pattern and outcome in paracetamol-induced acute severe hepatotoxicity. *Br J Clin Pharmacol* 2011;71:273–82.
- [19] Tomishima Y, Ishitsuka Y, Matsunaga N, Nagatome M, Furusho H, Irikura M, et al. Ozagrel hydrochloride, a selective thromboxane A2 synthase inhibitor, alleviates liver injury induced by acetaminophen overdose in mice. *BMC Gastroenterol* 2013;13:21.
- [20] Hanawa N, Shinohara M, Saberi B, Gaarde WA, Han D, Kaplowitz N. Role of JNK translocation to mitochondria leading to inhibition of mitochondria bioenergetics in acetaminophen-induced liver injury. *J Biol Chem* 2008;283:13565–77.
- [21] Hinson JA, Pike SL, Pumford NR, Mayeux PR. Nitrotyrosine-protein adducts in hepatic centrilobular areas following toxic doses of acetaminophen in mice. *Chem Res Toxicol* 1998;11:604–7.
- [22] LoGuidice A, Boelsterli UA. Acetaminophen overdose-induced liver injury in mice is mediated by peroxynitrite independently of the cyclophilin d-regulated permeability transition (Hoboken, NJ, U S) *Hepatology* 2011;54:969–78.
- [23] Masubuchi Y, Sugiyama S, Horie T. Th1/Th2 cytokine balance as a determinant of acetaminophen-induced liver injury. *Chem Biol Interact* 2009;179:273–9.
- [24] North TE, Babu IR, Vedder LM, Lord AM, Wishnok JS, Tannenbaum SR, et al. PGE2-regulated wnt signaling and N-acetylcysteine are synergistically hepatoprotective in zebrafish acetaminophen injury. *Proc Natl Acad Sci U S A* 2010;107:17315–20. S11–S19.
- [25] Iohara D, Hirayama F, Higashi K, Yamamoto K, Uekama K. Formation of stable hydrophilic  $C_{60}$  nanoparticles by 2-hydroxypropyl- $\beta$ -cyclodextrin. *Mol Pharm* 2011;8:1276–84.
- [26] Kogure K, Goto S, Abe K, Ohiwa C, Akasu M, Terada H. Potent antiperoxidation activity of the bisbenzylisoquinoline alkaloid cepharanthine: the amine moiety is responsible for its pH-dependent radical scavenge activity. *Gen Subj Biochim Biophys Acta* 1999;1426:133–42.
- [27] Leelarungrayub N, Rattanapanone V, Chanarat N, Gebicki JM. Quantitative evaluation of antioxidant properties of garlic and shallot preparations. *N Y, NY, U S Nutrition* 2006;22:266–74.
- [28] Stadtman ER, Berlett BS. Fenton chemistry. Amino acid oxidation. *J Biol Chem* 1991;266:17201–11.
- [29] Shimizu D, Ishitsuka Y, Miyata K, Tomishima Y, Kondo Y, Irikura M, et al. Protection afforded by pre- or post-treatment with 4-phenylbutyrate against liver injury induced by acetaminophen overdose in mice. *Pharmacol Res* 2014;87:26–41.
- [30] Yoshida Z, Takekuma H, Takekuma S, Matsubara Y. Molecular recognition of  $C_{60}$  with  $\gamma$ -cyclodextrin. *Angew Chem Int Engl* 1994;33:1597–9.
- [31] Andersson T, Nilsson K, Sundahl M, Westman G, Wennerstrom O.  $C_{60}$  embedded in  $\gamma$ -cyclodextrin: a water-soluble fullerene. *J Chem Soc Chem Commun* 1992:604–6.
- [32] Jaeschke H, Williams CD, Ramchandran A, Bajt ML. Acetaminophen hepatotoxicity and repair: the role of sterile inflammation and innate immunity. *Liver Int* 2012;32:8–20.
- [33] Cover C, Mansouri A, Knight TR, Bajt ML, Lemasters JJ, Pessayre D, et al. Peroxynitrite-induced mitochondrial and endonuclease-mediated nuclear DNA damage in acetaminophen hepatotoxicity. *J Pharmacol Exp Ther* 2005;315:879–87.
- [34] Knight TR, Kurtz A, Bajt ML, Hinson JA, Jaeschke H. Vascular and hepatocellular peroxynitrite formation during acetaminophen toxicity: role of mitochondrial oxidant stress. *Toxicol Sci* 2001;62:212–20.
- [35] Lauterburg BH, Corcoran GB, Mitchell JR. Mechanism of action of N-acetylcysteine in the protection against the hepatotoxicity of acetaminophen in rats in vivo. *J Clin Invest* 1983;71:980–91.
- [36] Corcoran GB, Wong BK. Role of glutathione in prevention of acetaminophen-induced hepatotoxicity by N-acetyl-L-cysteine in vivo: studies with N-acetyl-D-cysteine in mice. *J Pharmacol Exp Ther* 1986;238:54–61.
- [37] Mitchell JR, Jollow DJ, Potter WZ, Davis DC, Gillette JR, Brodie BB. Acetaminophen-induced hepatic necrosis. I. Role of drug metabolism. *J Pharmacol Exp Ther* 1973;187:185–94.
- [38] Foley S, Crowley C, Smaih M, Bonfils C, Erlanger BF, Seta P, et al. Cellular localization of a water-soluble fullerene derivative. *Biochem Biophys Res Commun* 2002;294:116–9.
- [39] Misirkic MS, Todorovic-Markovic BM, Vucicevic LM, Janjetovic KD, Jokanovic VR, Dramicanin MD, et al. The protection of cells from nitric oxide-mediated apoptotic death by mechanochemically synthesized fullerene ( $C_{60}$ ) nanoparticles. *Biomaterials* 2009;30:2319–28.

