Risk factor profiles, drug usage, and prevalence of aspirin- associated gastroduodenal injuries among high-risk cardiovascular Japanese patients: the results from the MAGIC study	Uemura N, Sugano K, Hiraishi H, Shimada K, Goto S, Uchiyama S, Okada Y, Origasa H, Ikeda Y; MAGIC Study Group.	J Gastroenterol	2014/5月	国外
Recombinant Tissue-type Plasminogen Activator (rt-PA) Therapy in an Acute Stroke Patient Taking Dabigatran Etexilate: A Case Report and Literature Review	Tabata E,Yasaka M,Wakugawa Y, Okada Y	Internal Medicine	2014/6月	国外
Which should Be the Essential Components of Stroke Centers in Japan? A Survey by Questionnaires Sent to the Directors of Facilities Certified by the Japan Stroke Society	Uehara T, Yasui N, Okada Y, Hasegawa Y, Nagatsuka K, Minematsu K	Cerebrovascular Diseases	2014/8月	国外
Randomized clinical trial: prevention of recurrence of peptic ulcers by rabeprazole in patients taking low-dose aspirin	Iwakiri R, Higuchi K, Kato M, Fujishiro M, Kinishita Y, Watanabe T, Takeuchi T, Yamauchi M, Sanomura M, Nakagawa H, Sugisaki N, Okada Y, Ogawa H, Arakawa T, Fujimoto K	Aliment Pharmacol Ther	2014/10月	国外

抗凝固療法と頭蓋内出血	外山祐一郎、 矢坂正弘、齋 藤正樹、下濱 俊、岡田靖	脳と循環	2014/5月	国内
心原性脳塞栓症のチーム医 療	岡田靖	Cardio-Coagulation	2014/6月	国内
			·	
日本人における脳梗塞再発 抑制戦略を考える	田中耕太郎、岡田靖、高木誠、橋本洋一	Cardio Vascular Contemporary	2014/6月	国内
	郎、長谷川泰弘、松本昌泰			,
		:		
心房細動に対するワルファ	外山祐一郎、	血栓と循環	2014/9月	国内
リンと新規抗凝固薬:脳梗塞・脳出血の発症リスクは 異なるのか?	矢坂正弘、齋藤正樹、下濱 俊、岡田靖			
	,	·		

TOAST (The trial of Org 10172 in Acute Stroke Treatment) 分 類と脳卒中病型診断の現状	城貴弘、岡田靖		2014/7月	国内
Cautionプレゼンテーション の禁句「もともと」「フォ ローアップ」	岡田靖	日本医事新報	2014/8月	国内
IST, CAST	岡田靖、齋藤 正樹、下濱 俊、矢坂正弘	Clinical Neuroscience	2014/10月	国内
血栓塞栓症のリスク評価と 抗凝固療法に伴う出血のリ スク評価	坂正弘、岡田 靖	Medical Practice	2014/10月	国内
プラタナス 医療者としての4段階目標	岡田靖	日本医事新報	2014/11月	国内
脳卒中 顔・腕・言葉です ぐ受診	岡田靖	Branch	2014/11月	国内

日本脳卒中協会 福岡県支	岡田靖	脳梗塞と心房細動	2014/11月	国内
部の活動について				
İ				
1				

書籍

著者氏名	タイトル	書籍全体の編集者名	書籍名	出版社名	出版	出版年	ページ
Okada Y	Transient ischemic attack as a medical emergency	Uchiyama S, Amarenco P,Minematsu K, Wong K.S.L.	TIA as Acute Cerebrovascular Syndrome	KARGER	Basel	2014	19~29
岡田靖	脳卒中の前触れ「皇を中のがまち」 警報」を見逃さいる さずれば脳卒中の発き で発る できる	『健康』編集部	高血圧が自分で 治せる101のワ ザ	株式会社 主婦の友社	東京	2014	62~63
岡田靖	脳卒中は顔、 腕、言葉で判 断。周囲に疑 わしいくが かしいぐ を呼ぶ!	『健康』編集部	高血圧が自分で 治せる101のワ ザ	株式会社 主婦の友社	東京	2014	64~65

IV. 研究成果の刊行物・別刷

The Impact of Comprehensive Stroke Care Capacity on the Hospital Volume of Stroke Interventions: A Nationwide Study in Japan: J-ASPECT Study

Koji Iihara, MD, PhD,* Kunihiro Nishimura, MD, PhD,† Akiko Kada, MPH,‡
Jyoji Nakagawara, MD,§ Kazunori Toyoda, MD, PhD,|| Kuniaki Ogasawara, MD, PhD,¶
Junichi Ono, MD, PhD,# Yoshiaki Shiokawa, MD, PhD,** Toru Aruga, MD, PhD,††
Shigeru Miyachi, MD, PhD,‡‡ Izumi Nagata, MD, PhD,§§ Shinya Matsuda, MD, PhD,||||
Koichi B. Ishikawa, PhD,¶¶ Akifumi Suzuki, MD, PhD,## Hisae Mori, MD,*
and Fumiaki Nakamura, MD,***and the J-ASPECT Study Collaborators

Background: The association between comprehensive stroke care capacity and hospital volume of stroke interventions remains uncertain. We performed a nationwide survey in Japan to examine the impact of comprehensive stroke care capacity on the hospital volume of stroke interventions. Methods: A questionnaire on hospital characteristics, having tissue plasminogen activator (t-PA) protocols, and 25 items regarding personnel, diagnostic, specific expertise, infrastructure, and educational components recommended for comprehensive stroke centers (CSCs) was sent to 1369 professional training institutions. We examined the effect of hospital characteristics, having a t-PA protocol, and the number of fulfilled CSC items (total CSC score) on the hospital volume of t-PA infusion, removal of intracerebral hemorrhage, and coiling and clipping of intracranial aneurysms performed in 2009. Results: Approximately 55% of hospitals responded to the survey. Facilities with t-PA protocols (85%) had a significantly higher likelihood of having 23 CSC items, for example, personnel (eg, neurosurgeons: 97.3% versus 66.1% and neurologists: 51.3% versus 27.7%), diagnostic (eg, digital cerebral angiography: 87.4% versus 43.2%), specific expertise (eg, clipping and coiling: 97.2% and 54% versus 58.9% and 14.3%, respectively), infrastructure (eg, intensive care unit: 63.9% versus 33.9%), and education (eg, professional education: 65.2% versus 20.7%). On multivariate analysis adjusted for hospital characteristics, total CSC score, but not having a t-PA protocol, was associated with the volume of all types of interventions with a clear increasing trend (P for trend < .001). Conclusion: We demonstrated a significant association between

From the *Department of Neurosurgery, National Cerebral and Cardiovascular Center, Osaka; †Department of Preventive Medicine, National Cerebral and Cardiovascular Center, Osaka; ‡Advanced Medical Technology Development, National Cerebral and Cardiovascular Center, Osaka; §Integrative Stroke Imaging Center, National Cerebral and Cardiovascular Center, Osaka; ¶Department of Cerebrovascular Medicine, National Cerebral and Cardiovascular Center, Osaka; ¶Department of Neurosurgery, Iwate Medical University, Iwate; #Chiba Cardiovascular Center, Chiba; **Department of Neurosurgery, Kyorin University, Tokyo; ††Showa University Hospital, Tokyo; ‡†Department of Neurosurgery, Nagoya University, Nagoya; §§Department of Neurosurgery, Nagasaki University, Nagasaki; ||||Department of Preventive Medicine and Community Health, School of Medicine, University of Occupational and Environmental Health, Fukuoka; ¶¶Center for Cancer Control and Informa

tion Services, National Cancer Center, Tokyo; ##Research Institute for Brain and Blood Vessels, Akita; and ***Department of Healthcare Epidemiology, Faculty of Medicine, Kyoto University, Kyoto, Japan.

Received May 16, 2013: revision received August 8, 2013: accepted

Received May 16, 2013; revision received August 8, 2013; accepted August 21, 2013.

Conflict of interest: None declared.

This study was supported by Grants-in-Aid from the Ministry of Health, Labor, and Welfare of Japan (principal investigator: KI).

Address correspondence to Koji Iihara, MD, PhD, Department of Neurosurgery, National Cerebral and Cardiovascular Center, 5-7-1 Fujishiro-dai, Suita, Osaka 565-8565, Japan. E-mail: kiihara@hsp.ncvc.go.jp.

1052-3057/\$ - see front matter © 2014 by National Stroke Association http://dx.doi.org/10.1016/j.jstrokecerebrovasdis.2013.08.016 comprehensive stroke care capacity and the hospital volume of stroke interventions in Japan. **Key Words:** Stroke facilities—stroke units—ischemic stroke—intracerebral hemorrhage—subarachnoid hemorrhage—acute stroke therapy. © 2014 by National Stroke Association

Introduction

Stroke is the third leading cause of death worldwide and a leading cause of long-term disability. In 2000, the Brain Attack Coalition discussed the concept of stroke centers and proposed 2 types of centers: primary and comprehensive.2 Most stroke patients can be treated appropriately at primary stroke centers (PSCs), and the Joint Commission has established programs for the certification of, and measurement of performance in, PSCs. The concept of a comprehensive stroke center (CSC) enables intensive care and the use of specialized techniques, which are not available at most PSCs.2 A set of metrics and associated data elements that cover the major types of care that distinguish CSCs from PSCs have been published recently.3 At present, when the certification of CSCs has been initiated worldwide, there are no nationwide reports on the associations between primary and comprehensive stroke care capacity and impact of comprehensive stroke care capacity on hospital volume of stroke interventions.

Aims

We performed a nationwide survey in Japan to determine the association between primary and comprehensive stroke care capacity and the impact of comprehensive stroke care capacity on the annual hospital volume of tissue plasminogen activator (t-PA) infusion, removal of intracerebral hemorrhage (ICH), and coiling and clipping of intracranial aneurysms (IAs).

Methods

Eligible Hospitals

The J-ASPECT study (Nationwide survey of Acute Stroke care capacity for Proper dEsignation of Comprehensive stroke cenTer in Japan) group developed a 49-question survey regarding hospital characteristics (eg, bed number, academic status, geographic location, and participation in the diagnosis procedure combination [DPC] payment system), primary and comprehensive stroke care capacity, and hospital volume of stroke interventions. The questionnaire was mailed on February 2011 to the 1369 certified training institutions of the Japan Neurosurgical Society, the Japanese Society of Neurology, and the Japan Stroke Society. In the case of nonresponders, a second mail was sent to the medical directors to ensure that the questionnaire was completed and returned.

Assessment of Comprehensive and Primary Stroke Care Capacities and Hospital Volume of Stroke Interventions

This survey included 25 items related to the 5 major components of CSCs (personnel, diagnostic programs, specific expertise, infrastructure, and educational components) and 5 items related to PSC certification (Tables 1 and 2). Personnel were assessed according to 7 categories (eg, board-certified neurologists, board-certified neurosurgeons, and board-certified endovascular physicians). Because the original questions were highly specific, they were modified if necessary. Six advanced neuroimaging capabilities (eg, magnetic resonance imaging [MRI] with diffusion-weighted imaging [DWI] and digital cerebral angiography [DSA]) were investigated based on their availability 24 h/d, 7 d/wk (24/7). The availability of specific expertise for the stroke interventions was examined according to 5 categories (eg, carotid endarterectomy [CEA] and clipping and coiling of IAs). Regarding infrastructure, the availability of 5 items (eg, stroke unit and intensive care unit [ICU]) was surveyed.

Table 1. Characteristics of the responding hospitals and items of primary stroke care capacity

Variables	Category	n	%
Number of beds	<50	20	2.7
	50-99	30	4
	100-299	232	31
	300-499	260	34.7
	≥500	207	27.6
Academic hospital		90	12
DPC hospital		553	73.8
Geographic	MEA-central	381	50.9
locations	MEA-outlying	239	31.9
	McEA-central	90	12
	McEA-outlying	12	1.6
	Unclassified	27	3.6
PSC component	t-PA protocol	637	85
-	t-PA physician*	688	92
	NIHSS*	529	70.7
	Acute stroke team*	198	26.5
	Direct phone with EMS*	435	58.5

Abbreviations: DPC, diagnosis procedure combination; EMS, emergency medical services; MEA, metropolitan employment area; McEA, micropolitan employment area; NIHSS, National Institutes of Health Stroke Scale; PSC, primary stroke center; t-PA, tissue plasminogen activator.

*Data missing: t-PA physician, 1; NIHSS, 1; acute stroke team, 1; direct phone with EMS, 5.

Table 2. Number (percentage) of responding hospitals (n = 749) with the recommended items of comprehensive stroke care capacity

Components	Items	n	%
Personnel	Neurologists	358	47.8
	Neurosurgeons	694	92.7
	Endovascular physicians	272	36.3
	Critical care medicine	162	21.6
	Physical medicine and	113	15.1
	rehabilitation		
	Rehabilitation therapy	742	99.1
	Stroke rehabilitation	102	13.8
	nurses*		
Diagnostic	CT*	742	99.2
(24/7)	MRI with diffusion	647	86.4
	Digital cerebral angiography*	602	80.8
	CT angiography*	627	84
	Carotid duplex	257	34.5
	ultrasound*		
	TCD*	121	16.2
Specific	Carotid endarterectomy*	603	80.6
expertise	Clipping of IA	685	91.5
	Hematoma removal/ draining	689	91.9
	Coiling of IA	360	48.1
	Intra-arterial reperfusion therapy	498	66.5
Infrastructure	Stroke unit*	132	17.6
	Intensive care unit	445	59.4
	Operating room staffed 24/7*	451	60.4
	Interventional services coverage 24/7	279	37.3
	Stroke registry*	235	31.7
Education	Community education*	369	49.4
	Professional education*	436	58.6

Abbreviations: CT, computed tomography; IA, intracranial aneurysm; MRI, magnetic resonance imaging; TCD, transcranial Doppler.

*Data missing: stroke rehabilitation nurse, 9; CT, 1; digital cerebral angiography, 4; CT angiography, 3; carotid endarterectomy, 1; carotid duplex, 3; TCD, 3; stroke unit, 1; operating room staffed, 2; stroke registry, 7; community education, 2; professional education, 5.

Educational/research programs were assessed according to 2 items (community and professional education). Overall organizational and staffing levels of the hospitals in terms of CSC capacity were scored on the basis of the results of a questionnaire referring to 25 items originally recommended by the Brain Attack Coalition (total CSC score). A score of 1 was assigned for meeting each recommended item, and the maximum total CSC score was 25. Hospital volume of stroke interventions (eg, t-PA infusion, removal of ICH, and clipping and coiling of IAs) performed in 2009 was assessed.

Other Hospital Characteristics

Hospital characteristics pertaining to the number of beds, academic status (university/nonuniversity hospital), participation in the DPC-based payment system, and geographic location were described. The DPC database is a nationwide database in Japan comprising discharge abstract and administrative claim data.4,5 In 2010, approximately 1388 acute care hospitals in Japan, representing approximately 18% and 50% of the number of hospitals and hospital beds, respectively, adopted the DPC-based payment system.⁶ The geographic location of the hospitals was classified according to urban employment areas,7 which are divided into Metropolitan Employment Areas (MEAs) and Micropolitan Employment Areas (McEAs). The MEAs and McEAs are further classified into central and outlying areas based on the commuting pattern of their inhabitants. These classifications are based on the 2005 population census.8

Selection Bias of the Responding Hospitals

To overcome selection bias, we compared the proportion of hospitals that adopted the Japanese DPC-based payment system (DPC hospital) between the responding and nonresponding hospitals. Among the eligible DPC hospitals, we compared the proportion of hospitals that treated more than 10 cases of ischemic stroke (IS), ICH, and subarachnoid hemorrhage between July and December 2008, by using the DPC data from the responding and nonresponding hospitals.

Statistical Analyses

We used Fisher's exact test to detect significant differences in the proportions of hospitals. Kruskal-Wallis and Wilcoxon tests were used to determine differences in categorical and total CSC scores. Multiple linear regression analysis was used to examine the impact of total CSC score and availability of t-PA protocol adjusted for other hospital characteristics on hospital volume of stroke interventions. We also calculated *P* values (2 tailed) for trend across total CSC score categorized into quartiles, and *P* less than .05 was considered significant. Percentages were calculated excluding missing data. SAS version 9.2 (SAS Institute, Inc., Cary, NC) and STATA version 12 (STATA Corp., College Station, TX) were used for all statistical analyses.

Ethics

This research was approved by the Institutional Review Board of the National Cerebral and Cardiovascular Center, which waived the requirement for individual informed consent.

1004 K. IIHARA ET AL.

Results

Response Rate and Responding-Hospital Characteristics

In total, 55% (749 hospitals) of the eligible institutions completed the questionnaire. The characteristics of the responding hospitals are shown in Table 1. DPC hospitals represented 65.3% of the eligible hospitals and 73.8% and 55% of the responding and nonresponding hospitals, respectively (P < .0001). The response rate of the DPC hospitals was significantly higher than that of the remaining hospitals (61.9% versus 41.3%, P < .0001). Among the DPC hospitals, a greater proportion of responding hospitals, rather than nonresponding hospitals, treated more than 10 cases of ICH (65.6% versus 49.9%, P < .0001) and subarachnoid hemorrhage (30.7% versus 17%, P < .0001) per 6 months but not of IS (79% versus 76.8%, P = .454).

Primary Stroke Care Capacity Recommended for PSCs

Written t-PA protocols were available in 85% of hospitals, and the National Institutes of Health Stroke Scale score was routinely documented in 70.7% of hospitals; however, an acute stroke team was available only in 26.5% of hospitals (Table 1).

Comprehensive Stroke Care Capacity Recommended for CSCs

Regarding the recommended personnel components of CSCs, 92.7% of hospitals had a neurosurgeon, 47.8% had a neurologist, and 36.3% had an endovascular physician (Table 2). The proportion of hospitals with critical care medicine physicians and physical medicine and stroke rehabilitation nurses was 21.6% and 15.1% and 13.8%, respectively. The availability of the personnel component ranged from 0 to 7 (median, 3; interquartile range [IQR], 2-4).

Computed tomography (CT), MRI with DWI, DSA, and CT angiography (CTA) were available 24/7 in 99.2%, 86.4%, 80.8%, and 84% of institutions, respectively, whereas carotid duplex ultrasonography and transcranial Doppler (TCD) were available only in 34.5% and 16.2% of hospitals, respectively. The availability of diagnostic components ranged from 0 to 6 (median, 4; IQR, 4-5).

CEA, clipping of IAs, and removal of ICH were available in 80.6%, 91.5%, and 91.9% of hospitals, respectively, whereas coiling of IAs and intra-arterial thrombolysis were available in 48.1% and 66.5% of hospitals, respectively. The availability of surgical and interventional components ranged from 0 to 5 (median, 4; IOR, 3-5).

A stroke unit and an ICU were available in 17.6% and 59.4% of hospitals, respectively. The availability of interventional service coverage on a 24-hour basis was observed in 37.3% of hospitals, whereas an operating room staffed 24/7 was available in 60.4% of hospitals. The availability of infrastructure components ranged from 0 to 5 (median, 2; IQR, 1-3).

Professional and community education were available in 58.6% and 49.4% of institutions, respectively (Table 2). The availability of the educational component ranged from 0 to 2 (median, 1; IQR, 0-2).

Geographical Disparity of Primary and Comprehensive Stroke Care Capacity

Among certified personnels, the proportion of neurologists, interventional physicians, and stroke nurses available at MEA-central institutes was significantly higher (P < .001, P < .001, and P = .045, respectively), whereas neurosurgeons, critical care medicine, and physical medicine were in place, irrespective of the location. Imaging techniques, such as MRI, diffusion-weighted MRI, CTA, digital subtraction angiography, and carotid duplex ultrasonography, were available on a 24-hour basis in similar proportions irrespective of the location, whereas TCD was available in a significantly higher proportion at MEA-central institutes except for TCD (P = .003). Specific surgical and interventional services for CEA (P = .002), coiling of IAs (P < .001), and intra-arterial reperfusion therapy (P = .04) were available in a significantly higher proportion at MEA-central institutes. The availability of stroke unit (P < .001), operating room staffed, and interventional services on a 24/7 basis (P < .001) was significantly more established at MEA-central institutes, whereas no significant differences were found with regard to the ICU. Community (P = .011) and professional education (P = .003) were more established at MEAcentral institutes (Appendix Tables 2 and 3).

Associations between the Implementation of a t-PA Protocol and Comprehensive Stroke Care Capacity

Facilities with t-PA protocols had a higher likelihood of having a neurosurgeon (97.3% versus 66.1%, P < .0001), neurologist (51.3% versus 27.7%, P < .0001), endovascular physician (41.1% versus 8.9%, P < .0001), and physicians in critical care medicine (23.7% versus 9.8%, P = .0007) and physical medicine and rehabilitation (16.5% versus 7.1%, P = .006) (Table 3). There were no differences in the availability of personnel in rehabilitation therapy (P = .355) or of stroke rehabilitation nurses (P = .695). Facilities with t-PA protocols had a higher percentage of 24/ 7 availability of CT (99.5% versus 97.3%, P = .042), MRI with DWI (89.6% versus 67.9%, P < .0001), DSA (87.4% versus 43.2%, P < .0001), carotid duplex ultrasonography (37% versus 19.8%, P = .0003), and TCD (18.1% versus 5.4%, P = .0002). Moreover, hospitals with a t-PA protocol had greater availability of CEA (87.1% versus 43.8%, P < .0001), clipping of IAs (97.2% versus 58.9%, P < .0001), removal of ICH (97.5% versus 60.7%, P < .0001), coiling of IAs (54% versus 14.3%, P < .0001), and intra-arterial thrombolysis (73% versus 29.5%, P < .0001) and greater availability of a stroke unit (20.1% versus 3.6%, P < .0001), an ICU (63.9% versus

Table 3. Characteristics of comprehensive stroke care capacity according to the presence or absence of a t-PA protocol

Variables	t-PA proto	col (+) n, %	t-PA protoc	col (-) n, %	OR	95% CI	P valu
Personnel							
Neurologists	327	51.3	31	27.7	2.8	1.8-4.3	<.000
Neurosurgeons	620	97.3	74	66.1	18.7	10.1-34.8	<.000
Endovascular physicians	262	41.1	10	8.9	7.1	3.7-13.9	<.000
Critical care medicine	151	23.7	11	9.8	2.9	1.5-5.5	.000
Physical medicine and rehabilitation	105	16.5	8	7.1	2.3	1.2-5.4	.006
Rehabilitation therapy	632	99.2	110	98.2	1.14	0.4-12	.355
Stroke rehabilitation nurses	88	14	14	12.6	1.1	0.6-2.1	.695
Diagnostic							
CT*	634	99.5	108	97.3	5.9	1.2-29.5	.042
MRI with diffusion	571	89.6	76	67.9	4.1	2.6-6.6	<.000
Digital cerebral angiography*	554	87.4	48	43.2	9.1	5.8-14.2	<.000
CT angiography*	566	89.1	61	55	6.7	4.3-10.5	<.000
Carotid duplex ultrasound*	235	37	22	19.8	2.4	1.5-3.9	.000
TCD*	115	18.1	6	5.4	3.9	1.7-9.1	.000
Specific expertise							
Carotid endarterectomy*	554	87.1	49	43.8	8.7	5.6-13.5	<.000
Clipping of IA	619	97.2	66	58.9	24	13.1-43.7	<.000
Hematoma removal/draining	621	97.5	68	60.7	25.1	13.4-46.9	<.000
Coiling of IA	344	54	16	14.3	7	4.1-12.2	<.000
Intra-arterial reperfusion therapy	465	73	33	29.5	6.5	4.2-10.1	<.000
Infrastructure							
Stroke unit*	128	20.1	4	3.6	6.8	2.5-18.8	<.000
Intensive care unit	407	63.9	38	33.9	3.4	2.3-5.3	<.000
Operating room staffed 24/7	426	67.1	25	22.3	7.1	4.4-11.4	<.000
Interventional services coverage 24/7	268	42.1	11	9.8	6.7	3.5-12.7	<.000
Stroke registry*	229	36.3	6	5.4	10	4.3-23.1	<.000
Education							
Community education*	348	54.8	21	18.8	5.3	3.2-8.7	<.000
Professional education*	413	65.2	23	20.7	7.2	4.4-11.7	<.000

Abbreviations: CI, confidence interval; CT, computed tomography; IA, intracranial aneurysm; MRI, magnetic resonance imaging; OR, odds ratio; TCD, transcranial Doppler.

*Data missing: CT, 1; digital cerebral angiography, 4; carotid U/S, 3; TCD, 3; carotid endarterectomy, 1; stroke unit, 1; stroke registry, 7; community education, 2; professional education, 5.

33.9%, P < .0001), an operating room staffed 24/7 (67.1% versus 22.3%, P < .0001), interventional service coverage 24/7 (42.1% versus 9.8%, P < .0001), and a stroke registry (36.3% versus 5.4%, P < .0001). Finally, hospitals with a t-PA protocol had greater availability of professional (65.2% versus 20.7%, P < .0001) and community (54.8% versus 18.8%, P < .0001) education (Table 3).

Impact of Total CSC Score on Hospital Volume of Stroke Interventions

The total CSC score, which ranged from 0 to 24 (median, 14; IQR, 11-18), was significantly higher in facilities with a t-PA protocol than in other facilities (median, IQR: 15, 12.5-18 versus 9, 5-11; P < .0001). On univariate analysis, the implementation of a t-PA protocol and total CSC score categorized into quartiles (Q1, 0-10; Q2, 11-13; Q3, 14-17; and Q4, 18-24) were each significantly associated with hospital volume of stroke interventions, as shown in Table 4. The impact of total CSC score (Q1-Q4) and the implemen-

tation of a tPA protocol on the annual case volume of stroke interventions in the participating hospitals was shown in Figure 1. On multiple linear regression analysis, the availability of a t-PA protocol and total CSC score each had a significant impact on the hospital volume of the stroke interventions after adjustment for other hospital characteristics (Appendix Tables 4-6). The inclusion of total CSC score, availability of a t-PA protocol, and other hospital characteristics in the model revealed that total CSC score, but not availability of a t-PA protocol, was significantly associated with the hospital volume of stroke interventions (Table 5).

Discussion

This study demonstrated a significant impact of comprehensive stroke care capacity represented by the total CSC score on the hospital volume of stroke interventions and unique aspects of comprehensive stroke care capacity in Japan.

1006 K. IIHARA ET AL.

Table 4. Impact of the availability of a t-PA protocol and of total CSC score on hospital volume of stroke interventions performed in 2009: hospital volume of stroke interventions based on the availability of a t-PA protocol and on total CSC score (univariate analysis)

	t-1	t-PA protocol			Total CSC score			
	(+)	(-)	P value	Q1	Q2	Q3	Q4	P value
t-PA infusion	5 (2-11)	0 (0-1)	<.0001	1 (0-3)	4 (1-7)	6 (4-10.5)	10 (5-16)	<.0001
Removal of ICH	7 (3-13)	0 (0-3)	<.0001	2 (0-4)	5 (3-9)	7.5 (3-15)	11.5 (6-18)	<.0001
Clipping of IA Coiling of IA	17 (8-29) 4 (0-13)	0 (0-10) 0 (0-1)	<.0001 <.0001	3 (0-11.5) 0 (0-1)	12 (6-21.25) 1 (0-4)	18 (10-30) 5.5 (1.25-13)	27 (18-43.75) 13 (5-22)	<.0001 <.0001

Abbreviations: CSC, comprehensive stroke center; IA, intracranial aneurysm; ICH, intracerebral hemorrhage; t-PA, tissue plasminogen

Unique Aspects of Comprehensive Stroke Care Capacity in Japan

This study illustrated several unique aspects of comprehensive stroke care capacity in Japan, for example, higher availability of neurosurgeons (92.7% versus 24%-54% in the United States)^{9,10} and endovascular surgeons (36.3% versus 15%-22% in the United States),⁹⁻¹¹ which was in sharp contrast with the relative shortage of neurologists (47.8% versus 31%-73% in the United States) and other personnel.⁹⁻¹¹ In the United States, only 7% of neurosurgeons play an active role in nontraumatic cranial emergencies,¹² whereas in Japan, 59% of the board-certified neurosurgeons are engaged in stroke care. The proportion of Japanese hospitals offer-

Table 5. Impact of the availability of a t-PA protocol and of total CSC score on hospital volume of stroke interventions performed in 2009: multivariate linear regression analysis of the impact of total CSC score and availability of a t-PA protocol on hospital volume of stroke interventions

	β	P value	95%	CI
tPA				
Total CSC score	.83	<.001	.67	.98
tPA protocol (+)	-1.31	.117	-2.95	.33
ICH				
Total CSC score	.97	<.001	.76	1.18
tPA protocol (+)	-1.17	.297	-3.36	1.03
Clipping				
Total CSC score	2.23	<.001	1.68	2.79
tPA protocol (+)	-4.33	.146	-10.17	1.51
Coiling				
Total CSC score	1.20	<.001	.92	1.48
tPA protocol (+)	-1.01	.496	-3.92	1.90

Abbreviations: CI, confidence interval; CSC, comprehensive stroke center; IA, intracranial aneurysm; ICH, intracerebral hemorrhage; t-PA, tissue plasminogen activator; Total CSC score, availability of a t-PA protocol, and other hospital characteristics were included in the model as independent variables.

ing MRI with DWI and CTA corresponded with the gradual increase in the availability of certain special diagnostic tests in the United States, whereas the availability of DSA (80.8%) was in contrast with the temporal decrease in the availability of catheter angiography observed in North Carolina (from 38% in 1998 to 30% in 2008) because of declines in the proportion of hospitals with neurointerventionalists.¹¹

A stroke unit/stroke care unit as the critical infrastructure for acute stroke, which has been proven to reduce the number of deaths and long-term dependency,¹³ was available in only 17.6% of hospitals in Japan; this was comparable with the proportion observed in hospitals in the United States (6.6%-28%).^{9-11,14}

To determine the relationship between primary and comprehensive stroke care capacity, we used the implementation of a written t-PA protocol as a key item of primary stroke care capacity as it is a key step in reducing t-PA-related complications. 15 The availability of a t-PA protocol (85%) recorded in the present study was comparable with that reported in 2 US statewide studies performed in Illinois in 2000 (72.8%)⁹ and in North Carolina between 1998 (43%) and 2008 (69%). 11 Notably, facilities with a t-PA protocol in Japan had a higher availability of nearly all (92%) recommended items of comprehensive stroke care capacity, with the exception of personnel in rehabilitation therapy and stroke rehabilitation nurses. In a previous study performed in the United States,⁹ however, no significant differences were noted in the availability of a larger number of critical items corresponding to the CSC items in this study (eg, endovascular physicians, CTA, conventional cerebral angiography, carotid duplex ultrasound, intraarterial thrombolysis, stroke unit, and community stroke awareness program) according to the presence or absence of a t-PA protocol. Therefore, comprehensive stroke care capacity tended to coexist in hospitals with primary stroke care capacity in Japan than in the United States. A relatively larger commitment of neurosurgeons in acute stroke care (eg, t-PA infusion) in Japan may explain such a coexistence.




Figure 1. A bar graph showing the impact of total CSC score (Q1-Q4) and the implementation of a tPA protocol on the annual case volume of stroke interventions (A, t-PA infusion; B, ICH removal; C, clipping; and D, coiling of intracranial aneurysms) in the participating hospitals. Abbreviations: CSC, comprehensive stroke center; t-PA, tissue plasminogen activator.

Geographical Disparity of Primary and Comprehensive Stroke Care in Japan

These findings could assist in identifying underserved, high-population density areas that may benefit from professional and hospital education by national organizations, such as the Japan Stroke Society and Japan Neurosurgical Society. In light of insufficient advanced acute stroke capacities, these findings represent fundamental information for the establishment of a spoke-and-hub stroke care system.² Such a system could maximize use of acute reperfusion therapies for IS and surgical/endovascular management of hemorrhagic stroke, depending on the personnel and specific expertise of the hospital and the geographical access of the potential population in Japan.

The Impact of Comprehensive Stroke Care Capacity on the Hospital Volume of Stroke Interventions

The present study demonstrated a significant impact of comprehensive stroke care capacity, as represented by the total CSC score, but not the availability of a t-PA protocol, on the hospital volume of stroke interventions performed in Japan in 2009, after adjustment for other hospital characteristics. High volume is associated with better outcomes across a wide range of procedures and conditions; however, the magnitude of the association varies greatly. High annual hospital volume was reported to be consistently associated with lower IS mortality. In the field of comprehensive stroke care, in-hospital mortality increases when the annual number of craniotomies for aneurysms is lower than 30¹⁸ and when CEA is performed by less experienced surgeons (<5 procedures/y). Conversely, a previous Japanese nationwide study

found no correlation between case volume and outcome after cerebral aneurysm clipping.²⁰ The mechanism via which volume influences outcomes remains uncertain. Specific processes of care, which are correlated with volume, are the most likely explanatory factors.¹⁶ The strong associations between comprehensive stroke care capacity and hospital volume of the stroke interventions observed in this study may support this notion. Further studies are necessary to establish the association between comprehensive stroke care capacity and in-hospital mortality after all types of stroke. If such association is established, as recently reported for PSC,²¹ the total CSC score may be used as a quality indicator of comprehensive stroke care capacity for benchmarking purposes.

Limitations

This study was likely to have included a potential information bias (self-report, recall, and nonresponse). Hospitals actively working to improve stroke care are more likely to respond to the questionnaire. The fact that a greater proportion of the responding DPC hospitals treated more than 10 cases of hemorrhagic stroke in the latter half of the previous year than did the nonresponding DPC hospitals may support such selection bias. We did not attempt to assess the self-report bias as no official information was available that could have been used to verify the data.

Conclusion

The present study demonstrated a significant impact of comprehensive stroke care capacity, as represented by the total CSC score on the hospital volume of stroke

1008 K. IIHARA ET AL.

interventions and unique aspects of comprehensive stroke care capacity in Japan.

References

- Alberts MJ, Hademenos G, Latchaw RE, et al. Recommendations for the establishment of primary stroke centers. Brain Attack Coalition. JAMA 2000;283:3102-3109.
- Alberts MJ, Latchaw RE, Selman WR, et al. Recommendations for comprehensive stroke centers: a consensus statement from the Brain Attack Coalition. Stroke 2005; 36:1597-1616.
- Leifer D, Bravata DM, Connors JJ III, et al. Metrics for measuring quality of care in comprehensive stroke centers: detailed follow-up to Brain Attack Coalition comprehensive stroke center recommendations: a statement for healthcare professionals from the American Heart Association/American Stroke Association. Stroke 2011;42:849-877.
- Kadono Y, Yasunaga H, Horiguchi H, et al. Statistics for orthopedic surgery 2006-2007: data from the Japanese Diagnosis Procedure Combination database. J Orthop Sci 2010;15:162-170.
- Shoda N, Yasunaga H, Horiguchi H, et al. Risk factors affecting in-hospital mortality after hip fracture: retrospective analysis using the Japanese Diagnosis Procedure Combination Database. BMJ Open 2012;2. http://dx.doi.org/10.1136/bmjopen-2011-000416.
- Current status of DPC hospitals [online]. Available at: http://www.mhlw.go.jp/stf/shingi/2r9852000000bartatt/2r9852000000bc5v.pdf. Accessed January 19, 2013.
- Kanemoto Y, Kurima R. Urban employment areas: defining Japanese metropolitan areas and constructing the statistical database for them. In: Okabe A, ed. GIS-based studies in the humanities and social sciences. Boca Raton, FL: CRC Taylor & Francis 2006:85-97.
- Population census [online]. Available at: http://www.st at.go.jp/english/data/kokusei/2005/outline.htm. Accessed September 15, 2013.
- Ruland S, Gorelick PB, Schneck M, et al. Acute stroke care in Illinois: a statewide assessment of diagnostic and treatment capabilities. Stroke 2002;33:1334-1339.

 Shultis W, Graff R, Chamie C, et al. Striking rural-urban disparities observed in acute stroke care capacity and services in the pacific northwest: implications and recommendations. Stroke 2010;41:2278-2282.

- 11. Goldstein LB. Statewide hospital-based stroke services in North Carolina: changes over 10 years. Stroke 2010; 41:778-783.
- 12. Seaver MJ. Baseline ER survey explores system's cracks: 2004 AANS/CNS neurosurgical emergency and trauma services survey [online]. Available at: http://www.aans.org/Media/Article.aspx?ArticleId=26367. Accessed January 23, 2013.
- Organised inpatient (stroke unit) care for stroke. Cochrane Database Syst Rev 2007. CD000197.
- Kidwell CS, Shephard T, Tonn S, et al. Establishment of primary stroke centers: a survey of physician attitudes and hospital resources. Neurology 2003;60:1452-1456.
- 15. Albers GW, Bates VE, Clark WM, et al. Intravenous tissue-type plasminogen activator for treatment of acute stroke: the Standard Treatment with Alteplase to Reverse Stroke (STARS) study. JAMA 2000;283:1145-1150.
- Halm EA, Lee C, Chassin MR. Is volume related to outcome in health care? A systematic review and methodologic critique of the literature. Ann Intern Med 2002;137:511-520.
- Saposnik G, Baibergenova A, O'Donnell M, et al. Hospital volume and stroke outcome: does it matter? Neurology 2007;69:1142-1151.
- Solomon RA, Mayer SA, Tarmey JJ. Relationship between the volume of craniotomies for cerebral aneurysm performed at New York state hospitals and in-hospital mortality. Stroke 1996;27:13-17.
- Hannan EL, Popp AJ, Tranmer B, et al. Relationship between provider volume and mortality for carotid endarterectomies in New York state. Stroke 1998;29: 2292-2297.
- Hattori N, Katayama Y, Abe T. Case volume does not correlate with outcome after cerebral aneurysm clipping: a nationwide study in Japan. Neurol Med Chir (Tokyo) 2007;47:95-100. discussion 1.
- 21. Lichtman JH, Jones SB, Wang Y, et al. Outcomes after ischemic stroke for hospitals with and without Joint Commission-certified primary stroke centers. Neurology 2011;76:1976-1982.

Appendix Table 1. List of the responding hospitals

Appendix Table 1. (Continued)

Abashiri Neurosurgical Rehabilitation Hospital

Ageo Central General Hospital Aichi Medical University Hospital

Aidu Chuo Hospital Aizawa Hospital Akiba Hospital

Akiru Municipal Medical Center

Akita General Hospital

Akita Kumiai General Hospital

Akita Rosai Hospital Akita University Hospital Ako Central Hospital Ako City Hospital

Almeida Memorial Hospital Amakusa Medical Center Anji Kosei Hospital Aomori Kyoritsu Hospital Aomori Municipal Hospital Aomori Rosai Hospital Aoyama Hospital

Araki Neurosurgical Hospital

Arao City Hospital Arita Kyouritsu Hospital

Asahikawa Medical University Hospital

Asahikawa Red Cross Hospital Asao General Hospital Ashikaga Red Cross Hospital

Atsumi Hospital Ayabe City Hospital

Azuma Neurosurgical Hospital Azumi General Hospital Azumino Red Cross Hospital Baba Memorial Hospital Bellland General Hospital

Beppu Medical Center, National Hospital Organization

Bizen Municipal Hospital

Central Gunma Neurosurgical Hospital

Chiba Cardiovascular Center Chiba Central Medical Center Chiba Emergency Medical Center Chiba Neurosurgical Clinic Chiba University Hospital Chichibu City Hospital Chigasaki City Hospital Chikamori Hospital

Chubu-Tokusyukai Hospital Chugoku Rousai Hospital Chuno Kosei Hospital Daiichi Hospital

Daini Okamoto General Hospital

Daiwa Hospital

Dohtoh Neurosurgical Hospital Dokkyo Medical University Hospital Ehime Prefectural Central Hospital Ehime Prefectural Imabari Hospital

Ehime University Hospital

Enshu Hospital Fuchu Hospital

Fuji Brain Institute and Hospital

Fuji City General Hospital Fujii Neurosurgical Hospital Fujimoto Hayasuzu Hospital Fujisawa City Hospital

Fujisawa Neurosurgical Hospital

Fujita General Hospital

Fujita Health University Hospital Fujiyoshida Municipal Medical Center

Fukaya Red Cross Hospital Fukui Red Cross Hospital Fukui General Hospital Fukui Kosei Hospital

Fukui Social Insurance Hospital Fukui University Hospital Fukuoka City Hospital Fukuoka Kieikai Hospital Fukuoka Seishukai Hospital Fukuoka Shin Mizumaki Hospital Fukuoka Tokushukai Medical Center

Fukuoka University Hospital Fukuroi Municipal Hospital

Fukushima Medical University Hospital Fukushima Prefectural Aizu General Hospital

Fukushima Red Cross Hospital

Fukushima Takanori Memorial Hospital

Furukawa Seiryo Hospital

Fussa Hospital

Gamagohri City Hospital

Geriatrics Research Institute and Hospital

Gifu Central Hospital Gifu Municipal Hospital

Gifu Prefectural General Medical Center

Gifu Prefectural Tajimi Hospital

Gifu University Hospital

Goshi Hospital

Gunma University Hospital Hachinohe Heiwa Hospital Hachisuga Hospital

Hakodate Neurosurgical Hospital Hakodate Shintoshi Hospital

Hakuai Hospital

Hakuo-kai Sumi Hospital Hamamatsu Medical Center

Hamamatsu University School of Medicine, University

Hospital

Hamamatu Rosai Hospital Hamanomachi Hospital Hanwa Memorial Hospital

Harada Hospital Hashima City Hospital Hata Kenmin Hospital Hayashi Hospital

Health Insurance Amakusa Chuo General Hospital

Health Insurance Nankai Hospital Health Insurance Naruto Hospital

Heart Life Hospital Heisei Memorial Hospital

(Continued)

Appendix Table 1. (Continued)

Heisei Neurosurgical Hospital

Hibino Hospital

Hidaka General Hospital

Higashimatsuyama Medical Association Hospital

Higashiyamato Hospital Hikone Municipal Hospital Himeji Central Hospital Himi Municipal Hospital Hiratsuka City Hospital Hirosaki University Hospital Hiroshima City Asa Hospital Hiroshima General Hospital

Hiroshima Prefectural Hospital

Hiroshima Red Cross Hospital and Atomic-bomb

Survivors Hospital

Hiroshima University Hospital

Hokkaido Neurosurgical Memorial Hospital

Hokkaido University Hospital

Hokushin General Hospital Nagano Prifectural Fideration

of Agricultural Cooperatives for Health and Welfare

Hokushinkai Megumono Hospital

Hokuto Hospital

Hoshigaoka Koseinenkin Hospital

Houetsu Hospital

Hskinan Municipal Hospital Hyogo Brain and Heart Center

Hyogo Prefectural Amagasaki Hospital Hyogo Prefectural Awaji Hospital Hyogo Prefectural Nishinomiya Hospital Ibaraki Prefectural Central Hospital Ibaraki Seinan Medical Center Hospital

Ibi Kousei Hospital

Ichinomiya Neurosurgery Hospital

Ichinomiya West Hospital Ichinose Hospital Iida Municipal Hospital

Iizuka Hospital

Ikeda Neurosurgical Center Hospital

Ikuwakai Memorial Hospital Imakiire General Hospital Imamura Bun-in Hospital Inagi Municipal Hospital

International University of Health and Welfare ATAMI

HOSPITAL

Inuyama Chuo Hospital

Iseikai Hospital

Iseikai Yahata Central Hospital Isesaki Municipal Hospital

Ishikawa Prefectural Central Hospital Ishinkai Yao General Hospital Ishinomaki City Hospital Ishioka Dai-Ichi Hospital Isogo Central Hospital

Itami Kousei Neurosurgical Hospital

Itoigawa General Hospital Itsukaichi Memorial Hospital Iwaki Kyoritsu Hospital

Iwamizawa Municipal General Hospital

Iwata City Hospital

Iwate Medical University Hospital Iwate Prefectural Iwai Hospital Iwate Prefectural Ninohe Hospital Iwate Prefecturel Isawa Hospital Izumi General Medical Center

Izumino Hospital

Izumu Municipal Hospital

JA Kochi Hospital

JA Toride Medical Center

Japan Medical Alliance Higashi Saitama General Hospital

Japanese Red Cross Hadano Hospital Japanese Red Cross Kitami Hospital Japanese Red Cross Kobe Hospital Japanese Red Cross Kochi Hospital Japanese Red Cross Koga Hospital Japanese Red Cross Kyoto Daini Hospital Japanese Red Cross Medical Center

Japanese Red Cross Nagasaki Genbaku Hospital

Japanese Red Cross Ogawa Hospital

Japanese Red Cross Society Hachinohe Medical Center

Japanese Red Cross Society Himeji Hospital

Jiaikai Kajiura Hospital

Jichi Medical University Hospital JR Tokyo General Hospital

Junshin Hospital

Juntendo University Hospital Juntendo University Nerima Hospital Juntendo University Urayasu Hospital

Junwakai Kinen Hospital Juzenkai Hospital Jyuzen General Hospital Kaga City Hospital

Kagawa Prefectural Central Hospital

Kagawa Rosai Hospital Kagawa University Hospital Kagoshima City Hospital

Kagoshima prefectural Oshima Hospital Kagoshima Tokushukai Hospital

Kagoshima University Medical and Dental Hospital

Kainan Hospital Kaiseikai Onishi Hospital

Kakegawa Municipal General Hospital

Kakizoe Hospital Kakogawa City Hospital Kameda Medical Center Kameoka Shimizu Hospital Kamiichi General Hospital Kamitsuga General Hospital

Kanazawa Medical University Hospital

Kanazawa Municipal Hospital Kanazawa Neurosurgical Hospital Kanazawa University Hospital

Kanetsu Hospital Kanoya Medical Center

Kansai Medical University Takii Hospital

Kanto Central Hospital for Public School Teachers

Kantoh Neurosurgical Hospital

(Continued)

Appendix Table 1. (Continued)

Karatsu Red Cross Hospital Kariya Toyota General Hospital

Kasai Cardiology Neurosurgery Hospital

Kasaoka Daiichi Hospital Kashima Rosai Hospital

Kashiwaba Neurosurgical Hospital

Kashiwazaki General Hospital and Medical Center

Kasugai Municipal Hospital Kasukabe Chuo General Hospital Katagi Neurosurgical Hospital

Katano Hospital Katsuta Hospital

Kawachi General Hospital Kawakita General Hospital Kawano Neurosurgical Hospital

Kawasaki Hospital

Kawasaki Medical School Hospital Kawasaki municipal Hospital Kawasaki Municipal Tama Hospital

Kawasaki Kamai Hospital
Kazuno Kosei Hospital
Keijin Kai Kawasaki Hospital
Keiwakai Ebetsu Hospital
Kenwakai Otemachi Hospital
Kimitsu Chuo Hospital
Kinan Hospital

Kinki University Hospital Kiryuu Kohsei General Hospital Kishiwada Tokushukai Hospital Kitakyushu Municipal Medical Center

Kitamurayama Hospital Kitano Hospital

Kizawa Memorial Hospital KKR Sapporo Medical Center Kobari General Hospital Kobe University Hospital Kochi Health Sciences Center Kochi Medical School Hospital

Kofu Jonan Hospital
Kofu Municipal Hospital
Kofu Neurosurgical Hospital
Kohka Public Hospital
Kohnan Hospital

Kokubu Neurosurgical Clinic Kokura Memorial Hospital Komaki City Hospital Komatsu Municipal Hospital Komono Kosei Hospital Konan Kosei Hospital Konan Tobu General Hospital

Kosei Hospital

Koshigaya Municipal Hospital Koto Memorial Hospital Kousei General Hospital Kouseikai Takai Hospital Kouseiren Murakami Hospital

Kugayama Hospital Kumamoto City Hospital Kumamoto Red Cross Hospital Kumamoto Rousai Hospital Kumamoto Takumadai Hospital Kumamoto University Hospital Kurashiki Central Hospital Kurashiki Heisei Hospital Kure Kyosai Hospital Kurobe City Hospital

Kurose City Hospital
Kurosihi City Hospital
Kurosawa Hospital
Kurosu Hospital

Kurume University Hospital Kusatsu General Hospital

Kushiro Kojinkai Memorial Hospital

Kushiro Rosai Hospital Kuwana Hospital

Kuwana West Medical Center Kyorin University Hospital Kyoritsu General Hospital Kyosai Tachikawa Hospital Kyoto City Hospital Kyoto Kujo Hospital

Kyoto Min-iren Chuo Hospital Kyushu University Hospital Machida Municipal Hospital Maebashi Red Cross Hospital Makita General Hospital Masu Memorial Hospital Matsudo City Hospital Matsunami General Hospital Matsushita Memorial Hospital Matsuyama Red Cross Hospital Matsuyama Shimin Hospital Mattoh-Ishikawa Central Hospital

Mazda Hospital

Medical Corporation Kawamura Society Kubokawa

Hospital Meisei Hospital Meitetsu Hospital Midorigaoka Hospital

Mie Prefectural General Medical Center

Mie University Hospital Mihara Memorial Hospital

Minamata City General Hospital and Medical Center

Minamisoma City General Hospital

Minase Hospital
Mine City Hospital
Minei Daiichi Hospital
Mino Municipal Hospital
Minoh City Hospital
Mishuku Hospital
Mishuku Hospital
Mishuku Hospital
Mito Medical Center

Mito Saiseikai General Hospital Mitoyo General Hospital Mitsugi General Hospital

Miyakonojo Regional Medical Center

Miyoshi Central Hospital

(Continued)

Mizushima Central Hospital Morioka JRC Hospital

Moriya-daiichi General Hospital

Murakami Memorial Hospital Asahi University

Murata Hospital

Muroran City General Hospital

Musashino General Hospital

Nadogaya Hospital

Naga District Hospital

Nagahama City Hospital

Nagahama Red Cross Hospital

Nagano Municipal Hospital

Nagano Prefectural Kiso Hospital

Nagano Prefectural Suzaka Hospital

Nagano Red Cross Hospital

Nagaoka Chuo General Hospital

Nagaoka Red Cross Hospital

Nagareyama Central Hospital

Nagasaki Kawatana Medical Center

Nagasaki Municipal Hospital

Nagasaki Prefecture Shimabara Hospital

Nagasaki University Hospital

Nagatomi Neurosurgical Hospital

Nagoya Daini Red Cross Hospital

Nagoya Ekisaikai Hospital

Nagoya Medical Center

Nagoya Memorial Hospital

Nagoya University Hospital

Naha City Hospital

Nakamura Memorial Hospital

Nakamura Memorial South Hospital

Nakano General Hospital

Nakatsu Gastrointestinal Hospital

Namegata District General Hospital

Nanbu Tokushukai Hospital

Nantan General Hospital

Nantan General Hospital

Nara City Hospital

Nara Medical University Hospital

Nara Prefectural Hospital

Nara Prefectural Mimuro Hospital

Narita Red Cross Hospital

Nasu Neurosurgical Hospital

Nasu Red Cross Hospital

National Cancer Center Hospital

National Center for Child Health and Development

National Center for Global Health and Medicine

National Cerebral and Cardiovascular Center

National Disaster Medical Center

National Fukuoka-Higashi Medical Center

National Hospital Organization Chiba Medical Center

National Hospital Organization Hamada Medical Center

National Hospital Organization Himeji Medical Center

National Hospital Organization Kanazawa Medical

National Hospital Organization Kanmon Medical Center

National Hospital Organization Kobe Medical Center

(Continued)

Appendix Table 1. (Continued)

National Hospital Organization Kumamoto Medical

National Hospital Organization Kure Medical Center

National Hospital Organization Maizuru Medical Center

National Hospital Organization Minami Wakayama

Medical Center

National Hospital Organization Nagasaki Medical Center

National Hospital Organization Nara Medical Center

National Hospital Organization Okayama Medical Center

National Hospital Organization Osaka Minami Medical Center

National Hospital Organization Saitama National

Hospital

National Hospital Organization Shizuoka Medical Center

National Hospital Organization Takasaki General

Medical Center

National Hospital Organization Ureshino Medical Center

National Hospital Organization Utano Hospital

National Hospital Organization Yokohama Medical

Center

National Kyushu Medical Center

Nayoro City General Hospital

Nihon University Itabashi Hospital

Niigata City General Hospital

Niigata Minami Hospital

Niigata Neurosurgical Hospital and Brain Research

Center

Niigata Prefectural Cancer Center Hospital

Niigata Prefectural Central Hospital

Niigata Prefectural Shibata Hospital

Niigata Prefectural Tokamachi Hospital

Niigata Rosai Hospital

Niigata University Medical and Dental Hospital

Nippon Medical School Chiba Hokusoh Hospital

Nippon Medical School Hospital

Nippon Medical School Musashi Kosugi Hospital

Nippon Medical School Tama-Nagayama Hospital

Nipponbashi Hospital

Nishi-Agatsuma Welfare Hospital

Nishi-Kobe Medical Center

Nishinara Tyuou Hospital

Nishinomiya Kyoritsu Neurosurgical Hospital

Nishio Municipal Hospital

Nishisaitama-chuo National Hospital Nishiwaki Municipal Hospital

North Fukushima Medical Center

North Osaka Police Hospital

NTT Medical Center Tokyo

Obara Hospital

Obihiro-Kosei General Hospital

Odate Municipal General Hospital

Odawara Municipal Hospital Oe Kyodou Hospital

Ofunato Hospital

Ogachi Central Hospital

Ogaki Municipal Hospital

Ogori Daiichi General Hospital

Appendix Table 1. (Continued)

Ohda Municipal Hospital Ohkawara Neurosurgical Hospital Ohmiya Chuo General Hospital Ohnishi Neurological Center Ohta General Hospital Oita Prefectural Hospital Oita University Hospital Oita-Oka-Hospital

Okayama East Neurosurgical Clinic

Okayama Kyokuto Hospital

Okayama City Hospital

Okayama Kyoritsu General Hospital

Okayama Rosai Hospital Okayama University Hospital Okazaki City Hospital Okinawa Chubu Hospital Okinawa Hokubu Hospital Okinawa Kyodo Hospital

Okinawa Prefectural Miyako Hospital

Okinawa Prefectural Nanbu Medical Center and

Children's Medical Center

Okinawa Prefectural Yaeyama Hospital Okitama Public General Hospital Okyama East Neurosurgical clinic Omihachiman Community Medical Center

Omori Red Cross Hospital Omuta City Hospital Ooi-Byouin Ookuma Hospital Ota Memorial Hospital

Osaka City General Hospital Osaka City University Hospital Osaka General Medical Center Osaka Kosei-Nenkin Hospital

Osaka Medical Center

Osaka Medical College Hospital

Osaka Mishima Critical Care Medical Center

Osaka Neurological Institute Osaka Neurosurgical Hospital Osaka Police Hospital

Osaka Prefectural Senshu Critical Care Medical Center

Osaka Red Cross Hospital Osaka Rosai Hospital Osaka University Hospital

Otemae Hospital Otsu Municipal Hospital

Research Institute for Brain and Blood Vessels Akita

Rumoi Central Clinic Sadamoto Hospital

Saga Prefectural Hospital Koseikan Saga Social Insurance Hospital Sagamihara Kyodo Hospital Sagamihara-chuo Hospital

Saisei-kai Yokohama-shi Nanbu Hospital

Saiseikai Central Hospital

Saiseikai Fukuoka General Hospital

Saiseikai Gose Hospital Saiseikai Hita Hospital

Saiseikai Hyuga Hospital Saiseikai Imabari Hospital Saiseikai Kumamoto Hospital Saiseikai Kurihashi Hospital Saiseikai Kyoto Hospital

Saiseikai Matsusaka General Hospital Saiseikai Matsuyama Hospital Saiseikai Nagasaki Hospital Saiseikai Nakatsu Hospital Saiseikai Nara Hospital Saiseikai Noe Hospital Saiseikai Saijo Hospital

Saiseikai Shimonoseki General Hospital

Saiseikai Toyama Hospital Saiseikai Utsunomiya Hospital Saiseikai Yahata General Hospital Saiseikai Yamaguchi Hospital Saiseikai Yokohanashi Tobu Hospital

Saiseikai-Chuwa Hospital Saitama Cancer Center Hospital

Saitama Cardiovascular and Respiratory Center

Saitama Medical Center

Saitama Medical Center Jichi Medical University

Saitama Medical University Hospital

Saitama Medical University International Medical Center

Saitama Municipal Hospital Saito Memorial Hospital Sakai City Hospital

Sakai Hospital Kinki University Faculty of Medicine

Sakaide Municipal General Hospital

Saku Central Hospital Sakura General Hospital Sakurakai Hospital Sanda City Hospital Sankoukai Miyazaki Hospital

Sannocho Hospital

Sano Kousei General Hospital

Sanyudo Hospital

Sapporo City General Hospital Sapporo Higashi-Tokushukai Hospital

Sasebo Chuo Hospital Sasebo City General Hospital

Sayama Hospital

Seguchi Neurosurgery Hospital Seirei Hamamatsu General Hospital

Seirei Memorial Hospital

Seirei Mikatagahara General Hospital

Seirei Yokohama Hospital Seiyu Memorial Hospital Sendai City Hospital

Sendai Medical Association Hospital

Sendai Open Hospital

Senpo Tokyo Takanawa Hospital

Senseki Hospital

Shakaihoken Kobe Central Hospital Shakaihoken Shimonoseki Kosei Hospital

Shiga Medical Center for Adults

Shiga University of Medical Science Hospital

(Continued)

Appendix Table 1. (Continued)

Shimada City Hospital

Shimane Prefectural Central Hospital

Shimane University Hospital Shimonoseki City Hospital Shimotsuga General Hospital

Shin Koga Hospital Shin Yukuhashi Hospital Shin-Tokyo Hospital

Shingu Municipal Medical Center

Shinko Hospital

Shinoda General Hospital Shinonoi General Hospital Shinrakuen Hospital

Shinseikai Toyama Hospital Shinshu Ueda Medical Center

Shinsuma Hospital

Shirahama Hamayu Hospital Shirakawa Kosei General Hospital Shiroishi neurosurgical Hospital

Shiroyama Hospital Shiseikai Daini Hospital Shizuoka Children's Hospital Shizuoka City Hospital Shizuoka General Hospital Shobara Red Cross Hospital Shonai Hospital

Shonan Kamakura General Hospital

Showa General Hospital Showa Inan General Hospital Showa University Fujigaoka Hospital

Showa University Hospital Social Insurance Chukyo Hospital Social Insurance Chuo General Hospital Social Insurance Takahama Hospital

Soseikai General Hospital South Miyagi Medical Center Southern Tohoku General Hospital

St Marianna University School of Medicine Toyoko Hospital

St Francisco Hospital

St Marianna University School of Medicine Hospital

Steeel Memorial Hirohata Hospital Steel Memorial Yawata Hospital Suiseikai Kajikawa Hospital Suita Municipal Hospital Suwa Central Hospital Suwakohan Hospital Suzuka Kaisei Hospital Tachikawa Medical Center Takada Chuo Hospital Takamatsu Municipal Hospital

Takamatsu Municipal Hospital Takamatsu Red Cross Hospital Takarazuka City Hospital Takarazuka Daiichi Hospital Takashima Municipal Hospital Takatsuki General Hospital Takeda General Hospital

Takikawa Neurosurgery Hospital

Tama-Hokubu Medical Center Tama-Nanbu Chiiki Hospital Tamana Central Hospital Tane General Hospital Tano Hospital

Tanushimaru Central Hospital

Tatebayashi Kosei Hospital Teiko University Chiba Medical Center

Teikyo University School of Medicine Hospital,

Mizonokuchi Tekeda Hospital

Tenshindo Hetsugi Hospital

The Taiju-Kai Foundation Social Medical Corporation

Kaisei General Hospital Tochigi National Hospital

Toho University Ohashi Medical Center

Tohoku KoseiNenkin Hospital

Tokai Central Hospital of the Mutual Aid Association of

Public School Teachers

Tokai University Hachioji Hospital

Tokai University Hospital
Tokai University Oiso Hospital

Toki General Hospital

Tokuda Neurosurgical Hospital

Tokushima Prefectural Central Hospital Tokushima Prefectural Kaifu Hospital Tokushima University Hospital Tokuyama Central Hospital

Tokuyama Central Hospita Tokyo Kyosai Hospital

Tokyo Medical And Dental University Hospital Faculity of Medicine

Tokyo Medical University Hospital

Tokyo Medical University Ibaraki Medical Center

Tokyo Metropalitan Hiroo Hospital

Tokyo Metropolitan Health and Medical Corporation

Toshima Hospital

Tokyo Metropolitan Ohtsuka Hospital Tokyo Metropolitan Police Hospital

Tokyo Women's Medical University Hospital

Tokyo Women's Medical University Yachiyo Medical Center

Tokyo Women's University Medical Center East

Tokyo-Teishin Hospital

Tokyo-West Tokushukai Hospital

Tokyu Medical University Hachioji Medical Center

Tomakomai Neurosurgical Hospital Tomakomai Nissho Hospital Tomei Atsugi Hospital Tominaga Hospital

Tominaga Hospital
Tomioka General Hospital
Tonami General Hospital
Tone Central Hospital
Tosei General Hospital
Tottori Pref.Kousei Hospital

Tottori Red Cross Hospital Tottori Seikyo Hospital Tottori University Hospital

Toyama City Hospital

(Continued)

Toyama Prefectural Central Hospital

Toyama Rosai Hospital

Toyohashi Medical Center

Toyokawa City Hospital

Toyooka Chuo Hospital

Toyooka Public Hospitals' association Toyooka Hospital

Toyota Kosei Hospital

Tsuchiura Kyodo General Hospital

Tsukazaki Hospital

Tsukuba Medical Center Hospital

Tsushima City Hospital

Tsuyama Central Hospital

Ube Industries, ltd. Central Hospital

Ueyama Hospital.

Ugadake Hospital

University Hospital of the Ryukyus

University of Miyazaki Hospital

University of Tokyo Hospital

University of Tsukuba Hospital

University of Yamanashi Hospital

Urasoe General Hospital

Ushiku Aiwa General Hospital

Ushioda General Hospital

Uwajima City Hospital

Uwajima Tokushukai Hospital

Veritas Hospital

Wada Hospital

Wakakusa Dai-ichi Hospital

Wakayama Co-operative Hospital

Wakayama Medical University Hospital

Wakayama Medical University Kihoku Hospital

Wakayama Rosai Hospital

Wakayama Saiseikai Hospital

Yaentoge Neurosurgery Hospital

Yagi Neurosurgical Hospital

Yaizu City General Hospital

Yamachika Memorial Hospital

Yamada Kinen Hospital

Yamada Red Cross Hospital

Yamagata City Hospital Saiseikan

Yamagata Prefectural Kahoku Hospital

Yamagata Prefectural Shinjo Hospital

Yamagata University Hospital

Yamaguchi Grand Medical Center

Yamaguchi Red Cross Hospital

Yamaguchi Rousai Hospital

Yamaguchi University Hospital

Yamamoto Memorial Hospital

Yamanashi Kosei Hospital

Yamanashi Prefectural Central Hospital

Yamanashi Red Cross Hospital

Yamashiro Public Hospital

Yamato Municipal Hospital

Yao Tokushukai General Hospital

Yasugi municipal Hospital

Yatsuo General Hospital

Yatsushiro Health Insurance General Hospital

Yahata General Hospital

(Continued)

Appendix Table 1. (Continued)

Yayoigaoka Hospital

Yodogawa Christian Hospital

Yokkaichi Municipal Hospital

Yokohama Central Hospital

Yokohama City Minato Red Cross Hospital

Yokohama City University Hospital

Yokohama City University Medical Center

Yokohama General Hospital

Yokohama Rosai Hospital

Yokohama Sakae Kyosai Hospital

Yokohama Shin-midori General Hospital

Yokohama Stroke and Brain Center

Yokohamashintoshi Neurosurgical Hospital

Yokosuka General Hospital Uwamachi

Yomeikai Obase Hospital

Yonabaru Chu-ou Hospital

Yonago Medical Center

Yonezawa City Hospital

Yuaikai Hospital

Yukioka Hospital