


Fig. 1 Inclusion and exclusion criteria



Stata 12 (Stata Corp, College Station, TX, USA) was used for all statistical analyses. In all analyses, $P < 0.05$ was taken to indicate statistical significance.

Results

Participant selection and baseline characteristics

A total of 7,449 participants undergoing PFTs and 16,743 participants undergoing eGFR and urinary protein measurement were enrolled between 1 April 2005, and 30 June 2010 at Kanazawa University Hospital. A total of 562 participants underwent both PFTs and renal function tests. We excluded 252 participants with major respiratory disease and abnormal CT and 177 participants with smoking history. Thus, 133 participants were finally included in this study (Fig. 1).

Table 1 shows the characteristics of the subjects. Neither age, body mass index (BMI), Hb, brain natriuretic peptide (BNP) nor the presence of diabetes and hypertension were significantly different between categories. As we excluded participants with smoking history, females were predominant. The percentages of males in category 2 (2.3 %) and category 4 (0 %) were significantly lower than those in category 1 (32.4 %) and category 3 (22.2 %).

Changes in pulmonary function according to eGFR and urinary protein excretion

Table 2 shows a comparison between PFT parameters in each category. Hb-adjusted %DL_{CO} in category 4 (46.2 ± 7.5) and category 2 (63.6 ± 17.8) were significantly lower than in category 1 (75.8 ± 18.9) ($P < 0.05$). Other PFT markers (%VC, %FEV1, FEV1/FVC, %TLC,

and %RV) were not significantly different between the categories.

Association between Hb-adjusted %DL_{CO} and eGFR and urinary protein excretion

Figure 2 shows the results of single regression analysis between Hb-adjusted %DL_{CO} and kidney function. Hb-adjusted %DL_{CO} was significantly positively correlated with eGFR ($R = 0.36, P < 0.001$, Fig. 2a), but was significantly negatively correlated with urinary protein ($R = -0.25, P = 0.005$, Fig. 2b).

Figure 3 shows the results of single regression analysis between Hb-adjusted %DL_{CO} and eGFR. Hb-adjusted %DL_{CO} was weakly correlated with eGFR in patients with urinary protein <0.3 g/gCr ($R = 0.30, P = 0.001$, Fig. 3a), but was strongly correlated with eGFR in patients with urinary protein ≥0.3 g/gCr ($R = 0.81, P < 0.001$, Fig. 3b).

To examine the precise relationships between kidney function and Hb-adjusted %DL_{CO}, multiple linear regression analysis was conducted in all patients examined in this study. As shown in Table 3, eGFR was associated with Hb-adjusted %DL_{CO} ($P = 0.023$) after adjusting for physical features (age and gender), expiratory minute volume (VE), and the presence or absence of DM and Hypertension. Urinary protein tended to be associated with Hb-adjusted %DL_{CO} ($P = 0.095$) after adjusting for physical features (age and gender), VE, and the presence or absence of DM, and hypertension.

Discussion

This study was performed to examine the associations between kidney function or urinary protein excretion and

Table 1 Characteristics of subjects

Category	1	2	3	4	Total	<i>P</i>
eGFR (ml/min/1.73 m ²)	≥60	<60	≥60	<60		
Urinary protein (g/gCr)	<0.3	<0.3	≥0.3	≥0.3		
<i>n</i> (% of category)	74 (55.6)	43 (32.3)	9 (6.8)	6 (4.5)	133	
Age (years)	52.7 ± 17.9	56.8 ± 18.5	49.0 ± 17.3	64.7 ± 9.2	54.4 ± 17.9	N.S.
Sex (% male)	24 (32.4)	1 (2.3)	2 (22.2)	0 (0)	27 (20.3)	<0.001
BMI (kg/m ²)	22.5 ± 3.8	22.2 ± 5.5	21.2 ± 4.2	21.8 ± 3.7	22.3 ± 4.4	N.S.
Albumin (g/dl)	3.97 ± 0.54	4.11 ± 1.45	3.86 ± 0.89	3.28 ± 0.82	3.97 ± 0.97	N.S.
Creatinine (mg/dl)	0.66 ± 0.14	1.02 ± 0.31	0.69 ± 0.09	1.37 ± 0.92	0.81 ± 0.34	<0.001
eGFR (ml/min/1.73 m ²)	86.2 ± 24.7	48.7 ± 9.3	81.3 ± 22.7	40.5 ± 17.9	71.6 ± 28.2	<0.001
Urinary protein (g/gCr)	0.06 ± 0.06	0.07 ± 0.06	1.22 ± 0.07	0.94 ± 1.39	0.18 ± 0.5	<0.001
Hb (g/dl)	12.6 ± 2.3	11.6 ± 1.9	11.0 ± 2.6	9.27 ± 2.2	12.0 ± 2.3	N.S.
BNP (pg/ml) (<i>n</i> of >20 pg/ml)	21.4 ± 49.2 (8)	27.2 ± 68.5 (7)	10.2 ± 5.8 (0)	18.4 ± 16.8 (1)	22.9 ± 55.0 (16)	N.S.
The presence of comorbidity						
Diabetes (%)	12 (17.6)	4 (9.3)	1 (11.1)	1 (16.7)	18 (13.5)	
Hypertension (%)	6 (8.1)	2 (4.7)	1 (11.1)	1 (16.7)	10 (7.5)	
Connective tissue disease (%)	25 (33.8)	19 (44.2)	2 (22.2)	1 (16.7)	47 (35.3)	
Haematologic disease (%)	13 (17.6)	3 (7.0)	2 (22.2)	2 (33.3)	20 (15.0)	
Cardiac disease (%)	5 (6.8)	2 (4.7)	0 (0)	0 (0)	7 (5.3)	
Cancer (%)	14 (18.9)	8 (18.6)	3 (33.3)	1 (16.7)	26 (19.5)	
Other diseases ^a (%)	8 (10.8)	10 (23.3)	2 (22.2)	1 (16.7)	21 (15.8)	

Data are presented as means ± standard deviation (SD)

N.S. not significant

^a Depression, vitreous haemorrhage, cataract, tonsillitis, sleep apnoea syndrome, thyroiditis, chronic hepatitis B, chronic hepatitis C, ileus, condyloma acuminatum, toxic eruption

Table 2 Comparison between PFT parameters in each category (renal function)

Category	1	2	3	4	Total
eGFR (ml/min/1.73 m ²)	≥60	<60	≥60	<60	
Urinary protein (g/gCr)	<0.3	<0.3	≥0.3	≥0.3	
%VC	109.2 ± 18.6	107.2 ± 15.7	111.3 ± 17.8	101.5 ± 14.2	108.3 ± 17.4
%FEV1	109.5 ± 20.4	114.2 ± 24.6	110.7 ± 12.9	105.9 ± 10.4	110.9 ± 21.1
FEV1/FVC	81.2 ± 7.2	80.1 ± 7.6	84.1 ± 7.7	76.7 ± 6.1	80.8 ± 7.4
%TLC	99.4 ± 14.0	99.2 ± 13.9	102.2 ± 12.1	96.1 ± 10.9	99.4 ± 13.6
%RV	94.2 ± 22.6	95.2 ± 20.4	94.3 ± 22.9	93.7 ± 15.9	94.6 ± 21.4
Hb-adjusted %DL _{CO}	75.8 ± 18.9	63.6 ± 17.8*	62.5 ± 15.7	46.2 ± 7.5*	69.6 ± 18.9
VE (l/min)	9.37 ± 3.33	8.35 ± 2.53	8.53 ± 2.57	8.86 ± 1.67	9.00 ± 3.00

Data are presented as means ± standard deviation (SD)

* *P* < 0.05, compared with category 1

pulmonary function in participants undergoing examination for eGFR, proteinuria, and PFTs. Our data indicated that decreased eGFR (<60 ml/min/1.73 m²) along with increased urinary protein (≥0.3 g/gCr) were associated with decreased Hb-adjusted %DL_{CO}. These results suggest that patients with CKD may have impaired pulmonary capacity.

Our data indicated that increased urinary protein was associated with decreased Hb-adjusted %DL_{CO}. Urinary

protein is one of the surrogate markers for vascular damage in the kidney [16]. In addition, urinary protein is also a surrogate marker of interstitial fibrosis in the kidney [17]. Various studies demonstrated that increased serum levels of cytokines and chemokines in patients with end-stage renal disease (ESRD) promote kidney fibrosis [10, 17–19]. These increased serum cytokine and chemokine levels would affect lung fibrosis in a similar way to the kidney. Vascular damage (impaired capillary vessels in lung

Fig. 2 Relationships between Hb-adjusted %DL_{CO} and kidney function (eGFR and urinary protein)




Fig. 3 Relationships between Hb-adjusted %DL_{CO} and eGFR depending on urinary protein




Table 3 Multiple linear regression analysis with Hb-adjusted %DL_{CO} as the dependent variable and age, gender, VE, eGFR, urinary protein, and the presence or absence of DM and hypertension as independent variables in all patients examined

Coefficient	Partial regression		Standard partial	
	Coefficient		Regression coefficient	
Model	B	SE (B)	β	P
Constant	71.643	9.300	-0.142	
Age (years)	-0.149	0.111	0.037	0.182
Gender (male)	1.643	5.196	0.037	0.753
VE (l/min)	14.253	11.040	0.135	0.200
eGFR (<60 ml/min/1.73 m ²)	-10.260	4.422	-0.260	0.023*
Urinary protein (≥0.3 g/gCr)	-10.019	5.922	-0.169	0.095
Diabetes (yes)	-0.111	6.085	-0.001	0.985
Hypertension (yes)	13.609	7.764	0.182	0.083

Regression analysis was adjusted for each factor

* P < 0.05

alveoli) and interstitial fibrosis (impaired diffusion from lung alveoli to the lumens of capillary vessels) are associated with decreased Hb-adjusted %DL_{CO} [20–27]. These mechanisms may be associated with the correlation between urinary protein and Hb-adjusted %DL_{CO}.

Our data also indicated that decreased eGFR was associated with decreased Hb-adjusted %DL_{CO}. Decreased eGFR enhances the risk of atherosclerosis [28–30]. Patients with CKD are more likely to be complicated by whole-body atherosclerosis caused by many factors, such as elevation of blood pressure, impaired glucose tolerance, activation of the renin–angiotensin system, abnormalities of calcium homeostasis, or deposition of calcium in the vessels [31–33]. Furthermore, Kumar et al. [34] reported that patients with CKD and ESRD are more likely to be complicated by pulmonary embolism. Vascular damage (impaired capillary vessels in lung alveoli) is associated with decreased Hb-adjusted %DL_{CO} [20–25]. These findings may have an impact on the contribution to the correlation between eGFR and Hb-adjusted %DL_{CO}.

Our data further revealed that both urinary protein and decreased eGFR were associated with decreased Hb-adjusted %DL_{CO}. The coexistence of urinary protein and decreased eGFR may result in a greater increase in vascular damage (impaired capillary vessels in lung alveoli) [20–25] and more severe interstitial fibrosis (impaired diffusion from lung alveoli to the lumens of capillary vessels) [26, 27] compared to the presence of urinary protein or decreased eGFR alone. Thus, they can have a synergetic effect on decreasing Hb-adjusted %DL_{CO}. The mechanisms underlying the association between Hb-adjusted %DL_{CO} and decreased eGFR with increased urinary protein may be complicated. Therefore, further studies, such as longitudinal studies and basic research, are needed to clarify the detailed mechanisms underlying this association between the lungs and kidneys.

This study had some limitations, including sampling bias. The participants included in this study were limited to patients treated at Kanazawa University Hospital. As this is an advanced medical institution, our participants had an increased likelihood of having various complications. Therefore, it is possible that participants with normal pulmonary function did not undergo PFTs in this study, leading to overestimation of the relationships between renal dysfunction and pulmonary dysfunction. The other major limitation of this study may be confounding factors. Other factors have been reported to affect %DL_{CO}. Reduction of haemoglobin concentration and impairment of lung alveolus are two major factors that affect %DL_{CO} [15, 35–38]. Reduction of haemoglobin concentration was not excluded in this study, because the influence of anaemia on %DL_{CO} is adjusted by haemoglobin. Impairment of lung alveoli is mainly associated with smoking habit and chronic obstructive pulmonary disease. Therefore, these patients were excluded from the study population. Moreover, diffusing capacity of the lung is influenced not only by Hb and inflammatory status (e.g., profibrotic cytokines), but also by oedematous condition (the presence of heart failure even in occult status) [39]. CKD patients with advanced stages may be complicated with mild or sub-clinical heart failure. Further studies are required to overcome these limitations.

In conclusion, the present study indicated that eGFR and urinary protein excretion were associated with pulmonary function (Hb-adjusted %DL_{CO}). Further studies, such as longitudinal studies and basic research, are needed to confirm the present results and to clarify the detailed mechanisms underlying these observations.

Acknowledgments We wish to thank the staff of the Clinical Laboratory at Kanazawa University Hospital for their technical assistance in this study. This study was supported in part by Grant-in-Aids for Diabetic Nephropathy Research and for Diabetic Nephropathy and Nephrosclerosis Research, from the Ministry of Health,

Labour and Welfare of Japan and by the Ministry of Education, Science, Sports and Culture, Japan.

Conflict of interest All authors declare that they have no conflicts of interest associated with this study.

References

1. Bouchi R, Babazono T, Yoshida N, Nyumura I, Toya K, Hayashi T, et al. Association of albuminuria and reduced estimated glomerular filtration rate with incident stroke and coronary artery disease in patients with type 2 diabetes. *Hypertens Res*. 2010;33:1298–304.
2. Hamaguchi S, Tsuchihashi-Makaya M, Kinugawa S, Yokota T, Ide T, Takeshita A, et al. Chronic kidney disease as an independent risk for long-term adverse outcomes in patients hospitalized with heart failure in Japan. Report from the Japanese Cardiac Registry of Heart Failure in Cardiology (JCARE-CARD). *Circ J*. 2009;73:1442–7.
3. Ko GJ, Rabb H, Hassoun HT. Kidney-lung crosstalk in the critically ill patient. *Blood Purif*. 2009;28:75–83.
4. Klein CL, Hoke TS, Fang WF, Altmann CJ, Douglas IS, Faubel S. Interleukin-6 mediates lung injury following ischemic acute kidney injury or bilateral nephrectomy. *Kidney Int*. 2008;74:901–9.
5. Ishii T, Doi K, Okamoto K, Imamura M, Dohi M, Yamamoto K, et al. Neutrophil elastase contributes to acute lung injury induced by bilateral nephrectomy. *Am J Pathol*. 2010;177:1665–73.
6. Grigoryev DN, Liu M, Hassoun HT, Cheadle C, Barnes KC, Rabb H. The local and systemic inflammatory transcriptome after acute kidney injury. *J Am Soc Nephrol*. 2008;19:547–58.
7. Lie ML, White LE, Santora RJ, Park JM, Rabb H, Hassoun HT. Lung T lymphocyte trafficking and activation during ischemic acute kidney injury. *J Immunol*. 2012;189:2843–51.
8. Ahuja N, Andres-Hernando A, Altmann C, Bhargava R, Bacalja J, Webb RG, et al. Circulating IL-6 mediates lung injury via CXCL1 production after acute kidney injury in mice. *Am J Physiol Renal Physiol*. 2012;303:864–72.
9. Furuichi K, Shintani H, Sakai Y, Ochiya T, Matsushima K, Kaneko S, et al. Effects of adipose-derived mesenchymal cells on ischemia-reperfusion injury in kidney. *Clin Exp Nephrol*. 2012;16:679–89.
10. Furuichi K, Kaneko S, Wada T. Chemokine/chemokine receptor-mediated inflammation regulates pathologic changes from acute kidney injury to chronic kidney disease. *Clin Exp Nephrol*. 2009;13:9–14.
11. American Thoracic Society. Standardization of spirometry. *Eur Respir J*. 2005;26:319–38.
12. Yamamoto R, Nagasawa Y, Shoji T, Iwatani H, Hamano T, Kawada N, et al. Cigarette smoking and progression of IgA nephropathy. *Am J Kidney Dis*. 2010;56:313–24.
13. Soriano JB, Zielinski J, Price D. Screening for and early detection of chronic obstructive pulmonary disease. *Lancet*. 2009;374:721–32.
14. American Thoracic Society. Single-breath carbon monoxide diffusing capacity (transfer factor). Recommendations for a standard technique—1995 update. *Am J Respir Crit Care Med*. 1995;152:2185–98.
15. Macintyre N, Crapo RO, Viegi G, Johnson DC, van der Grinten CP, Brusasco V, et al. Standardisation of the single-breath determination of carbon monoxide uptake in the lung. *Eur Respir J*. 2005;26:720–35.
16. Deckert T, Feldt-Rasmussen B, Borch-Johnsen K, Jensen T, Kofoed-Enevoldsen A. Albuminuria reflects widespread vascular damage: the Steno hypothesis. *Diabetologia*. 1989;32:219–26.

17. Wei CC, Li HH, Hsu YH, Hsing CH, Sung JM, Chang MS. Interleukin-20 targets renal cells and is associated with chronic kidney disease. *Biochem Biophys Res Commun*. 2008;374:448–53.
18. Wada T, Sakai N, Sakai Y, Matsushima K, Kaneko S, Furuichi K. Involvement of bone-marrow-derived cells in kidney fibrosis. *Clin Exp Nephrol*. 2011;15:8–13.
19. Furuichi K, Wada T, Kaneko S, Murphy PM. Roles of chemokines in renal ischemia/reperfusion injury. *Front Biosci*. 2008;13:4021–8.
20. Steenhuis LH, Groen HJ, Koëter GH, van der Mark TW. Diffusion capacity and haemodynamics in primary and chronic thromboembolic pulmonary hypertension. *Eur Respir J*. 2000;16:276–81.
21. Fernandez-Bonetti P, Lupi-Herrera E, Martinez-Guerra ML, Barrios R, Seoane M, Sandoval J. Peripheral airways obstruction in idiopathic pulmonary artery hypertension (primary). *Chest*. 1983;83:732–8.
22. Tashkin DP, Clements PJ, Wright RS, Gong H Jr, Simmons MS, Lachenbruch PA, et al. Interrelationships between pulmonary and extrapulmonary involvement in systemic sclerosis. A longitudinal analysis. *Chest*. 1994;105:489–95.
23. Hills EA, Geary M. Membrane diffusing capacity and pulmonary capillary volume in rheumatoid disease. *Thorax*. 1980;35:851–5.
24. Songür N, Songür Y, Tüzün M, Doğan I, Tüzün D, Ensari A, et al. Pulmonary function tests and high-resolution CT in the detection of pulmonary involvement in inflammatory bowel disease. *J Clin Gastroenterol*. 2003;37:292–8.
25. Herrlinger KR, Noftz MK, Dalhoff K, Ludwig D, Stange EF, Fellermann K. Alterations in pulmonary function in inflammatory bowel disease are frequent and persist during remission. *Am J Gastroenterol*. 2002;97:377–81.
26. Zompatori M, Calabrò E, Chetta A, Chiari G, Marangio E, Olivieri D. Chronic hypersensitivity pneumonitis or idiopathic pulmonary fibrosis? Diagnostic role of high resolution Computed Tomography (HRCT). *Radiol Med*. 2003;106:135–46.
27. Watters LC, King TE, Schwarz MI, Waldron JA, Stanford RE, Cherniack RM. A clinical, radiographic, and physiologic scoring system for the longitudinal assessment of patients with idiopathic pulmonary fibrosis. *Am Rev Respir Dis*. 1986;133:97–103.
28. Ishizaka N, Ishizaka Y, Toda E, Koike K, Seki G, Nagai R, et al. Association between chronic kidney disease and carotid intima-media thickening in individuals with hypertension and impaired glucose metabolism. *Hypertens Res*. 2007;30:1035–41.
29. Wada T, Shimizu M, Toyama T, Hara A, Kaneko S, Furuichi K. Clinical impact of albuminuria in diabetic nephropathy. *Clin Exp Nephrol*. 2012;16:96–101.
30. Gansevoort RT, Matsushita K, van der Velde M, Astor BC, Woodward M, Levey AS, et al. Lower estimated GFR and higher albuminuria are associated with adverse kidney outcomes in both general and high-risk populations. A collaborative meta-analysis of general and high-risk population cohorts. *Kidney Int*. 2011;80:93–104.
31. Bakris GL, Williams M, Dworkin L, Elliott WJ, Epstein M, Toto R, et al. Preserving renal function in adults with hypertension and diabetes: a consensus approach. National Kidney Foundation Hypertension and Diabetes Executive Committees Working Group. *Am J Kidney Dis*. 2000;36:646–61.
32. Ogihara T, Saruta T, Rakugi H, Fujimoto A, Ueshima K, Yasuno S, et al. Relationship between the achieved blood pressure and the incidence of cardiovascular events in Japanese hypertensive patients with complications: a sub-analysis of the CASE-J trial. *Hypertens Res*. 2009;32:248–54.
33. Pedrinelli R, Giampietro O, Carmassi F, Melillo E, Dell’Omo G, Catapano G, et al. Microalbuminuria and endothelial dysfunction in essential hypertension. *Lancet*. 1994;344:14–8.
34. Kumar G, Sakhuja A, Taneja A, Majumdar T, Patel J, Whittle J, et al. Pulmonary Embolism in Patients with CKD and ESRD. *Clin J Am Soc Nephrol*. 2012;7:1584–90.
35. Riepl G. Effects of abnormal hemoglobin concentration in human blood on membrane diffusing capacity of the lung and on pulmonary capillary blood volume. *Respiration*. 1978;36:10–8.
36. Klings ES, Wyszynski DF, Nolan VG, Steinberg MH. Abnormal pulmonary function in adults with sickle cell anemia. *Am J Respir Crit Care Med*. 2006;173:1264–9.
37. Morrison NJ, Abboud RT, Ramadan F, Miller RR, Gibson NN, Evans KG, et al. Comparison of single breath carbon monoxide diffusing capacity and pressure-volume curves in detecting emphysema. *Am Rev Respir Dis*. 1989;139:1179–87.
38. Gould GA, Redpath AT, Ryan M, Warren PM, Best JJ, Flenley DC, et al. Lung CT density correlates with measurements of airflow limitation and the diffusing capacity. *Eur Respir J*. 1991;4:141–6.
39. Cattadori G, Wasserman K, Meloni C, Mustaq S, Contini M, Apostolo A, et al. Alveolar membrane conductance decreases as BNP increases during exercise in heart failure. Rationale for BNP in the evaluation of dyspnea. *J Card Fail*. 2009;15:136–44.

Clinical impact of albuminuria and glomerular filtration rate on renal and cardiovascular events, and all-cause mortality in Japanese patients with type 2 diabetes

Takashi Wada · Masakazu Haneda · Kengo Furuichi · Tetsuya Babazono · Hiroki Yokoyama · Kunitoshi Iseki · Shin-ichi Araki · Toshiharu Ninomiya · Shigeo Hara · Yoshiki Suzuki · Masayuki Iwano · Eiji Kusano · Tatsumi Moriya · Hiroaki Satoh · Hiroyuki Nakamura · Miho Shimizu · Tadashi Toyama · Akinori Hara · Hirofumi Makino · The Research Group of Diabetic Nephropathy, Ministry of Health, Labour, and Welfare of Japan

Received: 3 August 2013 / Accepted: 21 September 2013 / Published online: 17 October 2013
© Japanese Society of Nephrology 2013

Abstract

Background The number of patients suffering from diabetic nephropathy resulting in end-stage kidney disease is increasing worldwide. In clinical settings, there are limited data regarding the impact of the urinary albumin-to-creatinine ratio (UACR) and reduced estimated glomerular filtration rate (eGFR) on renal and cardiovascular outcomes and all-cause mortality.

Methods We performed a historical cohort study of 4328 Japanese participants with type 2 diabetes from 10 centers. Risks for renal events (requirement for dialysis or

transplantation, or half reduction in eGFR), cardiovascular events (cardiovascular death, nonfatal myocardial infarction, or nonfatal stroke), and all-cause mortality were assessed according to UACR and eGFR levels.

Results During follow-up (median 7.0 years, interquartile range 3.0–8.0 years), 419 renal events, 605 cardiovascular events and 236 deaths occurred. The UACR levels increased the risk and the adjusted hazard ratios for these three events. In addition to the effects of UACR levels, eGFR stages significantly increased the adjusted hazard ratios for renal events and all-cause mortality, especially in

T. Wada · K. Furuichi · M. Shimizu · T. Toyama · A. Hara
Division of Nephrology, Kanazawa University Hospital,
Kanazawa, Japan

T. Wada (✉)
Division of Nephrology, Department of Laboratory Medicine,
Institute of Medical, Pharmaceutical and Health Sciences,
Faculty of Medicine, Kanazawa University, Kanazawa,
13-1 Takara-machi, Kanazawa 920-8641, Japan
e-mail: twada@m-kanazawa.jp

M. Haneda
Department of Medicine, Asahikawa Medical University,
Asahikawa, Japan

K. Furuichi · M. Shimizu · T. Toyama · A. Hara
Department of Disease Control and Homeostasis, Institute of
Medical, Pharmaceutical and Health Sciences, Kanazawa
University, Kanazawa, Japan

T. Babazono
Division of Nephrology and Hypertension, Diabetes Center,
Tokyo Women's Medical University School of Medicine,
Tokyo, Japan

H. Yokoyama
Jiyugaoka Medical Clinic, Internal Medicine, Obihiro, Japan

K. Iseki
Dialysis Unit, University Hospital of the Ryukyus, Nishihara,
Okinawa, Japan

S. Araki
Department of Medicine, Shiga University of Medical Science,
Otsu, Shiga, Japan

T. Ninomiya
Department of Medicine and Clinical Science, Graduate School
of Medical Sciences, Kyushu University, Fukuoka, Japan

S. Hara
Center of Health Management, Toranomon Hospital, Tokyo,
Japan

Y. Suzuki
Health Administration Center, Niigata University, Niigata, Japan

M. Iwano
Division of Nephrology, Department of General Medicine,
University of Fukui, Fukui, Japan

E. Kusano
Division of Nephrology, Department of Internal Medicine, Jichi
Medical University, Tochigi, Japan

patients with macroalbuminuria. Diabetic nephropathy score, based on the prognostic factors, well predicted incidence rates per 1000 patient/year for each event.

Conclusions Increased UACR levels were closely related to the increase in risks for renal, cardiovascular events and all-cause mortality in Japanese patients with type 2 diabetes, whereas the association between high levels of UACR and reduced eGFR was a strong predictor for renal events.

Keywords Diabetic nephropathy · Chronic kidney disease · Albuminuria · Cardiovascular disease · Mortality · Glomerular filtration rate

Introduction

Diabetic nephropathy is a leading cause of end-stage kidney (renal) disease (ESKD or ESRD) worldwide [1]. In addition, cardiovascular diseases and deaths increase in patients with diabetic nephropathy before and after dialysis [2–4]. Therefore, to determine and manage risk factors for progression of renal and cardiovascular outcomes and mortality is of importance to prolong the life expectancy of diabetic patients.

A high urinary albumin-to-creatinine ratio (UACR) and low estimated glomerular filtration rate (eGFR) have been believed to be predictors for diabetic ESKD and death [5–7]. Kidney Disease Improving Global Outcomes (KDIGO) provided a new classification for chronic kidney disease (CKD) by adding stages that stratified urinary albumin excretion as well as eGFR and emphasizing clinical diagnosis [8]. This new classification, mainly based on the collaborative meta-analysis of general population cohorts [8], has shed light on prognosis assigned by clinical diagnosis, stage, and other key factors relevant to renal and cardiovascular outcomes. However, the clinical impact of UACR levels in combination with eGFR on outcomes in

Japanese patients with diabetic nephropathy needs to be confirmed. Therefore, deeper clinical insights of UACR along with GFR are required to provide a key for the pathogenesis and outcomes of progressive renal complications, and associated cardiovascular events in type 2 diabetic patients.

Here we examined the prognostic value of UACR and eGFR for renal events, cardiovascular events and all-cause mortality in Japanese patients with type 2 diabetes. Furthermore, we proposed a diabetic nephropathy score for predicting prognosis in diabetic patients.

Subjects and methods

Subjects

This study was a historical cohort consisting of 4814 Japanese patients with type 2 diabetes who were treated by trained physicians at 10 centers between 1985 and 2010. Four hundred and fifty-nine patients were excluded because of age <18 ($n = 6$), follow-up <1 year ($n = 151$) and no measurement of urinary albumin or HbA1c or blood pressure (BP) ($n = 329$), leaving 4328 Japanese patients to be enrolled in this study. Patients with secondary diabetes, renal transplantation or dialysis were also excluded.

This study was conducted according to the ethical guidelines for epidemiological research designed by the Japanese Ministry of Education, Culture, Sports, Science and Technology and Ministry of Health, Labour, and Welfare (http://www.lifescience.mext.go.jp/files/pdf/n796_01.pdf).

The study design was included in a comprehensive protocol of retrospective study at the Division of Nephrology, Kanazawa University Hospital approved by Kanazawa University ethical committee (approval number 815).

Follow-up and assessments

Type 2 diabetes was defined according to the Japan Diabetes Society (JDS) criteria [9]. In this historical cohort study, UACR and eGFR were also determined. Measurement of UACR, by a turbidimetric immunoassay at each laboratory, was performed on spot urine samples at baseline. Serum creatinine was measured at baseline, at subsequent yearly intervals, and at the end of follow-up. Serum and urinary concentrations of creatinine were measured by an enzymatic method, and eGFR was estimated using the equation proposed by the Japanese Society of Nephrology [10]. Both UACR and serum creatinine were measured at local laboratories. At each study visit, blood pressure (BP) was measured in the sitting position. Hypertension was defined as $BP \geq 140/90$ mmHg or

T. Moriya
Health Care Center, Kitasato University, Sagami-hara, Japan

H. Satoh
Department of Nephrology, Hypertension, Diabetology,
and Metabolism, Fukushima Medical University,
Fukushima, Japan

H. Nakamura
Department of Environmental and Preventive Medicine, Institute
of Medical, Pharmaceutical and Health Sciences, Kanazawa
University, Kanazawa, Japan

H. Makino
Department of Medicine and Clinical Science, Okayama
University Graduate School of Medicine, Dentistry and
Pharmaceutical Sciences, Okayama, Japan

current use of antihypertensive drugs. Non-fasting blood samples were obtained for measurements of HbA1c and lipid levels at local laboratories. HbA1c was measured and standardized by the JDS (normal range 4.3–5.8 %) and certified by the US National Glycohemoglobin Standardization Program (National Glycohemoglobin Standardization Program, NGSP; NGSP = JDS + 0.4) [9].

UACR and GFR categories

Based on the new classification of CKD [8], the albuminuria category was classified at baseline as normoalbuminuria (<30 mg/g), microalbuminuria (≥ 30 and <300 mg/g), and macroalbuminuria (≥ 300 mg/g). In addition, baseline eGFR levels were divided into six categories: ≥ 90 , 60–89, 45–59, 30–44, 15–29 and <15 ml/min per 1.73 m². Patients examined in this study were categorized and assessed based on the above classifications.

Outcomes

The main outcomes of this study were renal events (requirement for dialysis or transplantation, or half reduction in eGFR), cardiovascular events (cardiovascular death, nonfatal myocardial infarction, or nonfatal stroke), and all-cause mortality death. These conditions corresponded to the International Classification of Diseases, 11th version (<http://www.who.int/classification/icd/en/>). Definitions for nonfatal myocardial infarction and nonfatal stroke are given elsewhere [11]. Patients were referred to cardiologists, neurosurgeons, neurologists or else to confirm diagnoses. Only the first event of the relevant outcome type was included in each analysis and the last day of the observation period was also noted if there were no incidences.

Statistical analysis

Data are expressed as mean \pm SD or median (interquartile range). Incidence rates of renal events, cardiovascular events, and all-cause death for different categories were calculated. Cox proportional hazards analysis was used to compute hazard ratios and 95 % confidence intervals (CI) to assess the impact of albuminuria and eGFR on the outcomes by using the group with eGFR ≥ 60 ml/min per 1.73 m² and/or the group with normoalbuminuria (<30 mg/g) as the reference [8]. In multivariate analysis, adjustment for risk factors for renal events, cardiovascular events, or all-cause mortality included age, gender, HbA1c, and systolic BP. A *p* value <5 % was considered significant. *p* values for trend tests examined whether UACR and eGFR levels were associated with increased hazard ratios. Trend tests across increasing risks for renal, cardiovascular

events and all-cause mortality are stratified by factors for diabetic nephropathy score.

All analyses were performed with the statistical software package SPSS (SPSS Japan, Tokyo, Japan).

Results

Baseline characteristics

Table 1 shows the baseline characteristics of patients examined in this study. The 4328 patients were distributed according to CKD stage and were followed until the onset of the first event or the end of the observation period (Table 1).

Incidence of numbers of patients of each event

During a median follow-up of 7.0 years (interquartile range 3.0–8.0 years), 419 renal events, 605 cardiovascular events and 236 deaths occurred, which were stratified by stages of renal function and levels of UACR with each event (Table 2). The incidence rates of each outcome per 1000 person-years were 19.8 for renal events, 23.3 for cardiovascular events and 8.4 for all-cause mortality. The incidence of each event increased with worsening of UACR levels and eGFR stages. Of importance, high incidence rates were noted in patients with macroalbuminuria plus reduced eGFR, especially for renal events (Table 2).

Risk for renal events, cardiovascular events, and all-cause mortality stratified by albuminuria and eGFR

Risks for renal events, cardiovascular events and all-cause mortality were evaluated by Cox proportional hazards analysis. The estimates were adjusted for age, gender, HbA1c, and systolic BP. The adjusted hazard ratios for renal events were 3.21 (95 % CI 2.31–4.47) for microalbuminuric patients and 21.86 (95 % CI 16.15–29.59) for macroalbuminuric patients as compared to normoalbuminuric patients as reference. Similarly, the adjusted hazard ratios for cardiovascular events and all-cause mortality were 1.38 (95 % CI 1.14–1.67) and 1.37 (95 % CI 0.99–1.89) for microalbuminuric patients and 2.05 (95 % CI 1.61–2.58) and 3.60 (95 % CI 2.53–5.20) for macroalbuminuric patients as compared to reference, respectively. Interestingly, UACR levels had the most significant impact on renal events. In addition to the effects of UACR levels, eGFR stages significantly increased the adjusted hazard ratios for renal events in patients with macroalbuminuria (Table 3). In Table 3, hazard ratios for cardiovascular events increased in patients with a higher UACR. In

Table 1 Baseline characteristics of participants ($n = 4328$)

Variable	All	Normoalbuminuria	Microalbuminuria	Macroalbuminuria	p For trend
N	4328	2679	1115	534	
Age (years; mean [SD])	60.2 (11.6)	59.5 (11.4)	61.9 (11.5)	59.8 (12.0)	<0.001
Male (n [%])	2546 (58.8)	1531 (57.1)	656 (58.8)	359 (67.2)	<0.001
Kidney factors					
UACR (mg/g; median [IQR])	18.2 (8.6–66.6)	10.2 (6.5–16.4)	66.6 (42.6–121.3)	994.8 (518.5–2272.5)	<0.001
eGFR (ml/min/1.73 m ² ; mean [SD])	77.0 (25.9)	81.3 (23.8)	76.4 (25.3)	56.9 (28.0)	<0.001
eGFR ≥ 90 (n [%])	1201 (27.7)	839	297	65	
eGFR 60–89 (n [%])	2051 (47.4)	1371	530	150	
eGFR 45–59 (n [%])	642 (14.8)	345	174	123	
eGFR 30–44 (n [%])	311 (7.2)	109	100	102	
eGFR 15–29 (n [%])	117 (2.7)	15	14	88	
eGFR <15 (n [%])	6 (0.1)	0	0	6	
BP (mmHg)					
Systolic BP (mean [SD])	131.0 (18.6)	127.2 (16.6)	134.2 (18.4)	143.0 (22.0)	<0.001
Diastolic BP (mean [SD])	74.3 (18.0)	73.3 (20.7)	74.8 (11.8)	78.1 (13.1)	<0.001
Other major risk factors					
HbA1c (%; mean [SD])	7.6 (1.7)	7.5 (1.7)	7.8 (1.7)	7.9 (1.8)	<0.001
Total cholesterol (mg/dL; mean [SD])	205.2 (35.9)	205.3 (34.1)	202.2 (33.9)	214.2 (50.2)	0.925
Body mass index (kg/m ² ; mean [SD])	25.3 (4.2)	25.1 (4.2)	25.5 (4.2)	25.4 (4.8)	0.098

Table 2 Number of patients and incidence rates of each outcome stratified by stages of eGFR and albuminuria

UACR	eGFR (ml/min/1.73 m ²)					
	>90	60–89	45–59	30–44	15–29	<15
Renal events (RRT or halving reduced eGFR)						
Normoalbuminuria	58 (4.2)		4 (2.3)	3 (6.9)	1 (21.3)	0
Microalbuminuria	31 (17.2)	41 (13.1)	15 (18.0)	10 (21.0)	1 (25.6)	0
Macroalbuminuria	20 (59.5)	62 (87.7)	56 (126.7)	54 (193.5)	61 (309.6)	2 (250.0)
Cardiovascular events						
Normoalbuminuria	229 (16.2)		40 (23.1)	14 (33.0)	1 (18.9)	0
Microalbuminuria	31 (16.1)	95 (28.7)	33 (32.6)	28 (56.1)	2 (43.5)	0
Macroalbuminuria	7 (17.2)	41 (44.6)	30 (44.3)	30 (64.2)	23 (57.9)	1 (100.0)
All-cause mortality						
Normoalbuminuria	70 (4.7)		26 (13.8)	4 (8.2)	4 (67.8)	0
Microalbuminuria	11 (5.4)	32 (8.9)	13 (11.9)	5 (8.5)	6 (117.6)	0
Macroalbuminuria	6 (14.4)	13 (12.3)	19 (24.6)	13 (23.6)	12 (26.8)	2 (142.9)

Number of patients (incidence rates per 1000 person-years)

RRT renal replacement therapy

addition, our results showed that there was a slight increase in the hazard ratios of cardiovascular events based on UACR levels plus co-existing reduced eGFR, especially in patients with microalbuminuria based on p for trend. In contrast, all-cause mortality was strongly associated with reduced eGFR <30 ml/min per 1.73 m², and the presence of macroalbuminuria even with preserved eGFR. The present study also revealed that normoalbuminuric renal insufficient diabetic patients did not have relatively poor

outcomes for renal events. Table 4 highlights the impact of low GFR and/or UACR on three distinct outcomes.

The clinical significance of diabetic nephropathy score in predicting the prognoses of renal events, cardiovascular events, and all-cause mortality

Considering the results of univariable and multivariable analyses, weighted arbitrary scores were allocated to each

Table 3 Hazard ratios based on CKD stages for each outcome

UACR	eGFR (ml/min/1.73 m ²)						<i>p</i> for trend (eGFR)
	>90	60–89	45–59	30–44	15–29	<15	
Renal events (RRT or halving reduced eGFR)							
Normoalbuminuria	1.00 (Reference)	1.00 (Reference)	0.69 (0.24–1.98)	1.83 (0.53–6.31)	11.59 (1.43–93.78)	NA	0.85
Microalbuminuria	3.31 (2.07–5.28)	3.04 (1.98–4.68)	3.36 (1.63–6.93)	3.10 (1.41–6.83)	3.60 (0.42–31.28)	NA	0.60
Macroalbuminuria	11.14 (5.87–21.17)	15.64 (10.30–23.74)	33.37 (20.58–50.91)	41.36 (25.09–68.16)	71.58 (40.41–126.80)	NA	<0.01
<i>p</i> for trend (albuminuria)	<0.01	<0.01	<0.01	<0.01	0.06	NA	
Cardiovascular events							
Normoalbuminuria	1.00 (Reference)	1.00 (Reference)	1.05 (0.73–1.49)	1.30 (0.74–2.28)	0.42 (0.06–3.06)	NA	0.46
Microalbuminuria	1.01 (0.69–1.49)	1.48 (1.15–1.90)	1.33 (0.89–2.00)	1.85 (1.20–2.85)	0.47 (0.11–1.97)	NA	0.04
Macroalbuminuria	1.28 (0.56–2.94)	2.10 (1.46–3.02)	1.85 (1.23–2.78)	2.37 (1.55–3.63)	2.09 (1.26–3.45)	12.76 (0.95–171.19)	0.20
<i>p</i> for trend (albuminuria)	0.81	<0.01	0.09	0.45	0.17	NA	
All-cause mortality							
Normoalbuminuria	1.00 (Reference)	1.00 (Reference)	1.67 (1.02–2.74)	1.22 (0.43–3.46)	8.19 (2.65–25.34)	NA	<0.01
Microalbuminuria	1.51 (0.78–2.95)	1.44 (0.92–2.24)	1.22 (0.63–2.35)	0.84 (0.31–2.26)	8.36 (2.81–24.90)	NA	0.04
Macroalbuminuria	4.37 (1.70–11.24)	1.92 (0.97–3.79)	4.84 (2.72–8.62)	4.09 (2.00–8.34)	6.16 (2.80–13.56)	70.57 (3.65–1363.68)	0.06
<i>p</i> for trend (albuminuria)	0.01	0.01	0.01	0.02	0.80	NA	

The estimates are adjusted for age, gender, HbA1c, systolic BP
RRT renal replacement therapy, *NA* not available

Table 4 Hazard ratios based on levels of UACR and eGFR for each outcome

UACR	eGFR (ml/min/1.73 m ²)		
	>60	30–59	<30
Renal events (RRT or halving reduced eGFR)			
Normoalbuminuria	1.00 (Reference)		49.82 (29.9–83.0)
Microalbuminuria	3.26 (2.34–4.55)		
Macroalbuminuria	13.6 (9.3–20.0)	33.0 (22.7–48.2)	
Cardiovascular events			
Normoalbuminuria	1.00 (Reference)		1.54 (1.00–2.39)
Microalbuminuria	1.40 (1.16–1.69)		
Macroalbuminuria	1.90 (1.36–2.65)	2.09 (1.54–2.84)	
All-cause mortality			
Normoalbuminuria	1.00 (Reference)		7.08 (4.16–12.05)
Microalbuminuria	1.30 (0.93–1.81)		
Macroalbuminuria	2.34 (1.35–4.04)	4.59 (2.90–7.25)	

The estimates are adjusted for age, gender, HbA1c, systolic BP

selected variable on the basis of each odds ratio (OR), and we defined a summation of scores as a new risk scoring system as the diabetic nephropathy score. We evaluated the diabetic nephropathy score for predicting the prognoses of renal events, cardiovascular events, and all-cause mortality. Each prognostic factor has a score and the maximum score is 6—microalbuminuria = 1, macroalbuminuria = 2, eGFR <45 ml/min per 1.73 m² = 1, age ≥60 years = 1, systolic BP >130 mmHg = 1, and HbA1c (NGSP) ≥6.9 % = 1. This score put stress on amounts of UACR.

Importantly, this simple score well predicted the incidence rates per 1000 patient/year for each event (Table 5).

Discussion

In this study we examined the clinical impact of UACR as well as the evaluation of GFR on outcomes in diabetic patients. We now report that increased urinary albumin excretion was strongly associated with risks for renal

Table 5 Diabetic nephropathy score reflects diabetic outcomes

Score	Renal events (RRT or halving reduced eGFR)				Cardiovascular events				All-cause mortality			
	Number of patients	Number of incidents	Rate per 1000 patient-years	95 % CI	Number of patients	Number of incidents	Rate per 1000 patient-years	95 % CI	Number of patients	Number of incidents	Rate per 1000 patient-years	95 % CI
	0	204	0	0.0		204	2	1.4	0.2–5.1	204	0	0.0
1	902	18	3.1	1.9–5.0	954	58	9.7	7.4–12.6	954	13	2.1	1.1–3.6
2	1228	54	6.9	5.2–9.0	1,310	174	21.4	18.4–24.9	1310	71	8.1	6.3–10.3
3	952	96	16.6	13.4–20.0	1,017	161	26.2	22.3–30.6	1017	62	9.4	7.2–12.0
4	471	118	46.6	38.5–56.0	532	116	39.0	32.2–46.8	532	45	13.4	9.8–18.0
5	213	93	103.8	83.8–127.0	238	71	60.5	47.3–76.3	238	35	25.5	17.7–35.4
6	61	40	216.2	154.5–294.0	73	23	80.4	51.0–120.7	73	10	28.1	13.5–51.7
<i>p</i> for trend			<0.001				<0.001				<0.001	

events, cardiovascular events and deaths in Japanese patients with type 2 diabetes. Of note, eGFR stages significantly increased the adjusted hazard ratios for renal events, especially when co-existing with macroalbuminuria, while patients with normoalbuminuria had relatively low risks for renal events. All-cause mortality was strongly associated with reduced eGFR <30 ml/min per 1.73 m² and the presence of macroalbuminuria even with preserved eGFR. However, the association between normoalbuminuria and reduced eGFR showed relatively low risks for cardiovascular events in the cohort of the Japanese population with type 2 diabetes. These findings suggested that diabetic patients with macroalbuminuria and low GFR had risks for adverse outcomes, even though UACR levels and eGFR had distinct clinical impacts on each event, respectively. Finally, the diabetic nephropathy score based on our present study may be useful for predicting the prognoses of outcomes in diabetic patients.

The present study has clearly shown that renal insufficiency plus the presence of macroalbuminuria accelerated risks for adverse outcomes, especially renal events. Recently, KDIGO reported the definition, classification and prognosis of CKD based both on estimated GFR and urinary levels of albumin excretion, emphasizing that a decrease in GFR as well as macroalbuminuria is important for renal outcomes of CKD [8]. In addition, the Action in Diabetes and Vascular disease: preterAx and diamicroN-MR Controlled Evaluation (ADVANCE) study reported that reduced eGFR with macroalbuminuria was associated with a higher risk for renal events [6]. Interestingly, the Casale Monferrato study revealed that macroalbuminuria was the main predictor of mortality, independently of both eGFR and cardiovascular risk factors [12]. In contrast, reduced eGFR did not increase the adjusted hazard ratio for renal events even in patients with microalbuminuria. This may be partly because the number of microalbuminuric patients with reduced GFR having renal events was relatively small as shown in Table 2. Collectively, these findings suggest that the assessment of macroalbuminuria as well as levels of eGFR may enable us to predict high risk for renal events.

The association between UACR and reduced eGFR showed relatively low risks for cardiovascular events, even though the incidence rate of cardiovascular events was 23.3, which was almost comparable to that observed in the Japan Diabetes Complications Study (JDACS) [13]. Our results also demonstrated that UACR was closely associated with cardiovascular events in patients with eGFR 60–89 ml/min per 1.73 m² and that reduced eGFR was important in microalbuminuric patients based on *p* values for trend. Of note, Yokoyama et al. recently reported that the risk for cardiovascular events was associated with progression of UACR stage in type 2 Japanese diabetic

patients [14]. In contrast, reduced eGFR was a high risk for developing cardiovascular endpoints (cardiovascular death, new admissions due to angina, myocardial infarction, stroke, revascularization or heart failure) and all-cause mortality independent of UACR [15]. Interestingly, the Second Nord-Trøndelag Health (HUNT II) study [16] reported that reduced eGFR with higher UACR was associated with a higher risk for cardiovascular events. This discrepancy compared to our present study may be partly because the number of patients with cardiovascular events in the present study was relatively small as shown in Table 2. Further studies will be required to examine this discrepancy.

This study also revealed that normoalbuminuric renal insufficient diabetic patients did not have relatively poor renal outcomes. In fact, the percentage of diabetic patients with normoalbuminuria and low eGFR is supposed to be relatively common in clinical settings. In this aspect, Yokoyama et al. [17] described that the proportion of subjects with low eGFR (<60 ml/min per 1.73 m²) and normoalbuminuria was 11.4 % of type 2 diabetic patients examined (262/2,298). Supporting our notion, Rigalleau et al. [18] reported that risk for renal progression in such patients with type 1 or type 2 diabetes is lower. On the contrary, all-cause mortality, not cardiovascular events, was strongly associated with reduced eGFR <30 ml/min per 1.73 m² in normoalbuminuric diabetic patients in this present study. Supporting this notion, hazard ratios for all-cause mortality as well as cardiovascular mortality increased in normoalbuminuric diabetic patients with low GFR [19]. The FIELD study also revealed that normoalbuminuric patients with eGFR 30–59 ml/min per 1.73 m² had a higher risk of cardiovascular events, cardiovascular death, non-coronary heart disease deaths, death from any cause than normoalbuminuric patients with eGFR ≥ 60 ml/min per 1.73 m² [7]. Interestingly, in the ADVANCE study, patients with normoalbuminuria and eGFR <60 ml/min/ 1.73 m² had a 3.95-fold higher risk for renal events, a 1.33-fold higher risk for cardiovascular events and a 1.85-fold higher risk for cardiovascular death [6]. In contrast, Vlek et al. [20] reported that eGFR <60 ml/min per 1.73 m² without UACR mainly influenced the risk of vascular events (hazard ratio 1.50; 1.05–2.15), but did not affect all-cause mortality. Furthermore, in type 2 diabetic patients, eGFR provided no further information for all-cause mortality and cardiovascular mortality in normoalbuminuric patients [14]. Therefore, further studies are needed to determine renal outcomes as well as all-cause mortality in normoalbuminuric diabetic patients with low eGFR.

We proposed a novel diabetic nephropathy score to predict incidence rates per 1000 patient/year for each event. To date, few studies have addressed individual prognostic factors/scores to predict outcomes of diabetic

complications in clinical settings. Couchoud et al. [21] reported development and validation of a prognostic score for 6-month mortality in elderly patients starting dialysis for ESKD. Nine risk factors were selected and points assigned for the score were body mass index <18.5 kg/m² (2 points), diabetes (1 point), congestive heart failure stages III to IV (2 points), peripheral vascular disease stages III to IV (2 points), dysrhythmia (1 point), active malignancy (1 point), severe behavioral disorder (2 points), total dependency for transfers (3 points) and unplanned dialysis (2 points). These scores effectively predict short-term prognosis among elderly patients, in which approximately 20 % of the patients had diabetic nephropathy. In contrast to this previous study, our simple prognostic scoring system may clearly predict cardiovascular events and all-cause mortality as well as renal events for patients of any age. Even though validation of this score system will be required for other cohorts, this system seems simple and useful for predicting clinical aspects.

To date, UACR levels and reduced eGFR have independently been reported to predict cardiovascular and renal outcomes in diabetes [6]. Previously, diabetic patients with microalbuminuria/macroalbuminuria had a risk for adverse outcomes, including cardiovascular events, cardiovascular death, and renal events as reported by the ADVANCE study [6]. Importantly, the present study, consisting of 4328 Japanese patients with type 2 diabetes, was critically different from the ADVANCE study in terms of (1) being a historical cohort study consisting of 10 centers, (2) longer observation period (median 7.0 years), (3) including the assessment of all-cause mortality, (4) including assessment of each event based on the new classification CKD stages, and (5) providing a diabetic nephropathy score to predict the prognoses of renal events, cardiovascular events, and all-cause mortality. Therefore, our present study further revealed the clinical significance of UACR and eGFR on adverse outcomes in diabetic patients.

There are several limitations to this study. First, the lack of histologically proven diabetic nephropathy should be discussed, even though diabetic nephropathy is clinically diagnosed by the presence of microalbuminuria. Second, the low incidence of cardiovascular events may result in a relatively weak statistical power. Furthermore, the lack of data regarding whether enrolled patients have predisposing cardiovascular diseases must be considered. However, this multicenter observational study of 4328 diabetic patients over 7 years may strengthen the present results and increase the accuracy of risk estimation and establishment of a prognostic diabetic nephropathy score.

In conclusion, these results conclude that the presence of microalbuminuria/macroalbuminuria is closely related to the increase in risks for adverse outcomes in Japanese diabetic patients, whereas the association between

macroalbuminuria and reduced eGFR was a strong predictor for renal events. Further studies will be required to validate the prognostic factors and related diabetic nephropathy score by using other cohorts together with future perspectives.

Acknowledgments The authors thank Dr. Yukinari Yamaguchi (Nara Medical University), Dr. Mitsuhiro Yoshimura (Noto General Hospital), and Miyuki Murakami (Kanazawa University) for supporting this study. This study was supported by Grant-in-Aids for Diabetic Nephropathy Research and for Diabetic Nephropathy and Nephrosclerosis Research, from the Ministry of Health, Labour and Welfare of Japan and by the Ministry of Education, Science, Sports and Culture, Japan.

Conflict of interest H. Makino is a consultant for AbbVie, Astellas and Teijin, receives speaker honoraria from Astellas, MSD, Takeda, and Tanabe Mitsubishi, and receives grant support from Astellas, Daiichi Sankyo, Dainippon Sumitomo, MSD, Novo Nordisk and Takeda.

References

1. Parving HH, Mauer M, Fioretto P, Rossing P, Ritz E. Diabetic nephropathy. In: Taal MW, Chertow GM, Marsden PA, Skorecki K, Yu ASL, Brenner BM, editors. *The kidney*. Philadelphia: Elsevier Saunders; 2012. p. 1411–54.
2. Nakayama M, Sato T, Sato H, Yamaguchi Y, Obara K, Kurihara I, et al. Different clinical outcomes for cardiovascular events and mortality in chronic kidney disease according to underlying renal disease: the Gonryo study. *Clin Exp Nephrol*. 2010;14:333–9.
3. Foley RN, Culeton BF, Parfrey PS, Harnett JD, Kent GM, Murray DC, et al. Cardiac diseases in diabetic end-stage renal disease. *Diabetologia*. 1997;40:1307–12.
4. Wada T, Shimizu M, Toyama T, Hara A, Kaneko S, Furuichi K. Clinical impact of albuminuria in diabetic nephropathy. *Clin Exp Nephrol*. 2012;16:96–101.
5. Berhane AM, Weil EJ, Knowler WC, Nelson RG, Hanson RL. Albuminuria and estimated glomerular filtration rate as predictors of diabetic end-stage renal disease and death. *Clin J Am Soc Nephrol*. 2011;6:2444–51.
6. Ninomiya T, Perkovic V, de Galan BE, Zoungas S, Pillai A, Jardine M, et al. Albuminuria and kidney function independently predict cardiovascular and renal outcomes in diabetes. *J Am Soc Nephrol*. 2009;20:1813–21.
7. Drury PL, Ting R, Zannino D, Ehnholm C, Flack J, Whiting M, et al. Estimated glomerular filtration rate and albuminuria are independent predictors of cardiovascular events and death in type 2 diabetes mellitus: the Fenofibrate Intervention and Event Lowering in Diabetes (FIELD) study. *Diabetologia*. 2011;54:32–43.
8. Levey AS, de Jong PE, Coresh J, El Nahas M, Astor BC, Matsushita K, et al. The definition, classification and prognosis of chronic kidney disease: a KDIGO Controversies Conference report. *Kidney Int*. 2011;80:17–28.
9. Report of the Committee on the classification and Diagnostic Criteria of Diabetes Mellitus: The committee of the Japan Diabetes Society on the Diagnostic Criteria of Diabetes Mellitus. *J Diabetes Invest*. 2010;1:212–28.
10. Matsuo S, Imai E, Horio M, Yasuda Y, Tomita K, Nitta K, et al. Revised equations for estimated GFR from serum creatinine in Japan. *Am J Kidney Dis*. 2009;53:982–92.
11. Yokoyama H, Matsushima M, Kawai K, Hirao K, Oishi M, Sugimoto H, et al. Low incidence of cardiovascular events in Japanese patients with type 2 diabetes in primary care settings; a prospective cohort study. *Diabetic Med*. 2011;28:1221–8.
12. Bruno G, Merletti F, Bargero G, Novelli G, Melis D, Soddu A, et al. Estimated glomerular filtration rate, albuminuria and mortality in type 2 diabetes: the Casale Monferrato study. *Diabetologia*. 2007;50:941–8.
13. Sone H, Tanaka S, Tanaka S, Iimuro S, Oida K, Yamasaki Y, et al. Serum level of triglycerides is a potent risk factor comparable to LDL cholesterol for coronary heart disease in Japanese patients with type 2 diabetes: subanalysis of the Japan Diabetes Complications Study (JDCS). *J Clin Endocrinol Metab*. 2011;96:3448–56.
14. Yokoyama H, Araki S, Haneda M, Matsushima M, Kawai K, Hirao K, et al. Chronic kidney disease categories and renal-cardiovascular outcomes in type 2 diabetes without prevalent cardiovascular disease: a prospective cohort study (JDDM 25). *Diabetologia*. 2012;55:1911–8.
15. So WY, Kong AP, Ma RC, Ozaki R, Szeto CC, Chan NN, et al. Glomerular filtration rate, cardiorenal end points, and all-cause mortality in type 2 diabetic patients. *Diabetes Care*. 2006;29:2046–52.
16. Hallan S, Astor B, Romundstad S, Assarød K, Kvenild K, Coresh J. Association of kidney function and albuminuria with cardiovascular mortality in older vs. younger individuals: The HUNT II Study. *Arch Intern Med*. 2007;167:2490–6.
17. Yokoyama H, Kawai K, Kobayashi M, Japan Diabetes Clinical Data Management Study Group. Microalbuminuria is common in Japanese type 2 diabetic patients: a nationwide survey from the Japan Diabetes Clinical Data Management Study Group (JDDM 10). *Diabetes Care*. 2007;30:989–92.
18. Rigalleau V, Lasseur C, Raffaitin C, Beauvieux MC, Barthe N, Chauveau P, et al. Normoalbuminuric renal-insufficient diabetic patients: a lower-risk group. *Diabetes Care*. 2007;30:2034–9.
19. Fox CS, Matsushita K, Woodward M, Biló HJ, Chalmers J, Heerspink HJ, et al. Association of kidney disease measures with mortality and end-stage renal disease in individuals with and without diabetes: a meta-analysis. *Lancet*. 2013;381:374.
20. Vlek AL, van der Graaf Y, Spiering W, Algra A, Visseren FL, SMART study group. Cardiovascular events and all-cause mortality by albuminuria and decreased glomerular filtration rate in patients with vascular disease. *J Intern Med*. 2008;264:351–60.
21. Couchoud C, Labeeuw M, Moranne O, Allot V, Esnault V, Frimat L, et al. A clinical score to predict 6-month prognosis in elderly patients starting dialysis for end-stage renal disease. *Nephrol Dial Transplant*. 2009;24:1553–61.

Kidney lesions in diabetic patients with normoalbuminuric renal insufficiency

Miho Shimizu · Kengo Furuichi · Hitoshi Yokoyama ·
Tadashi Toyama · Yasunori Iwata ·
Norihiko Sakai · Shuichi Kaneko · Takashi Wada

Received: 17 July 2013 / Accepted: 9 September 2013 / Published online: 1 October 2013
© Japanese Society of Nephrology 2013

Abstract Diabetic nephropathy is a leading cause of end-stage renal disease in Japan. Microalbuminuria has been considered as the first clinical sign of diabetic nephropathy. However, recent studies demonstrated that *normoalbuminuric renal insufficiency is not uncommon* for diabetic patients, especially in type 2 diabetes. Although the pathogenesis of normoalbuminuric renal insufficiency in diabetic nephropathy remains to be fully elucidated, distinct clinical and pathological features of diabetic patients with this finding have been reported as compared to those in diabetic patients with a typical clinical course. In type 1 diabetes, more advanced glomerular lesions were found in patients with normoalbuminuric renal insufficiency than in patients with normoalbuminuric preserved renal function. In contrast,

disproportionately advanced tubulointerstitial and vascular lesions, despite minor diabetic glomerular lesions, which denote the presence of diabetic kidney lesions as well as nephrosclerosis, were likely to be related to the development of *normoalbuminuric renal insufficiency* in some type 2 diabetic patients. In addition, long-term outcomes of diabetic patients with normoalbuminuric renal insufficiency remain controversial. Further studies to gain a better understanding of the structural–functional relationships and natural history of diabetic patients with normoalbuminuric renal insufficiency may improve the benefits of therapeutic interventions for diabetic nephropathy.

Keywords Diabetic nephropathy · Normoalbuminuric renal insufficiency · Kidney lesions · Nephrosclerosis · Comprehensive medicine · Humans

Presented at the 42nd Eastern Regional Meeting of the Japanese Society of Nephrology.

M. Shimizu · K. Furuichi · T. Toyama · Y. Iwata · N. Sakai ·
T. Wada
Division of Nephrology, Kanazawa University Hospital,
Kanazawa, Japan

M. Shimizu · K. Furuichi · T. Toyama · Y. Iwata · N. Sakai ·
S. Kaneko
Department of Disease Control and Homeostasis, Kanazawa
University Graduate School of Medical Sciences, Kanazawa,
Japan

H. Yokoyama
Division of Nephrology, Kanazawa Medical University,
Uchinada, Japan

T. Wada (✉)
Division of Nephrology, Department of Laboratory Medicine,
Kanazawa University Graduate School of Medical Sciences,
13-1 Takara-machi, Kanazawa, Ishikawa 920-8641, Japan
e-mail: twada@m-kanazawa.jp

Introduction

Diabetic nephropathy is a leading cause of end-stage renal disease in Japan [1]. The percentage of patients with diabetic nephropathy was 44.1 % among new dialysis patients, and 37.1 % among all dialysis patients at the end of 2012 based on the latest annual report of the Japanese Society for Dialysis Therapy.

Albuminuria and glomerular filtration rate (GFR) are recommended for use as clinical markers of diabetic nephropathy, and microalbuminuria has been considered as the first clinical sign of diabetic nephropathy [2–5]. However, recent studies demonstrated that reduction in GFR may precede the development of microalbuminuria in some diabetic patients [6–14].

On the other hand, pathological markers of diabetic nephropathy are complicated because a variety of renal lesions can be found in diabetic nephropathy. In addition, factors such as obesity, hypertension, dyslipidemia, and aging, which are frequent complications in type 2 diabetes, cause a wide variety of pathological changes [15–17]. Currently, renal biopsy is not always applicable for diabetic patients with a typical clinical course. The latest committee report by the Japan Renal Biopsy Registry (J-RBR) and the Japan Kidney Disease Registry (J-KDR) indicated that the percentage of diabetic nephropathy was 5.1 % (376/7442) in the pathological diagnoses as classified by pathogenesis in J-RBR 2009 and 2010 [18]. Therefore, the structural–functional relationships of diabetic nephropathy remain to be fully investigated.

Previously, Takazakura et al. [19] reported the clinical factors related to the development and progression of renal lesions in diabetic nephropathy by evaluation of serial renal biopsies or autopsy. In this study, significant relationships were found between progression of diabetic glomerulosclerosis and clinical factors such as the control of blood glucose, type of diabetes, age at onset, type of treatment, and degree of obesity. Subsequently, we conducted a long-term retrospective study to evaluate the structural–functional relationships and prognostic impacts of clinicopathological parameters for renal events, cardiovascular events, and all-cause mortality among 260 Japanese type 2 diabetic patients with biopsy-proven diabetic nephropathy [20]. Our study suggested that the characteristic pathological lesions and macroalbuminuria (severe proteinuria) were closely related to the long-term outcomes of diabetic nephropathy in type 2 diabetes. Based on our results, it is reasonable to predict renal prognosis of diabetic nephropathy by a combination of clinical and pathological parameters. Furthermore, we speculate that the evaluation of renal pathology provides a key note to have deeper insights for ‘Comprehensive Medicine in Humans’ including renal events and cardiovascular events in patients with diabetic nephropathy.

Here, we focus on the clinical characteristics, renal lesions, and outcomes of diabetic nephropathy with normoalbuminuric renal insufficiency, and describe future perspectives for clinical research on diabetic nephropathy.

Prevalence of normoalbuminuric renal insufficiency in diabetes

The prevalence of normoalbuminuria among type 1 diabetic patients with low GFR (<60 ml/min/ 1.73 m²) was 23.6 % (21/89) in the Diabetes Control and Complications Trial (DCCT) and Epidemiology of Diabetes Interventions

and Complications (EDIC) study conducted in 1982–2006 [6].

In contrast, the prevalence of normoalbuminuria among type 2 diabetic patients with low GFR (<60 ml/min/ 1.73 m²) was 32 % in the Developing Education on Microalbuminuria for Awareness of Renal and Cardiovascular Risk in Diabetes (DEMAND) study conducted in 2002–2005 [7], 35.1 % (60/171) in the Third National Health and Nutrition Examination Survey (NHANES III) conducted in 1988–1994 [8], 50.8 % (575/1132) in the United Kingdom Prospective Diabetes Study (UKPDS)-74 study conducted in 1977–1991 [9], 51.8 % (262/506) in the Japan Diabetes Clinical Data Management (JDDM) study conducted in 2004–2005 [10], 55.0 % (506/920) in the National Evaluation of the Frequency of Renal Impairment co-existing with Non-insulin-dependent diabetes (NEFRON) study conducted in 2005 [11], 56.5 % (1673/2959) in the Renal Insufficiency And Cardiovascular Events (RIACE) study conducted in 2007–2008 [12], 61.6 % (1252/2033) in the Action in Diabetes and Vascular disease: preterAx and diamicroN-MR Controlled Evaluation (ADVANCE) study conducted in 2001–2003 [13], and 71.3 % (290/407) in the Swedish National Diabetes Register conducted in 2002–2007 [14]. These results suggest that normoalbuminuric renal insufficiency is not uncommon among diabetic patients, especially in type 2 diabetes.

Pathogenesis of normoalbuminuric renal insufficiency in diabetes

There are several possible pathogenic mechanisms that may account for the development of normoalbuminuric renal insufficiency in diabetes. One possibility is that renal ischemia due to intrarenal arteriosclerosis may be related to the development of normoalbuminuric renal insufficiency. A negative correlation between GFR and the intrarenal arterial resistance index was found in type 2 diabetes, regardless of albuminuria stage [21, 22]. In addition, carotid intimal medial thickness, carotid stiffness, and silent cerebral infarction were also reported to be associated with impaired kidney function in type 2 diabetes, independent of microalbuminuria [21, 23].

The other possibility is that genetic susceptibility may contribute to the development of normoalbuminuric renal insufficiency. Polymorphisms of the protein kinase C- β gene were reported to be associated with accelerated decline of estimated GFR (eGFR) in type 2 diabetes without overt proteinuria [24].

Although the increasing use of renin–angiotensin system (RAS) blockade may be related to the increasing prevalence of normoalbuminuric renal insufficiency, the RIACE study showed that the use of RAS blockade was

more common in patients with albuminuric renal insufficiency than in those with normoalbuminuric renal insufficiency [12].

Clinical characteristics associated with normoalbuminuric renal insufficiency in diabetes

The reported clinical characteristics of diabetic patients with normoalbuminuric renal insufficiency include a higher proportion of females, a shorter duration of diabetes, lower prevalence of hypertension, smoking, retinopathy, neuropathy, previous cardiovascular disease, and antihypertensive agents including RAS blockade, lower levels of hemoglobin A1c and triglycerides, and higher levels of hemoglobin and high-density lipoprotein cholesterol, as compared to patients with albuminuric renal insufficiency [10, 12, 25]. In addition, compared to patients with normoalbuminuric preserved renal function, those with normoalbuminuric renal insufficiency are older, more frequently females and non-smokers, and have a higher prevalence of hypertension, dyslipidemia, metabolic syndrome, and previous cardiovascular disease, and higher levels of homeostasis model assessment of insulin resistance [10, 26].

Although our study of 260 type 2 diabetic patients with biopsy-proven diabetic nephropathy (96 females, 106 males; age 58.2 ± 11.4 years) included negative proteinuria as well as trace proteinuria in the normoalbuminuria (normal proteinuria) category based on the new classification of chronic kidney disease [3, 4], 154 (59.2 %) of the 260 patients showed low eGFR (<60 ml/min/1.73 m²), and 15 (9.7 %) of the 154 patients with low eGFR were not associated with albuminuria (proteinuria) [20]. Table 1 presents the baseline clinical and pathological findings of fifteen patients with normoalbuminuria (normal proteinuria) and low eGFR [7 males, 8 females; an average age 62.5 years (range 49–72 years)] at the time of renal biopsy.

The mean eGFR was 46.0 ml/min/1.73 m² (range 25.6–57.2 ml/min/1.73 m²), the mean duration of diabetes was 7.5 years (range 0–21 years), and the mean hemoglobin A1c level (NGSP) was 7.8 % (range 4.5–11.9 %). The prevalence of diabetic retinopathy, hypertension, dyslipidemia, and history of cardiovascular disease were 50.0 % (6/12), 35.7 % (5/14), 27.3 % (3/11), and 14.3 % (2/14), respectively. Our study demonstrated lower prevalence of hematuria and retinopathy, shorter duration of diabetes, lower systolic blood pressure, and higher hemoglobin level in patients with normoalbuminuria (normal proteinuria) and low eGFR as compared to patients with micro/macroulbuminuria (mild/severe proteinuria) and low eGFR. In addition, when compared to patients with normoalbuminuria (normal proteinuria) and preserved eGFR,

those with normoalbuminuria (normal proteinuria) and low eGFR were older. Our findings were consistent with previous reports, and provided additional information on the prevalence of hematuria. Furthermore, our study showed that aging was associated with low eGFR regardless of albuminuria (proteinuria) category, and that hematuria, diabetic retinopathy, and low hemoglobin were associated with the progression of albuminuria (proteinuria) regardless of eGFR category [20].

Pathological characteristics associated with normoalbuminuric renal insufficiency in diabetes

There have been few studies regarding the pathological characteristics of diabetic nephropathy with normoalbuminuric renal insufficiency. A study of 8 long-standing type 1 diabetic women with normoalbuminuria and low GFR (75 ± 10 ml/min/1.73 m²), as measured by creatinine clearance, found that these patients had more advanced glomerular lesions, such as higher volume fraction of mesangium, greater index of arteriolar hyalinosis, and higher percentage of global glomerular sclerosis than 19 normoalbuminuric women with preserved GFR (115 ± 15 ml/min/1.73 m²) [27]. Similarly, another study indicated that 23 long-standing type 1 diabetic patients with normoalbuminuria and low GFR (<90 ml/min/1.73 m²), as measured by either iothalamate clearance or creatinine clearance, had more advanced glomerular lesions, such as more increased glomerular basement membrane width and greater fractional volume of glomerulus occupied by mesangium, when compared with 82 normoalbuminuric patients with preserved GFR (≥ 90 ml/min/1.73 m²) [28]. In addition, the Cohen diabetic rat, an animal model of type 2 diabetes which shows progressive depression of renal function without proteinuria or hypertension, was reported to show classical diabetic glomerulosclerosis, including mesangial matrix expansion, thickened basement membranes, and increased deposition of type IV collagen [29].

Our study of 260 type 2 diabetic patients also demonstrated that diffuse lesions, nodular lesions, tubulointerstitial lesions, and vascular lesions in patients with normoalbuminuria (normal proteinuria) and low eGFR (<60 ml/min/1.73 m²) were more advanced compared to those in patients with normoalbuminuria (normal proteinuria) and preserved eGFR (≥ 60 ml/min/1.73 m²) in type 2 diabetes [20].

Fioretto et al. [15, 16] proposed a classification system including three major categories based on renal biopsy lesions in type 2 diabetic patients with microalbuminuria and proteinuria—category I with normal or near-normal renal structure (35 % of patients with microalbuminuria

Table 1 Clinicopathological findings at the time of renal biopsy and outcomes in type 2 diabetic patients with normoalbuminuria (normal proteinuria) and low eGFR <60 ml/min/1.73 m²

Case no.	Gender	Clinical and pathological findings at the time of renal biopsy								Outcomes				
		Age (years)	eGFR (ml/min/1.73 m ²)	Diabetes duration (years)	Hemoglobin A1c (%) (NGSP)	Diabetic retinopathy	Hypertension	Dyslipidemia	History of CVD	Category of renal lesions	Follow-up (years)	Renal events	CVD events	All-cause mortality
1	Male	69	25.6	2	n.d. ^a	n.d.	n.d.	n.d.	n.d.	III	n.d.	n.d.	n.d.	n.d.
2	Female	71	31.7	4	4.5	(-)	(-)	(-)	(-)	II	4.6	(-)	(-)	(-)
3	Male	56	32.1	15	7.1	n.d.	(-)	(-)	(-)	II	16.4	(-)	(-)	(-)
4	Female	61	36.1	1	6.5	(-)	(+)	(-)	(-)	III	5.1	(-)	(-)	(-)
5	Female	64	39.2	21	8.0	(-)	(+)	n.d.	(-)	II	14.5	(+)	(-)	(-)
6	Female	53	41.3	6	7.5	(+)	(+)	(-)	(+)	III	7.0	(-)	(-)	(-)
7	Female	61	49.4	10	6.5	(-)	(-)	(+)	(+)	III	21.8	(-)	(+)	(-)
8	Male	72	51.2	17	11.9	(+)	(-)	(-)	(-)	III	3.6	(-)	(-)	(-)
9	Male	65	52.7	5	5.9	(-)	(-)	(-)	(-)	III	5.3	(-)	(-)	(+)
10	Male	64	53.0	2	7.0	(-)	(-)	(-)	(-)	III	5.6	(-)	(-)	(-)
11	Male	61	53.7	0 ^b	6.0	(-)	(-)	n.d.	(-)	III	13.7	(-)	(-)	(-)
12	Female	64	55.5	10	11.8	(+)	(+)	(-)	(-)	III	2.7	(-)	(+)	(-)
13	Female	63	55.7	10	9.9	(+)	(-)	(+)	(-)	II	5.3	(-)	(-)	(-)
14	Female	65	56.0	1	8.0	(+)	(+)	(+)	(-)	II	2.1	(-)	(-)	(-)
15	Male	49	57.2	8	8.3	(+)	(-)	n.d.	(-)	II	13.4	(-)	(+)	(-)

Presence of diabetic retinopathy, hypertension, dyslipidemia, and history of CVD are indicated by (+). Category of renal lesions was evaluated according to the classification proposed by Fioretto et al. [15, 16]

eGFR estimated glomerular filtration rate, CVD cardiovascular disease, n.d. no data, NGSP National Glycohemoglobin Standardization Program

^a This patient had poorly controlled type 2 diabetes, which was followed at another hospital

^b This patient was diagnosed for the first time with type 2 diabetes by a random blood glucose test and an oral glucose tolerance test at the time of renal biopsy




Fig. 1 Representative microscopic findings in type 2 diabetic patients with normoalbuminuria (normal proteinuria) and low eGFR (<60 ml/min/1.73 m²). **a** Severe diffuse lesions in a patient classified as typical lesions of diabetic nephropathy (category II) (Case no. 3, Table 1; periodic acid-Schiff (PAS) stain $\times 200$). **b** Mild diffuse

lesions (*asterisk*) associated with global glomerular sclerosis (*arrow*) and disproportionately advanced arteriosclerosis, which denotes the presence of diabetic kidney lesions as well as nephrosclerosis, in a patient classified as atypical patterns (category III) (Case no.12, Table 1; PAS stain $\times 100$)

and 10 % of patients with proteinuria); category II with typical diabetic lesions with an approximately balanced severity of glomerular, tubulointerstitial, and arteriolar changes similar to type 1 diabetes (30 % of patients with microalbuminuria and 55 % of patients with proteinuria); and category III with disproportionately advanced tubulointerstitial lesions, vascular lesions, and global glomerulosclerosis, despite minor diabetic glomerulopathy (35 % of patients with microalbuminuria and 35 % of patients with proteinuria). We categorized the renal lesions of 15 patients with normoalbuminuria (normal proteinuria) and low eGFR according to this classification (Table 1). As a result, no patients were classified as having almost normal biopsies (category I), 6 patients (40 %) were classified as having typical lesions of diabetic nephropathy (category II), and 9 patients (60 %) were classified as having atypical patterns (category III). Representative microscopic findings of renal biopsies from type 2 diabetic patients with normoalbuminuria (normal proteinuria) and low eGFR are shown in Fig. 1. Figure 1a shows severe diffuse lesions in a patient classified as category II (Case no. 3, Table 1). Figure 1b shows mild diffuse lesions associated with global glomerular sclerosis and disproportionately advanced arteriosclerosis, which denotes the presence of diabetic kidney lesions as well as nephrosclerosis, in a patient classified as category III (Case no. 12, Table 1). Another study of type 2 diabetic patients with low eGFR (<60 ml/min/1.73 m²) by research biopsy also indicated that the typical glomerular changes of diabetic nephropathy were less common in normoalbuminuric (3/8) patients than in microalbuminuric (5/6) or macroalbuminuric (17/17) patients [30].

These results suggest that multifactorial pathogenesis in addition to diabetic conditions may contribute to the kidney lesions in type 2 diabetic patients with normoalbuminuric renal insufficiency.

Outcomes of diabetic patients with normoalbuminuric renal insufficiency

Although the outcome of diabetic patients with normoalbuminuric renal insufficiency remains controversial, it is likely to be better than that of diabetic patients with albuminuria, even with preserved renal function. In a study of 89 diabetic patients with eGFR <60 ml/min/1.73 m² (22 type 1 diabetes, 67 type 2 diabetes), 15 (17 %), 36 (40 %), and 38 (43 %), were found to be normoalbuminuric, microalbuminuric, and macroalbuminuric at baseline, respectively. During the 38-month follow-up, none of the normoalbuminuric patients started dialysis (microalbuminuric patients 2/36, macroalbuminuric patients 10/38), and their albumin excretion rates and serum creatinine levels were stable. Furthermore, none of the normoalbuminuric patients died (microalbuminuric patients 3/36, macroalbuminuric patients 7/38) during the follow-up period [25]. In a population-based study of 1,538 people with type 2 diabetes, 51, 32, and 17 % were found to be normoalbuminuric, microalbuminuric, and macroalbuminuric at baseline, respectively. During the 11-year follow-up, increasing trends of hazard rate ratios for all-cause mortality and cardiovascular mortality by decreasing eGFR values were observed in macroalbuminuric people only [31]. In addition, our study also showed similar results [20]. In our study, the outcomes were defined as the first occurrence of renal events (requirement of dialysis, or a 50 % decline in eGFR from baseline), cardiovascular events (cardiovascular death, nonfatal myocardial infarction, coronary interventions, or nonfatal stroke), and all-cause mortality. The mean follow-up period was 8.1 years (range 5–9,739 days), and there were 118 renal events, 62 cardiovascular events, and 45 deaths during the follow-up period. The hazard ratios for renal events, cardiovascular events, and all-cause mortality were increased by




Fig. 2 Event-free rates of **a** renal events, **b** cardiovascular events, and **c** all-cause mortality in type 2 diabetic patients with normoalbuminuria (normal proteinuria) stratified by eGFR categories according to Kaplan–Meier method. Differences between groups were

compared by the log-rank test. *Solid line* normoalbuminuria (normal proteinuria) and eGFR ≥ 60 ml/min/ 1.73 m² group, *dotted line* normoalbuminuria (normal proteinuria) and eGFR < 60 ml/min/ 1.73 m² group

decreasing eGFR levels in patients with macroalbuminuria (severe proteinuria) only, and macroalbuminuria (severe proteinuria) was a major clinical determinant of renal events and all-cause mortality. Fifteen patients with normoalbuminuria (normal proteinuria) and low eGFR had 1 renal event, 3 cardiovascular events, and 1 death during the follow-up period (Table 1). One patient who developed end-stage renal disease was positive for anti-hepatitis C virus (HCV) antibody, although renal biopsy showed typical diabetic glomerular lesions, such as diffuse and exudative lesions, without HCV-related glomerulonephritis (Case no. 5, Table 1). Event-free rates of renal events, cardiovascular events, and all-cause mortality in patients with normoalbuminuria (normal proteinuria) were not significantly different between eGFR categories (Fig. 2).

In contrast to the previous results outlined above, including our report, several studies showed that higher levels of urinary albumin excretion within the normal range predict faster decline in GFR and higher incidence of cardiovascular disease in diabetic patients [32, 33]. Furthermore, some studies demonstrated that albuminuria and renal function independently or respectively predict renal events, cardiovascular events, and death in diabetic patients [13, 34–36]. Therefore, further studies on clinical impacts of low GFR without albuminuria and new biomarkers for early and definitive diagnosis of diabetic nephropathy are required in clinical settings.

In this sense, the newly established prospective registry system, the Japan Diabetic Nephropathy Cohort Study (JDNCs), may provide key insights for future perspectives [37]. The Japanese Society of Nephrology established a nationwide, web-based, and prospective registry system including two basic registries—the J-RBR and the J-KDR [18, 38]. In addition to the two basic registries, the JDNCs

enrolled Japanese diabetic patients with broad-ranging albuminuria (proteinuria) stage and GFR levels. The aims of the JDNCs are to obtain clinical data and urine samples for revising the clinical staging of diabetic nephropathy, and to develop new diagnostic biomarkers for early detection or prediction of diabetic nephropathy. The prevalence of normoalbuminuria in patients with low eGFR (< 60 ml/min/ 1.73 m²) at baseline was 19.8 % in the JDNCs. Interestingly, the JDNCs is now prospectively collecting clinical data annually. Moreover, participants with diabetes are also enrolled in the J-RBR and the J-KDR [37]. Therefore, the combined data of the three registries will allow evaluation of the natural course and long-term outcomes of diabetic nephropathy, including patients with normoalbuminuric renal insufficiency, in the near future.

Conclusion

We focused on the structural–functional relationships and outcomes of diabetic nephropathy with normoalbuminuric renal insufficiency. Although the pathogenesis of normoalbuminuric renal insufficiency in diabetic nephropathy remains to be fully elucidated, disproportionately advanced tubulointerstitial lesions, vascular lesions, and global glomerulosclerosis, despite minor diabetic glomerular lesions, which denote the presence of diabetic kidney lesions as well as nephrosclerosis, are likely to be related to the development of normoalbuminuric renal insufficiency in some type 2 diabetic patients. However, other processes may contribute to the development of normoalbuminuric renal insufficiency with advanced diabetic glomerular lesions. Furthermore, long-term outcomes of diabetic patients with normoalbuminuric renal insufficiency remain

controversial. Further studies to gain a better understanding of the structural–functional relationships and natural history of diabetic patients with normoalbuminuric renal insufficiency may provide insights for the development of therapeutic strategies for diabetic nephropathy.

Acknowledgments This study was supported in part by a Grant-in-Aid for Diabetic Nephropathy Research, and for Diabetic Nephropathy and Nephrosclerosis Research from the Ministry of Health, Labour and Welfare of Japan. T.W. is also a recipient of a Grant-in-Aid from the Ministry of Education, Science, Sports and Culture in Japan.

Conflict of interest No potential conflicts of interest relevant to this article were reported.

References

- Nakai S, Iseki K, Itami N, Ogata S, Kazama JJ, Kimata N, et al. An overview of regular dialysis treatment in Japan (as of 31 December 2010). *Ther Apher Dial*. 2012;16:483–521.
- American Diabetes Association. Standards of medical care in diabetes—2013. *Diabetes Care*. 2013;36(Suppl. 1):S11–66.
- Levey AS, de Jong PE, Coresh J, et al. The definition, classification, and prognosis of chronic kidney disease: a KDIGO Controversies Conference report. *Kidney Int*. 2011;80:17–28.
- Japanese Society of Nephrology. Clinical practice guidebook for diagnosis and treatment of chronic kidney disease 2012. Tokyo:yoigakusha; 2012.
- Wada T, Shimizu M, Toyama T, Hara A, Kaneko S, Furuichi K. Clinical impact of albuminuria in diabetic nephropathy. *Clin Exp Nephrol*. 2012;16:96–101.
- Molitch ME, Steffes M, Sun W, Rutledge B, Cleary P, de Boer IH, et al. Development and progression of renal insufficiency with and without albuminuria in adults with type 1 diabetes in the diabetes control and complications trial and the epidemiology of diabetes interventions and complications study. *Diabetes Care*. 2010;33:1536–43.
- Dwyer JP, Parving HH, Hunsicker LG, Ravid M, Remuzzi G, Lewis JB. Renal dysfunction in the presence of normoalbuminuria in type 2 diabetes: results from the DEMAND Study. *Cardiorenal Med*. 2012;2:1–10.
- Kramer HJ, Nguyen QD, Curhan G, Hsu CY. Renal insufficiency in the absence of albuminuria and retinopathy among adults with type 2 diabetes mellitus. *JAMA*. 2003;289:3273–7.
- Retnakaran R, Cull CA, Thorne KI, Adler AI, Holman RR, UKPDS Study Group. Risk factors for renal dysfunction in type 2 diabetes: U.K. Prospective Diabetes Study 74. *Diabetes*. 2006;55:1832–9.
- Yokoyama H, Sone H, Oishi M, Kawai K, Fukumoto Y, Kobayashi M, et al. Prevalence of albuminuria and renal insufficiency and associated clinical factors in type 2 diabetes: the Japan Diabetes Clinical Data Management study (JDDM15). *Nephrol Dial Transplant*. 2009;24:1212–9.
- Thomas MC, Macisaac RJ, Jerums G, Weekes A, Moran J, Shaw JE, et al. Nonalbuminuric renal impairment in type 2 diabetic patients and in the general population (national evaluation of the frequency of renal impairment co-existing with NIDDM [NEFRON] 11). *Diabetes Care*. 2009;32:1497–502.
- Penno G, Solini A, Bonora E, Fondelli C, Orsi E, Zerbini G, et al. Clinical significance of nonalbuminuric renal impairment in type 2 diabetes. *J Hypertens*. 2011;29:1802–9.
- Ninomiya T, Perkovic V, de Galan BE, Zoungas S, Pillai A, Jardine M, et al. Albuminuria and kidney function independently predict cardiovascular and renal outcomes in diabetes. *J Am Soc Nephrol*. 2009;20:1813–21.
- Afghahi H, Cederholm J, Eliasson B, Zethelius B, Gudbjörnsdóttir S, Hadimeri H, et al. Risk factors for the development of albuminuria and renal impairment in type 2 diabetes—the Swedish National Diabetes Register (NDR). *Nephrol Dial Transplant*. 2011;26:1236–43.
- Fioretto P, Mauer M, Brocco E, Velussi M, Frigato F, Muollo B, et al. Patterns of renal injury in NIDDM patients with microalbuminuria. *Diabetologia*. 1996;39:1569–76.
- Fioretto P, Caramori ML, Mauer M. The kidney in diabetes: dynamic pathways of injury and repair. The Camillo Golgi Lecture 2007. *Diabetologia*. 2008;51:1347–55.
- Gambara V, Mecca G, Remuzzi G, Bertani T. Heterogeneous nature of renal lesions in type II diabetes. *J Am Soc Nephrol*. 1993;3:1458–66.
- Sugiyama H, Yokoyama H, Sato H, Saito T, Kohda Y, Nishi S, et al. Japan renal biopsy registry and Japan kidney disease registry: Committee report for 2009 and 2010. *Clin Exp Nephrol*. 2013;17:155–73.
- Takazakura E, Nakamoto Y, Hayakawa H, Kawai K, Muramoto S. Onset and progression of diabetic glomerulosclerosis; a prospective study based on serial renal biopsies. *Diabetes*. 1975;24:1–9.
- Shimizu M, Furuichi K, Toyama T, Kitajima S, Hara A, Kitagawa K, et al. Long-term outcomes of Japanese type 2 diabetic patients with biopsy-proven diabetic nephropathy. *Diabetes Care*. 2013. doi:10.2337/dc13-0298.
- Taniwaki H, Nishizawa Y, Kawagishi T, Ishimura E, Emoto M, Okamura T, et al. Decrease in glomerular filtration rate in Japanese patients with type 2 diabetes is linked to atherosclerosis. *Diabetes Care*. 1998;21:1848–55.
- MacIsaac RJ, Panagiotopoulos S, McNeil KJ, Smith TJ, Tsalamandris C, Hao H, et al. Is nonalbuminuric renal insufficiency in type 2 diabetes related to an increase in intrarenal vascular disease? *Diabetes Care*. 2006;29:1560–6.
- Uzu T, Kida Y, Shirahashi N, Harada T, Yamauchi A, Nomura M, et al. Cerebral microvascular disease predicts renal failure in type 2 diabetes. *J Am Soc Nephrol*. 2010;21:520–6.
- Araki S, Haneda M, Sugimoto T, Isono M, Isshiki K, Kashiwagi A, et al. Polymorphisms of the protein kinase C-beta gene (PRKCB1) accelerate kidney disease in type 2 diabetes without overt proteinuria. *Diabetes Care*. 2006;29:864–8.
- Rigalleau V, Lasseur C, Raffaitin C, Beauvieux MC, Barthe N, Chauveau P, et al. Normoalbuminuric renal-insufficient diabetic patients: a lower-risk group. *Diabetes Care*. 2007;30:2034–9.
- Kramer CK, Leitão CB, Pinto LC, Silveiro SP, Gross JL, Canani LH. Clinical and laboratory profile of patients with type 2 diabetes with low glomerular filtration rate and normoalbuminuria. *Diabetes Care*. 2007;30:1998–2000.
- Lane PH, Steffes MW, Mauer SM. Glomerular structure in IDDM women with low glomerular filtration rate and normal urinary albumin excretion. *Diabetes*. 1992;41:581–6.
- Caramori ML, Fioretto P, Mauer M. Low glomerular filtration rate in normoalbuminuric type 1 diabetic patients: an indicator of more advanced glomerular lesions. *Diabetes*. 2003;52:1036–40.
- Yagil C, Barak A, Ben-Dor D, Rosenmann E, Bernheim J, Rosner M, et al. Nonproteinuric diabetes-associated nephropathy in the Cohen rat model of type 2 diabetes. *Diabetes*. 2005;54:1487–96.
- Ekinci EI, Jerums G, Skene A, Crammer P, Power D, Cheong KY, et al. Renal structure in normoalbuminuric and albuminuric patients with type 2 diabetes and impaired renal function. *Diabetes Care*. 2013;. doi:10.2337/dc12-2572.
- Bruno G, Merletti F, Bargero G, Novelli G, Melis D, Soddu A, et al. Estimated glomerular filtration rate, albuminuria and