

Int J Vasc Med, 2012;2012:162545. Review.

- Ikeda Y, Aihara K, Yoshida S, Akaike M, Matsumoto T. et al.
Effects of androgens on cardiovascular remodeling.
J Endocrinol, 2012;214:1-10. Review.

研究分担者：戸口田淳也

- Matsumoto Y, Ikeya M, Hino K, Horigome K, Fukuta M, Watanabe M, Nagata S, Yamamoto T, Otsuka T, Toguchida J.
New protocol to optimize iPS cells for genome analysis of fibrodysplasia ossificans progressive.
Stem Cells, in press.
- Yokoyama K, Ikeya M, Umeda K, Oda H, Nodomi S, Nasu A, Matsumoto Y, Izawa K, Horigome K, Kusaka T, Tanaka T, Saito MK, Yasumi T, Nishikomori R, Ohara O, Nakayama N, Nakahata T, Heike T, Toguchida J.
Enhanced chondrogenesis of iPS cells from neonatal-onset multisystem inflammatory disease occurs via the caspase-1-independent cAMP/PKA/CREB pathway.
Arthritis Rheumatol. 2015;67(1):302-314.
- Fukuta M, Nakai Y, Kirino K, Nakagawa M, Sekiguchi K, Nagata S, Matsumoto Y, Yamamoto T, Umeda K, Heike T, Okumura N, Koizumi N, Sato T, Nakahata T, Saito M, Otsuka T, Kinoshita S, Ueno M, Ikeya M, Toguchida J.
Derivation of mesenchymal stromal cells from pluripotent stem cells through a neural crest lineage using small molecule compounds with defined media.
PLOS ONE, 2014;9(12):e112291.
- Aoyama T, Fujita Y, Madoba K, Nankaku M, Yamada M, Tomita M, Goto K, Ikeguchi R, Kakinoki R, Matsuda S, Nakamura T, Toguchida J.
Rehabilitation program after mesenchymal stromal cell transplantation augmented by vascularized bone grafts for idiopathic osteonecrosis of the femoral head: A preliminary study.
Arch Phys Med Rehabil. 2014; pii: S0003-9993(14)01210-6.
- Totoki Y, Yoshida A, Hosoda F, Nakamura H, Hama N, Ogura K, Yoshida A, Fujiwara T, Arai Y, Toguchida J, Tsuda H, Miyano S, Kawai A, Shibata T.
Unique mutation portraits and frequent COL2A1 gene alteration in chondrosarcoma.
Genome Res, 2014;24(9):1411-1420.
- Aoyama T, Goto K, Kakinoki R, Ikeguchi R, Ueda M, Kasai Y, Maekawa T, Tada H, Teramukai S, Nakamura T, Toguchida J.
An exploratory clinical trial for idiopathic osteonecrosis of femoral head by cultured autologous multipotent mesenchymal stromal cells augmented with vascularized bone grafts.
Tissue Eng Part B Rev, 2014;20(4):233-242.
- Husain SR, Ohya Y, Toguchida J, Puri RK.
Current Status of Multipotent Mesenchymal Stromal Cells.
Tissue Eng Part B Rev, 2014;20(3):189.

- Kajita Y, Kato T Jr, Tamaki S, Furu M, Takahashi R, Nagayama S, et al. The transcription factor Sp3 regulates the expression of a metastasis-related marker of sarcoma, actin filament-associated protein 1-like 1 (AFAP1L1). PLoS One, 2013;8:e49709.
- Yamada K, Ohno T, Aoki H, Semi K, Watanabe A, Moritake H, et al. EWS/ATF1 expression induces sarcomas from neural crest-derived cells in mice. J Clin Invest, 2013;123:600–610.
- Hayakawa K, Ikeya M, Fukuta M, Woltjen K, Tamaki S, Takahara N, et al. Identification of target genes of synovial sarcoma-associated fusion oncoprotein using human pluripotent stem cells. Biochem Biophys Res Commun, 2013;432:713–719.
- Nasu A, Ikeya M, Yamamoto T, Watanabe A, Jin Y, Matsumoto Y, et al. Genetically matched human iPS cells reveal that propensity for cartilage and bone differentiation differs with clones, not cell type of origin. PLoS One, 2013;8:e53771.
- Okita Y, Tatematsu N, Nagai K, Nakayama T, Nakamata T, Okamoto T, et al. Characteristics of flexed knee gait and functional outcome of a patient who underwent knee reconstruction with a hingeless prosthesis for bone tumor resection: a case report with gait analysis and comparison with healthy subjects. Eur J Phys Rehabil Med, 2013 Jul 9. [Epub ahead of print]
- Okita Y, Tatematsu N, Nagai K, Nakayama T, Nakamata T, Okamoto T, et al. Compensation by nonoperated joints in the lower limbs during walking after endoprosthetic knee replacement following bone tumor resection. Clin Biomech (Bristol, Avon), 2013;28:898–903.
- Matsumoto Y, Hayashi Y, Schlieve CR, Ikeya M, Kim H, Nguyen TD, et al. Induced pluripotent stem cells from patients with human fibrodysplasia ossificans progressiva show increased mineralization and cartilage formation. Orphanet J Rare Dis, 2013;8:190.
- Masago Y, Hosoya A, Kawasaki K, Kawano S, Nasu A, Toguchida J, et al. Molecular chaperone Hsp47 is essential for cartilage and endochondral bone formation. J Cell Sci, 2012;125:1118–1128.
- Kawaguchi S, Tsukahara T, Ida K, Kimura S, Murase M, Kano M, et al. SYT-SSX breakpoint peptide vaccines in patients with synovial sarcoma: A study from the Japanese Musculoskeletal Oncology Group. Cancer Sci. 2012;103:1625–1630.
- Kajiwara M, Aoi T, Okita K, Takahashi R, Inoue H, Takayama N, et al. Donor-dependent variations in hepatic differentiation from human-induced pluripotent stem cells. Proc Natl Acad Sci U S A. 2012;109:12538–12543.

- ・ 戸口田淳也
骨再生医療(間葉系幹細胞) 最新の骨粗鬆症学-骨粗鬆症の最新知見.
日本臨床, 2013;71:535-538.
- ・ 戸口田淳也
疾患特異的 iPS 細胞を用いた難治性骨・軟骨疾患の病態解析.
Bio Clinica, 2013;28:37-41.
- ・ 渡辺真、横山宏司、佐藤孝明、戸口田淳也
疾患特異的 iPS 細胞を用いた遺伝性軟骨疾患マーカー探索.
Medical Scinece Digest, 2013;39:530-533.
- ・ 松本佳久、福田誠、池谷真、戸口田淳也
iPS 細胞の基礎と応用 (第 3 回) iPS 細胞を使う-骨の研究へ.
整形外科, 2013;64:1017-1019.
- ・ 松本佳久、池谷真、戸口田淳也
難治性骨軟骨疾患罹患者由来 iPS 細胞を用いた病態再現と治療薬開発の試み. 最新疾患モデルと病態解明、創薬応用研究、細胞医薬創製研究の最前線
遺伝子医学 MOOK, 2012;22:190-194.

研究分担者 : 松田秀一

- ・ Kuroda Y, Akiyama H, Nankaku M, So K, Goto K, Matsuda S.
A report on three consecutive cases using computer tomography 3D preoperative planning for conversion of arthrodesed hips to total hip replacements.
HSSJ. in press, 2014.
- ・ Goto K, So K, Kuroda Y, Kawanabe K, Matsuda S.
Two-staged reconstruction for ankylosed hip with severe limb shortening: a case report.
Hip Int. 2014;5:24(6):660-663.
- ・ Tsukagoshi R, Tateuchi H, Fukumoto Y, Ibuki S, Akiyama H, So K, Kuroda Y, Okumura H, Ichihashi N. Functional performance of female patients more than 6 months after total hip arthroplasty shows greater improvement with weight-bearing exercise than with non-weight-bearing exercise: a randomized controlled trial. Eur J Phys Rehabil Med. 2014;50(6):665-675.
- ・ Tateuchi H, Tsukagoshi R, Fukumoto Y, Akiyama H, So K, Kuroda Y, Ichihashi N. Compensatory turning strategies while walking in patients with hip osteoarthritis. Gait Posture. 2014;39(4):1133-1137.
- ・ Fukumoto Y, Tateuchi H, Ikezoe T, Tsukagoshi R, Akiyama H, So K, Kuroda Y, Ichihashi N. Effects of high-velocity resistance training on muscle function, muscle properties, and physical performance in individuals with hip osteoarthritis: a randomized controlled trial. Clin Rehabil. 2014;28(1):48-58.
- ・ So K, Kuroda Y, Matsuda S, Akiyama H.

- Revision total hip replacement with a cemented long femoral component: minimum 9-year follow-up results. Arch Orthop Trauma Surg. 2013;133(6):869–874.
- Fukumoto Y, Ikezoe T, Tateuchi H, Tsukagoshi R, Akiyama H, So K, Kuroda Y, Yoneyama T, Ichihashi N; Muscle mass and composition of the hip, thigh and abdominal muscles in women with and without hip osteoarthritis. Ultrasound Med Biol. 2012;38:1540–1545.
 - Akiyama H, Kuroda Y, So K, Kakinoki R, Nakamura T. Study of a retrieved cementless acetabular cup with an AW-GC bottom coating. J Orthop Sci. 2013;18(6):1037–1041.
 - Tateuchi H, Tsukagoshi R, Fukumoto Y, Akiyama H, So K, Kuroda Y, Ichihashi N. Pelvic instability and trunk and hip muscle recruitment patterns in patients with total hip arthroplasty. J Electromyogr Kinesiol. 2013;23(1):151–8.
 - So K, Kanatani KT, Kuroda Y, Nakamura T, Matsuda S, Akiyama H. Good short-term outcome of primary total hip arthroplasty with cementless bioactive glass ceramic bottom-coated implants. Acta Orthop, 2012;83(6):599–603.
 - 黒田 隆
生医療の現状と展望：細胞を用いない再生医療 FGF-2ゼラチンハイドロゲルを用いた大腿骨頭壊死症の治療 一家兔動物モデルを用いて一，整形・災害外科 第56巻2013年4月臨時増刊号，金原出版，2013;625–633.
 - 黒田 隆
大腿骨頭壊死動物モデルを用いた細胞増殖因子治療の取り組みと臨床応用研究，遺伝子MOOK22号 最新疾患モデルと病態解明-創薬応用研究 細胞医薬創製研究の最前線-，メディカルドゥ，2012;195–200.

研究分担者：今井祐記

- Omata Y, Yasui T, Hirose J, Izawa N, Imai Y, Matsumoto T, Masuda H, Tokuyama N, Nakamura S, Tsutsumi S, Yasuda H, Okamoto K, Takayanagi H, Hikita A, Imamura T, Matsuo K, Saito T, Kadono Y, Aburatani H, Tanaka S. Genome-wide comprehensive analysis reveals critical cooperation between Smad and c-Fos in RANKL-induced osteoclastogenesis. J Bone Miner Res, 2014; [Epub ahead of print]
- Yano K, Yasuda H, Takaoka K, Takahashi M, Nakamura H, Imai Y, Wakitani S. Fate, origin and roles of cells within free bone grafts. J Orthop Sci, 2014;[Epub ahead of print]
- Inoue K and Imai Y. Identification of Novel Transcription Factors in Osteoclast Differentiation using Genome-wide

- Analysis of Open Chromatin Determined by DNase-seq.
J Bone Miner Res, 2014;29(8):1823–1832.
- Shimizu K, Uematsu A, Imai Y, Sawasaki T.
Pctairel/Cdk16 promotes skeletal myogenesis by inducing myoblast migration and fusion.
FEBS Lett, 2014;25;588(17):3030–3037.
 - Ipulan LA, Suzuki K, Sakamoto Y, Murashima A, Imai Y, Omori A, Nakagata N, Nishinakamura R, Valasek P, Yamada G.
Non-myocytic androgen receptor regulates the sexually dimorphic development of the embryonic bulbocavernosus muscle.
Endocrinology, 2014;155(7):2467–2479.
 - Yu TY, Kondo T, Matsumoto T, Fujii-Kuriyama Y, Imai Y.
Aryl hydrocarbon receptor catabolic activity in bone metabolism is osteoclast dependent in vivo.
Biochem Biophys Res Commun, 2014;450(1):416–422.
 - Enoki Y, Sato T, Tanaka S, Iwata T, Usui M, Takeda S, Kokabu S, Matsumoto M, Okubo M, Nakashima K, Yamato M, Okano T, Fukuda T, Chida D, Imai Y, Yasuda H, Nishihara T, Akita M, Oda H, Okazaki Y, Suda T, Yoda T.
Netrin-4 derived from murine vascular endothelial cells inhibits osteoclast differentiation in vitro and prevents bone loss in vivo.
FEBS Lett, 2014;588(14):2262–2269.
 - Kondoh S, Inoue K, Igarashi K, Sugizaki, H, Shirode-Fukuda Y, Inoue E, Yu T, Takeuchi J, Kanno J, Bonewald L, Imai Y.
Estrogen receptor α in osteocytes regulates trabecular bone formation in female mice.
Bone, in press, 2014.
 - Okuno Y, Inoue K, Imai Y.
Novel insights into histone modifiers in Adipogenesis.
Adipocyte, 2013;2(4):285–288.
 - Si Y, Inoue K, Igarashi K, Kanno J, Imai Y.
Autoimmune regulator, Aire, is a novel regulator of chondrocyte differentiation.
Biochem Biophys Res Commun, 2013;437(4):579–584.
 - Imai Y, Youn MY, Inoue K, Takada I, Kouzmenko A, Kato S.
Nuclear receptors in bone physiology and diseases.
Physiol Rev, 2013;93(2):481–523. doi: 10.1152/physrev.00008.2012.
 - Inoue K, Inoue E, Imai Y.
Female sex hormones ameliorate arthritis in SKG mice.
Biochem Biophys Res Commun, 2013;434(4):740–745. doi: 10.1016/j.bbrc.2013.03.111.
 - Okuno Y, Otake F, Igarashi K, Kanno J, Matsumoto T, Takada I, et al.
Epigenetic regulation of adipogenesis by PHF2 histone demethylase.
Diabetes, 2013;62(5):1426–1434.

- Yamamoto Y, Yoshizawa T, Fukuda T, Shirode-Fukuda Y, Yu T, Sekine K, Sato T, Kawano H, Aihara K, Nakamichi Y, Watanabe T, Shindo M, Inoue K, Inoue E, Tsuji N, Hoshino M, Karsenty G, Metzger D, Chambon P, Kato S, Imai Y.
Vitamin D receptor in osteoblasts is a negative regulator of bone mass control.
Endocrinology, 2013;154(3):1008–1020.
- You MY, Yokoyama A, Fujiyama-Nakamura S, Ohtake F, Minehata KI, Yasuda H, Suzuki T, Kato S, Imai Y.
JMJD5, a JmjC-domain-containing protein, negatively regulates osteoclastogenesis by facilitating NFATc1 protein degradation.
J Biol Chem, 2012;287(16):12994–3004.

研究分担者：高橋大介

- Takahashi D, Majima T, Onodera T, Kasahara Y, Inoue M, Irie T, Kasemura T.
Celecoxib does not affect the release of hyaluronic acid in end stage osteoarthritic joints.
Mod Rheumatol, 2013;23(5):934–938.
- Onodera T, Majima T, Nishiike O, Kasahara Y, Takahashi D.
Posterior femoral condylar offset after total knee replacement in the risk of knee flexion contracture.
J Arthroplasty, 2013;28(7):1112–1116.
- Tanabe A, Majima T, Onodera T, Sawaguchi N, Kasahara Y, Takahashi D.
Sagittal alignment of the first metatarsophalangeal joint after arthrodesis for rheumatoid forefoot deformity.
J Foot Ankle Surg, 2013;52(3):343–347.
- Nishio Y, Onodera T, Kasahara Y, Takahashi D, Iwasaki N, Majima T.
Intraoperative medial pivot affects deep knee flexion angle and patient-reported outcomes after total knee arthroplasty.
J Arthroplasty, in press 2013.
- Majima T, Inoue M, Kasahara Y, Onodera T, Takahashi D, Minami A.
Effect of the Japanese herbal medicine, Boiogito, on the osteoarthritis of the knee with joint effusion.
Sports Med Arthrosc Rehabil Ther Technol, 2012;10, 3.
- Ito H, Tanino H, Yamanaka Y, Nakamura T, Takahashi D, Minami A, et al.
Long-term results of conventional varus halfwedge proximal femoral osteotomy for the treatment of osteonecrosis of the femoral head.
J Bone Joint Surg Br, 2012;94-B:308–314.
- Matsuoka M, Majima T, Onodera T, Ieko M, Souri M, Takahashi D.
Hemorrhagic-acquired factor XIII deficiency associated with tocilizumab for treatment of rheumatoid arthritis.
Int J Hematol, 2012;96(6):781–785.

- Onodera T, Majima T, Kasahara Y, Takahashi D, Yamazaki S, Ando R, et al. Outcome of transfibular ankle arthrodesis with Ilizarov apparatus. Foot Ankle Int, 2012;33(11):964–968.
- 高橋大介、本宮 真、岩崎倫政
北海道大学整形外科で行っている外傷の臨床と教育～創外固定治療とマイクロサージャリー～.
北海道整形災害外科学会雑誌, 2013;55(1):50–56.
- 校條祐輔、笠原靖彦、小野寺智洋、高橋大介、総村俊之、入江 徹、他
膝関節発生びまん型色素性絨毛結節性滑膜炎に対する滑膜全切除術の治療成績.
JOSKAS, 2013;38(2):318–319.
- 岩崎浩司、笠原靖彦、小野寺智洋、高橋大介、総村俊之、入江 徹、他
高度内反変形膝に対する全人工膝関節置換術の治療成績.
JOSKAS, 2013;38(2):230–231.
- 小野寺智洋、眞島任史、笠原靖彦、高橋大介、入江 徹、総村俊之、他
距骨骨軟骨病変に対する骨軟骨柱移植術の治療成績.
JOSKAS, 2013;38(1):146–147.
- 清水智弘、高橋大介、紺野拓也、井上正弘、田辺明彦、入江 徹、他
股関節後方脱臼骨折の治療成績.
Hip Joint, 2013;39:506–510.
- 入江 徹、眞島任史、高橋大介、笠原靖彦、小野寺智洋、門司順一
先天性膝関節脱臼の治療成績
日本小児整形外科学会雑誌, 2013;22(1):94–99.
- 馬場力哉、寺島尚志、笠原靖彦、小野寺智洋、高橋大介、岩崎倫政、他
リウマチ性膝関節症に対する人工膝関節全置換術の臨床成績 生物学的製剤使用群と非使用群の違い.
北海道整形災害外科学会雑誌, 2013;55(1):73–77.
- 高橋大介、眞島任史、井上正弘、笠原靖彦、小野寺智洋、紺野拓也、他
S-ROM THA と S-ROM-A THA の中期成績の比較.
日本人工関節学会誌, 2013;43:125–126.
- 高橋大介、眞島任史、山田勝久、渡部琢哉、沢口直弘、三浪明男、他
難治性の両側先天性膝関節脱臼の治療経験.
日小整会誌, 2012;21(1):81–85.
- 高橋大介、眞島任史、井上正弘、安部聰弥、菅野大己、増田武志
THA 術後ステム周囲骨折の治療経験.
Hip Joint, 2012;38:707–710.
- 遠藤香織、加藤浩仁、菅野忠博、高橋大介、眞島任史
特発性大腿骨頭壊死症に対する CT・MRI fusion による 3 次元術前シミュレーションプログラム.
Hip Joint, 2012;38:1112–1116.
- 遠藤香織、高橋大介、眞島任史、井上正弘
特発性大腿骨頭壊死症の関節温存手術の術前計画における MRI 画像の再構築技術の有用性.

- Hip Joint, 2012;38:793–797.
- ・ 紹村俊之、眞島任史、笠原靖彦、小野寺智洋、高橋大介、入江 徹
TKA 術後成績 PFC Σ RP-F を使用した TKA の短期成績.
日本人工関節学会誌, 2012;42:219–220.
 - ・ 高橋大介、眞島任史、山田勝久、渡部琢哉、沢口直弘、三浪明男、他
難治性の両側先天性膝関節脱臼の治療経験.
日本小児整形外科学会雑誌, 2012;21(1):81–85.
 - ・ 遠藤香織、加藤浩仁、菅野忠博、高橋大介、眞島任史
特発性大腿骨頭壞死症に対する CT・MRI fusion 画像による三次元術前シミュレーションプログラム.
Hip Joint, 2012;38:1112–1116.
 - ・ 笹沢史生、高橋大介、井上正弘、大浦久典、伊藤 浩、眞島任史
ペルテス病に対する大腿骨内反骨切り術の術後成績に影響を与える因子.
Hip Joint, 2012;38:838–842.
 - ・ 遠藤香織、高橋大介、眞島任史、井上正弘
特発性大腿骨頭壞死症の関節温存手術の術前計画における MRI 画像の再構築技術の有用性.
Hip Joint, 2012;38:793–797.
 - ・ 入江 徹、井上正弘、大浦久典、高橋大介、眞島任史
大腿骨転子下骨切り併用セメントレス S -ROM THA の中期成績.
Hip Joint, 2012;38:432–436.

研究分担者：菅野伸彦

- ・ Tamura S, Takao M, Sakai T, Nishii T, Sugano N.
Spinal Factors Influencing Change in Pelvic Sagittal Inclination From Supine Position to Standing Position in Patients Before Total Hip Arthroplasty.
J Arthroplasty, 2014; 29(12):2294–2297.
- ・ Nakahara I, Takao M, Bandoh S, Sugano N.
Fixation strength of taper connection at head-neck junction in retrieved carbon fiber-reinforced PEEK hip stems. J Artif Organs. 2014; 17(4):358–363.
- ・ Takahashi Y, Sugano N, Puppulin L, Zhu W, Pezzotti G.
Raman spectroscopic study of remelting and annealing-induced effects on microstructure and compressive deformation behavior of highly crosslinked UHMWPE for total hip arthroplasty.
J Biomed Mater Res B Appl Biomater, 2014; 102(8):1762–1770.
- ・ Takao M, Nishii T, Sakai T, Sugano N.
Navigation-aided visualization of lumbosacral nerves for anterior sacroiliac plate fixation: a case report. Int J Med Robot, 2014; 10(2):230–236.
- ・ Miki H, Kyo T, Kuroda Y, Nakahara I, Sugano N.
Risk of edge-loading and prosthesis impingement due to posterior pelvic tilting after total hip arthroplasty. Clin Biomech (Bristol, Avon), 2014; 29(6):607–613.

- Takao M, Nishii T, Sakai T, Yoshikawa H, Sugano N. Iliosacral screw insertion using CT-3D-fluoroscopy matching navigation. Injury, 2014; 45(6):988–994.
- Nakahara I, Takao M, Sakai T, Miki H, Nishii T, Sugano N. Three-dimensional morphology and bony range of movement in hip joints in patients with hip dysplasia. Bone Joint J, 2014; 96-B(5):580–589.
- Zhu W, Puppulin L, Leto A, Takahashi Y, Sugano N, Pezzotti G. In situ measurements of local temperature and contact stress magnitude during wear of ceramic-on-ceramic hip joints. J Mech Behav Biomed Mater, 2014; 31:68–76.
- Puppulin L, Leto A, Wenliang Z, Sugano N, Pezzotti G. Innovative tribometer for in situ spectroscopic analyses of wear mechanisms and phase transformation in ceramic femoral heads. J Mech Behav Biomed Mater, 2014; 31:45–54.
- Puppulin L, Sugano N, Zhu W, Pezzotti G. Structural modifications induced by compressive plastic deformation in single-step and sequentially irradiated UHMWPE for hip joint components. J Mech Behav Biomed Mater, 2014; 31:86–99.
- Takahashi Y, Sugano N, Takao M, Sakai T, Nishii T, Pezzotti G. Raman spectroscopy investigation of load-assisted microstructural alterations in human knee cartilage: Preliminary study into diagnostic potential for osteoarthritis. J Mech Behav Biomed Mater, 2014; 31:77–85.
- Sakai T, Hanada T, Murase T, Kitada M, Hamada H, Yoshikawa H, Sugano N. Validation of patient specific surgical guides in total hip arthroplasty. Int J Med Robot, 2014; 10(1):113–120.
- Sugano N, Iida H, Akiyama H, Takatori Y, Nagoya S, Hasegawa M, Kabata T, Hachiya Y, Yasunaga Y. Nationwide investigation into adverse tissue reactions to metal debris after metal-on-metal total hip arthroplasty in Japan. J Orthop Sci, 2014; 19:85–89.
- Abe H, Sakai T, Ando W, Takao M, Nishii T, Nakamura N, Hamasaki T, Yoshikawa H, Sugano N. Synovial joint fluid cytokine levels in hip disease. Rheumatology (Oxford), 2014; 53(1):165–172.
- Abe H, Sakai T, Nishii T, Takao M, Nakamura N, Sugano N. Jogging After Total Hip Arthroplasty. Am J Sports Med, 2014; 42(1):131–137.
- Nakasone S, Takao M, Sakai T, Nishii T, Sugano N. Does the extent of osteonecrosis affect the survival of hip resurfacing? Clin Orthop Relat Res 2013;471(6):1926–1934.

- Sakai T, Iwana D, Nakamura N, Nishii T, Takao M, Watanabe T, Yoshikawa H, Sugano N. Histological characteristics of the human femoral head in patients with femoral neck fracture. *Virchows Arch*, 2012;461(6):705–711.
- Nakasone S, Takao M, Nishii T, Sakai T, Sugano N. Incidence and natural course of initial polar gaps in Birmingham Hip Resurfacing cups. *J Arthroplasty*, 2012;27(9):1676–1682.
- Nishii T, Sakai T, Takao M, Yoshikawa H, Sugano N. Ultrasound screening of periarticular soft tissue abnormality around metal-on-metal bearings. *J Arthroplasty*, 2012;27:895–900.
- Takao M, Sakai T, Nishii T, Yoshikawa H, Takahara S, Sugano N. Incidence and predictors of osteonecrosis among cyclosporin- or tacrolimus-treated renal allograft recipients. *Rheumatol Int*, 2011;31:165–170.
- 仲宗根哲、高尾正樹、西井 孝、坂井孝司、中村宣雄、岩名大樹、北田 誠、菅野伸彦 特発性大腿骨頭壞死症に対する表面置換型人工股関節全置換術の中期成績-従来型 THAとの比較- *Hip Joint*, 2012;38:810–811.
- 坂井孝司、中村宣雄、安藤 渉、西井 孝、高尾正樹、阿部裕仁、田村 理、大園健二、吉川秀樹、菅野伸彦 大腿骨頭軟骨下骨折と急速破壊型股関節症の組織学的検討 *Hip joint*, 2012;38:1005–1009.
- 西井 孝 特発性大腿骨頭壞死症 今日の治療指針, 2012;908–909.

研究分担者：赤司浩一

- Oryoji K, Kiyohara C, Horiuchi T, Tsukamoto H, Niiro H, Shimoda T, Akashi K, Yanase T. Reduced carotid intima-media thickness in systemic lupus erythematosus patients treated with cyclosporine A. *Mod Rheumatol*, 2014; 24: 86–92.
- Chong Y, Shimoda S, Yakushiji H, Ito Y, Aoki T, Miyamoto T, Kamimura T, Shimono N, Akashi K. Clinical impact of fluoroquinolone-resistant *Escherichia coli* in the fecal flora of hematological patients with neutropenia and levofloxacin prophylaxis. *PLoS One*, 2014; 9: e85210.
- Yabuuchi H, Matsuo Y, Tsukamoto H, Horiuchi T, Sunami S, Kamitani T, Jinnouchi M, Nagao M, Akashi K, Honda H. Evaluation of the extent of ground-glass opacity on high-resolution CT in patients with interstitial pneumonia associated with systemic sclerosis: Comparison between quantitative and qualitative analysis. *Clin Radiol*, 2014;69: 758–764.
- Shima T, Miyamoto T, Kikushige Y, Yuda J, Tochigi T, Yoshimoto G, Kato K, Takenaka K, Iwasaki H,

Mizuno S, Goto N, Akashi K.

The ordered acquisition of Class II and Class I mutations directs formation of human t(8;21) acute myelogenous leukemia stem cell.

Exp Hematol, 2014; 42: 955-965.

- Kato K, Ohno Y, Kamimura T, Kusumoto H, Tochigi T, Jinnouchi F, Kohno K, Kuriyama T, Henzan H, Takase K, Kawano I, Kadokami M, Nawata R, Muta T, Eto T, Iwasaki H, Ohshima K, Miyamoto T, Akashi K.

Long-term remission after high-dose chemotherapy followed by auto-SCT as consolidation for intravascular large B-cell lymphoma.

Bone Marrow Transplant, 2014; 1543-1544.

- Yonekawa A, Saijo S, Hoshino Y, Miyake Y, Ishikawa E, Suzukawa M, Inoue H, Tanaka M, Yoneyama M, Oh-Hora M, Akashi K, Yamasaki S.

Dectin-2 Is a Direct Receptor for Mannose-Capped Lipoarabinomannan of Mycobacteria.

Immunity, 2014; 41: 402-413.

- Miyawaki K, Arinobu Y, Iwasaki H, Kohno K, Tsuzuki H, Iino T, Shima T, Kikushige Y, Takenaka K, Miyamoto T, Akashi K.

CD41 marks the initial myelo-erythroid lineage specification in adult mouse hematopoiesis: Redefinition of murine common myeloid progenitor.

Stem Cells, 2014 (in press)

- Iwamoto C, Takenaka K, Urata S, Yamauchi T, Shima T, Kuriyama T, Daitoku S, Saito Y, Miyamoto T, Iwasaki H, Kitabayashi I, Itoh K, Kishimoto J, Kohda D, Matozaki T, Akashi K.

The BALB/c-specific polymorphic SIRPA enhances its affinity for human CD47, inhibiting phagocytosis against human cells to promote xenogeneic engraftment.

Exp Hematol, 2013; 42: 163-171.

- Yamauchi T, Takenaka K, Urata S, Shima T, Kikushige Y, Tokuyama T, Iwamoto C, Nishihara M, Iwasaki H, Miyamoto T, Honma N, Nakao M, Matozaki T, Akashi K.

Polymorphic sirpa is the genetic determinant for nod-based mouse lines to achieve efficient human cell engraftment.

Blood, 2013; 121: 1316-1325.

- Shima T, Miyamoto T, Kikushige Y, Mori Y, Kamezaki K, Takase K, Henzan H, Numata A, Ito Y, Takenaka K, Iwasaki H, Kamimura T, Eto T, Nagafuji K, Teshima T, Kato K, Akashi K.

Quantitation of hematogones at the time of engraftment is a useful prognostic indicator in allogeneic hematopoietic stem cell transplantation.

Blood, 2013; 121: 840-848.

- Shima T, Forraz N, Sato N, Yamauchi T, Iwasaki H, Takenaka K, Akashi K, McGuckin C, Teshima T. A novel filtration method for cord blood processing using a polyester fabric filter.

International journal of laboratory hematology, 2013; 35: 436-446.

- Chong Y, Shimoda S, Yakushiji H, Ito Y, Miyamoto T, Kamimura T, Shimono N, Akashi K.

Antibiotic rotation for febrile neutropenic patients with hematological malignancies: Clinical significance of antibiotic heterogeneity.

PloS one, 2013;8:e54190.

- Furukawa M, Kiyohara C, Horiuchi T, Tsukamoto H, Mitoma H, Kimoto Y, Uchino A, Nakagawa M, Oryoji K, Shimoda T, Harada M, Akashi K.

Prevalence and risk factors of vertebral fracture in female Japanese patients with systemic lupus erythematosus.

Modern rheumatology, 2013;23:765–773.

- Ueda N, Tsukamoto H, Mitoma H, Ayano M, Tanaka A, Ohta S, Inoue Y, Arinobu Y, Niro H, Akashi K, Horiuchi T. The cytotoxic effects of certolizumab pegol and golimumab mediated by transmembrane tumor necrosis factor alpha.

Inflammatory bowel diseases, 2013;19:1224–1231.

- Sumida K, Shimoda S, Iwasaka S, Hisamoto S, Kawanaka H, Akahoshi T, Ikegami T, Shirabe K, Shimono N, Maehara Y, Selmi C, Gershwin ME, Akashi K.

Characteristics of splenic CD8(+) T cell exhaustion in patients with hepatitis C.

Clinical and experimental immunology, 2013;174:172–178.

- Fukata M, Ishikawa F, Najima Y, Yamauchi T, Saito Y, Takenaka K, Miyawaki K, Shimazu H, Shimoda K, Kanemaru T, Nakamura K, Odashiro K, Nagafuji K, Harada M, Akashi K.

Contribution of bone marrow-derived hematopoietic stem/progenitor cells to the generation of donor-marker(+) cardiomyocytes in vivo.

PloS one, 2013;8:e62506.

- Welner RS, Bararia D, Amabile G, Czibere A, Benoukraf T, Bach C, Wansa KD, Ye M, Zhang H, Iino T, Hetherington CJ, Akashi K, Tenen DG.

C/ebpalpha is required for development of dendritic cell progenitors.

Blood, 2013;121:4073–4081.

- Niro H, Jabbarzadeh-Tabrizi S, Kikushige Y, Shima T, Noda K, Ota S, Tsuzuki H, Inoue Y, Arinobu Y, Iwasaki H, Shimoda S, Baba E, Tsukamoto H, Horiuchi T, Taniyama T, Akashi K.

CIN85 is required for Cbl-mediated regulation of antigen receptor signaling in human B cells.

Blood, 2012;119:2263–2273.

- Kumamaru H, Ohkawa Y, Saiwai H, Yamada H, Kubota K, Kobayakawa K, Akashi K, Okano H, Iwamoto Y, Okada S. Direct isolation and RNA-seq reveal environment-dependent properties of engrafted neural stem/progenitor cells.

Nat Commun 2012;3:1140.

- Kuriyama T, Takenaka K, Kohno K, Yamauchi T, Daitoku S, Yoshimoto G, Kikushige Y, Kishimoto J, Abe Y, Harada N, Miyamoto T, Iwasaki H, Teshima T, Akashi K.

Engulfment of hematopoietic stem cells caused by down-regulation of CD47 is critical in the pathogenesis of hemophagocytic lymphohistiocytosis,

Blood, 2012;120:4058–4067.

- Eriguchi Y, Takashima S, Oka H, Shimoji S, Nakamura K, Uryu H, Shimoda S, Iwasaki H, Shimono N, Ayabe T, Akashi K, Teshima T.
Graft-versus-host disease disrupts intestinal microbial ecology by inhibiting Paneth cell production of α -defensins.
Blood, 2012;120:223-231.
- Yuki H, Ueno S, Tatetsu H, Niiro H, Iino T, Endo S, Kawano Y, Komohara Y, Takeya M, Hata H, Okada S, Watanabe T, Akashi K, Mitsuya H, Okuno Y.
PU.1 is a potent tumor suppressor in classical Hodgkin lymphoma cells.
Blood, 2012;121:962-970.
- Maehara K, Odawara J, Harada A, Yoshimi T, Nagao K, Obuse C, Akashi K, Tachibana T, Sakata T, Ohkawa Y.
A co-localization model of paired ChIP-seq data using a large ENCODE data set enables comparison of multiple samples.
Nucleic Acids Res, 2012;41:54-62.
- Harada A, Okada S, Konno D, Odawara J, Yoshimi T, Yoshimura S, Kumamaru H, Saiwai H, Tsubota T, Kurumizaka H, Akashi K, Tachibana T, Imbalzano AN, Ohkawa Y.
Chd2 interacts with H3.3 to determine myogenic cell fate.
EMBO J, 2012;31:2994-3007.
- Mizuochi C, Fraser ST, Biasch K, Horio Y, Kikushige Y, Tani K, Akashi K, Tavian M, Sugiyama D.
Intra-aortic clusters undergo endothelial to hematopoietic phenotypic transition during early embryogenesis.
PLoS One, 2012;7:35763.
- Harris AC, Ferrara JL, Braun TM, Holler E, Teshima T, Levine JE, Choi SW, Landfried K, Akashi K, Lugt MV, Couriel DR, Reddy P, Paczesny S.
Plasma biomarkers of lower gastrointestinal and liver acute GVHD.
Blood, 2012;119:2960-2963.

研究分担者：天野宏一

- Tanaka Y, Takeuchi T, Amano K, Saito K, Hanami K, Nawata M, Fukuyo S, Kameda H, Kaneko Y, Kurasawa T, Nagasawa H, Hoshi D, Sato E, Yamanaka H.
Effect of interleukin-6 receptor inhibitor, tocilizumab, in preventing joint destruction in patients with rheumatoid arthritis showing inadequate response to TNF inhibitors.
Mod Rheumatol, 2014; 24: 399-404.
- Kurasawa T, Nagasawa H, Kishimoto M, Amano K, Takeuchi T, Kameda H.
Addition of another disease-modifying anti-rheumatic drug to methotrexate reduces the flare rate within 2 years after infliximab discontinuation in patients with rheumatoid arthritis: An open, randomized, controlled trial.
Mod Rheumatol, 2014; 24: 561-566.

- Sada KE, Amano K, Uehara R, Yamamura M, Arimura Y, Nakamura Y, Makino H. Research Committee on Intractable Vasculitides, the Ministry of Health, Labour, Welfare of Japan. : A nationwide survey on the epidemiology and clinical features of eosinophilic granulomatosis with polyangiitis (Churg-Strauss) in Japan. *Mod Rheumatol*, 2014; 24: 640–644.
- Tanaka Y, Kubo S, Tanaka Y, Kubo S, Yamanaka H, Amano K, Hirata S, Tanaka E, Nagasawa H, Yasuoka H, Takeuchi T. Efficacy and safety of abatacept in routine care of patients with rheumatoid arthritis: Orenicia® as Biological Intensive Treatment for RA (ORBIT) study. *Mod Rheumatol*, 2014 ; 24: 754–762.
- Terai C, Tsutsumi T, Sakurai T, Moriguchi M, Azuma T, Kaneko M, Kawagoe M, Hoshi K, Yoshida H, Matsui T, Nakajima K, Okuyama A, Nishi E, Amano K, Ota M, Mimura T, Chino K, Aoki K, Hand Y, Hirose T, Kida I, Kobayashi S, Suzuki K, Matsuzaki T, Kuga Y. The efficacy of mizoribine for the treatment of rheumatoid arthritis and its correlation with renal function. *Mod Rheumatol*, 2014; 24: 892–896.
- Sakai R, Cho SK, Nanki T, Koike R, Watanabe K, Yamazaki H, Nagasawa H, Amano K, Tanaka Y, Sumida T, Ihata A, Yasuda S, Nakajima A, Sugihara T, Tamura N, Fujii T, Dobashi H, Miura Y, Miyasaka N, Harigai M. For the REAL study group. : The risk of serious infection in patients with rheumatoid arthritis treated with tumor necrosis factor inhibitors decreased over time: a report from the registry of Japanese rheumatoid arthritis patients on biologics for long-term safety (REAL) database. *Rheumatol Int*, 2014; 34: 1729–1736.
- Kurasawa T, Nagasawa H, Nishi E, Takei H, Okuyama A, Kondo T, Nishimura K, Sakai R, Shibata A, Chino K, Ogawa H, Ito T, Amano K, Kato H. Successful treatment of class IV+V lupus nephritis with combination therapy of high-dose corticosteroids, tacrolimus and intravenous cyclophosphamide. *Intern Med*, 2013;52(10):1125–30.
- Nishi E, Kameda H, Ogawa H, Nagasawa H, Takei H, Okuyama A, Kurasawa T, Kondo T, Nishimura K, Shirai Y, Sakai R, Ito T, Takeuchi T, Amano K. Efficacy of weekly mizoribine pulse therapy in refractory lupus nephritis. *Mod Rheumatol*, 2013; 23(1): 97–103.

研究分担者：黒田 肇

- 黒田 肇、佐藤弘恵、中枝武司、和田庸子、村上修一、中野正明、成田一衛
全身性エリテマトーデスの初回ステロイド投与におけるスタチンの大腿骨頭壊死予防効果の検討.
中部リウマチ, 2014;44(1):11–13.
- 黒田 肇、若松彩子、佐藤弘恵、中枝武司、和田庸子、中野正明、成田一衛

全身性エリテマトーデスの初回ステロイド投与における大腿骨頭壊死症の発生に関する背景因子の検討.

中部リウマチ（印刷中）

- Kuroda T, Wada Y, Nakano M.

Diagnosis and treatment of AA amyloidosis with rheumatoid arthritis: State of the art.

In Dali Feng (ed), Amyloidosis, In Tech. Croatia, 2013, 187-212.

研究分担者：竹内 勤

- Mok CC, Yap DY, Navarra SV, Liu Z-H, Zao M-H, Lu L, Takeuchi T, Avihingsanon Y, Yu X-Q, Lapid EA, Lugue-Lizardo LR, Smethkul V, Shen N, Chen S-L, and Chan TM, for the Asian Lupus Nephritis Network Overview of lupus nephritis management guidelines and perspective from Asia. Nephrology, 2014;19:11-20.
- Takeuchi T, Matsubara T, Ohta S, Mukai M, Amano K, Tohma S, Tanaka Y, Namanaka H, Miyasaka N. Biologic-free remission of established rheumatoid arthritis after discontinuation of abatacept: a prospective multi-center, observational study in Japan. Rheumatology, online September 24, 2014.
- Takeuchi T, Suzuki K. CD247 variants and single-nucleotide polymorphisms observed in systemic lupus erythematosus patients. Rheumatology (Oxford). 2013;52(9):1551-1555.
- Takeuchi T, Suzuki K, Kondo T, Yoshimoto K, Tsuzaka K. CD3 z defects in systemic lupus erythematosus. Ann Rheum Dis, 2012 ; 71:i78-i81.
- 倉沢隆彦、鈴木勝也、竹内 勤
特集 自己免疫疾患の最近の話題と展望 マイクロアレイを用いた自己免疫疾患の解析
Pharma Medica, 2013: vol. 31 No. 1 メディカルレビュー

研究分担者：多田芳史

- Koarada S, Sadanaga Y, Nagao N, Tashiro S, Suematsu R, Ohta A, Tada Y. In Vitro Analysis of Proliferating CD4+ T Cells in Patients with Rheumatoid Arthritis as a Possible Marker of TNF-alpha Blockers. Clinical Anti-Inflammatory & Anti-Allergy Drugs, 2014; 1:1-7.
- Washio M, Fujii T, Kuwana M, Kawaguchi Y, Mimori A, Horiuchi T, Tada Y, Takahashi H, Mimori T, and Japan MCTD Study group. Lifestyle and other related factors for the development of mixed connective tissue disease among Japanese females in comparison with systemic lupus erythematosus. Mod Rheumatol, 2014:24:788-792.
- Kiyohara C, Washio M, Horiuchi T, Asami T, Ide S, Atsumi T, Kobashi G, Takahashi H, Tada Y, and The Kyusyu Sapporo SLE (KYSS) STUDY GROUP.

- Modifying effect of N-acetyltransferase 2 genotype on the association between systemic lupus erythematosus and consumption of alcohol and caffeine-rich beverages.
Arthritis Care Res, 2014; 66:1049–1056.
- Takahashi H, Washio M, Kiyohara C, Tada Y, Asami T, Ide S, Atsumi T, Kobashi G, Yamamoto M, Horiuchi T; Kyushu Sapporo SLE (Kyss) Study Group.
 Psychological stress in a Japanese population with systemic lupus erythematosus: finding from KYSS study.
Mod Rheumatol, 2014;24:448–452.
 - Ono N, Niiro H, Ueda Ak, Sawabe T, Nishizaka H, Furugo I, Yoshizawa S, Yoshizawa S, Tsukamoto H, Kiyohara C, Tada Y, Horiuchi T.
 Characteristics of MPO-ANCA-positive granulomatosis with polyangiitis: a retrospective multi-center study in Japan.
Rheumatol Int, 2014; ePub ahead of print.
 - Koarada S, Tashiro S, Tokuda Y, et al.
 Subsets of RP105-negative plasmablasts in IgG4-related disease.
Ann Rheum Dis, 2014;73: e65.
 - Kumano T, Ishibashi H, Kubota Y, Ono Y, Tada Y, Nagasawa K, et al.
 Adult-onset Still's disease in an elderly woman with atypical presentation.
Case Report in Internal Medicine, 2014;1:158–162.
 - Koarada S, Tashiro S, Nagao N, Suematsu R, Ohta A, Tada Y.
 Increased RP105-negative B cells in IgG4-related disease.
The Open Rheumatology Journal, 2013;7:55–57.
 - Koarada S, Tada Y.
 Autoantibody-producing RP105-negative B cells in human and a murine model of lupus erythematosus.
 Systemic lupus erythematosus (SLE): Prevalence, pathophysiology and prognosis.
 Koarada S (Eds) Nova Science publishers Inc, 2013:159–166.
 - Koarada S, Sadanaga Y, Nagao N, Tashiro S, Suematsu R, Tada Y. et al.
 Illustrated overview of the prevalence and clinical symptoms of systemic lupus erythematosus.
 Systemic lupus erythematosus (SLE): Prevalence, pathophysiology and prognosis.
 Koarada S (Eds) Nova Science publishers Inc, 2013:1–48.
 - Kiyohara C, Washio M, Horiuchi T, Asami T, Ide S, Atsumi T, et al.
 Risk modification by CYP1A1 and GSTM1 polymorphisms in the association of cigarette smoking and systemic lupus erythematosus in a Japanese population.
Scand J Rheumatol, 2012;41:103–109.
 - Kiyohara C, Washio M, Horiuchi T, Asami T, Ide S, Atsumi T, et al.
 Cigarette smoking, alcohol consumption, and risk of systemic lupus erythematosus: a case-control study in a Japanese population.
J Rheumatol, 2012;39:1363–1370.

- Tada Y, Suematsu E, Ueda A, Nagano S, Sawabe T, Nishizaka H, et al. Clinical factors to predict a poor prognosis and refractory disease in patients with polymyositis and dermatomyositis associated with interstitial lung disease. Clin Exp Rheumatol, 2012;30:450.
- Koarada S, Tada Y, Suematsu R, Soejima S, Inoue H, Ohta A, et al. Phenotyping of P105-negative B cell subsets in patients with systemic lupus erythematosus. Clin Dev Immunol, 2012;2012:198206.
- Koarada S, Tada Y. RP105-negative B cells in systemic lupus erythematosus. Clin Dev Immunol, 2012;2012:259186.
- 小荒田秀一、多田芳史
Polyangiitis overlap 症候群
日本臨床新領域別症候群シリーズ, 2014 : 27:663-666.
- 小野伸之、多田芳史
ANCA 関連血管炎の最新の診断と治療 -治癒を目指した治療へ-
九州リウマチ, 2014;34:73-78.
- 永尾奈津美、貞永裕梨、末松梨絵、小荒田秀一、田代知子、小野伸之、多田芳史、大田明英
本邦における強皮症に続発した顕微鏡的多発血管炎の検討.
九州リウマチ, 2014 ; 34:86-92.
- 多田芳史
全身性エリトマトーデスとインターフェロン制御因子.
臨床リウマチ, 2014 ; 26:70-72.
- 杉 元子、末松梨絵、小荒田秀一、大田明英、多田芳史
医療秘書における臨床研究支援業務.
医療秘書実務論集, 2014; 4:63-68.
- 杉 元子、末松梨絵、小荒田秀一、大田明英、多田芳史
医療秘書におけるリウマチ薬処方実態調査.
Medical secretary : 日本医療秘書機関誌, 2014;11(2) : 17-19.
- 多田芳史
膠原病患者の下肢病変. 総論.
下肢救済マニュアル pp200-203, 秀潤社、東京、2014.
- 多田芳史
膠原病患者の下肢病変. 膠原病患者の下肢病変と治療.
下肢救済マニュアル. pp204-211, 秀潤社、東京、2014.
- 多田芳史
全身性エリテマトーデスとIRF5.
臨床免疫・アレルギー科, 2013;59:104-109.
- 近江雅代、鷲尾昌一、堀内孝彦、塚本 浩、多田芳史、他

- 全身性エリトマトーデス発症に関連する食事因子～栄養素等摂取状況および食品群別摂取について～
 日本病態栄養学会誌, 2013; 16: 99–106.
- 多田芳史
 成人Still病に対するアナキンラの効果
 リウマチ科 2013;50:233-237.
 - 本池 悠、田代知子、永尾奈津美、末松梨絵、小荒田秀一、多田芳史、他
 後腹膜線維症の所見を呈した洞組織球症の1例
 九州リウマチ, 2013;33:119–124.
 - 水田和孝、末松梨絵、貞永裕梨、永尾奈津美、田代知子、多田芳史、他
 後腹膜神経鞘腫を合併したRS3PE症候群の一例
 九州リウマチ, 2013;33:108–113.
 - 多田芳史
 成人スタイル病 免疫疾患イラストレイテッド 田中良哉編集 羊土社、東京、2013:127–130.
 - 小荒田秀一、田代知子、末松梨絵、井上久子、大田明英、多田芳史
 自己抗体産生RP105陰性B細胞上に発現するB-Cell maturation antigenは、全身性エリテマトーデスの治療標的となりうるか。
 日臨免誌, 2012;35:38–45.
 - 近江雅代、鶯尾昌一、堀内孝彦、塙本 浩、赤司浩一、多田芳史、他
 全身性エリトマトーデス発症に関連する食事因子。
 臨床と研究, 2012;89:74–78.
 - 多田芳史
 免疫抑制薬の臨床応用実践論 RS3PE症候群。
 炎症と免疫, 2012;20:523–527.

研究分担者：田中良哉

- Kondo M, Yamaoka K, Sakata K, Sonomoto K, Lin L, Nakano K, Tanaka Y.
 IL-6/STAT3 signaling pathway contributes to chondrogenic differentiation of human mesenchymal stem cells.
Arthritis Rheum (in press).
- Iwata S, Nakayamada S, Fukuyo S, Kubo S, Yunoue N, Wang S-P, Yoshikawa M, Saito K, Tanaka Y.
 Activation of Syk in peripheral blood B cells in patients with rheumatoid arthritis: A potential target for abatacept therapy.
Arthritis Rheum (in press).
- Tanaka Y, Hirata S, Kubo S, Fukuyo S, Hanami K, Sawamukai N, Nakano K, Nakayamada S, Yamaoka K, Sawamura F, Saito K.
 Discontinuation of adalimumab after achieving remission in patients with established rheumatoid arthritis: 1-year outcome of the HONOR study.
Ann Rheum Dis, 2013 Nov 28. doi: 10.1136/annrheumdis-2013-204016.
- Zhang X, Yamaoka K, Sonomoto K, Kaneko H, Satake M, Yamamoto Y, Kondo M, Zhao J, Miyagawa I,

- Yamagata K, Fukuyo S, Okada Y, Tanaka Y.
 Local delivery of mesenchymal stem cells with poly-lactic-co-glycolic acid nano-fiber scaffold suppress arthritis in rats.
Plos ONE, 2014;9(12):e114621.
- Kubo S, Yamaoka K, Kondo M, Yamagata K, Zhao J, Iwata S, Tanaka Y.
 The JAK inhibitor tofacitinib reduces the T cell stimulatory capacity of human monocyte-derived dendritic cells.
Ann Rheum Dis, 2014;73: 2192–2198.
 - Tanaka Y, Martin Mola E.
 Can IL-6-targeting catch up TNF-targeting in rheumatoid arthritis: from studies of olokizumab, sarilumab and sirukumab.
Ann Rheum Dis, 2014;73:1395–1397.
 - Wang S-P, Iwata S, Nakayamada S, Sakata K, Yamaoka K, Tanaka Y.
 Tofacitinib, a Jak inhibitor, inhibits human B cell activation in vitro.
Ann Rheum Dis, 2014;73:2213–2215.
 - Kondo M, Yamaoka K, Tanaka Y.
 Acquiring chondrocyte phenotype from human mesenchymal stem cells under inflammatory conditions.
Int J Mol Sci, 2014;15:21270–21285.
 - Fukuyo S, Yamaoka K, Sonomoto K, Oshita K, Okada Y, Saito K, Yoshida Y, Kanazawa T, Minami Y, Tanaka Y.
 IL-6-accelerated calcification by induction of ROR2 in human adipose tissue-derived mesenchymal stem cells is STAT3-dependent.
Rheumatology, 2014;53:1282–1290.
 - Tanaka Y.
 Next stage of RA treatment: TNF-inhibitor-free remission will be a possible treatment goal?
Ann Rheum Dis, 2013;72:ii124–ii127.
 - Kondo M, Yamaoka K, Sonomoto K, Fukuyo S, Oshita K, Okada Y, Tanaka Y.
 IL-17 inhibits chondrogenic differentiation of human mesenchymal stem cells.
Plos ONE, 2013;8 (11): e79463.
 - Iwata S, Saito K, Tokunaga M, Tanaka Y.
 B-cell or T-cell-dominant recurrence after rituximab therapy in patients with SLE.
Ann Rheum Dis, 2012;71:1749–1750.
 - Tanaka Y, Maeshima Y, Yamaoka K.
 In vitro and in vivo analysis of a Jak inhibitor in rheumatoid arthritis.
Ann Rheum Dis, 2012;71:i70–i74.
 - Maeshima K, Yamaoka K, Kubo S, Nakano K, Iwata S, Saito K, Ohishi M, Miyahara H, Tanaka S, Ishi K, Yoshimatsu H, Tanaka Y.
 A JAK inhibitor tofacitinib regulates synovitis through inhibition of IFN- γ and IL-17 production by

human CD4+ T cells.

Arthritis Rheum, 2012;64:1790–1798.

- Sonomoto K, Yamaoka K, Oshita K, Fukuyo S, Zhang X, Nakano K, Okada Y, Tanaka Y. IL-1 β induces differentiation of human mesenchymal stem cells into osteoblasts via the Wnt5a/Ror2 pathway.
Arthritis Rheum, 2012;64:3353–3363.

研究分担者：山路 健

- Gorai M, Ogasawara M, Matsuki Y, Yamada Y, Murayama G, Sugisaki N, Nemoto T, Ando S, Minowa K, Kon T, Tada K, Matsushita M, Yamaji K, Tamura N, Takasaki Y. Weighting with the Lansbury articular index improves the correlation of ultrasound score with serum matrix metalloproteinase-3 level in rheumatoid arthritis patients. Mod Rheumatol, 2014;24(6): 915–919.
- Doe K, Nozawa K, Okada T, Tada K, Yamaji K, Tamura N, Takasaki Y. Usefulness of minor salivary gland biopsy in the diagnosis of IgG4-related disease: a case report. Int J Clin Exp Pathol, 2014;15: 7(5): 2673–7.
- Murayama G, Ogasawara M, Matsushita M, Yamaji K, Tamura N, Takasaki Y, et al. Clinical miscount of involved joints denotes the need for ultrasound complementation in usual practice for patients with rheumatoid arthritis. Clin Exp Rheumatol, 2013;31(4):506–514.
- Inami Y, Yamaji K, Gohda T, Io H, Takasaki Y, Tomino Y, et al. Effects of dialysis on the pharmacokinetics of salazosulfapyridine. Rheumatol Int, 2013;33(2):535–539.
- Yamada R, Sekiya F, Kusaoi M, Yamaji K, Tsuda H, Takasaki Y, et al. Reduction in bradykinin generation during leukocytapheresis using novel cellsorbaTM CS-180S: Effects of changing the filling solution. J Clin Apher, 2013 [Epub ahead of print]
- Nemoto T, Ogasawara M, Sekiya F, Yamaji K, Tamura N, Takasaki Y, et al. Can routine clinical measures predict ultrasound-determined synovitis and remission in rheumatoid arthritis patients? Clin Exp Rheumatol, 2013 Oct 17. [Epub ahead of print]
- Kusaoi M, Yamaji K, Ogasawara M, Sekiya F, Tsuda H, Takasaki Y, et al. Gene expression analysis using a high-resolution DNA microarray of peripheral whole blood immediately before and after leukocytapheresis for rheumatoid arthritis. Ther Apher Dial, 2012;16(5):456–466.
- Ogasawara M, Kageyama M, Kusaoi M, Yamaji K, Tamura N, Takasaki Y, et al. Recent trends in the use of non-biologic DMARDs and an evaluation of their continuation rates in single and dual combination therapies in rheumatoid arthritis patients in Japan. Mod Rheumatol, 2012;22(6):831–836.