

**Figure 1** Phenotypic stratification of early-onset collagen VI-related myopathies; modification from Quijano-Roy *et al*<sup>10</sup> and Briñas *et al*<sup>11</sup> (A). Typical distal hyperlaxity and protruding calcanei in a patient with Ullrich congenital muscular dystrophy (UCMD) (B,C). Diagnostic muscle MRI findings of the thigh (D): fatty regeneration along the fascia in the centre of the rectus femoris (arrow) and in the periphery of the vastus lateralis with relative sparing of its central part (arrowheads). Early-onset and rapidly progressive scoliosis in UCMD (E–H). E, 3.7; F, 6.1; G, 6.9; H, 7.9 years of age at assessment. (Images courtesy of Drs Ikuya Nonaka and Akihiko Ishiyama).

(kyphoscoliosis) and ankles. Arthrogryposis multiplex are seen in 20.0% of patients.<sup>19</sup> Some transient feeding difficulties or poor sucking might occur in the neonatal period.<sup>19 20</sup>

The most common presentations are delayed motor milestone and proximal muscle weakness. Patients with UCMD usually become able to sit independently with or without delay.<sup>19</sup> The majority of patients with typical UCMD achieve the ability to walk by 2 years of age. The age at independent ambulation is reported to be  $1.7 \pm 0.8$  and  $1.7 \pm 0.5$  years in English and Japanese natural history studies, respectively.<sup>19 20</sup> This group is referred to as having ‘moderate progressive’ disease in the phenotypic stratification of early-onset collagen VI-related myopathy (figure 1A).<sup>10 11</sup> In the remaining patients, ambulation is never achieved. Those with the most severe presentation, accounting for 19.4–25.7% of UCMD patients,<sup>11 19</sup> are referred to as having ‘early-severe’ disease (figure 1A). However, even these severely affected children can usually learn to roll, crawl and maintain a sitting position. Patients who never walk due to severe contractures that prevent an upright posture may walk on their knees for a certain period of time. Achievement of speaking phrases is not delayed, ranging from 12 to 25 months of age.<sup>19</sup>

### Clinical manifestations

Patients show generalised muscle weakness predominantly in trunk and proximal limbs. Neck flexors are weak. Facial weakness is reported respectively in 24.1% and 30.8% of patients in two different studies.<sup>19 20</sup> The most striking feature is hyperlaxity of the distal joints (figure 1B), although it can be absent in severely affected individuals. Spinal rigidity, scoliosis and various proximal joint contractures develop with progression of the disease. Typically, initially flexible distal joints, such as fingers, wrists and ankles, eventually become contractured with time. Interestingly, calcanei are often protruded posteriorly (figure 1C). Distal joint hyperlaxity, proximal joint contractures, scoliosis and protruding calcaneus are observed in >50% patients in a Japanese cohort.<sup>19</sup> Many patients have a characteristic facial appearance with round-shaped face with slight drooping of the lower lid and prominent ears.<sup>15</sup> Intelligence is normal. Other features include follicular hyperkeratosis over the extensor surfaces of upper and lower limbs, a softer consistency of the skin in the palms and soles and the tendency to keloid formation.<sup>15</sup> Respiratory insufficiency is not common at birth, although it becomes a critical complication of the disease as the condition progresses. Cardiac involvement is not documented to date.<sup>21 22</sup>



## Neuromuscular

Serum creatine kinase (CK) activity in patients with UCMD is usually within normal range or only mildly elevated,<sup>19–21</sup> which is unusual in other (congenital) muscular dystrophies. Electromyography shows action potentials of low amplitude and short duration.<sup>13</sup> Muscle MRI shows a characteristic pattern on transverse T1-weighted images—diffuse involvement of the thigh muscles with relative sparing of the medial muscles (sartorius, gracilis and adductor longus).<sup>23</sup> Rectus femoris is variably involved with a typical central area of high signal, called ‘central shadow’.<sup>23</sup> In vastus lateralis, the peripheral part is mainly involved and signal intensity is markedly increased while the central part is relatively spared (figure 1D).<sup>23</sup>

### Muscle pathology and collagen VI immunohistochemistry

Variable degrees of histological changes can be observed in muscle biopsies from patients with UCMD. The spectrum includes fibre size variation affecting both fast and slow fibres, type 1 fibre predominance, increased endomysial connective tissue or adipose tissue, increased number of internal nuclei and mild necrotic and regenerating process along with indirect evidence of regenerating fibres such as the presence of fibres containing fetal myosin.<sup>21–24</sup> One report described that, early in the disease, UCMD presents as a non-dystrophic myopathy with predominant fibre atrophy, showing a bimodal size distribution of type 1 fibres or a diagnostic pattern of congenital fibre-type disproportion.<sup>25</sup> Unlike other muscular dystrophies, interstitial fibrosis seems disproportionately prominent considering the relative paucity of necrotic and regenerating fibres in UCMD.

Collagen VI is widely distributed throughout ECMs in various tissues. In skeletal muscle, collagen VI is found in the epimysial, perimysial and endomysial interstitium, but it is concentrated in particular in and adjacent to basement membranes of myofibers, blood vessels and intramuscular nerves. Muscle biopsies from UCMD patients can show anything from mild reduction of endomysial or basal lamina collagen VI staining to complete deficiency (CD) of collagen VI in the ECM. We previously showed that, in the majority of patients with UCMD, collagen VI is present in the interstitium but is absent from the sarcolemma by using double immunostaining for collagen IV and VI (figure 2), and named it ‘sarcolemma specific collagen VI deficiency (SSCD)’.<sup>26</sup> Electron microscopic findings support a lack of connection between collagen VI microfibrils in the interstitium and the basal lamina, leading to SSCD.<sup>26</sup> Space is observed between muscle fibres and connective tissue that are normally closely attached. Basal lamina appear intact even in degenerating muscle fibres with disorganised myofibrils. These findings suggest a loose connection between the basal lamina and other ECM collagens in UCMD.<sup>27</sup> Collagen VI is deficient also in capillaries in muscle.<sup>13</sup> The absence or alteration of collagen VI can be demonstrated by immunocytochemistry of cultured skin fibroblasts, although this analysis is available only in limited laboratories. In skin, collagen VI expression is decreased in the papillary dermis and skin hair follicles, but not in vessels, peripheral nerves, smooth muscle and sweat glands.<sup>13</sup> One report described that collagen VI levels were greatly decreased in peripheral blood macrophages from three patients with UCMD.<sup>28</sup>

### Natural history of disease

Muscle weakness is slowly progressive. Most affected children become able to walk independently but eventually lose ambulation often by early teenage years. Loss of ambulation is reported to occur at  $10.7 \pm 4.8$ ,  $10.1 \pm 4.4$  and  $8.8 \pm 2.9$  years of age in English, French and Japanese group of patients with UCMD,

respectively.<sup>11 19 20</sup> However, this can be widely variable: some patients never walk while others can still walk even beyond late teens. After loss of ambulation, progression of muscle weakness becomes less prominent. In contrast, the contractures can still be progressive, particularly in the ankles, knees, hips and elbows, aggravating physical disability. These clinical features may well be in line with strikingly progressive interstitial fibrosis on muscle pathology.


Respiratory insufficiency usually occurs after the loss of ambulation. It is noteworthy, however, some patients have impending respiratory dysfunction while they are still ambulant. Progressive decline in the predicted forced vital capacity (FVC) or vital capacity (VC) is observed from the preschool age to the early teens.<sup>19 20</sup> Of note, restrictive respiratory dysfunction develops rapidly in the first decade of life; indeed, %predicted FVC declines by  $6.6 \pm 1.9\%$ /year from 6 to 10 years of age compared to  $0.4 \pm 3.0\%$ /year from 11 to 15 years of age.<sup>20</sup> Similarly, VC declines exponentially with a sharp decrease by 10 years of age.<sup>19</sup> This may well be associated with proximal joint and vertebral contractures together with weakness of the diaphragm. The introduction of non-invasive ventilation (NIV) is usually sufficient to treat this situation effectively for many years. The percentage of patients with NIV increases with age; half require NIV by age 11–12 years.<sup>12 19</sup> Natural history study from UK reported that age at initiation of NIV was  $14.3 \pm 4.7$  years, with a mean FVC of 20%.<sup>20</sup> The other two studies have recently reported similar findings: an estimated predicted VC of 36% at the time of initiation of NIV at  $11.2 \pm 3.6$  years in a Japanese cohort<sup>19</sup> and an average FVC of 34% just before NIV initiation at  $11.3 \pm 4.0$  years in a large international cohort,<sup>12</sup> demonstrating a remarkable consistency of the pulmonary function declining in patients with UCMD among different cohorts, regardless of different approaches to data acquisition.

Scoliosis, which may require surgical correction, is a common complication.<sup>15 20</sup> Substantial scoliosis appears as early as preschool years and its onset precedes loss of ambulation.<sup>19 20</sup> Development of scoliosis in Duchenne muscular dystrophy (DMD) is strongly related to the loss of walking ability—scoliosis is not typically evident in ambulatory patients and develops after they become wheelchair dependent. In contrast, in UCMD, scoliosis develops even when patients are still ambulant and is progressive from early stage. In our cohort, a maximum progression rate of Cobb angle was  $16.2 \pm 10.0^\circ$ /year.<sup>19</sup> Importantly, scoliosis progresses rapidly within years, once it starts (figure 1E–H). The early-onset and rapidly progressive scoliosis in UCMD may well accelerate physical disability, such as difficulty sitting, standing and walking, and cause pain. More importantly, scoliosis can aggravate respiratory function by reducing the rib cage compliance in combination with other proximal joint contractures.

### Differential diagnosis

Differential diagnosis includes wide range of neuromuscular disorders in infants and children, such as other forms of muscular dystrophy, congenital myopathies, spinal muscular atrophy (SMA), especially type 2 and 3, and other diseases of connective tissue such as Ehlers–Danlos syndrome (EDS). However, combination of normal or minimally elevated CK levels, lack of cardiac manifestation and specific pattern of thigh muscle involvement is often suggestive of UCMD. Brain MRI is usually normal unlike other CMD forms such as MDC1A, Walker–Warburg syndrome, muscle-eye-brain disease and Fukuyama CMD, which may show structural abnormalities or white matter





**Figure 2** Double immunostaining for collagen IV and VI. (A–C) Patient with non-diagnostic muscle pathology; (D–F) UCMD patient with CD; (G–I) UCMD patient with SSCD. (A,D,G) Immunostaining for collagen IV. Collagen IV is present in the sarcolemma. (B,E,H) Immunostaining for collagen VI. Collagen VI is absent in a patient with CD, while it is present in the interstitium but markedly reduced in a patient with SSCD. (C,F,I) Merged images. Both collagen IV and VI are present in the sarcolemma in a patient with non-diagnostic pathology, as indicated by yellow; in contrast, collagen VI is absent in the sarcolemma in a patient with SSCD, as indicated by only red. UCMD, Ullrich congenital muscular dystrophy; CD, complete collagen VI deficiency; SSCD, sarcolemma-specific collagen VI deficiency, bar 20  $\mu$ m.

changes. *RYR1*-related core myopathy, including central core disease (CCD) and multimincore disease, may also show similar clinical features as they can multiple joint contractures and spinal deformity. Progressive respiratory muscle involvement, often disproportionate to the skeletal muscle weakness, is also seen in multimincore disease. However, significant respiratory insufficiency is unusual in CCD. Interestingly, a recent report demonstrated that muscle pathology in a patient with a heterozygous *COL6A3* gene mutation included cores, rods and lobulated fibres.<sup>29</sup> SMA is characterised by fasciculation and diagnosed by identifying mutations in the *SMN* gene. EDS may mimic UCMD in terms of joint laxity. Furthermore, various types of EDS are commonly associated with mild to moderate muscle weakness. Muscle pathology can demonstrate mild myopathic features but necrotic fibres and fibrosis are absent and collagen VI staining is normal.<sup>30</sup>

Of particular importance is rigid spine with muscular dystrophy type 1 (RSMD1) in forms of CMD, which results from

mutations in the *SEPN1* gene, because it can show a significant clinical overlap in the late stage of the disease. RSMD1 patients typically show combination of mild or moderate proximal muscle weakness, Achilles tendon tightness, spinal rigidity and scoliosis, and require ventilator assistance in the first decade of life,<sup>15</sup> similarly to UCMD.

### Management

Treatment for patients with UCMD is supportive for symptoms such as respiratory insufficiency and scoliosis, and is dependent on age of the individual and severity of symptoms. Respiratory failure is a common complication of UCMD and careful follow-up with regular assessments of respiratory function, including spirometry and nocturnal oximetry studies, is important to detect asymptomatic decline in patients. In a recent review based on expert consensus on the standard care for CMD, cough assistance using mechanical insufflation-exsufflation is generally accepted as the method to improve cough efficiency.<sup>22</sup> Other methods such as



## Neuromuscular

breath stacking with Ambu bag maintain thoracic compliance and reduce the risk of chronic atelectasis.<sup>22</sup> Respiratory support with nocturnal NIV usually becomes necessary in the first or second decade of life for patients<sup>12 19</sup> and might be effective in reducing symptoms and promoting quality of life. There are times when chronic ventilation can require an invasive application via tracheostomy.

For scoliosis, conservative management including standing frame, positioning and bracing is widely used, although controversial whether those approaches are preventive.<sup>22</sup> However, scoliosis may require active management including spinal surgery to prevent progression, although there have been no formal studies on the efficacy of scoliosis surgery. The choice of the instrumentation such as growing rods depends on the age of the individual, his/her ability to grow and the severity of scoliosis.<sup>22</sup> Suggested contraindications to spinal surgery includes family decision, very poor or deteriorating cardiac status and/or respiratory status, very young age, potential loss of function after spinal fixation and severe scoliosis.<sup>22</sup> Severe respiratory insufficiency may not be a contraindication in certain high specialised centres. In fact, one study on surgical correction of spinal deformity from Japan reported that scoliosis surgery was successfully performed in three patients with UCMD at 11, 13 and 17 years of age, respectively<sup>31</sup>; spinal surgery, however, did not prevent deterioration of respiratory function in these patients, suggesting that at such older ages pulmonary and chest wall compliance might be too severely compromised for patients to benefit from scoliosis surgery, and earlier surgical intervention may be more beneficial. Indeed, a single case report described that slower decline of predicted VC in a patient after scoliosis surgery performed at 5 years of age compared with another patient who had undergone surgical correction of scoliosis at 9 years of age.<sup>19</sup> Further studies are necessary to conclude the efficacy of early scoliosis surgery.

Equipment recommended for assistance in standing, ambulation and/or other forms includes walking frames, standing frames, swivel walker, knee-ankle-foot orthoses, ankle-foot orthoses, scooters and wheelchairs.<sup>22</sup> The joint contractures of patients with UCMD in particular seem to be progressive and regular stretching is recommended to maintain a certain level of mobility of the joints. In addition, feeding and swallowing difficulties can be encountered in UCMD.<sup>20</sup> Issues of feeding and nutrition are multifactorial and closely related to other areas of care; for example, nocturnal hypoventilation can affect appetite and growth and respiratory insufficiency can result in easy fatigue and difficulty in swallowing.<sup>22</sup> Consultation with a nutrition specialist is often helpful to boost energy intake. Some children may need a temporary or permanent gastric feeding tube support to maintain an adequate nutritional and fluid intake.<sup>20 21</sup> Survival has not been fully documented under the current standards of medical care, but failure to introduce adequate respiratory support might lead to the death of teenagers with UCMD.<sup>11 21</sup> With the availability of effective respiratory interventions, patients commonly survive into adulthood to date, and other potential aspects of the disease could surface.

### MOLECULAR DIAGNOSIS, PATHOGENESIS AND THERAPEUTIC AVENUES

Collagen VI is a ubiquitously expressed ECM protein composed of three  $\alpha$ -chains,  $\alpha 1$ ,  $\alpha 2$  and  $\alpha 3$ . The three chains are encoded by the genes *COL6A1* and *COL6A2* on chromosome 21q22.3 and *COL6A3* on chromosome 2q37.<sup>13</sup> The basic monomer, made up of two globular domains connected by a triple helical structure, is composed of one of each of the these  $\alpha$ -chain

subunits (1:1:1 ratio). Prior to secretion into the extracellular space, the two basic monomers assemble into dimers (two anti-parallel, overlapping monomers) and such dimers associate in a staggered parallel orientation to form tetramers (four monomers) in the cytoplasm.<sup>13 32</sup> Outside the cell, these tetramers, the secreted form of collagen VI, associate in an end-to-end fashion to form collagen VI microfibrillar structures, which interacts with collagen IV and other ECM components including proteoglycans decorin and biglycan, collagen I, hyaluronan, heparin and integrin.<sup>13 33</sup>

### Collagen VI gene mutations in UCMD

UCMD used to be regarded as an autosomal-recessive disorder. However, soon after the initial discovery of recessive mutations in the *COL6A2* gene,<sup>34 35</sup> a total of four patients with sporadic UCMD were found to carry *de novo* autosomal-dominant mutations in either *COL6A1*, *COL6A2* or *COL6A3*.<sup>36 37</sup> UCMD is now known to be caused by either recessive or dominant genetic mechanism, the latter most typically occur as *de novo* mutations.<sup>32</sup> This is most likely because these dominant mutations are associated with too severe phenotype to allow patients to produce their offspring. In contrast, in BM, the phenotype is typically mild enough for patients to produce children, resulting in autosomal-dominant inheritance condition.<sup>8 13</sup>

The most common types of mutations are point mutations, exon skipping and mutations leading to premature termination codons (PTCs).<sup>32</sup> Among point mutations, missense changes affecting glycine residues in the Gly-Xaa-Yaa motifs of the N-terminal triple helical domains are the most common and are often dominant *de novo*.<sup>32 33</sup> Splice mutations resulting in in-frame exon skipping are generally dominant *de novo* mutations.<sup>32 33</sup> These dominant mutations can result in secretion of some mutant-containing tetramers into the extracellular space; in the multistep collagen VI intracellular assembly process, only 1/16 of the tetramers produced by patients with dominant mutations could be composed entirely of normal  $\alpha$ -chains, thus exerting a dominant negative effect.<sup>37</sup> This leads to loss of normal localisation of collagen VI in the basement membrane and eventually results in a severe phenotype. Nonsense mutations and small deletions or insertions inducing PTCs with consequent nonsense-mediated mRNA decay (NMD), an mRNA quality control mechanism that degrades aberrant mRNAs containing PTCs, and loss of the mutated chain are mostly inherited as recessive mutations.<sup>32 33</sup> Patients with these mutations are unable to assemble or secrete functional collagen VI protein, as all three  $\alpha$ -chains are required to form a collagen VI monomer. Thus, such functional null alleles, which underlie typical UCMD, mostly lead to CD mode of collagen VI in skeletal muscles.<sup>16 26 33 34</sup> On the other hand, complete deletions of one copy of these genes also act in a recessive fashion. In support of this notion, carriers of the deletion are in fact clinically asymptomatic, indicating that complete haploinsufficiency of any of the three collagen VI genes does not cause the disease. Interestingly, two reports demonstrated that autosomal-recessive inheritance can also underlie BM, in which patients carried a truncating or null *COL6A2* mutation associated with missense changes in the partnering allele lying within the C2 domain of the  $\alpha 2$  chain.<sup>38 39</sup> Furthermore, myosclerosis syndrome was reported to be responsible for a homozygous nonsense *COL6A2* mutation.<sup>40</sup> Unlike nonsense mutations associated with UCMD, the mutated mRNA escaped NMD and was translated into a truncated  $\alpha 2$  chain, but secreted collagen VI was reduced and structurally abnormal and thus did not correctly localise in the basement membrane of myofibers. These facts suggest that the



severity of collagen VI gene mutations and the resulting functional abnormality of collagen VI in the ECM dictate a phenotypic spectrum of collagen VI-related myopathies, meaning that a fundamentally different genetic and biochemical mechanism among these myopathies can no longer be assumed.

We previously showed that there are two modes of collagen VI deficiency, CD and SSCD,<sup>26</sup> which respectively result from recessive and *de novo* dominant mutations in the collagen VI genes.<sup>16</sup> There is no straightforward correlation between protein levels and phenotypes; CD, however, is most likely to be associated with the more severe phenotype than SSCD.<sup>11 19</sup> Unlike patients with CD, a great heterogeneity in the maximal motor capacity was observed in patients with SSCD, ranging from no acquisition of walking ability to retaining ambulation throughout childhood.

### Properties of collagen VI and its pathological roles

Collagen VI is widely distributed throughout ECMs in various tissues, including muscle, skin, tendon, cartilage, intervertebral discs, lung, blood vessels and adipose tissue.<sup>4</sup> Given the clinical features seen in patients with collagen VI-related myopathies, the tissues in which collagen VI has the most important roles include muscle and tendon. In muscle, the cell source producing collagen VI is the interstitial mesenchymal cell.<sup>41 42</sup> In tendons, abundant collagen VI is present in immediate pericellular ECM of the resident tendon fibroblasts.<sup>43</sup>

Collagen VI contributes to the properties of the local ECM microenvironment by forming a discrete network of beaded microfilaments, which interact with a large number of matrix molecules and cell surface receptors. One possible molecule mediating its interaction would be collagen type IV, the most important collagenous component of basement membranes.<sup>13</sup> On the other hand, collagen VI might be indirectly linked to muscle cell surface receptors via biglycan and the dystrophin-associated protein complex, as collagen VI binds to biglycan,<sup>13</sup> which interacts with the sarcoglycan and dystroglycan complex.<sup>33</sup> The functions of collagen VI pertaining to various cell types also include the promotion of adhesion,<sup>1 2</sup> proliferation,<sup>3</sup> migration<sup>4</sup> and survival.<sup>5 44 45</sup> Attachment of cells to the ECM is important for preventing apoptosis,<sup>46</sup> which could be particularly relevant for muscle disorders that directly involve interactions between matrix and muscle, as is the case for high early implanted cell death, partially due to 'anoikis', in cell transplantation treatment of DMD.<sup>47</sup> Studies with cultured fibroblasts from patients with UCMD have shown that mutant cells or mutated collagen VI exhibit decreased adherence to their surroundings, emphasising that loss of cell ECM interactions is the key mechanism of collagen VI-related myopathies.<sup>2 48</sup> Collagen VI also has crucial roles in the regulation and differentiation of adipocytes.<sup>49</sup>

Studies on muscle fibres from *Col6a1*<sup>-/-</sup> mice, engineered by genetic ablation of the *Col6a1* gene,<sup>50</sup> and human myoblast cultures has suggested that collagen VI may be involved in preventing myofiber apoptosis, which seems to be mediated by regulating the mitochondrion-mediated cell death cascade<sup>5 51</sup>; a key event appears to be inappropriate opening of the mitochondrial permeability transition pore. These findings therefore link a defect of the ECM to mitochondrial dysfunction followed by apoptosis that is preventable by inactivation of cyclophilin D by using cyclosporine A, its derivative Debio025 or genetic inactivation of cyclophilin D.<sup>51-53</sup> However, there are contradictory reports in which researchers did not find evidence of myofiber apoptosis in biopsied muscles from UCMD patients or *Col6a3* mutant mice muscles,<sup>54 55</sup> suggesting that muscle cell death by apoptosis is not a universal phenomenon in all patients and collagen VI-deficient mice.

In addition, a study of the autophagic process in muscles of *Col6a1*<sup>-/-</sup> mice revealed that autophagy was not induced efficiently, which determines the presence of dysfunctional organelles in muscle fibres.<sup>56</sup> A similar alteration of autophagy was also detected in muscle biopsies derived from nine patients with UCMD or BM.<sup>56</sup> This defective autophagy provides the link between the previously described mitochondrial dysfunction and myofiber degeneration. These data thus provide a basis for novel therapeutic targets to reactivate of the autophagic flux by either nutritional approaches<sup>57</sup> or by pharmacological and genetic tools in collagen VI deficient skeletal muscle.

### Therapeutic advances

The events responsible for myofiber atrophy and loss might be different in UCMD than in other forms of muscular dystrophy with prominent membrane fragility such as DMD. To date, no single hypothesis can fully explain variation in fibre size, ongoing interstitial fibrogenesis and adipogenesis even in mild necrotic and regenerating process in UCMD or provide all targets for therapies, although important clues have been discovered.

Pathological hypotheses leading to myofiber degeneration in collagen VI-deficient skeletal muscle have been proposed and therapeutic targets have been suggested. There have been pilot studies on patients with UCMD based upon the theory of mitochondrial dysfunction or impaired autophagy.<sup>51-53 57</sup> A recent report has shown that collagen VI is a key component of satellite cell niche and lack of collagen VI causes impaired muscle regeneration and reduced satellite cell self-renewal capability after injury in *Col6a1*<sup>-/-</sup> mice.<sup>58</sup> Additionally, when normal collagen VI is supplied *in vivo* by grafting wild-type interstitial mesenchymal cells, the biochemical properties of collagen VI-deficient muscles are ameliorated and satellite cell defects also rescued.<sup>58</sup> These results can open new venues for a better understanding of the pathomechanism underlying collagen VI-related myopathies. Furthermore, multipotent mesenchymal stem cell (MSC) is the most common type of adult stem cells and is isolated from several sources such as bone marrow and adipose tissue. Another report has recently shown that transplanted human adipose-derived stem cells, with phenotypic and functional features of mesenchymal progenitors, secrete collagen VI protein in *Col6a1*<sup>-/-</sup> mice.<sup>59</sup> Thus, MSC-based therapy can be an attractive option as transplanted cells are able to self-renew and to differentiate into collagen VI-producing cells in skeletal muscle.<sup>41 42</sup>

Advances in molecular genetics provide gene-based therapies; that is, antisense oligonucleotide or small interfering RNA (siRNA) inhibition of mutant transcripts exerting dominant negative effects<sup>60-62</sup> and upregulation of mutant transcripts by specific inhibition of NMD.<sup>63</sup> As 60–80% of UCMD cases are attributed to dominant negative mutations,<sup>11 16</sup> the allele-specific antisense approach can be applied to the majority of patients. Recent reports have shown that siRNA-mediated knockdown of SMG-1 and Upf1, essential components for NMD, or SMG-8, a subunit of SMG-1 kinase, gives rise to the upregulation of mutant triple-helical collagen VI, thus ameliorating mutant phenotypes from UCMD fibroblasts with a homozygous frameshift mutation causing a PTC in the *COL6A2* gene.<sup>64 65</sup>

### CONCLUSION

UCMD is caused by mutations in either *COL6A1*, *COL6A2* or *COL6A3* gene, thereby leading to collagen VI deficiency in the ECM. We here presented the clinicopathological features, robust



natural history and the current supportive care for symptoms. Of special interest is progressive interstitial fibrosis even in very mild necrotic and regenerating process in muscle. Patients with UCMD have unique manifestations attributable to both muscle and connective tissue disorders. Collagen VI contributes to the properties of the local ECM microenvironment by forming a discrete network of beaded microfilaments, which interact with a large number of matrix molecules and cell surface receptors. Advanced researches have provide important clues to explain how collagen VI deficiency in the ECM can cause the development of muscle weakness in *Col6a1*<sup>-/-</sup> mice or patients with UCMD, although the link between the ECM defect and phenotype remains to be fully elucidated. Further studies are necessary to elucidate exactly how collagen VI deficiency in the ECM makes muscle cells vulnerable to apoptosis or interstitial fibrogenesis and adipogenesis strikingly progressive.

**Acknowledgements** The authors thank Drs Ikuya Nonaka, Hirofumi Komaki and Akihiko Ishiyama for their valuable assistance. This study was supported by Intramural Research Grant (23-5) for Neurological and Psychiatric Disorders of NCNP, Research on Rare and Intractable Diseases and Research on Applying Health Technology from the Ministry of Health, Labour and Welfare of Japan.

**Contributors** TY performed literature search and wrote the manuscript; IN was involved in literature search and preparation of the manuscript.

**Competing interests** None.

**Provenance and peer review** Commissioned; externally peer reviewed.

## REFERENCES

- Aumailley M, Specks U, Timpl R. Cell adhesion to type VI collagen. *Biochem Soc Trans* 1991;19:843–7.
- Kawahara G, Okada M, Morone N, et al. Reduced cell anchorage may cause sarcolemma-specific collagen VI deficiency in Ullrich disease. *Neurology* 2007;69:1043–9.
- Atkinson JC, Rühl M, Becker J, et al. Collagen VI regulates normal and transformed mesenchymal cell proliferation in vitro. *Exp Cell Res* 1996;228:283–91.
- Iyengar P, Espina V, Williams TW, et al. Adipocyte-derived collagen VI affects early mammary tumor progression in vivo, demonstrating a critical interaction in the tumor/stroma microenvironment. *J Clin Invest* 2005;115:1163–76.
- Irwin WA, Bergamin N, Sabatelli P, et al. Mitochondrial dysfunction and apoptosis in myopathic mice with collagen VI deficiency. *Nat Genet* 2003;35:367–71.
- Ullrich O. Kongenitale, atonisch-sklerotische Muskeldystrophie, ein weiterer Typus der hereditären Erkrankungen des neuromuskulären Systems. *Z Ges Neurol Psychiatr* 1930;126:171–201.
- Ullrich O. Kongenitale, atonisch-sklerotische Muskeldystrophie. *Monatsschr Kinderheilkd* 1930;47:502–10.
- Bethlem J, van Wijngaarden GK. Benign myopathy, with autosomal dominant inheritance. A report on three pedigrees. *Brain* 1976;99:91–100.
- Butterfield RJ, Foley AR, Dastgir J, et al. Position of glycine substitutions in the triple helix of COL6A1, COL6A2, and COL6A3 is correlated with severity and mode of inheritance in collagen VI myopathy. *Hum Mutat* 2013;34:1558–67.
- Quijano-Roy S, Allamand V, Riahi N, et al. Predictive factors of severity and management of respiratory and orthopaedic complications in 16 Ullrich CMD patients [abstract]. *Neuromuscul Disord* 2007;17:844.
- Briñas L, Richard P, Quijano-Roy S, et al. Early onset collagen VI myopathies: Genetic and clinical correlations. *Ann Neurol* 2010;68:511–20.
- Foley AR, Quijano-Roy S, Collins J, et al. Natural history of pulmonary function in collagen VI-related myopathies. *Brain* 2013;136:3625–33.
- Lampe AK, Bushby KM. Collagen VI related muscle disorders. *J Med Genet* 2005;42:673–85.
- Pepe G, Bertini E, Bonaldo P, et al. Bethlem myopathy (BETHLEM) and Ullrich scleroatonic muscular dystrophy: 100th ENMC International Workshop, 23–24 November 2001, Naarden, the Netherlands. *Neuromuscul Disord* 2002;12:984–93.
- Muntoni F, Voit T. The congenital muscular dystrophies in 2004: a century of exciting progress. *Neuromuscul Disord* 2004;14:635–49.
- Okada M, Kawahara G, Noguchi S, et al. Primary collagen VI deficiency is the second most common congenital muscular dystrophy in Japan. *Neurology* 2007;69:1035–42.
- Peat RA, Smith JM, Compson AG, et al. Diagnosis and etiology of congenital muscular dystrophy. *Neurology* 2008;71:312–21.
- Norwood FL, Harling C, Chinnery PF, et al. Prevalence of genetic muscle disease in Northern England: in depth analysis of a muscle clinic population. *Brain* 2009;132:3175–86.
- Yonekawa T, Komaki H, Okada M, et al. Rapidly progressive scoliosis and respiratory deterioration in Ullrich congenital muscular dystrophy. *J Neurol Neurosurg Psychiatry* 2013;84:982–9.
- Nadeau A, Kinali M, Main M, et al. Natural history of Ullrich congenital muscular dystrophy. *Neurology* 2009;73:25–31.
- Mercuri E, Yuva Y, Brown SC, et al. Collagen VI involvement in Ullrich syndrome: a clinical, genetic, and immunohistochemical study. *Neurology* 2002;58:1354–9.
- Wang CH, Bönnemann CG, Rutkowski A, et al. Consensus statement on standard of care for congenital muscular dystrophies. *J Child Neurol* 2010;25:1559–81.
- Mercuri E, Lampe A, Allsop J, et al. Muscle MRI in Ullrich congenital muscular dystrophy and Bethlem myopathy. *Neuromuscul Disord* 2005;15:303–10.
- Nonaka I, Ue Y, Ishihara T, et al. A clinical and histological study of Ullrich's disease (congenital atonic-sclerotic muscular dystrophy). *Neuropediatrics* 1981;12:197–208.
- Schessl J, Goemans NM, Magold AI, et al. Predominant fiber atrophy and fiber type disproportion in early Ullrich disease. *Muscle Nerve* 2008;38:1184–91.
- Ishikawa H, Sugie K, Murayama K, et al. Ullrich disease due to deficiency of collagen VI in the sarcolemma. *Ann Neurol* 2004;62:620–3.
- Ishikawa H, Sugie K, Murayama K, et al. Ullrich disease: collagen VI deficiency: EM suggests a new basis for muscular weakness. *Neurology* 2002;59:920–3.
- Gualandi F, Curci R, Sabatelli P, et al. Macrophages: a minimally invasive tool for monitoring collagen VI myopathies. *Muscle Nerve* 2011;44:80–4.
- Claeys KG, Schrading S, Bozkurt A, et al. Myopathy with lobulated fibers, cores, and rods caused by a mutation in collagen VI. *Neurology* 2012;79:2288–90.
- Voermans NC, van Alfen N, Pillen S, et al. Neuromuscular involvement in various types of Ehlers-Danlos syndrome. *Ann Neurol* 2009;65:687–97.
- Takaso M, Nakazawa T, Imura T, et al. Surgical correction of spinal deformity in patients with congenital muscular dystrophy. *J Orthop Sci* 2010;15:493–501.
- Allamand V, Briñas L, Richard P, et al. ColVI myopathies: where do we stand, where do we go? *Skelet Muscle* 2011;1:30–42.
- Bönnemann CG. The collagen VI-related myopathies: muscle meets its matrix. *Nat Rev Neurol* 2011;7:379–90.
- Camacho VO, Bertini E, Zhang RZ, et al. Ullrich congenital muscular dystrophy is caused by recessive mutations in collagen type VI. *Proc Natl Acad Sci USA* 2001;98:7516–21.
- Higuchi I, Shiraishi T, Hashiguchi T, et al. Frameshift mutation in the collagen VI gene causes Ullrich's disease. *Ann Neurol* 2001;50:261–5.
- Pan TC, Zhang RZ, Sudano DG, et al. New molecular mechanism for Ullrich congenital muscular dystrophy: a heterozygous in-frame deletion in the COL6A1 gene causes a severe phenotype. *Am J Hum Genet* 2003;73:355–69.
- Baker NL, Morgelin M, Peat R, et al. Dominant collagen VI mutations are a common cause of Ullrich congenital muscular dystrophy. *Hum Mol Genet* 2005;14:279–93.
- Gualandi F, Urciuolo A, Martoni E, et al. Autosomal recessive Bethlem myopathy. *Neurology* 2009;73:1883–91.
- Foley AR, Hu Y, Zou Y, et al. Autosomal recessive inheritance of classic Bethlem myopathy. *Neuromuscul Disord* 2009;19:813–17.
- Merlini L, Martoni E, Grumati P, et al. Autosomal recessive myosclerosis myopathy is a collagen VI disorder. *Neurology* 2008;71:1245–53.
- Zou Y, Zhang RZ, Sabatelli P, et al. Muscle interstitial fibroblasts are the main source of collagen VI synthesis in skeletal muscle: implications for congenital muscular dystrophy types Ullrich and Bethlem. *J Neuropathol Exp Neurol* 2008;67:144–54.
- Braghetta P, Ferrari A, Fabbro C, et al. An enhancer required for transcription of the Col6a1 gene in muscle connective tissue is induced by signals released from muscle cells. *Exp Cell Res* 2008;314:3508–18.
- Ritty TM, Roth R, Heuser JE. Tendon cell array isolation reveals a previously unknown fibrillin-2-containing macromolecular assembly. *Structure* 2003;11:1179–88.
- Rühl M, Sahin E, Johannsen M, et al. Soluble collagen VI drives serum-starved fibroblast through S phase and prevents apoptosis via down-regulation of Bax. *J Biol Chem* 1999;274:34361–8.
- Howell SJ, Doane KJ. Type VI collagen increases cell survival and prevents anti-β1 integrin-mediated apoptosis. *Exp Cell Res* 1998;241:230–41.
- Chiarugi P, Giannoni E. Anoikis: a necessary death program for anchorage-dependent cells. *Biochem Pharmacol* 2008;76:1352–64.
- Gerard C, Forest MA, Beauregard G, et al. Fibrin gel improves the survival of transplanted myoblasts. *Cell Transplant* 2012;21:127–37.
- Kawahara G, Ogawa M, Okada M, et al. Diminished binding of mutated collagen VI to the extracellular matrix surrounding myocytes. *Muscle Nerve* 2008;38:1192–5.
- Nakajima I, Muroya S, Tanabe R, et al. Extracellular matrix development during differentiation into adipocyte with a unique increase in type V and VI collagen. *Biol Cell* 2002;94:197–203.
- Bonaldo P, Braghetta P, Zanetti M, et al. Collagen VI deficiency induces early onset myopathy in the mouse: an animal model for Bethlem myopathy. *Hum Mol Genet* 1998;7:2135–40.
- Bernardi P, Bonaldo P. Dysfunction of mitochondria and sarcoplasmic reticulum in the pathogenesis of collagen VI muscular dystrophies. *Ann NY Acad Sci* 2008;1147:303–11.



- 52 Tiepolo T, Angelin A, Palma E, *et al*. The cyclophilin inhibitor Debio025 normalizes mitochondrial function, muscle apoptosis and ultrastructural defects in Col6a-/- myopathic mice. *Br J Pharmacol* 2009;157:1045–52.
- 53 Palma E, Tiepolo T, Angelin A, *et al*. Genetic ablation of cyclophilin D rescues mitochondrial defects and prevents muscle apoptosis in collagen VI myopathic mice. *Hum Mol Genet* 2009;18:2024–31.
- 54 Paco S, Ferrer I, Jou C, *et al*. Muscle fiber atrophy and regeneration coexist in collagen VI-deficient human muscle: role of calpain-3 and nuclear factor- $\kappa$ B signaling. *J Neuropathol Exp Neurol* 2012;71:894–906.
- 55 Pan TC, Zhang RZ, Markova D, *et al*. COL6A3 protein deficiency in mice leads to muscle and tendon defects similar to human collagen VI congenital muscular dystrophy. *J Biol Chem* 2013;288:14320–31.
- 56 Grumati P, Coletto L, Sabatelli P, *et al*. Autophagy is defective in collagen VI muscular dystrophies, and its reactivation rescues myofiber degeneration. *Nat Med* 2010;16:1313–20.
- 57 Low protein diet in patients with collagen VI related myopathies (LPD). <http://clinicaltrials.gov/ct2/show/NCT01438788> (accessed 3 Feb 2014).
- 58 Urciuolo A, Quarta M, Morbidoni V, *et al*. Collagen VI regulates satellite cell self-renewal and muscle regeneration. *Nat Commun* 2013;4:1964.
- 59 Alexeev V, Arita M, Donahue A, *et al*. Human adipose-derived stem cell transplantation as a potential therapy for collagen VI-related congenital muscular dystrophy. *Stem Cell Res Ther* 2014;5:21. <http://stemcellres.com/content/5/1/21>
- 60 Gualandi F, Manzati E, Sabatelli P, *et al*. Antisense-induced messenger depletion corrects a COL6A2 dominant mutation in Ullrich myopathy. *Hum Gene Ther* 2012;23:1313–8.
- 61 Bolduc V, Zou Y, Ko D, *et al*. siRNA-mediated allele-specific silencing of a COL6A3 mutation in a cellular model of dominant Ullrich muscular dystrophy. *Mol Ther Nucleic Acids* 2014;3:e147.
- 62 Noguchi S, Ogawa M, Kawahara G, *et al*. Allele-specific gene silencing of mutant mRNA restores cellular function in Ullrich congenital muscular dystrophy fibroblasts. *Mol Ther Nucleic Acids* (In press).
- 63 Usuki F, Yamashita A, Higuchi I, *et al*. Inhibition of nonsense-mediated mRNA decay rescues the phenotype in Ullrich's disease. *Ann Neurol* 2004;55:740–4.
- 64 Usuki F, Yamashita A, Kashima I, *et al*. Specific inhibition of nonsense-mediated mRNA decay components, SMG-1 or Upf1, rescues the phenotype of Ullrich disease fibroblasts. *Mol Ther* 2006;14:351–60.
- 65 Usuki F, Yamashita A, Shiraishi T, *et al*. Inhibition of SMG-8, a subunit of SMG-1 kinase, ameliorates nonsense-mediated mRNA decay-exacerbated mutant phenotypes without cytotoxicity. *Proc Natl Acad Sci USA* 2013;110:15037–42.



## Neuropathology Education

## Kyphoscoliosis and easy fatigability in a 14-year-old boy

Jantima Tanboon,<sup>1</sup> Yukiko K. Hayashi,<sup>2,3,4</sup> Ichizo Nishino<sup>3,4</sup> and Tumtip Sangruchi<sup>1</sup>

<sup>1</sup>Department of Pathology, Faculty of Medicine, Siriraj Hospital, Mahidol University, Bangkok, Thailand, <sup>2</sup>Department of Neurophysiology, Tokyo Medical University, <sup>3</sup>Department of Neuromuscular Research, National Institute of Neuroscience, and <sup>4</sup>Department of Clinical Development, Translational Medical Center, National Center of Neurology and Psychiatry, Tokyo, Japan

## CLINICAL COURSE

The patient was a 14-year-old Thai boy who presented at an orthopedic clinic with kyphoscoliosis first detected 1 year prior to medical attention. He was the first child of non-consanguineous parents; the pregnancy and delivery were uneventful. He had normal developmental milestones. The patient noticed easy fatigability after exercise which worsened during the past year. His younger brother was 10 years old and healthy. There was no history of neuromuscular disease in his family. His father passed away due to an unrelated incident. General examination revealed a body weight of 30 kg and a height of 149 cm, bilateral ptosis, high-arched palate, kyphoscoliosis, and asymmetrical chest wall. Proximal muscle weakness of grade 4 by Medical Research Council Scale in all extremities and areflexia were noted. The muscle tone was normal. Serum CK was 49 IU/L. Echocardiogram showed mild pulmonary and tricuspid regurgitation. Pulmonary function tests showed restrictive lung disease; the forced vital capacity was 1.56 L (51% of predicted). Sleep study revealed apnea-hypopnea index 13.6 per hour associated with severe oxygen desaturation (minimum SpO<sub>2</sub> 39.0%).

## PATHOLOGICAL FINDINGS

Muscle biopsy from the right quadriceps showed varying fiber sizes with type 1 hypotrophy and type 2 hypertrophy. The type 1 fibers were predominant (91%). Cap structures were seen in 32% of all fibers (Fig. 1). Ultrastructurally, the cap structures were composed of disorganized thin myofibrils and thickened Z-lines (Fig. 2). Nemaline rods were not present. Genetic analysis using genomic DNA

identified a heterozygous c.415\_417delGAG (p.Glu139del) in exon 4 of *TPM2*. The genetic tests were not performed in other family members.

## DIAGNOSIS

*TPM2*-related cap myopathy.

## DISCUSSION

Cap myopathy is a rare congenital myopathy characterized by presence of muscle fibers with cap structures on muscle biopsy and prominent respiratory muscle involvement. It is associated with dominant mutations in *TPM2*, *TPM3* and *ACTA1* genes, in descending order.<sup>1,2</sup>

Histologically, in cases with genetic results, cap structures range from 4% to 100% in muscle fibers.<sup>1</sup> Fidzianska reported two neonatal-onset cases with childhood death containing caps in 70–75% of fibers and two other childhood-onset cases with slowly progressive course containing caps in 20–30% of fibers, suggesting association of disease severity and the frequency of cap structure.<sup>3</sup> The number of caps seems to increase as the patients have aged.<sup>4,5</sup>


Onset of cap myopathy ranges from the neonatal period to childhood. Respiratory insufficiency is usually the major problem in patient management. The other common clinical manifestations include neonatal hypotonia, high-arched palate, myopathic facies and scoliosis. Muscle weakness is predominant in proximal muscles, which is non-specific. Serum CK is usually normal or slightly elevated. A case with distinctive abnormal cardiac function<sup>6</sup> and several cases with mild non-specific cardiac abnormalities have been reported.<sup>1</sup> Interestingly, one of the reported cases shows continuous gradual clinical improvement.<sup>5</sup>

Since all three causative genes in cap myopathy are also associated with nemaline myopathy (NM), it is not surprising to have concurrent caps and rods in the same patients<sup>2,7</sup>

Correspondence: Jantima Tanboon, MD, Department of Pathology, Faculty of Medicine, Siriraj Hospital, Mahidol University, Bangkok Noi, Bangkok 10700, Thailand. Email: jtanboon@gmail.com

Received 14 July 2014 and accepted 20 July 2014; published online 28 August 2014.






**Fig. 1** Cap structures at the periphery of the small fibers in HE stain (A). Caps are highlighted by modified Gomori trichrome stain (B) and nicotinamide adenine dehydrogenase tetrazolium reductase stain (C). The small fibers are type 1 demonstrated by ATPase stain (pH 4.4) and the cap areas are negatively stained (D). The caps are positive for desmin (E) and alpha-actinin (F). Bar = 50  $\mu$ m, inset bar = 10  $\mu$ m.

or have a family member with NM<sup>4,8</sup> although cap myopathy is much less common. Muscle biopsy findings of some cap myopathy patients were consistent with congenital fiber type disproportion (CFTD) at a younger age.<sup>5,8</sup> This may represent either a disease spectrum or insufficient tissue sampling since CFTD is also associated with *TPM2* and *TPM3* mutations.<sup>2</sup> Most cap myopathy patients are sporadic yet a few probable autosomal dominant familial cap myopathy cases have been reported.<sup>1,7</sup> It is likely that most cases are affected by *de novo* dominant mutations

causing too severe phenotypes to have offspring. Overlapping phenotypes between cap and nemaline myopathies include facial muscle weakness, high-arched palate and respiratory complications. The severity of both myopathies can range from mild to fatal.

The most common recurrent mutation in cap myopathy is p.Glu139del in exon 4 of *TPM2* gene and this mutation seems to be mostly related to cap myopathy, although six patients with different phenotypes have been reported.<sup>2,6,7,9</sup> Most patients with p.Glu139del, including our case, devel-





**Fig. 2** Electron microscopy study of the cap structures shows disorganized thin myofibrils and thickened Z-lines. Bar = 1  $\mu$ m.

oped symptoms during childhood, except for one case that presented in the neonatal period.<sup>9</sup> Our patient and two of the reported cases noticed developed scoliosis in a short period of time.<sup>6,7</sup> This feature has not been documented in cap myopathy patients with other mutations. It is possible yet inconclusive that p.Glu139del might affect axial muscles in later stage but progress at a faster rate compared to the other mutations.

In conclusion, we report a case of cap myopathy in a 14-year-old Thai boy. It is the first case reported from Asia and shares a common mutation, p.Glu139del, which was previously identified in patients with European ancestry.<sup>2,6,7,9</sup>

### ACKNOWLEDGMENTS

The authors thank Ms. Rangsim Nguenvattana for her secretarial work on the preparation of this manuscript. This work was supported partly by Research on Intractable Diseases, Comprehensive Research on Disability Health and Welfare, and Applying Health Technology from the Ministry of Health Labour and Welfare; partly by Intramu-

ral Research Grant 23-5 for Neurological and Psychiatric Disorders of National Center of Neurology and Psychiatry.

### DISCLOSURES

All authors report no disclosures.

### REFERENCES

- Schreckenbach T, Schroder JM, Voit T *et al*. Novel TPM3 mutation in a family with cap myopathy and review of the literature. *Neuromuscul Disord* 2014; **24**: 117–124.
- Marttila M, Lehtokari VL, Marston S *et al*. Mutation update and genotype-phenotype correlations of novel and previously described mutations in TPM2 and TPM3 causing congenital myopathies. *Hum Mutat* 2014; **35**: 779–790.
- Fidzianska A. “Cap disease” – a failure in the correct muscle fibre formation. *J Neurol Sci* 2002; **201**: 27–31.
- Cuisset JM, Maurage CA, Pellissier JF *et al*. “Cap myopathy”: case report of a family. *Neuromuscul Disord* 2006; **16**: 277–281.
- Ohlsson M, Quijano-Roy S, Darin N *et al*. New morphologic and genetic findings in cap disease associated with beta-tropomyosin (TPM2) mutations. *Neurology* 2008; **71**: 1896–1901.
- Clarke NF, Domazetovska A, Waddell L, Kornberg A, McLean C, North KN. Cap disease due to mutation of the beta-tropomyosin gene (TPM2). *Neuromuscul Disord* 2009; **19**: 348–351.
- Tasca G, Fattori F, Ricci E *et al*. Somatic mosaicism in TPM2-related myopathy with nemaline rods and cap structures. *Acta Neuropathol* 2013; **125**: 169–171.
- Tajsharghi H, Ohlsson M, Lindberg C, Oldfors A. Congenital myopathy with nemaline rods and cap structures caused by a mutation in the beta-tropomyosin gene (TPM2). *Arch Neurol* 2007; **64**: 1334–1338.
- Lehtokari VL, Ceuterick-de Groote C, de Jonghe P *et al*. Cap disease caused by heterozygous deletion of the beta-tropomyosin gene TPM2. *Neuromuscul Disord* 2007; **17**: 433–442.