

19. Watanabe M, Miyata H, Gotoh M, Baba H, Kimura W, Tomita N, et al. Total gastrectomy risk model: data from 20,011 Japanese patients in a nationwide internet-based database. *Ann Surg.* (in press).
20. Kobayashi H, Miyata H, Gotoh M, Baba H, Kimura W, Kitagawa Y, et al. Risk model for right hemicolectomy based on 19,070 Japanese patients in the National Clinical Database. *J Gastroenterol.* 2013. (Epub ahead of print).
21. Kenjo A, Miyata H, Gotoh M, Okubo S, Suzuki H, Kitagawa Y, et al. Risk stratification of 7732 hepatectomy cases in 2011 from National Clinical Database for Japan. *J Am Coll Surg.* 2014;218:412–22.
22. Khuri SF. The NSQIP: a new frontier in surgery. *Surgery.* 2005;138:837–43.
23. Miyata H, Hashimoto H, Horiguchi H, Matsuda S, Motomura N, Takamoto S. Performance of in-hospital mortality prediction models for acute hospitalization: hospital standardized mortality ratio in Japan. *BMC Health Serv Res.* 2008;8:229.
24. Riche FC, Dray X, Laisne MJ, Mateo J, Raskine L, Sanson-Le Pors MJ, et al. Factors associated with septic shock and mortality in generalized peritonitis: comparison between community-acquired and postoperative peritonitis. *Crit Care.* 2009;13:R99.
25. Chandra V, Nelson H, Larson DR, Harrington JR. Impact of primary resection on the outcome of patients with perforated diverticulitis. *Arch Surg.* 2004;139:1221–4.
26. Pisanu A, Reccia I, Deplano D, Porru F, Uccheddu A. Factors predicting in-hospital mortality of patients with diffuse peritonitis from perforated colonic diverticulitis. *Ann Ital Chir.* 2012;83:319–24.
27. Welcker K, Lederle J, Schorr M, Siebeck M. Surgery and adjuvant therapy in patients with diffuse peritonitis: cost analysis. *World J Surg.* 2002;26:307–13.
28. Trenti L, Biondo S, Golda T, Monica M, Kreisler E, Fraccalvieri D, et al. Generalized peritonitis due to perforated diverticulitis: Hartmann's procedure or primary anastomosis? *Int J Colorectal Dis.* 2011;26:377–84.
29. Arenal JJ, Bengoechea-Beeby M. Mortality associated with emergency abdominal surgery in the elderly. *Can J Surg.* 2003;46:111–6.
30. Wakefield H, Vaughan-Sarrazin M, Cullen JJ. Influence of obesity on complications and costs after intestinal surgery. *Am J Surg.* 2012;204:434–40.
31. Cook TM, Day CJ. Hospital mortality after urgent and emergency laparotomy in patients aged 65 yr and over. Risk and prediction of risk using multiple logistic regression analysis. *Br J Anaesth.* 1998;80:776–81.
32. Bader FG, Schroder M, Kujath P, Muhl E, Bruch HP, Eckmann C. Diffuse postoperative peritonitis—value of diagnostic parameters and impact of early indication for relaparotomy. *Eur J Med Res.* 2009;14:491–6.
33. Tokuda Y, Abe T, Ishimatsu S, Hinohara S. Ambulance transport of the oldest old in Tokyo: a population-based study. *J Epidemiol.* 2010;20:468–72.
34. Matzen LE, Jepsen DB, Ryg J, Masud T. Functional level at admission is a predictor of survival in older patients admitted to an acute geriatric unit. *BMC Geriatr.* 2012;12:32.
35. Nakazawa A, Nakamura K, Kitamura K, Yoshizawa Y. Association between activities of daily living and mortality among institutionalized elderly adults in Japan. *J Epidemiol.* 2012;22:501–7.

OPEN

# Risk Models of Operative Morbidities in 16,930 Critically Ill Surgical Patients Based on a Japanese Nationwide Database

Zenichiro Saze, MD, PhD, Hiroaki Miyata, PhD, Hiroyuki Konno, MD, PhD, Mitsukazu Gotoh, MD, PhD, Takayuki Anazawa, MD, PhD, Ai Tomotaki, RN, MS, Go Wakabayashi, MD, PhD, and Masaki Mori, MD, PhD

**Abstract:** The aim of the study was to evaluate preoperative variables predictive of lethal morbidities in critically ill surgical patients at a national level.

There is no report of risk stratification for morbidities associated with mortality in critically ill patients with acute diffuse peritonitis (ADP).

We examined data from 16,930 patients operated during 2011 and 2012 in 1546 different hospitals for ADP identified in the National Clinical Database of Japan. We analyzed morbidities significantly associated with operative mortality. Based on 80% of the population, we calculated independent predictors for these morbidities. The risk factors were validated using the remaining 20%.

The operative mortality was 14.1%. Morbidity of any grade occurred in 40.2% of patients. Morbidities correlated with mortality, including septic shock, progressive renal insufficiency, prolonged ventilation >48 hours, systemic sepsis, central nervous system (CNS) morbidities, acute renal failure and pneumonia, and surgical site infection (SSI), were selected for risk models. A total of 18 to 29 preoperative variables were selected per morbidity and yielded excellent C-indices for each (septic shock: 0.851; progressive renal insufficiency: 0.878; prolonged ventilation >48 h: 0.849; systemic sepsis: 0.839; CNS morbidities: 0.848; acute renal failure: 0.868; pneumonia: 0.830; and SSI: 0.688).

We report the first risk stratification study on lethal morbidities in critically ill patients with ADP using a nationwide surgical database. These risk models will contribute to patient counseling and help predict which patients require more aggressive surgical and novel pharmacological interventions.

(*Medicine* 94(30):e1224)

**Abbreviations:** ADL = activities of daily living, ADP = acute diffuse peritonitis, APACHE II = Acute Physiology and Chronic Health Evaluation II, ASA = American Society of Anesthesiologists, BMI = body mass index, CIs = confidence

Editor: Xiao-Dong Chen.

Received: April 20, 2015; revised: June 30, 2015; accepted: July 1, 2015. From the Japanese Society of Gastroenterological Surgery (JSGS) (ZS, TA, MM), JSGS Database Committee (HM, HK, MG, GW), and National Clinical Database (HM, AT), Tokyo, Japan.

Correspondence: Mitsukazu Gotoh, Department of Regenerative Surgery, Fukushima Medical University, 1-Hikarigaoka, Fukushima City 960-1295, Japan (e-mail: mgotoh@fmu.ac.jp).

This study was partially supported by a research grant from Ministry of Health, Labour and Welfare.

The authors have no conflicts of interest to disclose.

Copyright © 2015 Wolters Kluwer Health, Inc. All rights reserved.

This is an open access article distributed under the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0, where it is permissible to download, share and reproduce the work in any medium, provided it is properly cited. The work cannot be changed in any way or used commercially.

ISSN: 0025-7974

DOI: 10.1097/MD.0000000000001224

intervals, CNS = central nervous system, CVA = cerebrovascular accident, JSGS = Japanese Society of Gastroenterological Surgery, NCD = National Clinical Database, ROC = Receiver operating characteristic, SIRS = systemic inflammatory response syndrome, SSI = surgical site infection.

## INTRODUCTION

Acute diffuse peritonitis (ADP) is defined as the uncontained spread of intraabdominal infection, rapidly proceeding beyond the source of infection into multiple (2–4) quadrants of the intraabdominal cavity.<sup>1</sup> Most patients diagnosed with ADP are critically ill and therefore require emergency surgery, regardless of the source of infection.<sup>2–4</sup> A high incidence of severe postoperative complications such as septic shock, pneumonia, and organ failure has resulted in a high mortality rate of approximately 30%, even in modern case series.<sup>4</sup> Therefore, the identification of postoperative complications associated with mortality and their optimal treatment is necessary to improve outcomes. There have been risk models for mortality in critically ill patients. The Acute Physiology and Chronic Health Evaluation II (APACHE II) score,<sup>5</sup> Sequential Organ Failure Assessment score,<sup>6</sup> and Mannheim Peritonitis Index<sup>7</sup> have all been shown to be quite effective for predicting mortality in critically ill patients. However, there has been no risk model for the morbidity of critically ill patients using a nationwide database.

American College of Surgeons National Surgical Quality Improvement Program (ACS-NSQIP) risk models are powerful predictors of specific morbidities and mortality associated with gastrointestinal surgery.<sup>8–10</sup> However, there has been no nationwide analysis of critically ill surgical patients. In one regional report, Turner et al<sup>11</sup> showed that ACS-NSQIP criteria were associated with high APACHE II scores and poor outcomes in 340 surgical patients (mortality: 20.6%) treated in the intensive care unit of the University of Maryland Medical Center (Baltimore, MD). They found that APACHE II score predictions were consistent with ACS-NSQIP postoperative outcomes. This observation prompted us to hypothesize that ACS-NSQIP preoperative variables could be used to predict both postoperative morbidities and mortalities in ADP patients.

The National Clinical Database (NCD) in Japan, which commenced patient registration in January 2011, is a nationwide project linked to the surgical board certification system.<sup>12,13</sup> Submitting cases to the NCD is a prerequisite for all member institutions of both the Japan Surgical Society and Japanese Society of Gastroenterological Surgery, and only registered cases can be used for board certification. The NCD collaborates with the ACS-NSQIP<sup>10</sup>; they share the common goal of developing a standardized surgery database to achieve an improvement in treatment quality.<sup>14</sup>

Previously, we reported that patients with ADP are critically ill, most require emergency surgery, and their 30-day mortality and 90-day in-hospital mortality rates are 9% and 13.9%, respectively.<sup>15</sup> In this study, we used data from 16,930 patients with ADP treated in 2011 and 2012 and registered with the NCD to create risk models for postoperative morbidities associated with mortality.

## METHODS

### Patient Selection

The NCD is a nationwide project associated with the board certification system of surgery in Japan into which data from over 1,200,000 surgical cases treated at over 3500 hospitals are entered annually. We have created risk models of mortality for the 8 surgical procedures (esophagectomy, total gastrectomy, distal gastrectomy, right hemicolectomy, low anterior resection, hepatectomy, pancreaticoduodenectomy, and ADP) using NCD data sets, and the respective model was published separately,<sup>15–22</sup> and the results were summarized as a review article.<sup>13</sup> Thus, patient selection, preoperative and perioperative variables, and ethics consideration were quite consistent between the studies. The NCD continuously recruits individuals who approve these data, members of various departments in charge of cases, and data entry officers through a web-based data management system; thus, the traceability of the data is assured.<sup>12</sup> In addition, the project constantly validates the consistency of these data by the inspection of randomly chosen institutions. Current laws, ordinances, and guidelines regarding the confidentiality of data are observed. Patients agree for their data to be included in research projects by using presumed consent with opt-out through the Web page and/or a notice of each hospital.<sup>20</sup> The NCD project was approved on November 2010 by Japan Surgical Society Ethics Committee.

In this study, we focused on ADP in the Gastrointestinal Surgery section of the NCD. In the NCD, we identified 16,930 patients who underwent surgery for ADP in 2011 to 2012. Patients who declined to have their records entered in the NCD were excluded from our analysis. Records with missing data on patient age, sex, or status, 30 days after surgery were also excluded.

### Preoperative and Perioperative Variables

The preoperative and perioperative variables used by the NCD are almost identical to those used by the ACS-NSQIP ([http://site.acsnsqip.org/wp-content/uploads/2013/10/ACSNSQIP\\_PUF\\_UserGuide.2012.pdf#search=user+guide+for+the+2012+ACS+NSQIP](http://site.acsnsqip.org/wp-content/uploads/2013/10/ACSNSQIP_PUF_UserGuide.2012.pdf#search=user+guide+for+the+2012+ACS+NSQIP)). All variables, definitions, and inclusion criteria regarding the NCD are accessible to participating institutions on its website (<http://www.ncd.or.jp/>), which also features an E-learning system to instruct participants in how to input consistent data. The potential independent variables were previously described.<sup>13,15–22</sup> These included patient demographics, preexisting comorbidities, preoperative laboratory values, and perioperative data (Table 1).

### Outcome Measures (Mortality and Postoperative Occurrences)

We calculated the 30-day mortality and operative mortality. The former was defined as death within 30 days of surgery, regardless of the patient's geographical location, even if the patient had been discharged from the hospital. The latter was defined as death within the index hospitalization period,

regardless of the length of hospital stay (up to 90 days), as well as any death after discharge within 30 days of surgery.

The postoperative morbidities that occurred within 30 days of surgery included relaparotomy within 30 days of surgery; wound-related morbidities (superficial incisional surgical site infection [SSI], deep incisional SSI, organ/space SSI, wound disruption); respiratory morbidities (pneumonia, unplanned intubation, pulmonary embolism, ventilation >48 hours); urinary tract morbidities (progressive renal insufficiency, acute renal failure, urinary tract infection); central nervous system (CNS) morbidities (stroke/cerebrovascular accident [CVA], coma for <24 hours, peripheral nerve injury); cardiac morbidities (cardiac arrest, myocardial infarction); and other occurrences (bleeding 1–4 u or  $\geq 5$  u red blood cells, deep-vein thrombosis/thrombophlebitis, septic shock, severe sepsis, systemic inflammatory response syndrome [SIRS]).

### Statistical Analysis

We used IBM SPSS Statistics for Windows (Version 20; IBM Corp, Armonk, NY) for data analysis. Univariate analysis of the data was performed using Fisher exact test, the unpaired Student *t* test, and the Mann–Whitney *U* test. Correlations between each morbidity and operative mortality and between respective morbidities were analyzed using the Pearson product–moment correlation.

Data were randomly assigned into 2 subsets that were split 80/20: the first for model development and the second for validation. The 8 sets of logistic models (septic shock, systemic sepsis, progressive renal insufficiency, acute renal failure, ventilation >48 hours, pneumonia, CNS morbidities, and SSI) were constructed for dataset development using step-wise selection of the predictors with a probability (*P*) value for inclusion of 0.05. A “goodness-of-fit” test was performed to assess how well the model discriminated between patients with or without respective morbidities. Receiver operating characteristic (ROC) curves for respective morbidities were created for the validation dataset. A ROC curve is a plot of a test's true-positive rate (sensitivity) versus its false-positive rate (1–specificity).

## RESULTS

### Preoperative Risk Profiles and Laboratory Data of the Study Population

The demographic data and risk profile of 16,930 patients with ADP are shown in Table 1. The patient population had a mean age of  $64.9 \pm 18.6$  years (range: 0–106 years), and 60.5% ( $n = 10,248$ ) were male. In this population, 37.7% arrived at hospital by ambulance, and 92.9% required emergency surgery. Their original disease and associated operative mortalities were acute peritonitis (15.1%), appendicitis (1%), gastroduodenal ulcer/perforation (9.5%), intestinal perforation (18.4%), intestinal obstruction (18.9%), cholecystitis/cholangitis (13.3%), and vascular insufficiency (31.2%). These proportions and mortalities are consistent with findings from 2011.<sup>15</sup>

An abbreviated risk profile for the study population is also shown in Table 1. In brief, 58.4% of the patient population had an American Society of Anesthesiologists (ASA) classification of III–V, partial/total dependency for activities of daily living (ADL) was 41.2%, 0.5% of patients had body mass index (BMI) of  $>30 \text{ kg/m}^2$ , and 5.1% of patients had a weight loss of  $>10\%$ . With regard to preexisting comorbidities, failure of various organs occurred in a percentage of patients, including ventilator

**TABLE 1.** Preoperative Risk Profiles and Laboratory Data of the Study Population

Characteristics	Cases With Characteristics	% of Entire Population	No. of Death	Operative Mortality	Fisher
<b>Demographics</b>					
Age					
Under 60	5217	30.8%	236	4.5%	<0.001
61–65	1890	11.2%	185	9.8%	
66–70	1677	9.9%	236	14.1%	
71–75	1978	11.7%	349	17.6%	
76–80	2248	13.3%	435	19.4%	
80 and over	3920	23.2%	944	24.1%	
Males	10248	60.5%	1389	13.6%	0.014
Ambulance transportation	6375	37.7%	972	15.2%	<0.001
Emergency case	15731	92.9%	2231	14.2%	0.213
<b>Preoperative risk assessment</b>					
General					
ADL immediately before surgery					
Totally dependent	2278	13.5%	758	33.3%	<0.001
Partially dependent	4690	27.7%	1326	28.3%	<0.001
ASA classification					
Class 4 and 5	2431	14.4%	990	40.7%	<0.001
Class 3	7448	44.0%	1919	25.8%	<0.001
Body mass index $\geq 30$ kg/m <sup>2</sup>	452	0.5%	78	17.3%	0.052
Body mass index $\geq 26$ kg/m <sup>2</sup>	1873	1.5%	249	13.3%	0.307
Alcohol drinking (at times/occasional)	7106	42.0%	784	11.0%	<0.001
Brinkmann index $\geq 600$	2605	2.1%	358	13.7%	0.602
Brinkmann index $\geq 400$	3551	2.7%	456	12.8%	0.017
>10% loss body weight in last 6 months	861	5.1%	295	34.3%	<0.001
Respiratory					
Ventilator dependent	646	3.8%	283	43.8%	<0.001
Current pneumonia	637	3.8%	278	43.6%	<0.001
History of severe COPD	563	3.3%	150	26.6%	<0.001
Respiratory failure	1391	8.2%	545	39.2%	<0.001
Cardiovascular					
Congestive heart failure	447	2.6%	195	43.6%	<0.001
Hypertension requiring medication	5046	29.8%	901	17.9%	<0.001
Hypertension without treatment	521	3.1%	89	17.1%	0.052
Renal					
Acute renal failure	742	4.4%	321	43.3%	<0.001
Cerebral nervous system					
CVA/Stroke with neurological deficit	482	2.8%	111	23.0%	<0.001
Cerebrovascular disease within 14 days	142	0.8%	32	22.5%	0.006
Cerebrovascular disease	812	4.8%	202	24.9%	<0.001
Hematological					
Bleeding disorder without treatment	1086	6.4%	373	34.3%	<0.001
Bleeding disorder	1828	10.8%	592	32.4%	<0.001
Preop Transfusion of $\geq 1$ unit of RBCs	3487	20.6%	1028	29.5%	<0.001
Any blood transfused in the emergency room	702	4.1%	287	40.9%	<0.001
Infectious disorder					
Systemic sepsis	5233	30.9%	1266	24.2%	<0.001
Other					
Epidural anesthesia	3482	20.6%	224	0.064	<0.001
Open wound	450	2.7%	128	28.4%	<0.001
Steroid use for chronic condition	677	4.0%	197	29.1%	<0.001
Ascites without control	3742	22.1%	811	21.7%	<0.001
Esophageal varices without control	89	0.5%	29	32.6%	<0.001
Disease					
Acute peritonitis	8613	50.9%	1300	15.1%	<0.001
Appendicitis	2470	14.6%	24	1.0%	<0.001
Gastroduodenal ulcer/perforation	1742	10.3%	166	9.5%	<0.001
Intestinal perforation	2504	14.8%	461	18.4%	<0.001

Characteristics	Cases With Characteristics	% of Entire Population	No. of Death	Operative Mortality	Fisher
Intestinal obstruction	855	5.1%	162	18.9%	<0.001
Cholecystitis/cholangitis	451	2.7%	60	13.3%	0.676
Vascular insufficiency	253	1.5%	79	31.2%	<0.001
Oncological					
Other than cancer surgery	15202	89.8%	1899	12.5%	<0.001
Preoperative laboratory value					
WBC < 3500/mL	2717	3.3%	567	20.9%	<0.001
Hematocrit over 48% (male), 42% (female)	1056	0.7%	122	11.6%	0.015
Plate count < 150,000/mL	2798	4.7%	799	28.6%	<0.001
Plate count < 50,000/mL	199	0.6%	105	52.8%	<0.001
Serum albumin < 3.5 g/dL	8839	11.0%	1864	21.1%	<0.001
Serum albumin < 2.5 g/dL	3334	5.8%	977	29.3%	<0.001
Serum albumin < 2.0 g/dL	1293	2.8%	471	36.4%	<0.001
SGOT ≥ 40 U/L	3225	4.8%	819	25.4%	<0.001
SGOT ≥ 35 U/L	3848	5.5%	933	24.2%	<0.001
Bilirubin < 0.2 mg/dL	40	0.0%	8	20.0%	0.259
Serum creatinine ≥ 3.0 mg/dL	1104	2.2%	374	33.9%	<0.001
Serum creatinine ≥ 2.0 mg/dL	1980	3.7%	634	32.0%	<0.001
Serum creatinine ≥ 1.2 mg/dL	4378	6.9%	1176	26.9%	<0.001
BUN ≥ 60 mg/dL	905	2.0%	337	37.2%	<0.001
BUN ≥ 25 mg/dL	5458	8.5%	1435	26.3%	<0.001
BUN ≥ 20 mg/dL	7398	10.2%	1728	23.4%	<0.001
Serum sodium < 130 mEq/L	924	1.4%	236	25.5%	<0.001
Serum sodium ≥ 146 mEq/L	316	0.7%	120	38.0%	<0.001
Alkaline phosphatase < 110 mEq/L	372	0.4%	63	16.9%	0.111
CRP > 10 mg/dL	7934	7.3%	1240	15.6%	<0.001
INR of PT values ≥ 1.67	796	1.5%	248	31.2%	<0.001
PT < 10 s	1886	2.4%	398	21.1%	<0.001
PTT < 30 s	4330	2.5%	429	9.9%	<0.001

ADL = activities of daily living; ASA classification = American Society of Anesthesiologists Physical Status Classification; AST = aspartate amino transferase; BUN = blood urea nitrogen; COPD = chronic obstructive pulmonary disease; CRP = C-reactive protein; CVA = cerebrovascular accident; WBC = white blood cell.

dependence (3.8%), congestive heart failure (2.6%), and acute renal failure (4.4%). Signs of systemic sepsis were evident in 30.9% of patients. Blood transfusion was required in 4.1% of patients. An ASA classification of >IV and V and organ failure were associated with an operative mortality rate of >40%.

### Postoperative Occurrences in Patients with ADP

The 30-day mortality and operative mortality rates after surgery for ADP were 8.8% (1482) and 14.1% (2385), respectively. The incidences of various morbidities and percentage of consequent patient deaths are shown in Table 2. The postoperative morbidities that led to a high percentage of deaths (>40%) included transfusion (1–4 U: 43.5%; >5 U: 52.2%), prolonged ventilation (45.6%), unplanned intubation (51.4%), pneumonia (43%), cardiac and CNS morbidities (90.3% and 64.8%, respectively), acute renal failure (57.1%), progressive renal insufficiency (55.6%), any systemic sepsis (41%), and septic shock (55.8%). These morbidities occurred at a relatively high incidence (4.8%–15%) excepting cardiac morbidities (2.5%). SSI of any type, including organ space, deep incisional, and superficial incisional, occurred in 23.2% of patients and led to an operative mortality rate of 20.8%.

### Correlation Between Postoperative Morbidities and Operative Mortality

Correlation between 30-day operative mortality rates and postoperative morbidities were analyzed using the Pearson

product–moment correlation. The morbidities highly correlated with mortality (top 7) as well as SSI as the most representative complication of ADP were selected and are compared in Table 3. A better correlation with postoperative morbidities was found when operative rather than 30-day mortality was used. Among the postoperative morbidities, septic shock, progressive renal insufficiency, and ventilation >48 hours were highly correlated with each other ( $r > 0.5$ ). In contrast, SSI was only moderately correlated with systemic sepsis, and weakly correlated with ventilation >48 hours.

### Model Results and Performance

We developed risk models for postoperative morbidities with a relatively high incidence associated with high mortality (Table 4; Supplemental Table, <http://links.lww.com/MD/A344>, with 95% confidence intervals [CIs]). The postoperative morbidities selected correlated well with operative mortality. Septic shock, systemic sepsis (SIRS, sepsis, or septic shock), progressive renal insufficiency, acute renal failure, ventilation >48 hours, pneumonia, and CNS morbidities were selected, and SSI was also included as the most frequent morbidity.

The logistic models of these morbidities with odds ratios are shown in Table 4. The morbidities with a 95% CI showing statistical significance are shown in the Supplemental Table, <http://links.lww.com/MD/A344>. To evaluate the performance of the models, the C-index (a measure of model discrimination), which was the area under the ROC curve, was calculated for the

TABLE 2. Postoperative Occurrences After ADP Surgery

Postoperative Outcomes	Cases With the Outcome	% of Entire Population	No. of Death	% Death With the Outcome	% Death Without the Outcome	Fisher
General						
Any complication	6808	40.2	1828	26.9	5.5	<0.001
Bleeding transfusions	2353	13.9	1023	43.5	9.3	<0.001
Bleeding transfusions $\geq 5$ units	1337	7.9	698	52.2	10.8	<0.001
Reoperation within 30 d	1317	7.8	317	24.1	13.2	<0.001
Readmission within 30 d	340	2.0	14	4.1	14.3	<0.001
Respiratory						
On Ventilator >48 h	2592	15.3	1182	45.6	8.4	<0.001
Unplanned intubation	821	4.8	422	51.4	12.2	<0.001
Pneumonia	1693	10.0	728	43.0	10.9	<0.001
Cardiovascular						
Cardiac arrest/myocardial infarction	421	2.5	380	90.3	12.1	<0.001
Pulmonary embolism	55	0.3	16	29.1	14.0	<0.001
Cerebral nervous system						
CVA/Stroke	867	5.1	562	64.8	11.3	<0.001
Renal						
Acute renal failure	960	5.7	548	57.1	11.5	<0.001
Progressive renal insufficiency	1740	10.3	967	55.6	9.3	<0.001
Infectious disorder						
Systemic sepsis	3321	19.6	1361	41.0	7.5	<0.001
Septic shock	1786	10.5	996	55.8	9.2	<0.001
Sepsis	826	4.9	224	27.1	13.4	<0.001
SIRS	709	4.2	141	19.9	13.8	<0.001
SSI	3931	23.2	819	20.8	12.0	<0.001
Organ space SSI	1865	11.0	541	29.0	12.2	<0.001
Deep incisional SSI	1648	9.7	475	28.8	12.5	<0.001
Superficial SSI	3052	18.0	632	20.7	12.6	<0.001
Wound disruption	1179	7.0	403	34.2	12.6	<0.001
Urinary tract infection	440	2.6	124	28.2	13.7	<0.001

CVA = cerebrovascular accident, SIRS = systemic inflammatory response syndrome, SSI = surgical site infection.

validation sets (Figure 1). The C-indices and 95% CIs of each occurrence were 0.851 (0.841–0.860) for septic shock, 0.878 (0.870–0.887) for progressive renal insufficiency, 0.849 (0.841–0.858) for ventilation >48 hours, 0.848 (0.835–0.862) for CNS morbidities, 0.868 (0.856–0.880) for acute renal failure, 0.830 (0.819–0.840) for pneumonia, and 0.851 (0.841–0.860) for systemic sepsis. The C-index of SSI showed a weaker correlation (0.688 [0.677–0.698]) than other morbidities.

A total of 18 to 29 preoperative variables were selected as risk factors of each complication. Age, ASA classification, preoperative ventilation or pneumonia, acute renal failure, blood transfusion, and systemic sepsis, as well as selected preoperative laboratory values suggestive of severe infection and organ failure, were captured in the risk models as predictors of most of the complications.

## DISCUSSION

We hypothesized that ACS-NSQIP preoperative variables could be used to predict both postoperative morbidities and mortalities in ADP patients. In total, 93% of 16,930 patients with ADP included in this study required emergency surgery, and the overall operative mortality was 14.1%. This was comparable with the findings of a previous analysis using NCD data from 2011,<sup>15</sup> in which 93.1% of patients with




ADP required emergency surgery, and the overall operative mortality was 8.8%. This suggests that there is a consistent population of critically ill surgical patients who require emergency surgery in Japan. By examining the data of a large number of patients with ADP, we were able to identify the postoperative complications associated with mortality and create risk models for each complication. Septic shock, progressive renal insufficiency, ventilation >48 hours and systemic sepsis were moderately correlated ( $r > 0.36$ ) with operative mortality, whereas CNS morbidities, acute renal failure, and pneumonia were weakly ( $0.2 < r \leq 0.35$ ) correlated with operative mortality. For these complications, risk models showed excellent C-indices ( $> 0.830$ ) in the validation dataset. To our knowledge, this is the first report to successfully show and validate using a large-scale dataset that the preoperative variables of the ACS-NSQIP can predict postoperative morbidities in critical ill patients.

The prediction of postoperative complications is essential to the decision-making process before surgery, and useful to identify patients eligible for participation in the evaluation of novel pharmacologic interventions<sup>23,24</sup> or more aggressive surgical interventions. In the past, several scoring systems have been used to predict complications.<sup>25–31</sup> ASA score is a useful predictor for mortality,<sup>25,26</sup> but suffers from its reproducibility because of subjective parameters.<sup>26</sup> APACHE II was developed in a mixed group of medical and surgical patients.<sup>27</sup> It failed to

**TABLE 3.** Correlation Between Operative Mortality and Respective Postoperative Occurrences

Occurrences	thirtyday mortality	operative mortality	Septic shock	Progressive renal insufficiency	On Ventilator > 48 Hours	Any systemic sepsis	CVA/Stroke	Acute renal failure	Pneumonia	SSI
30-day mortality	1	.765	.398	.365	.327	.336	.328	.301	.187	.034
Operative mortality	.765	1	.411	.404	.385	.382	.339	.303	.277	.107
Septic shock	.398	.411	1	.526	.579	.695	.390	.465	.371	.268
Progressive renal insufficiency	.365	.404	.526	1	.554	.536	.411	.724	.390	.283
On Ventilator > 48 h	.327	.385	.579	.554	1	.621	.434	.444	.491	.329
Any systemic sepsis	.336	.382	.695	.536	.621	1	.367	.421	.439	.428
CVA/Stroke	.328	.339	.390	.411	.434	.367	1	.343	.265	.157
Acute renal failure	.301	.303	.465	.724	.444	.421	.343	1	.303	.195
Pneumonia	.187	.277	.371	.390	.491	.439	.265	.303	1	.285
SSI any	.034	.107	.268	.283	.329	.428	.157	.195	.285	1

The column mark indicates the following:

 0.3 ≤ r < 0.4    
  0.4 ≤ r < 0.5    
  0.5 ≤ r

CVA = cerebrovascular accident, SSI = surgical site infection.

predict the development of multiple organ failure syndrome or mortality with clinical utility in postoperative surgical patients.<sup>28</sup> Physiological and Operative Severity Score for the enUmeration of Mortality and Morbidity has been studied as a possible surgical audit system<sup>29</sup>; however, it seems to overestimate mortality, particularly for the low risk group.<sup>30,31</sup> A reliable model for predicting complications can only be based on the accurately recorded incidences of those complications. A comparison of the outcomes of patients with ADP registered with the NCD in 2011 with those registered in 2012 revealed that mortality and morbidities were highly correlated between these years ( $r = 0.9932$ ; Supplemental Figure, <http://links.lww.com/MD/A344>). The thorough data retrieval system of the NCD and clinically clear entity of ADP made it possible to create successful risk models for these morbidities.

Severe sepsis/septic shock, defined as the presence of acute organ dysfunction in the context of infection, has a mortality rate of approximately 25% to 35%,<sup>32,33</sup> but which can exceed 70%.<sup>34,35</sup> Anaya and Nathens<sup>36</sup> analyzed risk factors of severe sepsis in 11,202 patients using Washington State administrative hospital discharge data. They identified 11% with severe sepsis, which was present in 424 (62%) of the 686 decedents, and showed that source of infection, extent of peritonitis, increasing

age, and preexisting organ dysfunction were independently associated with severe sepsis. Our findings on the mortality of patients with ADP were consistent with their study. The mortality of patients with ADP as a result of appendicitis was low (1%) compared with that associated with other causes such as intestinal/gastroduodenal perforation (18.4%/9.5%), vascular insufficiency (31.2%), and cholecystitis/cholangitis (13.3%). Regarding peritonitis, when it is localized within an abscess, the operative mortality rate of cases registered with the NCD was relatively low (4.6%; 254 deaths/5470 cases) compared with that of patients with ADP (14.1%). This study provides more reliable information on clinical variables and laboratory data compared with the findings of Anaya and Nathens.<sup>36</sup> We were able to select significant variables to predict each complication, and discrimination and calibration using validation tests clearly showed the excellent performance of these models.

It is interesting to note that the risk models for morbidities moderately associated with mortality (septic shock, any systemic sepsis, renal failure, acute renal failure, prolonged ventilation, pneumonia, and CNS morbidities) picked up similar variables as risk factors—age, ADL status, ASA classification, blood transfusions, and systemic sepsis—to those found to be

**TABLE 4.** Risk Models of Postoperative Occurrences After ADP Surgery

Variable	Septic Shock	Any Systemic Sepsis	Progressive Renal Insufficiency	Acute Renal Failure	On Ventilator > 48 Hours	Pneumonia	CVA/Stroke	SSI Any
<b>Demographics</b>								
Age 60–75	1.144	1.095	1.105	1.144	1.16	1.214	1.174	1.04
Males		1.153			1.13	1.317		
<b>Preoperative risk assessment</b>								
<b>General</b>								
ADL totally dependent	1.178				1.399		1.426	
ADL partially dependent		1.175	1.23			1.278		
ASA class 4 and class 5	3.635	2.993	3.147	3.474	3.341	2.321	3.433	1.705
ASA class 3	1.77	1.888	1.957	1.922	2.066	1.837	1.691	1.347
Body mass index $\geq 30$ kg/m <sup>2</sup>					1.567			
Body mass index $\geq 26$ kg/m <sup>2</sup>			1.438	1.614	1.224			1.274
Alcohol drinking (at times/occasional)			1.181	1.256		1.206		1.118
Brinkmann index $\geq 600$	1.199				1.217			
Brinkmann index $\geq 400$		1.162						
>10% loss body weight in last 6 months								1.561
<b>Respiratory</b>								
Ventilator dependent	1.519	1.404	1.305		2.734		2.035	
Current pneumonia		1.35	1.667	1.704	1.89	4.994	1.599	
History of severe COPD	1.371				1.472	1.403		
Respiratory failure	1.236					1.292		
<b>Cardiovascular</b>								
Congestive heart failure			1.501		1.331			
Hypertension requiring medication		1.119	1.199		1.235			
Hypertension without treatment								
<b>Renal</b>								
Acute renal failure	1.471	1.258	2.975	3.869	1.26	1.504		
<b>Cerebral nervous system</b>								
CVA/Stroke		1.346		1.675	1.376	1.631	1.826	
Cerebrovascular disease within 14 days		1.933					3.406	
Cerebrovascular disease	1.373					1.421		
<b>Hematological</b>								
Bleeding disorder without treatment	1.437	1.494		1.471	1.377	1.289	1.92	
Bleeding disorder			1.361					
Blood transfusions	1.511	1.556	1.514	1.61	1.887	1.546	1.432	1.17
Preoperative transfusion of $\geq 1$ unit of RBCs			1.303		1.369			1.355
<b>Infectious disorder</b>								
Systemic Sepsis	2.821	4.086	1.974	2.035	2.092	1.901	1.776	1.824
<b>Oncological</b>								
Other than cancer surgery	0.734		0.803					
<b>Other</b>								
Open wound		1.469						2.186
Steroid use for chronic condition	1.486		1.585		1.586	1.545		1.507
Ascites without control		1.17						
Esophageal varices without control					1.846			
<b>Preoperative laboratory value</b>								
WBC < 3500/mL	1.989	1.462	1.318	1.55	1.553		1.428	1.225
Hematocrit over 48% (male), 42% (female)	1.441	1.334		1.52	1.493			
Plate count < 150,000/mL		1.175	1.192					
Plate count < 50,000/mL	1.741							
Serum albumin < 3.5 g/dL		1.286			1.153			1.162
Serum albumin < 2.5 g/dL	1.267					1.18	1.251	
Serum albumin < 2.0 g/dL		1.287	1.403		1.606	1.255		1.227



Variable	Septic Shock	Any Systemic Sepsis	Progressive Renal Insufficiency	Acute Renal Failure	On Ventilator > 48 Hours	Pneumonia	CVA/Stroke	SSI Any
SGOT ≥ 40 U/L							1.252	
SGOT ≥ 35 U/L	1.272	1.198	1.4	1.454	1.281			
Bilirubin < 0.2 mg/dL						2.611		
Serum creatinine ≥ 3.0 mg/dL							1.626	
Serum creatinine ≥ 2.0 mg/dL		1.233	1.637					
Serum creatinine ≥ 1.2 mg/dL	1.454		1.721	1.566	1.202		1.31	
BUN ≥ 60 mg/dL				1.388				
BUN ≥ 25 mg/dL			1.362	1.43				
BUN ≥ 20 mg/dL	1.355	1.357	1.344		1.404	1.278	1.415	1.156
Serum sodium < 130 mEq/L		1.233						
Serum sodium ≥ 146 mEq/L		1.482	1.432	1.586	1.68	1.501	1.499	
Alkaline phosphatase < 110 mEq/L								1.487
CRP > 10 mg/dL								1.353
INR of PT values ≥ 1.67	1.44	1.239						
PT < 10 s						1.232		1.157
PTT < 30 s	1.181							1.137

ADL = activities of daily living; ASA = American Society of Anesthesiologists Physical Status; AST = aspartate amino transferase; BUN = blood urea nitrogen; COPD = chronic obstructive pulmonary disease; CRP = C-reactive protein; CVA = cerebrovascular accident; WBC = white blood cell.

risk factors of mortality in patients with ADP.<sup>15</sup> Preoperative variables associated with organ dysfunction tended to be included as risk factors in most of the risk models: preoperative ventilation/pneumonia, acute renal failure, bleeding disorders, low white blood cell count, low albumin level, and elevation of blood urea nitrogen.<sup>15</sup> High serum sodium levels, indicative of

severe dehydration in patients, were also identified. In contrast, the risk model for SSI, which was poorly associated with mortality ( $r = 0.107$ ), showed a relatively low C-index (0.688) compared with the other risk models. Risk factors such as pulmonary, renal, and cerebral disorders were not included in the risk model. The key part of these risk models is that variables




FIGURE 1. Receiver operating characteristic (ROC) curves of each postoperative complication was shown with the C-indices and 95% CIs of each occurrence. ROC = receiver operating characteristic, CIs = confidence intervals.

that were not included as risk factors of mortality were picked up as predictors of morbidities leading to mortality. This will help to improve the postoperative management of patients with ADP.

There are several limitations to this study. First, although these risk models for morbidities effectively predicted their occurrence based on preoperative variables, the source of infection and degree of its control would affect mortality and morbidity. These intraoperative parameters will be evaluated in a future study. Second, in the NCD data-entry system, the final outcome of each morbidity, whether it improved, was unresolved, led to death, and was not recorded. It is not possible to relate each morbidity directly to mortality, although most fatal cases feature multiple organ failure at the end.

ADP is a clinically distinct entity requiring life-saving emergency surgery and intensive care. We created risk models for morbidities in critically ill patients with ADP, using variables recorded by the NCD comparable to those of the ACS-NSQIP, and these models performed well. These models could be formatted to feed information back to the NCD and can be expected to improve the quality of the surgical and postoperative care of patients with ADP.

#### ACKNOWLEDGMENTS

The authors thank all data managers and hospitals participating in this NCD project for their great efforts in entering the data, and also thank the members and the working members of the JSGS database committee.

#### REFERENCES

- Seiler CA, Brugger L, Forssmann U, et al. Conservative surgical treatment of diffuse peritonitis. *Surgery*. 2000;127:178–184.
- Ordonez CA, Puyana JC. Management of peritonitis in the critically ill patient. *Surg Clin North Am*. 2006;86:1323–1349.
- Blot S, De Waele JJ. Critical issues in the clinical management of complicated intra-abdominal infections. *Drugs*. 2005;65:1611–1620.
- Pieracci FM, Barie PS. Management of severe sepsis of abdominal origin. *Scand J Surg*. 2007;96:184–196.
- Knaus WA, Draper EA, Wagner DP, et al. APACHE II: a severity of disease classification system. *Crit Care Med*. 1985;13:818–829.
- Vincent JL, Moreno R, Takala J, et al. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. On behalf of the Working Group on Sepsis-Related Problems of the European Society of Intensive Care Medicine. *Intensive Care Med*. 1996;22:707–710.
- Linder MM, Wacha H, Feldmann U, et al. The Mannheim peritonitis index. An instrument for the intraoperative prognosis of peritonitis. *Chirurg*. 1987;58:84–92.
- Huffman KM, Cohen ME, Ko CY, et al. A Comprehensive Evaluation of Statistical Reliability in ACS NSQIP Profiling Models. *Ann Surg*. 2014.
- Raval MV, Cohen ME, Ingraham AM, et al. Improving American College of Surgeons National Surgical Quality Improvement Program risk adjustment: incorporation of a novel procedure risk score. *J Am Coll Surg*. 2010;211:715–723.
- Hall BL, Hamilton BH, Richards K, et al. Does surgical quality improve in the American College of Surgeons National Surgical Quality Improvement Program: an evaluation of all participating hospitals. *Ann Surg*. 2009;250:363–376.
- Turner PL, Ilano AG, Zhu Y, et al. ACS-NSQIP criteria are associated with APACHE severity and outcomes in critically ill surgical patients. *J Am Coll Surg*. 2011;212:287–294.
- Miyata H, Gotoh M, Hashimoto H, et al. Challenges and prospects of a clinical database linked to the board certification system. *Surg Today*. 2014;44:1991–1999.
- Gotoh M, Miyata H, Hashimoto H, et al. National Clinical Database feedback implementation for quality improvement of cancer treatment in Japan: from good to great through transparency. *Surg Today*. 2015.
- Viehl CT, Kraus R, Zurcher M, et al. The Acute Physiology and Chronic Health Evaluation II score is helpful in predicting the need of relaparotomies in patients with secondary peritonitis of colorectal origin. *Swiss Med Wkly*. 2012;142:w13640.
- Nakagoe T, Miyata H, Gotoh M, et al. Surgical risk model for acute diffuse peritonitis based on a Japanese nationwide database: an initial report on the surgical and 30-day mortality. *Surg Today*. 2014.
- Takeuchi H, Miyata H, Gotoh M, et al. A risk model for esophagectomy using data of 5354 patients included in a Japanese nationwide web-based database. *Ann Surg*. 2014;260:259–266.
- Watanabe M, Miyata H, Gotoh M, et al. Total gastrectomy risk model: data from 20,011 Japanese patients in a nationwide internet-based database. *Ann Surg*. 2014;260:1034–1039.
- Kurita N, Miyata H, Gotoh M, et al. Risk model for distal gastrectomy when treating gastric cancer on the basis of data from 33,917 Japanese patients collected using a nationwide web-based data entry system. *Ann Surg*. 2015.
- Kobayashi H, Miyata H, Gotoh M, et al. Risk model for right hemicolectomy based on 19,070 Japanese patients in the National Clinical Database. *J Gastroenterol*. 2014;49:1047–1055.
- Matsubara N, Miyata H, Gotoh M, et al. Mortality after common rectal surgery in Japan: a study on low anterior resection from a newly established nationwide large-scale clinical database. *Dis Colon Rectum*. 2014;57:1075–1081.
- Kenjo A, Miyata H, Gotoh M, et al. Risk stratification of 7,732 hepatectomy cases in 2011 from the National Clinical Database for Japan. *J Am Coll Surg*. 2014;218:412–422.
- Kimura W, Miyata H, Gotoh M, et al. A pancreaticoduodenectomy risk model derived from 8575 cases from a national single-race population (Japanese) using a web-based data entry system: the 30-day and in-hospital mortality rates for pancreaticoduodenectomy. *Ann Surg*. 2014;259:773–780.
- Artigas A, Niederman MS, Torres A, et al. What is next in sepsis: current trials in sepsis. *Expert Rev Anti Infect Ther*. 2012;10:859–862.
- Hutchins NA, Unsinger J, Hotchkiss RS, et al. The new normal: immunomodulatory agents against sepsis immune suppression. *Trends Mol Med*. 2014;20:224–233.
- Vacanti CJ, VanHouten RJ, Hill RC. A statistical analysis of the relationship of physical status to postoperative mortality in 68,388 cases. *Anesth Analg*. 1970;49:564–566.
- Prause G, Ratzenhofer-Comenda B, Pierer G, et al. Can ASA grade or Goldman's cardiac risk index predict peri-operative mortality? A study of 16,227 patients. *Anaesthesia*. 1997;52:203–206.
- Jones DR, Copeland GP, de Cossart L. Comparison of POSSUM with APACHE II for prediction of outcome from a surgical high-dependency unit. *Br J Surg*. 1992;79:1293–1296.
- Cerra FB, Negro F, Abrams J. APACHE II score does not predict multiple organ failure or mortality in postoperative surgical patients. *Arch Surg*. 1990;125:519–522.
- Copeland GP, Jones D, Walters M. POSSUM: a scoring system for surgical audit. *Br J Surg*. 1991;78:355–360.
- Whiteley MS, Prytherch DR, Higgins B, et al. An evaluation of the POSSUM surgical scoring system. *Br J Surg*. 1996;83:812–815.
- Prytherch DR, Whiteley MS, Higgins B, et al. POSSUM and Portsmouth POSSUM for predicting mortality. Physiological and Operative Severity Score for the enUmeration of Mortality and morbidity. *Br J Surg*. 1998;85:1217–1220.

32. Barie PS, Hydo LJ, Shou J, et al. Efficacy and safety of drotrecogin alfa (activated) for the therapy of surgical patients with severe sepsis. *Surg Infect*. 2006;7(Suppl 2):S77–S80.
33. Barie PS, Vogel SB, Dellinger EP, et al. A randomized, double-blind clinical trial comparing cefepime plus metronidazole with imipenem-cilastatin in the treatment of complicated intra-abdominal infections. Cefepime Intra-abdominal Infection Study Group. *Arch Surg*. 1997;132:1294–1302.
34. Farthmann EH, Schoffel U. Principles and limitations of operative management of intraabdominal infections. *World J Surg*. 1990;14:210–217.
35. Garcia-Sabrido JL, Tallado JM, Christou NV, et al. Treatment of severe intra-abdominal sepsis and/or necrotic foci by an 'open-abdomen' approach. *Arch Surg*. 1988;123:152–156.
36. Anaya DA, Nathens AB. Risk factors for severe sepsis in secondary peritonitis. *Surg Infect*. 2003;4:355–362.

## Effects of body mass index (BMI) on surgical outcomes: a nationwide survey using a Japanese web-based database

Motonari Ri<sup>1</sup> · Hiroaki Miyata<sup>2</sup> · Susumu Aikou<sup>1</sup> · Yasuyuki Seto<sup>1</sup> · Kohei Akazawa<sup>3</sup> · Masahiro Takeuchi<sup>4</sup> · Yoshiro Matsui<sup>5</sup> · Hiroyuki Konno<sup>6</sup> · Mitsukazu Gotoh<sup>6</sup> · Masaki Mori<sup>6</sup> · Noboru Motomura<sup>7</sup> · Shinichi Takamoto<sup>7</sup> · Yoshiki Sawa<sup>8</sup> · Hiroyuki Kuwano<sup>9</sup> · Norihiro Kokudo<sup>10</sup>

Received: 12 June 2015 / Accepted: 10 July 2015 / Published online: 12 August 2015  
© Springer Japan 2015

### Abstract

**Purpose** To define the effects of body mass index (BMI) on operative outcomes for both gastroenterological and cardiovascular surgery, using the National Clinical Database (NCD) of the Japanese nationwide web-based database.

**Methods** The subjects of this study were 288,418 patients who underwent typical surgical procedures between January 2011 and December 2012. There were eight gastroenterological procedures, including esophagectomy, distal gastrectomy, total gastrectomy, right hemicolectomy, low anterior resection, hepatectomy of >1 segment excluding the lateral segment, pancreaticoduodenectomy, and surgery for acute diffuse peritonitis ( $n = 232,199$ ); and five cardiovascular procedures, including aortic valve replacement, total arch replacement (TAR), descending thoracic aorta replacement (descending TAR), and on- or off-pump coronary artery bypass grafting ( $n = 56,219$ ). The relationships

of BMI with operation time and operative mortality for each procedure were investigated, using the NCD.

**Results** Operation times were longer for patients with a higher BMI. When a BMI cut-off of 30 was used, the operation time for obese patients was significantly longer than that for non-obese patients, for all procedures except esophagectomy ( $P < 0.01$ ). The mortality rate based on BMI revealed a U-shaped distribution, with both underweight and obese patients having high mortality rates for almost all procedures. **Conclusions** This Japanese nationwide study provides solid evidence to reinforce that both obesity and excessively low weight are factors that impact operative outcomes significantly.

**Keywords** National clinical database · Nationwide web-based database · Body mass index · Operation time · Operative mortality

On behalf of the Japan Surgical Society.

✉ Yasuyuki Seto  
seto-tyk@umin.ac.jp

<sup>1</sup> Department of Gastrointestinal Surgery, Graduate School of Medicine, The University of Tokyo, 7-3-1, Hongo, Bunkyo-ku, Tokyo 113-8655, Japan

<sup>2</sup> National Clinical Database (NCD), Tokyo, Japan

<sup>3</sup> Department of Medical Informatics, Niigata University Medical and Dental Hospital, Niigata, Japan

<sup>4</sup> Department of Clinical Medicine, School of Pharmacy, Kitasato University, Kanagawa, Japan

<sup>5</sup> Department of Cardiovascular and Thoracic Surgery, Hokkaido University Graduate School of Medicine, Hokkaido, Japan

<sup>6</sup> The Japanese Society of Gastroenterological Surgery, Tokyo, Japan

<sup>7</sup> Japan Cardiovascular Surgery Database Organization, Tokyo, Japan

<sup>8</sup> Department of Cardiovascular Surgery, Osaka University of Medicine, Osaka, Japan

<sup>9</sup> Department of General Surgical Science, Graduate School of Medicine, Gunma University, Gunma, Japan

<sup>10</sup> Hepato-Biliary-Pancreatic Surgery Division, Department of Surgery, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan

**Table 1** Summary of past studies on the effect of body mass index on surgical outcomes

Year	References	<i>n</i>	Disease	Operation	BMI cut-off values	Operation time (min) <i>P</i> value	Mortality (%) <i>P</i> value
2013	Bhayani et al. [7]	794	–	Subtotal or total esophagectomy	18.5–25 vs. 35≤	316 vs. 308 n.s.	3.3 vs. 2.3 ns
2013	Sugimoto et al. [8]	216	Gastric cancer	Laparoscopic distal gastrectomy	25> vs. 25≤	200 vs. 212 0.005	0 vs. 0 ns
2014	Makino et al. [9]	152	Colon cancer	Laparoscopic colectomy	30> vs. 30≤	157 vs. 182 0.008	0 vs. 0 ns
2012	Saunders et al. [10]	403	–	Liver resection	18.5–24.9 vs. 40≤	306 vs. 402 ns	6 vs. 27 0.05
2014	El Nakeeb et al. [11]	471	Some	Pancreaticoduodenectomy	25> vs. 25≤	300 vs. 321 0.003	0.8 vs. 7.1 0.001
2012	Zdichavsky et al. [12]	596	Some	Laparoscopic cholecystectomy	30> vs. 30≤	73.5 vs. 87 <0.001	–
2011	Alam et al. [13]	13,115	–	Coronary artery bypass graft	30> vs. 30≤	39.3 vs. 41.7* <0.0001	3.7 vs. 2.9 ns
2012	Smith et al. [14]	1066	Aortic stenosis	Aortic valve replacement	25> vs. 25–30 vs. 30<	72.9 vs. 74.9 vs. 76.5** ns	5.6 vs. 2.8 vs. 4.5 ns

*BMI* body mass index

\* Aortic clamp time, \*\* cross-clamp time

## Introduction

With the proportion of obese people in world populations increasing, obesity-related illnesses have become a major concern globally [1]. The World Health Organization (WHO) has documented that 34 % of adult men and 35 % of adult women have a body mass index (BMI) of more than 25, and that 10 % of adult men and 14 % of adult women worldwide are obese (BMI ≥ 30) [2]. In the United States, 34.9 % of adults are obese [3]. Japan is no exception to this trend, with a National Health and Nutrition Survey (2012) conducted by the Ministry of Health, Labour and Welfare revealing that 29.1 % of adult men and 19.4 % of adult women have a BMI of more than 25. A rising BMI and a growing obese population in Japan have been reported, especially among men, as well as middle-aged and older women [4–7]. Although there have been a number of studies on the relationship between obesity and operative outcomes, their conclusions are inconsistent (Table 1) [8–15]. Moreover, many of these studies were performed at individual institutions and there have been no large-scale surveys comparing different surgical areas. Ultimately, it has not yet been established whether obesity adversely affects operative outcomes, such as the operation time and risk. Thus, we conducted a cross-sectional investigation of the effects of BMI on operation time and operative mortality for both gastroenterological and cardiovascular surgery, using a Japanese nationwide database.

## Methods

### The nationwide database system

In January 2011, Japan's National Clinical Database (NCD) became accessible on-line, with the cooperation of some of the nation's surgical associations. The NCD is a large-scale nationwide database, in which data from over 1,200,000 surgical cases were collected from more than 3500 hospitals in 2011. The information about operations performed nationwide was registered in the NCD by data management departments from the participating institutions. The data were evaluated annually using a web-based data management system to assure data traceability. This system also validated data consistency by randomly inspecting the participating institutions. Several clinical studies conducted by various societies have used the NCD data [16–18].

### Patients

We analyzed data from a total 288,418 patients who underwent a typical procedure in the areas of gastroenterological and cardiovascular surgery between 1 January 2011 and 31 December 2012. There were 13 procedures in total: esophagectomy (Eso), distal gastrectomy (DG), total gastrectomy (TG), right hemicolectomy (RHC), low anterior resection (LAR), hepatectomy of more than one segment apart from the lateral segment (Hx),

pancreaticoduodenectomy (PD), surgery for acute diffuse peritonitis (ADP), aortic valve replacement (AVR), total arch replacement (TAR), descending thoracic aorta replacement (descending TAR), coronary artery bypass graft (CABG) on-pump, and CABG off-pump. Patients who did not consent to their data being used were excluded from this analysis. Records with missing data for age, sex, or status at postoperative day (POD) 30 were also excluded, as were data related to patients with a BMI of less than 10 or over 50 as these values were obviously erroneous.

### Study outcomes

The three outcomes examined using the NCD were BMI, operation time, and operative mortality. BMI is calculated as weight (in kilograms) divided by the square of height (in meters). Operative mortality included all deaths occurring within the index hospitalization period, regardless of the length of hospital stay (up to 90 days), or after hospital discharge (within 30 days after surgery). We evaluated influences of stratified BMI on operation time and operative mortality.

### Statistical analyses

All statistical analyses were conducted using SPSS (version 20). We performed univariate comparisons of BMI and operation time, using the unpaired Student's *t* test and Chi-square test. A *P* value of 0.05 was considered significant.

### Results

Table 2 summarizes the backgrounds of each procedure. There were 232,199 patients undergoing 1 or more of 8 gastroenterological surgery procedures and 56,219 patients undergoing 1 or more of 5 cardiovascular surgery procedures.

#### BMI distribution

Figure 1 shows the BMI distribution of all the patients whose data were analyzed. Approximately 60 % of the procedures (61.2–69.9 %) were performed on patients with a normal physique, with a BMI of 18.5–25. Of the remaining patients, 2.6 % were obese, 20.3 % were overweight, and 13.4 % were underweight. There were differences in BMI distributions between patients undergoing gastroenterological and cardiovascular procedures. The largest category of patients undergoing gastroenterological surgery had a BMI of 18.5–22, and the largest category of those undergoing cardiovascular surgery had a BMI of 22–25. The proportion of overweight patients (with a BMI of 25–30) among those

**Table 2** Background parameters

Operation	Number	Sex (male/female)	Age (mean $\pm$ SD)
Eso	10,825	9124/1701	66.1 $\pm$ 9.3
DG	63,650	42,438/21,212	69.1 $\pm$ 11.5
TG	37,817	27,877/9940	69.0 $\pm$ 11.0
RHC	37,750	18,878/18,872	71.7 $\pm$ 12.1
LAR	33,217	21,436/11781	66.2 $\pm$ 11.6
Hx	14,903	10,456/4447	67.0 $\pm$ 11.5
PD	17,485	10,841/6644	68.5 $\pm$ 10.2
ADP	16,552	10,020/6532	64.9 $\pm$ 18.6
AVR	14,827	7914/6913	72.4 $\pm$ 10.7
TAR	10,594	7421/3173	69.5 $\pm$ 11.6
Descending TAR	5605	4129/1476	69.6 $\pm$ 12.3
CABG on-pump	9221	7232/1989	67.7 $\pm$ 9.9
CABG off-pump	15,972	12,547/3425	69.1 $\pm$ 9.7


*Eso* esophagectomy, *DG* distal gastrectomy, *TG* total gastrectomy, *RHC* right hemicolectomy, *LAR* low anterior resection, *Hx* hepatectomy performed for >1 segment except for the lateral segment, *PD* pancreaticoduodenectomy, *ADP* surgery for acute diffuse peritonitis, *AVR* aortic valve replacement, *TAR* total arch replacement, *CABG* coronary artery bypass graft

undergoing gastroenterological surgery was less than 20 %, whereas the proportion of overweight patients among those undergoing cardiovascular surgery was more than 20 %. The proportion of obese patients did not exceed 3 % in any of the gastroenterological surgery groups, but it exceeded 4 % in all of the cardiovascular surgery groups, being especially low in the Eso (1 %) group and especially high in the on-pump (5 %) and off-pump (4.6 %) CABG groups. The proportion of underweight patients (with a BMI of <18.5) among those undergoing gastroenterological surgery, apart from Hx, was at least 10 %, whereas among those undergoing cardiovascular surgery, apart from AVR, it was no more than 10 %. This rate was high for ADP (23.3 %), Eso (21.1 %), and PD (14.9 %), but low for off-pump (5 %) and on-pump (5.1 %) CABG.

#### Operation time by BMI

Figure 2 shows the operation times for all procedures, which became longer as the BMI rose. When a BMI cut-off of 30 was used, the operation time for obese patients was significantly longer than that for those who were not obese, for all procedures except Eso (Table 3). Among the gastroenterological operations, the operation time for obese patients was prolonged for PD (56  $\pm$  8.3 min, mean  $\pm$  SE), TG (50  $\pm$  4.0 min), and LAR (50  $\pm$  3.9 min), while among the cardiovascular operations, the operation time for obese patients was prolonged for TAR (55  $\pm$  8.4 min) and descending TAR (54  $\pm$  13.8 min). The operation times were originally longer for men than for women, for all of

**Fig. 1 a** Body mass index (BMI) distribution in the gastroenterological surgery procedures (%). *Eso* esophagectomy, *DG* distal gastrectomy, *TG* total gastrectomy, *RHC* right hemicolectomy, *LAR* low anterior resection, *Hx* hepatectomy performed for >1 segment apart from the lateral segment, *PD* pancreaticoduodenectomy, *ADP* surgery for acute diffuse peritonitis. **b** BMI distribution in the cardiovascular surgery procedures (%). *AVR* aortic valve replacement, *TAR* total arch replacement, *CABG on pump* coronary artery bypass graft



the procedures (Table 4). The operation time was significantly longer for obese men than for men who were not obese, and this result was consistent among all procedures. The differences in the effects of obesity on operation times between men and women varied among the procedures. The prolongation of operation time by at least 60 min because of obesity was greater in men than in women undergoing LAR, TAR, and descending TAR. Among patients undergoing Hx, the impact of obesity on operation time was greater for women than for men.

**Influence of BMI on operative mortality**


Figure 3 shows the operative mortality rates. The mortality rate according to BMI had a U-shaped distribution for most procedures, with both underweight and obese patients having higher mortality rates. Overweight patients tended

to have the lowest mortality risk among those undergoing cardiovascular surgery, but not among those undergoing gastroenterological surgery. Obesity tended to have a small effect on mortality among patients undergoing Eso or DG. Among patients undergoing PD, those who were underweight had the lowest mortality rate.

**Discussion**

We conducted a large-scale trans-disciplinary study on the effects of BMI on gastroenterological vs. cardiovascular surgery, using NCD accumulated in Japan. Several previous studies have demonstrated a cause-and-effect relationship between BMI and operative outcomes, including operation time and operative mortality in various fields, including gastroenterological procedures for esophageal

**Fig. 2 a** Operation times for the gastroenterological surgery procedures (min). *Eso* esophagectomy, *DG* distal gastrectomy, *TG* total gastrectomy, *RHC* right hemicolectomy, *LAR* low anterior resection, *Hx* hepatectomy performed for >1 segment, apart from the lateral segment, *PD* pancreaticoduodenectomy, *ADP* surgery for acute diffuse peritonitis. **b** Operation times for the cardiovascular surgery procedures (min). *AVR* aortic valve replacement, *TAR* total arch replacement, *CABG* coronary artery bypass graft



cancer [8], gastric cancer [9], colon cancer [10], liver cancer [11], pancreatic cancer [12], and laparoscopic cholecystectomy [13], as well as cardiac procedures, including CABG and valve replacement [14, 15]. However, according to the literature, the influences of BMI on surgery vary among operative procedures and even for the same procedure, so the effects of BMI on operative outcomes remain controversial. To date, two studies involving a large number of cases have been reported. In a study of 30,765 patients undergoing surgery for gastric cancer or colorectal cancer in Japan, Yasanuga et al. found that obese patients had significantly higher rates of postoperative complications and that underweight patients had significantly higher rates of

both postoperative complications and operative mortality than those with a normal BMI [19]. In a study of 13,115 patients undergoing CABG, Alam et al. found that obese patients had significantly prolonged total circulatory bypass and aortic clamp times, as well as an increased risk of postoperative respiratory failure, postoperative renal failure, and surgical site infections, but not of operative mortality [14]. The number of patients in the present study was greater than that in either of the previous studies and we also examined data for various procedures in two major fields. We confirmed that BMI has similar effects on nearly all procedures in the fields of gastroenterological and cardiovascular surgery in this nationwide survey.



**Table 3** Operation times (min) according to a body mass index cut-off value of 30

Operation	Total number (BMI < 30:30 ≤ BMI)	Operation time (min)		Prolongation (min)	P value
		BMI < 30	30 ≤ BMI		
Eso	10,825 (10,727:98)	479	505	26	ns
DG	63,650 (62,391:1259)	248	288	40	<0.001
TG	37,817 (37,215:602)	279	329	50	<0.001
RHC	37,750 (36,850:900)	197	233	36	<0.001
LAR	33,217 (32,409:808)	262	312	50	<0.001
Hx	14,903 (14,459:444)	384	424	40	<0.001
PD	17,485 (17,229:256)	468	524	56	<0.001
ADP	16,552 (16,117:435)	127	148	21	<0.001
AVR	14,827 (14,257:570)	304	321	17	<0.001
TAR	10,594 (10,114:480)	426	481	55	<0.001
Descending TAR	5605 (5356:249)	274	328	54	<0.001
CABG on-pump	9221 (8760:461)	358	379	21	<0.001
CABG off-pump	15,972 (15,238:734)	303	328	25	<0.001

Student's *t* test was used to compare the mean operation time between obese (BMI ≥ 30) and non-obese (BMI < 30) patients. Statistical significance was set at *P* < 0.05

*Eso* esophagectomy, *DG* distal gastrectomy, *TG* total gastrectomy, *RHC* right hemicolectomy, *LAR* low anterior resection, *Hx* hepatectomy performed for >1 segment apart from the lateral segment, *PD* pancreaticoduodenectomy, *ADP* surgery for acute diffuse peritonitis, *AVR* aortic valve replacement, *TAR* total arch replacement, *CABG* coronary artery bypass graft, *BMI* body mass index

Operation times were longer for patients with a higher BMI. When a BMI cut-off of 30 was used, the operation time was significantly longer for obese patients for all procedures except Eso. The procedure associated with the greatest prolongation of operation time was PD, by 56 min, while AVR was prolonged by at least 17 min. The results of the previous studies were conflicting about the influences of obesity on operation time, but those of the present study allowed us to conclude that obesity prolongs the operation time for nearly all the procedures studied. Interestingly, for all procedures, the operation times were longer for men than for women, and they were significantly longer for men who were obese than men who were not. We also noted differences among procedures in the prolongation of operation time due to obesity between men and women. It is noteworthy that men typically have a body type characterized by visceral fat accumulation, whereas women tend to show subcutaneous fat accumulation. These body type differences may account for the differences in operative times and the prolongation of surgery because of obesity.

The mortality rates according to BMI showed a U-shaped distribution, with both underweight and obese patients having high mortality rates, except when PD was the procedure performed. Prior reports from Western countries showed similar mortality curves. These studies demonstrated the so-called 'obesity paradox' in that obesity exerts an advantageous effect on morbidity and mortality, compared with normal BMI, in surgical patients [20–22].

Davenport et al. [22] reported that patients with mild obesity, defined as a BMI over 30 but under 35, had a lower 30-day mortality rate than patients of normal weight following vascular surgery. The BMI category with the lowest mortality risk is lower in Asians than in Western populations because Asians have a higher percentage of body fat than Westerners at the same BMI [3, 23]. Deurenberg et al. [24] reported the BMI values of East Asians to be lower, by 1.9–3.2 %, than in Western people when the percentage of body fat was the same. The adverse effect of obesity on short-term prognosis in patients undergoing surgery for a malignant disease would be due to not only the high technical difficulty of the necessary surgical procedures in the obese patient, but to BMI-dependent differences in the severity of the cancer. However, the influence of BMI on malignancy remains controversial [25, 26]. Mullen et al. [27] reported that underweight patients had a significantly higher postoperative death rate, with an odds ratio (OR) of 5.24 and a 95 % confidence interval (CI) of 1.7–16.2, after major surgery for gastrointestinal cancer. As in previous studies, the underweight patients in the present series had higher mortality, which we speculate is related to a reduced capacity to tolerate surgery because of their underlying nutritional deficiencies and probable advanced stages of gastroenterological cancer.

This study has the advantages of being nationwide and including a very large number of cases; however, there are limitations because of its retrospective nature and cross-sectional design. Moreover, certain details were not

**Table 4** Operation times (min) according to a body mass index cut-off value of 30 and gender

Operation	Gender	Total number (BMI < 30:30 ≤ BMI)	Operation time (min)		Prolongation (min)	P value
			BMI < 30	30 ≤ BMI		
Eso	Male	9124 (9036:88)	488	521	33	<0.05
	Female	1701 (1691:10)	432	363	−69	ns
DG	Male	42438 (41674:764)	255	303	48	<0.001
	Female	21212 (20717:495)	234	265	31	<0.001
TG	Male	27877 (27484:393)	284	338	54	<0.001
	Female	9940 (9731:209)	264	313	49	<0.001
RHC	Male	18878 (18505:373)	206	251	45	<0.001
	Female	18872 (18345:527)	187	221	34	<0.001
LAR	Male	21,436 (21,013:423)	269	334	65	<0.001
	Female	11,781 (11,396:385)	248	287	39	<0.001
Hx	Male	10,456 (10,162:294)	392	428	36	<0.001
	Female	4447 (4297:150)	368	417	49	<0.001
PD	Male	10,841 (10,715:126)	480	540	60	<0.001
	Female	6644 (6514:130)	448	508	60	<0.001
ADP	Male	10,020 (9790:230)	127	153	26	<0.001
	Female	6532 (6327:205)	126	142	16	<0.001
AVR	Male	7914 (7687:227)	314	354	40	<0.001
	Female	6913 (6570:343)	292	300	8	ns
TAR	Male	7421 (7075:346)	427	494	67	<0.001
	Female	3173 (3039:134)	422	450	28	<0.05
Descending TAR	Male	4129 (3945:184)	275	343	68	<0.001
	Female	1476 (1411:65)	269	286	17	ns
CABG on-pump	Male	7232 (6885:347)	360	380	20	<0.001
	Female	1989 (1875:114)	351	375	24	<0.05
CABG off-pump	Male	12,547 (12,020:527)	305	334	29	<0.001
	Female	3425 (3218:207)	293	312	19	<0.01

Student's *t* test was used to compare the mean operation time between obese (BMI ≥ 30) and non-obese (BMI < 30) patients by sex. Statistical significance was set at  $P < 0.05$

*Eso* esophagectomy, *DG* distal gastrectomy, *TG* total gastrectomy, *RHC* right hemicolectomy, *LAR* low anterior resection, *Hx* hepatectomy performed for >1 segment apart from the lateral segment, *PD* pancreaticoduodenectomy, *ADP* surgery for acute diffuse peritonitis, *AVR* aortic valve replacement, *TAR* total arch replacement, *CABG* coronary artery bypass graft, *BMI* body mass index

assessed, such as the presence or absence of endoscopic surgery, emergency vs. non-urgent status, and co-existent illnesses, including hypertension and diabetes. Various factors other than BMI can affect operative outcomes, and these factors differ among operative procedures. It is important to analyze precise risks, including all relevant factors, for a broader range of operative procedures in the future. However, we should consider both obesity and very low body weight as the key factors impacting outcomes.


In the present study, a high BMI was shown to influence health economics and inflict excess burden on surgeons, considering the prolonged operation times for all surgical procedures evaluated. Yet, there are no additional fees in relation to obesity, in either Japan or, to our knowledge,

any other country. Given the added burden associated with this condition, it is anticipated that the medical economics of providing surgical treatment for obese patients will be reviewed.

## Conclusion

We investigated the effects of BMI on both gastroenterological and cardiovascular surgical procedures based on data from 288,418 patients, obtained from a Japanese nationwide database. The operation time was significantly longer for patients who were obese than for those who were not, for all procedures except Eso. Furthermore, both

**Fig. 3 a** Operative mortality for the gastroenterological surgery procedures (%). *Eso* esophagectomy, *DG* distal gastrectomy, *TG* total gastrectomy, *RHC* right hemicolectomy, *LAR* low anterior resection, *Hx* hepatectomy performed for >1 segment except for the lateral segment, *PD* pancreaticoduodenectomy, *ADP* surgery for acute diffuse peritonitis. **b** Operative mortality for the cardiovascular surgery procedures (%). *AVR* aortic valve replacement, *TAR* total arch replacement, *CABG* coronary artery bypass graft



obese and underweight patients had high mortality rates. This Japanese nationwide study provides further evidence that obesity and very low bodyweight are both factors with significant adverse effects on operative outcomes.

**Compliance with ethical standards**

**Conflict of interest** None of the authors has any conflicts of interest to disclose.

**Funding/support** This study was funded by the Japan Surgical Society.

**Role of the funders/sponsors** The NCD data used in this study were collected by the Japan Surgical Society. The Japan Surgical Society reviewed and approved this article before its submission.

**References**

1. Krishnamurthi RV, Feigin VL, Forouzanfar MH, Mensah GA, Connor M, Bennett DA, et al. Global and regional burden of first-ever ischaemic and haemorrhagic stroke during 1990–2010: findings from the Global Burden of Disease Study 2010. *Lancet Glob Health*. 2013. doi:10.1016/S2214-4109X(13)70089-5.
2. World Health Organization. Global status report on noncommunicable disease 2010. [http://www.who.int/nmh/publications/ncd\\_report\\_full\\_en.pdf](http://www.who.int/nmh/publications/ncd_report_full_en.pdf). Accessed 1 Oct 2014).
3. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of childhood and adult obesity in the United States, 2011–2012. *JAMA*. 2014;311:806–14.
4. Funatogawa I, Funatogawa T, Nakao M, Karita K, Yano E. Changes in body mass index by birth cohort in Japanese adults:

- results from the National Nutrition Survey of Japan 1956–2005. *Int J Epidemiol.* 2009;38:83–92.
5. Matsushita Y, Takahashi Y, Mizoue T, Inoue M, Noda M, Tsugane S. Overweight and obesity trends among Japanese adults: a 10-year follow-up of the JPHC Study. *Int J Obes (Lond).* 2008;32:1861–7.
  6. Yoshiike N, Seino F, Tajima S, Arai Y, Kawano M, Furuhashi T, et al. Twenty-year changes in the prevalence of overweight in Japanese adults: the National Nutrition Survey 1976–95. *Obes Rev.* 2002;3:183–90.
  7. Kouda K, Nakamura H, Nishio N, Fujita Y, Takeuchi H, Iki M. Trends in body mass index, blood pressure, and serum lipids in Japanese children: Iwata population-based annual screening (1993–2008). *J Epidemiol.* 2010;20:212–8.
  8. Bhayani NH, Gupta A, Dunst CM, Kurian AA, Halpin VJ, Swanstrom LL. Does morbid obesity worsen outcomes after esophagectomy? *Ann Thorac Surg.* 2013;95:1756–61.
  9. Sugimoto M, Kinoshita T, Shibasaki H, Kato Y, Gotohda N, Takahashi S, et al. Short-term outcome of total laparoscopic distal gastrectomy for overweight and obese patients with gastric cancer. *Surg Endosc.* 2013;27:4291–6.
  10. Makino T, Trencheva K, Shukla PJ, Rubino F, Zhuo C, Pavoors RS, et al. The influence of obesity on short- and long-term outcomes after laparoscopic surgery for colon cancer: a case-matched study of 152 patients. *Surgery.* 2014;156:661–8.
  11. Saunders JK, Rosman AS, Neihaus D, Gouge TH, Melis M. Safety of hepatic resections in obese veterans. *Arch Surg.* 2012;147:331–7.
  12. El Nakeeb A, Hamed H, Shehta A, Askr W, El Dosoky M, Said R, et al. Impact of obesity on surgical outcomes post-pancreaticoduodenectomy: a case-control study. *Int J Surg.* 2014;12:488–93.
  13. Zdichavsky M, Bashin YA, Blumenstock G, Zieker D, Meile T, Konigsrainer A. Impact of risk factors for prolonged operative time in laparoscopic cholecystectomy. *Eur J Gastroenterol Hepatol.* 2012;24:1033–8.
  14. Alam M, Siddiqui S, Lee VV, Elayda MA, Nambi V, Yang EY, et al. Isolated coronary artery bypass grafting in obese individuals: a propensity matched analysis of outcomes. *Circ J.* 2011;75:1378–85.
  15. Smith RL 2nd, Herbert MA, Dewey TM, Brinkman WT, Prince SL, Ryan WH, et al. Does body mass index affect outcomes for aortic valve replacement surgery for aortic stenosis? *Ann Thorac Surg.* 2012;93:742–6.
  16. Watanabe M, Miyata H, Gotoh M, Baba H, Kimura W, Tomita N, et al. Total gastrectomy risk model: data from 20,011 Japanese patients in a Nationwide Internet-Based Database. *Ann Surg.* 2014;260:1034–9.
  17. Kimura W, Miyata H, Gotoh M, Hirai I, Kenjo A, Kitagawa Y, et al. A pancreaticoduodenectomy risk model derived from 8575 cases from a national single-race population (Japanese) using a web-based data entry system: the 30-day and in-hospital mortality rates for pancreaticoduodenectomy. *Ann Surg.* 2014;259:773–80.
  18. Dohi M, Miyata H, Doi K, Okawa K, Motomura N, Takamoto S, et al. The off-pump technique in redo coronary artery bypass grafting reduces mortality and major morbidities: propensity score analysis of data from the Japan Cardiovascular Surgery Database. *Eur J Cardiothorac Surg.* 2015;47:299–308.
  19. Yasunaga H, Horiguchi H, Matsuda S, Fushimi K, Hashimoto H, Ayanian JZ. Body mass index and outcomes following gastrointestinal cancer surgery in Japan. *Br J Surg.* 2013;100:1335–43.
  20. Valentijn TM, Galal W, Tjeertes EK, Hoeks SE, Verhagen HJ, Stolker RJ. The obesity paradox in the surgical population. *Surg.* 2013;11:169–76.
  21. Mullen JT, Moorman DW, Davenport DL. The obesity paradox: body mass index and outcomes in patients undergoing nonbariatric general surgery. *Ann Surg.* 2009;250:166–72.
  22. Davenport DL, Xenos ES, Hosokawa P, Radford J, Henderson WG, Endean ED. The influence of body mass index obesity status on vascular surgery 30-day morbidity and mortality. *J Vasc Surg.* 2009;49:140–7.
  23. WHO Expert Consultation. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet.* 2004;363:157–63.
  24. Deurenberg P, Yap M, van Staveren WA. Body mass index and percent body fat: a meta analysis among different ethnic groups. *Int J Obes Relat Metab Disord.* 1998;22:1164–71.
  25. Seishima R, Okabayashi K, Hasegawa H, Sugiyama D, Ishii Y, Tsuruta M, et al. Obesity was associated with a decreased post-operative recurrence of rectal cancer in a Japanese population. *Surg Today.* 2014;44:2324–31.
  26. Kim SH, Park HS, Kim KH, Yoo H, Chae BJ, Bae JS, et al. Correlation between obesity and clinicopathological factors in patients with papillary thyroid cancer. *Surg Today.* 2015;45:723–9.
  27. Mullen JT, Davenport DL, Hutter MM, Hosokawa PW, Henderson WG, Khuri SF, et al. Impact of body mass index on perioperative outcomes in patients undergoing major intra-abdominal cancer surgery. *Ann Surg Oncol.* 2008;15:2164–72.