

working population. However, although practical methods, such as MetS screening, for a high-risk strategy exist, pragmatic approaches to a population strategy are still under development in Japan.

### Statement of Conflict of Interest

All authors declare that there are no conflicts of interest.

### Acknowledgments

This research was supported by Health and Labor Sciences Research Grants of the Ministry of Health, Labor and Welfare of Japan (Comprehensive Research on Life-Style Related Diseases including Cardiovascular Diseases and Diabetes Mellitus: H25-Junkankitou [Seishuu]-Ippan-013).

### REFERENCES

- Ministry of Health, Welfare and Labour. 2010 comprehensive survey of living conditions of the people on health and welfare. Tokyo Health and Welfare Statistics Association. 2012. <http://www.mhlw.go.jp/toukei/saikin/hw/k-tyosa/k-tyosa10/> (in Japanese).
- <http://law.e-gov.go.jp/htmldata/S47/S47HO057.html> (in Japanese).
- <http://law.e-gov.go.jp/htmldata/S47/S47F04101000032.html> (in Japanese).
- Sakurai H. Occupational safety and health in Japan: current situations and the future. *Ind Health*. 2012;50:253-260.
- Ikeda N, Saito E, Kondo N, et al. What has made the population of Japan healthy? *Lancet*. 2011;378:1094-1105.
- Saito I, Folsom AR, Aono H, Ozawa H, Ikebe T, Yamashita T. Comparison of fatal coronary heart disease occurrence based on population surveys in Japan and the USA. *Int J Epidemiol*. 2000;29:837-844.
- Lifestyle-Related Disease Control Bureau of the Ministry of Health, Labour and Welfare. Kokumin Eiyu no Genjyo 1998 [The National Nutrition Survey, Japan]. Tokyo: Daiichi Shuppan. 2000.
- Shimamoto T, Komachi Y, Inada H, et al. Trends for coronary heart disease and stroke and their risk factors in Japan. *Circulation*. 1989;79:503-515.
- Tanabe N, Iso H, Okada K, Nakamura Y, Harada A, Ohashi Y, Ando T, Ueshima H, Japan Arteriosclerosis Longitudinal Study Group. Serum total and non-high-density lipoprotein cholesterol and the risk prediction of cardiovascular events - the JALS-ECC -. *Circ J*. 2010;74:1346-1356.
- NIPPON DATA80 Research Group. Risk assessment chart for death from cardiovascular disease based on a 19-year follow-up study of a Japanese representative population. *Circ J*. 2006;70:1249-1255.
- Hirobe K, Terai T, Fujioka S, Goto K, Dohi S. 3M-Study Project Committee of the Japan Association of Occupational Physicians "San-yu-kai"; Morbidity of Myocardial Infarction Multi-center Study in Japan (3M Study): study design and event rates for myocardial infarction and coronary death by age category in Japanese workers. *Circ J*. 2005;69:767-773.
- Maruyama K, Hirobe K, Noda H, Iso H, Dohi S, Terai T, Fujioka S, Goto K, Horie S, Nakano S. Associations between blood lipid profiles and risk of myocardial infarction among Japanese male workers: 3M Study. *J Atheroscler Thromb*. 2009;16:714-721.
- Noda H, Maruyama K, Iso H, Dohi S, Terai T, Fujioka S, Goto K, Horie S, Nakano S, Hirobe K. 3M Study Project Committee of the Japan Association of Occupational Physicians "San-yu-kai"; Prediction of myocardial infarction using coronary risk scores among Japanese male workers: 3M Study. *J Atheroscler Thromb*. 2010;17:452-459.
- Rose G. The Strategy of Preventive Medicine. Oxford: Oxford University Press. 1992.
- Rose G. Sick individuals and sick populations. *Int J Epidemiol*. 2001;30:427-432.
- Okamura T, Tanaka T, Babazono A, et al. The high-risk and population strategy for occupational health promotion (HIPOP-OHP) study: study design and cardiovascular risk factors at the baseline survey. *J Hum Hypertens*. 2004;18:475-485.
- Okamura T, Tanaka T, Takebayashi T, et al. Methodological issues for a large-scale intervention trial of lifestyle modification: Interim assessment of the high-risk and population strategy for occupational health promotion (HIPOP-OHP) study. *Environ Health Prev Med*. 2004;9:137-143.
- Tanaka T, Okamura T, Yamagata Z, et al. Awareness and treatment of hypertension and hypercholesterolemia in Japanese workers: the High-risk and Population Strategy for Occupational Health Promotion (HIPOP-OHP) study. *Hypertens Res*. 2007;30:921-928.
- Kamon Y, Okamura T, Tanaka T, et al. Marital status and cardiovascular risk factors among middle-aged Japanese male workers: the High-risk and Population Strategy for Occupational Health Promotion (HIPOP-OHP) study. *J Occup Health*. 2008;50:348-356.
- Tamura U, Tanaka T, Okamura T, et al. Changes in weight, cardiovascular risk factors and estimated risk of coronary heart disease following smoking cessation in Japanese male workers: HIPOP-OHP study. *J Atheroscler Thromb*. 2010;17:12-20.
- Yoshita K, Tanaka T, Kikuchi Y, et al. The evaluation of materials to provide health-related information as a population strategy in the worksite: The high-risk and population strategy for occupational health promotion (HIPOP-OHP) study. *Environ Health Prev Med*. 2004;9:144-151.
- Naito M, Nakayama T, Okamura T, et al. Effect of a 4-year workplace-based physical activity intervention program on the blood lipid profiles of participating employees: the high-risk and population strategy for occupational health promotion (HIPOP-OHP) study. *Atherosclerosis*. 2008;197:784-790.
- Tanaka H, Yamato H, Tanaka T, et al. Effectiveness of a low-intensity intra-worksites intervention on smoking cessation in Japanese employees: a three-year intervention trial. *J Occup Health*. 2006;48:175-182.
- Hayashino Y, Fukuhara S, Okamura T, et al. A prospective study of passive smoking and risk of diabetes in a cohort of workers: the High-Risk and Population Strategy for Occupational Health Promotion (HIPOP-OHP) study. *Diabetes Care*. 2008;31:732-734.
- Funahashi T. Definition of metabolic syndrome in Japan—Concept and perspective. *Nippon Rinsho*. 2007;65(Suppl 7):84-90. (in Japanese).
- Kaplan NM. The deadly quartet: Upper-body obesity, glucose intolerance, hyper-triglyceridemia, and hypertension. *Arch Intern Med*. 1989;149:1514-1520.
- Kadota A, Hozawa A, Okamura T, et al. Relationship between metabolic risk factor clustering and cardiovascular mortality stratified by high blood glucose and obesity: NIPPON DATA90, 1990–2000. *Diabetes Care*. 2007;30:1533-1538.

28. Iso H, Sato S, Kitamura A, et al. Metabolic syndrome and the risk of ischemic heart disease and stroke among Japanese men and women. *Stroke*. 2007;38:1744-1751.
29. Hata J, Doi Y, Ninomiya T, et al. The effect of metabolic syndrome defined by various criteria on the development of ischemic stroke subtypes in a general Japanese population. *Atherosclerosis*. 2010;210:249-255.
30. Okamura T, Nakamura K, Kanda H, et al. Effect of combined cardiovascular risk factors on individual and population medical expenditures: a 10-year cohort study of national health insurance in a Japanese population. *Circ J*. 2007;71:807-813.
31. Okamura T, Kokubo Y, Watanabe M, et al. A revised definition of the metabolic syndrome predicts coronary artery disease and ischemic stroke after adjusting for low density lipoprotein cholesterol in a 13-year cohort study of Japanese: the Suita study. *Atherosclerosis*. 2011;217:201-206.
32. Miyake S. Long-term hypertension control in a community-comparison of stroke incidence and hypertension control between participants and nonparticipants in health examinations. *Nihon Koshu Eisei Zasshi*. 1993;40:606-623. (in Japanese).
33. Hirai T, Kusaka Y, Iki M, et al. Relationship of hypertension prevalence in companies to business type and scale—from an analysis of health examination results in Fukui prefecture. *Nihon Koshu Eisei Zasshi*. 1996;43:806-814. (in Japanese).
34. Yamataki H, Suwazono Y, Okubo Y, et al. Health status of workers in small and medium-sized companies as compared to large companies in Japan. *J Occup Health*. 2006;48:166-174.
35. Hoshuyama T, Hino Y, Kayashima K, et al. Inequality in the health status of workers in small-scale enterprises. *Occup Med (Lond)*. 2007;57:126-130.
36. Davey Smith G, Carroll D, Rankin S. Socio-economic differentials in mortality: evidence from Glasgow graveyards. *BMJ*. 1992;305:1554-1557.
37. Kagamimori S, Gaina A, Nasermoaddeli A. Socioeconomic status and health in the Japanese population. *Soc Sci Med*. 2009;68:2152-2160.
38. Hirokawa K, Tsutsumi A, Kayaba K, the Jichi Medical Cohort study group. Mortality risks in relation to occupational category and position among the Japanese working population: the Jichi Medical School (JMS) cohort study. *BMJ Open*. 2013;3:e002690.
39. Sakurai H. Healthy Japan 21. *JMAJ*. 2003;46:47-49.

