

引用文献リスト (採用論文、不採用論文リスト)

CQ1

採用論文

Wong 2014	"Wong D, Abdel-Latif M, Kent ",Antenatal steroid exposure and outcomes of very premature infants: a regional cohort study. ,Arch Dis Child Fetal Neonatal Ed,2014,99(1),F12-20
Romejko 2013	"Romejko-Wolniewicz E, Oleszczuk L, Zar?ba-Szczudlik J, Czajkowski K",Dosage regimen of antenatal steroids prior to preterm delivery and effects on maternal and neonatal outcomes.,Journal of maternal-fetal & neonatal medicine,2013,26(3),237-41
大関 2012	"大関 一裕, 松井 紗智子, 島村 直紀, 荻野 寛子, 高下 敦子, 岡田 真衣子, 大橋 祥子, 藤中 義史, 増永 健, 瀧川 逸朗",超早産児における出生前ステロイド投与と新生児予後,日本周産期・新生児医学会雑誌,2012,48(4),840-844
Smith 2012	"Smith PB, Ambalavanan N, Li L, Cotten CM, Laughon M, Walsh MC, Das A, Bell EF, Carlo WA, Stoll BJ, Shankaran S, Laptook AR, Higgins RD, Goldberg ",Approach to infants born at 22 to 24 weeks' gestation: relationship to outcomes of more-mature infants. ,Pediatrics,2012,129(6),e1508-16
Gulcan 2006	"Gulcan H, Gungor S, Tiker F, Kilicdag ",Effect of perinatal factors on time of first stool passage in preterm newborns: An open, prospective study. ",Curr Ther Res Clin Exp,2006,67(3),214-25
Attridge 2006	"Attridge JT, Clark R, Gordon ",New insights into spontaneous intestinal perforation using a national data set (3): antenatal steroids have no adverse association with spontaneous intestinal perforation. ,J Perinatol,2006,26(11),667-70
甲斐 2005	"甲斐 克秀, 池田 智明, 山下 理絵, 道方 香織, 金子 政時, 鮫島 浩, 茨 聡, 池ノ上 克",極低出生体重児と外科疾患 在胎22~27週で出生した新生児の特発性消化管穿孔に関する周産期因子の検討,日本周産期・新生児医学会雑誌,2005,41(4),816-820
Song 2005	"Song GA, Han ",Effect of antenatal corticosteroid and antibiotics in pregnancies complicated by premature rupture of membranes between 24 and 28 weeks of gestation. ,J Korean Med Sci,2005,20(1),88-92
Qublan 2001	"Qublan HS, Malkawi HY, Hiasat MS, Hindawi IM, Al-Taani MI, Abu-Khait SA, Al-Maaitah ",The effect of antenatal corticosteroid therapy on pregnancies complicated by premature rupture of membranes. ,Clin Exp Obstet Gynecol,2001,28(3),183-6
Wells 1999	"Wells LR, Papile LA, Gardner MO, Hartenberger CR, Merker",Impact of antenatal corticosteroid therapy in very low birth weight infants on chronic lung disease and other morbidities of prematurity. ,J Perinatol,1999,19(8 Pt 1),578-81

不採用論文

Rolnitsky 2015	"Rolnitsky A, Lee SK, Piedbouf B, Harrison A, Shah ",Prophylactic Interventions in Neonatology: How Do They Fare in Real Life? ,Am J Perinatol,2015,,
澤田 2014	澤田 祐季, 浅野 智子, 吉原 紘行, 小島 龍司, 小川 紫野, 松本 洋介, 井口 純子, 岡田 英幹, 松原 寛和, 大嶋 勉 出生前母体ステロイドの投与間隔による新生児の合併症発症率についての検討 東海産科婦人科学会雑誌 2014 50 49-52
Murthy 2014	"Murthy K, Yanowitz TD, DiGeronimo R, Dykes FD, Zaniletti I, Sharma J, Sullivan KM, Mirpuri J, Evans JR, Wadhawan R, Piazza A, Adams-Chapman I, Asselin JM, Short BL, Padula MA, Durand DJ, Palotto EK, Reber ",Short-term outcomes for preterm infants with surgical necrotizing enterocolitis. ,J Perinatol,2014,34(10),736-40
Dodd 2013	"Dodd JM, Jones L, Flenady V, Cincotta R, Crowther ",Prenatal administration of progesterone for preventing preterm birth in women considered to be at risk of preterm birth. ,Cochrane Database Syst Rev,2013,7,CD004947

Crowther 2012	"Crowther CA, Aghajafari F, Askie LM, Asztalos EV, Brocklehurst P, Bubner TK, Doyle LW, Dutta S, Garite TJ, Guinn DA, Hallman M, Hannah ME, Hardy P, Maurel K, Mazumder P, McEvoy C, Middleton PF, Murphy KE, Peltoniemi OM, Peters D, Sullivan L, Thom EA, Voysey M, Wapner RJ, Yelland L, Zhang", Repeat prenatal corticosteroid prior to preterm birth: a systematic review and individual participant data meta-analysis for the PRECISE study group (prenatal repeat corticosteroid international IPD study group: assessing the effects using the best level of evidence) – study protocol. ,Syst Rev,2012,1,12
Torrance 2009	Torrance HL, Derks JB, Scherjon SA, Wijnberger LD, Visser ,Is antenatal steroid treatment effective in preterm IUGR fetuses? ,Acta Obstet Gynecol Scand,2009,88(10),1068–73
Smrcek 2005	"Smrcek JM, Schwartau N, Kohl M, Berg C, Geipel A, Krapp M, Diedrich K, Ludwig", Antenatal corticosteroid therapy in premature infants. ,Arch Gynecol Obstet,2005,271(1),26–32
Harding 2001	"Harding JE, Pang J, Knight DB, Liggins ", Do antenatal corticosteroids help in the setting of preterm rupture of membranes? ,Am J Obstet Gynecol,2001,184(2),131–9
Smith 2000	"Smith LM, Qureshi N, Chao ", Effects of single and multiple courses of antenatal glucocorticoids in preterm newborns less than 30 weeks' gestation. ,J Matern Fetal Med,2000,9(2),131–5
Elimian 1999	"Elimian A, Verma U, Canterino J, Shah J, Visintainer P, Tejani", Effectiveness of antenatal steroids in obstetric subgroups. ,Obstet Gynecol,1999,93(2),174–9
Tucker 1995	"Tucker L, Hoff C, Peevy K, Brost B, Holland S, Calhoun ", The effects of antenatal steroid use in premature rupture of membranes. ,Aust N Z J Obstet Gynaecol,1995,35(4),390–2
Halac 1990	"Halac E, Halac J, Begue EF, Casanas JM, Indiveri DR, Petit JF, Figueroa MJ, Olmas JM, Rodriguez LA, Obregon RJ, et a", Prenatal and postnatal corticosteroid therapy to prevent neonatal necrotizing enterocolitis: a controlled trial. ,J Pediatr,1990,117(1 Pt 1),132–8
Bauer 1984	"Bauer CR, Morrison JC, Poole WK, Korones SB, Boehm JJ, Rigatto H, Zachman RD", A decreased incidence of necrotizing enterocolitis after prenatal glucocorticoid therapy.,Pediatrics,1984,73(5),682–8

## CQ2

### 採用論文

Herrmann 2014	"Herrmann K, Carroll",An exclusively human milk diet reduces necrotizing enterocolitis. ,Breastfeed Med,2014,9(4),184-90
Cristofao 2013	"Cristofalo EA, Schanler RJ, Blanco CL, Sullivan S, Trawoeger R, Kiechl-Kohlendorfer U, Dudell G, Rechtman DJ, Lee ML, Lucas A, Abrams S",Randomized trial of exclusive human milk versus preterm formula diets in extremely premature infants.,The Journal of pediatrics,2013,163(6),1592-1595.e1
Ganapathy 2012	"Ganapathy V, Hay JW, Kim ",Costs of necrotizing enterocolitis and cost-effectiveness of exclusively human milk-based products in feeding extremely premature infants. ,Breastfeed Med,2012,7(1),29-37
Maayan 2012	"Maayan-Metzger A, Avivi S, Schushan-Eisen I, Kuint",Human milk versus formula feeding among preterm infants: short-term outcomes. ,Am J Perinatol,2012,29(2),121-6
Ghandehari 2012	"Ghandehari H, Lee ML, Rechtman ",An exclusive human milk-based diet in extremely premature infants reduces the probability of remaining on total parenteral nutrition: a reanalysis of the data. ,BMC Res Notes,2012,5,188
Henderson 2009	"Henderson G, Craig S, Brocklehurst P, McGuire",Enteral feeding regimens and necrotising enterocolitis in preterm infants: a multicentre case-control study. ,Arch Dis Child Fetal Neonatal Ed,2009,94(2),F120-3
Sisk 2007	Sisk PM, Lovelady CA, Dillard RG, Gruber KJ, O'Shea ,Early human milk feeding is associated with a lower risk of necrotizing enterocolitis in very low birth weight infants. ,J Perinatol,2007,27(7),428-33
Henderson 2007	Henderson Ginny, Anthony Mary Y, McGuire William,Formula milk versus maternal breast milk for feeding preterm or low birth weight infants,Cochrane Database of Systematic Reviews,2007,-4,
Schanler 2005	Schanler RJ, Lau C, Hurst NM, Smith ,Randomized trial of donor human milk versus preterm formula as substitutes for mothers' own milk in the feeding of extremely premature infants. ,Pediatrics,2005,116(2),400-6
McGuire 2003	McGuire W, Anthony ,Donor human milk versus formula for preventing necrotising enterocolitis in preterm infants: systematic review. ,Arch Dis Child Fetal Neonatal Ed,2003,88(1),F11-4
McGuire 2001	McGuire W, Anthony ,Formula milk versus term human milk for feeding preterm or low birth weight infants. ,Cochrane Database Syst Rev,2001,-4,CD002971
McGuire 2001	McGuire W, Anthony ,Formula milk versus preterm human milk for feeding preterm or low birth weight infants. ,Cochrane Database Syst Rev,2001,-3,CD002972
Lucas 1996	Lucas A, Fewtrell MS, Morley R, Lucas PJ, Baker BA, Lister G, Bishop ,Randomized outcome trial of human milk fortification and developmental outcome in preterm infants. ,Am J Clin Nutr,1996,64(2),142-51

### 不採用論文

五味 2010	五味 明,窪田 昭男,野口 啓幸,奥山 直樹,石井 智浩,川原 央好,小室 広昭,中村 哲郎,増本 幸二,八木 実,田中 芳明,古村 眞,板橋 家頭夫,日本小児外科学会学術・先進医療検討委員会. 超低出生体重児における消化管穿孔の実態調査. 日本小児外科学会雑誌. 2010 46(4) 791-796 小児外科 2009 41(5) 500-503
奥山 2000	奥山 宏臣,佐々木 隆士,清水 義之,窪田 昭男,川原 央好,長谷川 利路,米田 光宏,野瀬 恵介,三谷 泰之,野村 元成,白石 淳,北島 博之【出生前診断の倫理と実際】子宮内発育遅延症例における胎便関連性腸閉塞 2000 36(3) 500-505 臨床小児医学 1997 45(2) 73-77

堀内 1987	堀内 勁, 宝樹 真理 新生児壊死性腸炎の発生状況 1982~1983年におけるアンケート調査 日本新生児学会雑誌 1987 23(2) 431-438
松田 1982	松田 博雄, 萩野 恵里子, 志村 浩二 新生児壊死性腸炎および極小未熟児の腹部レントゲン所見の検討 日本新生児学会雑誌 1982 18(4) 714-723
白井	白井 勝, 真鍋 博美, 石井 朋子, 立花 幸晃, 佐藤 敬, 小久保 雅代, 坂田 宏, 丸山 静男, 宮本 和俊 超低出生体重児の腸穿孔6例のまとめ 日本新生児学会雑誌
曾根	曾根 滋巳, 新飯田 裕一, 本間 真二郎, 他 極小未熟児の消化管穿孔14例についての臨床的検討
Colaizy 2015	Colaizy , "Donor human milk for very low birth weights: patterns of usage, outcomes, and unanswered questions. ", Curr Opin Pediatr, 2015, 27(2), 172-6
Pammi 2015	Pammi M, Abrams , Oral lactoferrin for the prevention of sepsis and necrotizing enterocolitis in preterm infants. , Cochrane Database Syst Rev, 2015, -2, CD007137
Rochow 2015	Rochow N, Landau-Crangle E, Fusch, Challenges in breast milk fortification for preterm infants. , Curr Opin Clin Nutr Metab Care, 2015, 18(3), 276-84
Salas 2015	Salas AA, Cuna A, Bhat R, McGwin G Jr, Carlo WA, Ambalavanan, A randomised trial of re-feeding gastric residuals in preterm infants. , Arch Dis Child Fetal Neonatal Ed, 2015, 100(3), F224-8
Saengtawesin 2014	Saengtawesin V, Tangpolkaiwalsak R, Kanjanapattankul, Effect of oral probiotics supplementation in the prevention of necrotizing enterocolitis among very low birth weight preterm infants. , J Med Assoc Thai, 2014, 97 Suppl 6, S20-5
Thoene 2014	Thoene M, Hanson C, Lyden E, Dugick L, Ruybal L, Anderson-Berry, Comparison of the effect of two human milk fortifiers on clinical outcomes in premature infants. , Nutrients, 2014, 6(1), 261-75
Bigger 2014	Bigger HR, Fogg LJ, Patel A, Johnson T, Engstrom JL, Meier , Quality indicators for human milk use in very low-birthweight infants: are we measuring what we should be measuring? , J Perinatol, 2014, 34(4), 287-91
Quigley 2014	Quigley M, McGuire, Formula versus donor breast milk for feeding preterm or low birth weight infants. , Cochrane Database Syst Rev, 2014, 4, CD002971
Morgan 2013	Morgan J, Bombell S, McGuire, Early trophic feeding versus enteral fasting for very preterm or very low birth weight infants. , Cochrane Database Syst Rev, 2013, 3, CD000504
Arslanoglu 2013	Arslanoglu S, Corpeleijn W, Moro G, Braegger C, Campoy C, Colomb V, Decsi T, Domellof M, Fewtrell M, Hojsak I, Mihatsch W, Molgaard C, Shamir R, Turck D, van Goudoever, Donor human milk for preterm infants: current evidence and research directions. , J Pediatr Gastroenterol Nutr, 2013, 57(4), 535-42
Menon 2013	Menon G, Williams , "Human milk for preterm infants: why, what, when and how? ", Arch Dis Child Fetal Neonatal Ed, 2013, 98(6), F559-62
Fallon 2012	Fallon EM, Nehra D, Potemkin AK, Gura KM, Simpser E, Compher C, Puder, A.S.P.E.N. clinical guidelines: nutrition support of neonatal patients at risk for necrotizing enterocolitis. , JPEN J Parenter Enteral Nutr, 2012, 36(5), 506-23
Downard 2012	Downard CD, Renaud E, St Peter SD, Abdullah F, Islam S, Saito JM, Blakely ML, Huang EY, Arca MJ, Cassidy L, Aspelund, Treatment of necrotizing enterocolitis: an American Pediatric Surgical Association Outcomes and Clinical Trials Committee systematic review. , J Pediatr Surg, 2012, 47(11), 2111-22

Adamkin 2012	Adamkin ,“Mother’s milk, feeding strategies, and lactoferrin to prevent necrotizing enterocolitis. ”,JPEN J Parenter Enteral Nutr,2012,36(1 Suppl),25S–9S
Thoene 2011	Thompson A, Bizzarro M, Yu S, Diefenbach K, Simpson BJ, Moss ,Risk factors for necrotizing enterocolitis totalis: a case–control study. ,J Perinatol,2011,31(11),730–8
Meinzen 2009	Meinzen–Derr J, Poindexter B, Wrage L, Morrow AL, Stoll B, Donovan ,Role of human milk in extremely low birth weight infants’ risk of necrotizing enterocolitis or death. ,J Perinatol,2009,29(1),57–62
Quigley 2007	Quigley MA, Henderson G, Anthony MY, McGuire,Formula milk versus donor breast milk for feeding preterm or low birth weight infants. ,Cochrane Database Syst Rev,2007,–4,CD002971
Maayan 2004	Maayan–Metzger A, Itzhak A, Mazkereth R, Kuint,Necrotizing enterocolitis in full–term infants: case–control study and review of the literature. ,J Perinatol,2004,24(8),494–9
Furman 2003	Furman L, Taylor G, Minich N, Hack,The effect of maternal milk on neonatal morbidity of very low–birth–weight infants. ,Arch Pediatr Adolesc Med,2003,157(1),66–71
Schanler 1994	Schanler RJ, Atkinson ,Effects of nutrients in human milk on the recipient premature infant. ,J Mammary Gland Biol Neoplasia,1999,4(3),297–307
Caplan 1993	Caplan MS, MacKendrick,Necrotizing enterocolitis: a review of pathogenetic mechanisms and implications for prevention. ,Pediatr Pathol,1993,13(3),357–69

### CQ 3

#### 採用論文

Lau 2015	Lau CS, Chamberlain RS. Probiotic administration can prevent necrotizing enterocolitis in preterm infants: A meta-analysis. <i>J Pediatr Surg.</i> 2015 Aug;50(8):1405-12
Totsu 2014	Satsuki Totsu,1 Chika Yamasaki,2 Masaki Terahara,3 Atsushi Uchiyama1 and Satoshi Kusuda,1 on behalf of Probiotics Study Group in Japan. 1Maternal and Perinatal Center, Tokyo Women's Medical University, 2Department of Neonatology, Aiiiku Hospital, Tokyo and 3Food Science Institute, Meiji Dairies Corporation, Odawara, Japan. Bifidobacterium and enteral feeding in preterm infants: Cluster-randomized trial. <i>Pediatrics International</i> (2014) 56, 714-719
Serce 2013	Serce O, Benzer D, Gursoy T, Karatekin G, Ovali F. Efficacy of <i>Saccharomyces boulardii</i> on necrotizing enterocolitis or sepsis in very low birth weight infants: a randomised controlled trial. <i>Early Hum Dev.</i> 2013 Dec;89(12):1033-6.
Demirel 2013	Demirel G1, Erdeve O, Celik IH, Dilmen U. <i>Saccharomyces boulardii</i> for prevention of necrotizing enterocolitis in preterm infants: a randomized, controlled study. <i>Acta Paediatr.</i> 2013 Dec;102(12):e560-5.
AI-Hosni 2012	M Al-Hosni, M Duenas, M Hawk, LA Stewart, RA Borghese, M Cahoon, L Atwood, D Howard, K Ferrelli and R Soll, 8. Probiotics-supplemented feeding in extremely low-birth-weight infants. <i>Journal of Perinatology</i> (2012) 32, 253-259
Mihatsch 2012	Mihatsch WA1, Braegger CP, Decsi T, Kolacek S, Lanzinger H, Mayer B, Moreno LA, Pohlandt F, Puntis J, Shamir R, Stadtmüller U, Szajewska H, Turck D, van Goudoever JB. Critical systematic review of the level of evidence for routine use of probiotics for reduction of mortality and prevention of necrotizing enterocolitis and sepsis in preterm infants. <i>Clin Nutr.</i> 2012 Feb;31(1):6-15
Mary 2012	Mary N Mugambi, Alfred Musekiwa, Martani Lombard, Taryn Young and Reneé Blaauw. Probiotics, prebiotics infant formula use in preterm or low birth weight infants: a systematic review. Mugambi et al. <i>Nutrition Journal</i> 2012, 11:58
Braga 2011	Braga TD, da Silva GA, de Lira PI, de Carvalho Lima M. Efficacy of <i>Bifidobacterium breve</i> and <i>Lactobacillus casei</i> oral supplementation on necrotizing enterocolitis in very-low-birth-weight preterm infants: a double-blind, randomized, controlled trial. <i>Am J Clin Nutr</i> 2011;93:81-6.
Sari 2011	FN Sari, EA Dizdar, S Oguz, O Erdeve, N Uras and U Dilmen. Oral probiotics: <i>Lactobacillus sporogenes</i> for prevention of necrotizing enterocolitis in very low-birth weight infants: a randomized, controlled trial. <i>European Journal of Clinical Nutrition</i> (2011) 65, 434-439

Carole 2009	Carole Rouge´, Hugues Piloquet, Marie-José´ Butel, Bernard Berger, Florence Rochat, Laurent Ferraris, Clotilde Des Robert, Arnaud Legrand, Marie-France de la Cochetie`re, Jean-Michel N' Guyen, Michel Vodovar, Marcel Voyer, Dominique Darmaun, and Jean-Christophe Roze´. Oral supplementation with probiotics in very-low-birth-weight preterm infants: a randomized, double-blind, placebo-controlled trial. Am J Clin Nutr 2009;89:1828-35.
Bin-Nun 2005	Bin-Nun A, Bromiker R, Wilschanski M, Kaplan M, Rudensky B, Caplan M, Hammerman C. Oral Probiotics prevent necrotizing enterocolitis in very low birth weight neonates. J Pediatr. 2005 Aug;147(2):192-6.
Dani 2002	Dani C, Biadaioli R, Bertini G, Martelli E, Rubaltelli FF. Probiotics feeding in prevention of urinary tract infection, bacterial sepsis and necrotizing enterocolitis in preterm infants. A prospective double-blind study. Biol Neonate. 2002 Aug;82(2):103-8.

不採用論文

## CQ4

## 採用論文

Livingston 2015	Livingston MH, Shawyer AC, Rosenbaum PL, Williams C, Jones SA, Walton. Glycerin enemas and suppositories in premature infants: a meta-analysis. Pediatrics 2015;135(6):1093-1106
Shinde 2014	Shinde S, Kabra NS, Sharma SR, Avasthi BS, Ahmed. Glycerin suppository for promoting feeding tolerance in preterm very low birthweight neonates: a randomized controlled trial. Indian Pediatr2014;51(5):367-370
Khadr 2011	Khadr SN, Ibhanebhor SE, Rennix C, Fisher HE, Manjunatha CM, Young D, Abara. Randomized controlled trial: impact of glycerin suppositories on time to full feeds in preterm infants. Neonatology2011;100(2):169-176
Haiden 2007	Haiden N, Jilma B, Gerhold B, Klebermass K, Prusa AR, Kuhle S, Rohrmeister K, Kohlhauser-Vollmuth C, Pollak J. Small volume enemas do not accelerate meconium evacuation in very low birth weight infants. Pediatr Gastroenterol Nutr2007;44(2):270-273

## 不採用論文

孫田 2008	孫田 みゆき, 岸 泰子, 横塚 太郎, 安田 彩子, 鬼頭 修, 鈴木 千鶴子, 津田 峰行. 胎便関連性腸閉塞の診断と治療 外科的介入を要した胎便関連性腸閉塞についての臨床学的検討. 日本周産期・新生児医学会雑誌 2008;44(4):939-942
---------	--



## CQ5

## 採用論文

Haiden 2012	Haiden N, Norooz F, Klebermass-Schrehof K, Horak AS, Jilma B, Berger A, Repa. The effect of an osmotic contrast agent on complete meconium evacuation in preterm infants. <i>Pediatrics</i> 2012; 130(6); 1600-6
Garza 2004	Garza-Cox S, Keeney SE, Angel CA, Thompson LL, Swischuk. Meconium obstruction in the very low birth weight premature infant. <i>Pediatrics</i> 2004; 114(1); 285-90
遠藤 2002	遠藤 大一, 山本 和歌子, 中島 やよひ, 与田 仁志, 川上 義, 吉野 浩之, 上井 義之, 横森 欣司. IUGRの超低出生体重児における胎便関連性腸閉塞症の増悪因子についての検討. <i>日本新生児学会雑誌</i> 2002; 38(3); 502-505

## 不採用論文

Cho 2015	Cho HH, Cheon JE, Choi YH, Lee SM, Kim WS, Kim IO, Shin SM, Kim EK, Kim HS, Choi JH, You. Ultrasound-guided contrast enema for meconium obstruction in very low birth weight infants: Factors that affect treatment success. <i>Eur J Radiol</i> 2015.
Koshinaga 2011	Koshinaga T, Inoue M, Ohashi K, Sugito K, Ikeda T, Tomita. Therapeutic strategies of meconium obstruction of the small bowel in very-low-birthweight neonates. <i>Pediatr Int</i> 2011; 53(3); 338-44.
Nakaoka 2009	Nakaoka T, Shiokawa C, Nishihara M, Tamai H, Funato M, Uemura. Iopamidol enema treatment for meconium obstruction of prematurity in extremely low-birth weight infants: a safe and effective method. <i>Pediatr Surg Int</i> 2009; 25(3); 273-6.
Shinohara 2007	Shinohara T, Tsuda M, Koyama. Management of meconium-related ileus in very low-birthweight infants. <i>Pediatr Int</i> 2007; 49(5); 641-4.
Shim 2007	Shim SY, Kim HS, Kim DH, Kim EK, Son DW, Kim BI, Choi. Induction of early meconium evacuation promotes feeding tolerance in very low birth weight infants. <i>Neonatology</i> 2007; 92(1); 67-72.
Krasna 1996	Krasna IH, Rosenfeld D, Salerno. Is it necrotizing enterocolitis, microcolon of prematurity, or delayed meconium plug? A dilemma in the tiny premature infant. <i>J Pediatr Surg</i> 1996; 31(6); 855-8.

今治 2014	今治 玲助, 加藤 怜子, 橋本 晋太郎, 向井 亘, 佐伯 勇, 秋山 卓士. 出生体重児における胎便関連性腸閉塞症の検討. 外科学会雑誌 2014; 50(6); 1005-1010.	超低 日本小児
春本 2011	春本 研, 塩川 智司, 権 英寿, 山道 拓, 辻本 嘉助. 出生体重児の重症胎便関連性腸閉塞症に対する外科的治療戦略. 児外科学会雑誌 2011; 47(1); 20-25	超低 日本小
河野 2008	河野 達夫. 児新生児学会雑誌第19巻第2号85(251)頁~88(254)頁2007年 寺田明佳他論文「極低出生体重児の胎便関連性腸閉塞症に対するガストログラフィン胃内投与の効果」について ガストログラフィン胃内投与の危険性. 日本未熟児新生児学会雑誌 2008; 20(1); 156.	日本未熟 児新生児
寺田2007	寺田 明佳, 市場 博幸, 郡山 健, 田中 裕子, 森 啓之, 大西 聡, 江原 英治. 極低出生体重児の胎便関連性腸閉塞症に対するガストログラフィン胃内投与の効果. 日本未熟児新生児学会雑誌 2007; 19(2); 251-254.	極 日本未熟児新生児学会雑誌
鈴木2007	鈴木 完, 仁科 孝子, 村越 孝次, 新井 真理, 佐藤 かおり, 近藤 昌敏, 柿沼 亮太, 岡崎 薫. 極低出生体重児に発症した胎便排泄遅延に対する造影剤治療の検討(早期に開始することの有効性について) 日本周産期・新生児医学会雑誌 2007; 43(1); 56-61.	
藤谷 2007	藤谷 しのぶ, 三浦 文宏, 水谷 佳世, 井上 真理, 水野 克巳, 竹内 敏雄, 板橋 家頭夫. 当院で経験した胎便関連性腸閉塞4症例における検討. 日本未熟児新生児学会雑誌 2007; 19(1); 70-75.	日
向井 2006	向井 基, 茨 聡, 丸山 有子, 徳久 琢也, 藤江 由夏, 松井 貴子, 佐々木 恒, 中澤 祐介, 野口 啓幸, 坂本 浩一, 高松 英夫.【極・超低出生体重児の外科】極・超低出生体重児における胎便性腸閉塞の治療と予後. 小児外科 2006; 38(1); 86-90.	
住田 2006	住田 裕, 藤村 正哲. に対するガストログラフィン注腸 その至適量に関する検討. 未熟児新生児学会雑誌 1998; 10(2); 191-195.	胎便病 日本
雨海 1992	雨海 照祥, 監物 久夫, 城 一也. 小児外科疾患とNon-Open Surgery ガストログラフィンの挟み撃ちによるmeconium diseaseの治療. 小児外科 1992; 24(5); 573-580.	小

## CQ6

### 採用論文

Rees 2008	Peritoneal drainage or laparotomy for neonatal bowel perforation?
Bonnard A 2008	The use of the score for neonatal acute physiology–perinatal extension in perforated necrotizing enterocolitis could it guide therapy in newborns less than 1500g?
Moss 2006	Laparotomy versus peritoneal drainage for necrotizing enterocolitis and perforation
Blakely M 2006	Laparotomy versus peritoneal drainage for necrotizing enterocolitis or isolated intestinal perforation in extremely low birth weight infants: outcomes through 18 months adjusted age
Sharma R 2004	Surgical management of bowel perforations and outcome in very low birth weight infants
Azarow KS	Laparotomy or drain for perforated necrotizing enterocolitis who gets what and why?

### 不採用論文

Downard 2012	Treatment of necrotizing enterocolitis : an american pediatric surgical association outcomes and clinical trials committee systematic review
Rao 2011	Peritoneal drainage versus laparotomy as initial surgical treatment for perforated necrotizing enterocolitis or spontaneous intestinal perforation in preterm low birth weight infants
Rees 2010	Peritoneal drainage does not stabilize extremely low birth weight infants with perforated bowel: data from the NET trial
Sola 2010	Peritoneal drainage versus laparotomy for necrotizing enterocolitis and intestinal perforation: a meta-analysis.

## Ⅱ資料

1. 疾患概要
2. 研究業績
  - (ア) 学会発表
  - (イ) 論文発表
3. 発表論文 PDF
4. ガイドライン外部評価結果
5. 研究者名簿

## 1. 疾患概要

### 消化器系疾患分野

#### 低出生体重児の消化管機能障害

1. 概要 近年の低出生体重児の出生数は増加傾向にある。周産期医療の進歩により低出生体重児の救命率は改善傾向にあるが、その一方で種々の臓器の未熟性に起因する合併症が周産期医療における大きな課題となってきた。なかでも壊死性腸炎、特発性腸穿孔、胎便関連性腸閉塞、胎便性腹膜炎といった消化管機能障害は、低出生体重児によくみられる重篤な消化管合併症であり、生命予後だけでなく長期予後を左右する重要な因子となっている。

2. 疫学 日本小児外科学会のアンケート(全国 NICU263 施設を対象、回答率 47%)では、2003-2007 年の5 年間で超低出生体重児 8282 例中消化管穿孔発生症は 444 例/5 年間(発生率 5.36%)という報告がある。消化管機能障害の多くが超低出生体重児に発生することやアンケートの回答率などを考慮すれば、本邦で年間 200 例前後の発生があると考えられる。

3. 原因 壊死性腸炎、特発性腸穿孔、胎便関連性腸閉塞、胎便性腹膜炎のほとんどは極低出生体重児に発生することから、腸管の未熟性を背景として、感染やストレスといった種々の周産期因子が関与して発症すると考えられている。しかし、個々の疾患の危険因子は明らかではなく、病態や病因も不明である。

4. 症状 ほとんどの場合、生後数日から生後 1-2 週間の新生児期に発症する。腸炎症状で発症する場合、腸閉塞症状で発症する場合、突然の腸穿孔で発症する場合など様々である。一旦腸穿孔を起こせば腹膜炎を併発して敗血症性ショックに陥り、全身状態は急速に悪化する。

5. 合併症 周産期管理の進歩とともに、本疾患の迅速な診断・治療により救命率は上昇してきた。しかし長期フォローに基づく最近の報告では、救命例の半数以上に精神運動発達遅延がみられることが明らかになってきた。そのため個々の疾患の周産期背景因子の解析から、その発症機序を明らかにして予防法を確立することが、低出生体重児全体の予後改善に不可欠であると考えられる。

6. 治療法 腸炎症状や腸閉塞症状が先行する場合は、絶食、抗生剤投与といった保存的治療が試みられる。保存的治療が有効でなく全身状態が悪化する場合や腸穿孔を併発した場合は手術適応となる。腸瘻造設術が一般的だが、全身状態が良ければ穿孔部の縫合閉鎖や腸吻合も行われる。

7. 研究班 低出生体重児消化管機能障害の疾患概念確立にむけた疫学調査研究  
( H26-難治等(難)一般-010 )

## 2. 研究業績

### (ア) 学会発表

Risk factors for surgical intestinal complications in very low birth weight infants: A multicenter case-control study

Hiroomi Okuyama

9<sup>th</sup> Annual Conference of Cambodian Society of Pediatric Surgery November 19, 2015 Phnom Penh, Cambodia

低出生体重児消化管機能障害における腸瘻閉鎖時期と術後経過

望月響子、奥山宏臣、早川昌弘、漆原直人、横井暁子、白石淳、藤永英志、大橋研介、永田公二、天江新太郎、古川泰三、矢内俊裕

1) 神奈川県立こども医療センター 外科

2) 低出生体重児消化管機能障害の疾患概念確立に向けた疫学調査研究班

第51回 日本周産期新生児医学会、福岡 2015. 7. 12-12 一般演題

極低出生体重児における MRI についての検討 NEC、FIP との差異

江角 元史郎, 永田 公二, 落合 正行, 田口 智章, 早川 昌弘, 漆原 直人, 望月 響子, 横井 暁子, 白石 淳, 藤永 英志, 大橋 研介, 大藤 さとこ, 天江 新太郎, 古川 泰三, 矢内 俊裕, 田附 裕子, 奥山 宏臣

低出生体重児の消化管機能障害に関する周産期背景因子の疫学調査研究班

第51回 日本周産期新生児医学会、福岡 2015. 7. 12-12 一般演題

SGA 児の長期予後改善にむけた周産期管理

SGA 児の消化管機能障害—胎便関連性腸閉塞—

奥山宏臣\*

低出生体重児消化管機能障害の疾患概念確立にむけた疫学調査研究班

\* 大阪大学大学院医学系研究科外科学講座小児成育外科学

第51回 日本周産期新生児医学会、福岡 2015. 7. 12-12 シンポジウム

消化管機能障害を有する低出生体重児の短腸症候群に対する戦略

横井暁子

低出生体重児消化管機能障害の疾患概念確立にむけた疫学調査研究班

兵庫県立こども病院 外科

第51回 日本周産期新生児医学会、福岡 2015. 7. 12-12 シンポジウム

小児外科におけるナショナルスタディと診断治療ガイドラインの構築

極低出生体重児の消化管機能障害に関する周産期背景因子ならびに予後の検討

奥山 宏臣, 大藤 さとこ, 早川 昌弘, 漆原 直人, 横井 暁子, 白石 淳, 望月 響子, 藤永 英志, 大橋 研介, 永田 公二, 天江 新太郎, 古川 泰三, 矢内 俊裕, 田附 裕子, 皆川 京子

低出生体重児消化管機能障害の疾患概念確立に向けた疫学調査研究班

第52回日本小児外科学会学術集会 神戸市 2016.5.28-30 シンポジウム

田附裕子、奥山宏臣、上原秀一郎、上野豪久、山中宏晃、白石 淳、漆原直人、武 浩志、藤永英志、横井暁子、大藤さとこ、大橋研介、早川昌弘、田口智章

手術適応となったMRIの予後改善をめざした小児外科的介入:多施設共同後方視的調査結果より

第51回 日本周産期新生児医学会、福岡 2015. 7. 12-12 一般演題

(イ) 論文発表

Hayakawa M, Taguchi T, Urushihara N, Yokoi A, Take H, Shiraishi J, Fujinaga H, Ohashi K, Oshiro M, Kato Y, Ohfuji S, Okuyama H.

Outcome in VLBW infants with surgical intestinal disorder at 18 months of corrected age. *Pediatr Int.* 2015,57(4):633-8.

Okuyama H, Ohfuji S, Hayakawa M, Urushihara N, Yokoi A, Take H, Shiraishi J, Fujinaga H, Ohashi K, Minagawa K, Misaki M, Nose S, Taguchi T.

Risk factors for surgical intestinal disorders in VLBW infants: Case-control study. *Pediatr Int.* 2016 58(1):34-9.

Saka R, Wakimoto T, Nishiumi F, Sasaki T, Nose S, Fukuzawa M, Oue T, Yanagihara I, Okuyama H.

Surfactant protein-D attenuates the lipopolysaccharide-induced inflammation in human intestinal cells overexpressing toll-like receptor 4. *Pediatr Surg Int.* 2016 Jan;32(1):59-63. doi: 10.1007/s00383-015-3812-y. Epub 2015 Oct 28.

Michinobu Ohno, Yutaka Kanamori, Kaori Sato, Misato Migita, Toshiko Takezoe, Masataka Takahashi, Toshihiko Watanabe, Yasushi Fuchimoto, Shoitiro Amari, Takuya Oshima, Yushi Ito, Rika Sugibayashi, Katsunori Ozawa, Seiji Wada,

Haruhiko Sago, Congenital diaphragmatic hernia in dizygotic twins. *J Ped Surg Case Reports* 2015; 3: 114-116.

第51回日本周産期新生児医学会学術集会シンポジウム2次抄録

SGA 児の長期予後改善にむけた周産期管理

SGA 児の消化管機能障害—胎便関連性腸閉塞—

奥山宏臣\*

低出生体重児消化管機能障害の疾患概念確立にむけた疫学調査研究班

\*大阪大学大学院医学系研究科外科学講座小児成育外科学

(日本周産期新生児学会雑誌 in press)

「改訂第2版 症例から学ぶ周産期診療ワークブック」


Ⅲ新生児編 10. 消化器 1) 新生児壊死性腸炎

和田和子\*、奥山宏臣\*\*

大阪大学大学院医学系研究科小児科\*、小児成育外科学\*\*

(メジカルビュー社 in press)





## Original Article

**Risk factors for surgical intestinal disorders in VLBW infants:  
Case-control study**

Hiroomi Okuyama,<sup>1</sup> Satoko Ohfuji,<sup>2</sup> Masahiro Hayakawa,<sup>3</sup> Naoto Urushihara,<sup>4</sup> Akiko Yokoi,<sup>5</sup> Hiroshi Take,<sup>6</sup> Jun Shiraiishi,<sup>7</sup> Hideshi Fujinaga,<sup>8</sup> Kensuke Ohashi,<sup>9</sup> Kyoko Minagawa,<sup>10</sup> Maiko Misaki,<sup>10</sup> Satoko Nose<sup>10</sup> and Tomoaki Taguchi<sup>11</sup>

<sup>1</sup>Department of Pediatric Surgery, Osaka University Graduate School of Medicine, Suita, <sup>2</sup>Department of Public Health, Faculty of Medicine, Osaka City University, Osaka, <sup>3</sup>Center for Maternal-Neonatal Care, Nagoya University Hospital, Nagoya, <sup>4</sup>Department of Pediatric Surgery, Shizuoka Children's Hospital, Shizuoka, <sup>5</sup>Department of Pediatric Surgery, Hyogo Children's Hospital, Kobe, <sup>6</sup>Department of Pediatric Surgery, Kanagawa Children's Hospital, Yokohama, <sup>7</sup>Department of Neonatology, Osaka Medical Center and Research Institute for Maternal and Child Health, Izumi, <sup>8</sup>Division of Neonatology, National Center for Child Health and Development, <sup>9</sup>Department of Pediatric Surgery, Nihon University School of Medicine, Tokyo, <sup>10</sup>Department of Pediatrics, Hyogo College of Medicine, Nishinomiya and <sup>11</sup>Department of Pediatric Surgery, Kyushu University, Fukuoka, Japan

**Abstract** **Background:** Very low-birthweight (VLBW) infants (VLBWI) are at increased risk for surgical intestinal disorders including necrotizing enterocolitis (NEC), focal intestinal perforation (FIP) and meconium-related ileus (MRI). The aim of this study was to identify disease-specific risk factors for surgical intestinal disorders in VLBWI.

**Methods:** A retrospective multicenter case-control study was conducted at 11 institutes. We reviewed VLBWI who underwent laparotomy for intestinal disorders including perforation and intractable bowel obstruction. The surgical disorders were classified into four categories (NEC, FIP, MRI, others) based on the macroscopic findings at operation. In order to identify risk factors, two matched controls for each subject were chosen based on gestational age and birthweight. OR and 95%CI were calculated using a conditional logistic regression model and a multivariate model.

**Results:** A total of 150 cases (NEC, n = 44; FIP, n = 47; MRI, n = 42; others, n = 17) and 293 controls were identified. The cases and controls were similar in terms of gestational age and birthweight (cases/controls, 26.7 ± 2.5/26.5 ± 2.6 weeks; 790 ± 256/795 ± 257 g). On multivariate modeling, disease-specific risk factors were as follows: female (OR, 0.23; 95% CI: 0.06–0.89), respiratory distress syndrome (OR, 35.7; 95%CI: 2.48–514) and patent ductus arteriosus (OR, 10.9; 95% CI: 1.51–79.3) for NEC; outborn delivery (OR, 5.47; 95%CI: 1.48–20.2) for FIP; and twin pregnancy (OR, 4.25; 95%CI: 1.06–17.1), PROM (OR, 6.85; 95%CI: 1.33–35.4) and maternal steroid (OR, 0.23; 95%CI: 0.07–0.79) for MRI.

**Conclusions:** Different risk factors were identified for NEC, FIP and MRI, suggesting that each disease has a different etiology, and that different strategies are required to prevent these diseases.

**Key words** case-control study, focal intestinal perforation, meconium-related ileus, necrotizing enterocolitis, very low-birthweight.

Despite recent advancements in neonatal care, surgical intestinal disorders remain the most devastating gastrointestinal diseases in very low-birthweight (VLBW) infants. In addition to being associated with high mortality rate, survivors are left with significant long-term morbidity.<sup>1,2</sup> Therefore, prevention of the intestinal complications is required to improve the overall outcome of VLBW infants. There are several diseases that cause surgical intestinal disorders without mechanical obstruction in VLBW infants, including necrotizing enterocolitis (NEC), focal intestinal perforation (FIP) and meconium-related ileus (MRI).

Necrotizing enterocolitis is one of the most common and serious intestinal disorders involving systemic inflammatory responses in VLBW infants. Although previous numerous studies showed multiple risk factors for NEC including immaturity of the intestine, formula feeding and microbial colonization, the case fatality rate with surgical intervention remains as high as 50%, and is highest for the smallest infants.<sup>3,4</sup> Therefore, risk factors for surgical NEC in VLBW infants should be analyzed separately from medical and term NEC.

Focal intestinal perforation, characterized by the presence of focal intestinal perforation with no or minimal adjacent bowel inflammation, has become an increasingly common intestinal disorder in VLBW infants. While the pathogenesis of FIP is poorly understood, recent research has confirmed FIP to be a separate disease entity from NEC based on pathological findings<sup>5</sup> and genome expression profiles.<sup>6</sup>

To date, several types of meconium obstruction in the absence of cystic fibrosis in VLBW infants have been described. In 1999, Kubota *et al.* proposed the term “meconium-related ileus” as a single pathophysiologic entity with a wide spectrum of symptoms

Correspondence: Hiroomi Okuyama, MD PhD, Department of Pediatric Surgery, Osaka University Graduate School of Medicine, 2-2, Yamadaoka Suita, Osaka 565-0871, Japan. Email: okuyama@pedsurg.med.osaka-u.ac.jp

Received 4 January 2015; revised 20 February 2015; accepted 10 June 2015.

caused by meconium obstruction in the absence of cystic fibrosis.<sup>7</sup> Although MRI responds to conservative management in most cases, delayed identification leads to increased risk of intestinal perforation. Thus, it is important to identify risk factors for the severe type of MRI requiring surgery.

Prematurity and low birthweight are the consistently recorded risk factors for the development of these diseases, but risk factors for surgical intestinal disorders in VLBW infants have not been well described. Because macroscopic findings of affected intestines are important indicators in the definitive diagnosis of NEC, FIP and MRI, the diseases should be categorized based on the operative findings in order to analyze the etiology of each disease. Therefore, the aim of this study was to identify disease-specific risk factors associated with surgical intestinal disorders in VLBW infants based on definitive diagnosis at surgery.

## Methods

A retrospective multicenter case-control study was conducted at 11 institutes in Japan. We reviewed VLBW infants who underwent laparotomy for intestinal disorders including perforation and intractable ileus between January 2003 and December 2012. In order to detect risk factors other than prematurity and low birthweight, two matched controls for every subject were chosen based on gestational age ( $\pm 1$  week) and birthweight ( $\pm 50$  g). The prenatal and preoperative data for each infant and their mothers were collected from the medical records. The data included disease category, gestational age, birthweight, maternal factors (i.e. outborn delivery, mode of delivery, multiple pregnancies, placental abruption, diabetes, pregnancy hypertension, maternal smoking, chorioamnionitis, premature rupture of membranes [PROM], maternal steroid treatment) and patient factors (i.e. sex, Apgar score, small for gestational age/appropriate for gestational age, umbilical artery catheter use, respiratory distress syndrome [RDS], surfactant treatment, steroid treatment, symptomatic patent ductus arteriosus [PDA], transfusions, breast/formula feeding, and probiotics treatment).

### Definition of surgical intestinal disorders

The surgical intestinal disorders observed in the VLBW infants were classified into the following four categories based on the operative findings.

Necrotizing enterocolitis is a condition of diffuse or segmental necrotic injury to the mucosal and submucosal layers of the bowel with systemic inflammatory responses. FIP is defined as an isolated

perforation of the bowel with no obvious inflammation in the adjacent site or the rest of the bowel.<sup>6</sup> MRI is characterized by functional ileus due to impaired meconium excretion not associated with cystic fibrosis and is usually diagnosed when laparotomy for intractable ileus indicates a microcolon or small-sized colon extending to the distal ileum with a dilated proximal ileum filled with sticky meconium.<sup>7</sup> The classification “others” was assigned to conditions not belonging to any of the aforescribed three categories.

### Statistical analysis

Comparison of subject characteristics between the cases and controls was done using chi-squared tests for categorical variables and Wilcoxon rank-sum test for continuous variables. Summarized data for continuous variables are presented as mean  $\pm$  SD. A conditional logistic regression model was used to calculate OR and 95% CI for surgical intestinal disorders. The trend of association was assessed by assigning ordinal scores to each level within the category. The following explanatory variables were used to construct the multivariate models: (i) crude OR on univariate analysis; (ii) an initial multivariate model involving variables with statistically significant crude OR for both surgical intestinal disorders and at least one disease; and (iii) a final multivariate model excluding variables with no association with the diseases in the initial model.

In order to examine disease-specific risk factors, a stratified analysis was conducted according to each disease (e.g. NEC, FIP and MRI).

All tests were two-sided, and  $P < 0.05$  was regarded as statistically significant. All analyses were performed using the SAS version 9.1.3 (SAS Institute, Cary, NC, USA).

### Ethics

This study was performed after being approved by the ethics committee of Hyogo College of Medicine (approval number: 1483) and the independent ethics committees of 10 other participating institutions: National Center for Child Health and Development, Kyushu University, Nagoya University Hospital, Japanese Red Cross Nagoya Daiichi Hospital, Anjyo-kousei Hospital, Osaka Medical Center and Research Institute for Maternal and Child Health, Hyogo Children's Hospital, Shizuoka Children's Hospital, Kanagawa Children's Hospital and Nihon University School of Medicine.

## Results

A basic description of the study sample is provided in Table 1. A total of 150 cases (NEC,  $n = 44$ ; FIP,  $n = 47$ ; MRI,  $n = 42$ ; others,

**Table 1** Disease categories in VLBW infants

Category	Cases			Controls		
	n	Gestational age (weeks) Mean $\pm$ SD	Birthweight (g) Mean $\pm$ SD	n	Gestational age (weeks) Mean $\pm$ SD	Birthweight (g) Mean $\pm$ SD
NEC	44	25.6 $\pm$ 1.9	750 $\pm$ 207	86	25.4 $\pm$ 1.9	745 $\pm$ 206
FIP	47	26.7 $\pm$ 2.3	811 $\pm$ 246	92	26.5 $\pm$ 2.3	818 $\pm$ 252
MRI	42	27.3 $\pm$ 2.9	756 $\pm$ 281	81	27.5 $\pm$ 3.0	771 $\pm$ 277
Others	17	27.1 $\pm$ 2.7	903 $\pm$ 297	34	27.8 $\pm$ 2.5	946 $\pm$ 275
Total	150	26.7 $\pm$ 2.5	790 $\pm$ 256	293	26.5 $\pm$ 2.6	795 $\pm$ 257

FIP, focal intestinal perforation; MRI, meconium-related ileus; NEC, necrotizing enterocolitis; VLBW very low-birthweight.

n = 17) and 293 controls were identified. "Others" consisted of nine cases of meconium peritonitis, three cases of torsion, one case of invagination, one case of adhesion ileus, one case of internal hernia and two cases of unclassified entities. The cases were similar to the controls in terms of both gestational age and birthweight (cases/controls gestational age,  $26.7 \pm 2.5/26.5 \pm 2.6$  weeks; birthweight,  $790 \pm 256/795 \pm 257$  g).

#### Disease-specific risk factors

The investigated maternal and patient risk factors for surgical NEC are listed in Table 2. The conditional logistic regression models indicated that twin pregnancy (OR, 5.21; 95%CI: 1.68–16.2), RDS (OR, 16.3; 95%CI: 2.09–128), surfactant treatment (OR, 11.6; 95%CI: 1.46–92.0) and PDA (OR, 2.33; 95%CI: 1.00–5.41) were significantly associated with the development of surgical NEC. In addition, female sex (OR, 0.40; 95%CI: 0.18–0.89) was a protective factor. On multivariate modeling, female sex, RDS and PDA remained significant risk factors associated with surgical NEC.

The investigated maternal and patient risk factors for FIP are given in Table 3. Condition logistic regression modeling indicated that outborn delivery (OR, 5.47; 95%CI: 1.48–20.2) was the only significant risk factor associated with FIP.

The investigated maternal and patient risk factors for surgical MRI are listed in Table 4. On conditional logistic regression modeling, PROM (OR, 3.10; 95%CI: 1.05–9.13) and surfactant treatment (OR, 4.83; 95%CI: 1.01–23.1) were significant risk factors associated with surgical MRI. In contrast, maternal steroid treatment (OR, 0.38; 95%CI: 0.15–0.96) was identified to be protective. Multivariate modeling showed that twin pregnancy, PROM, and maternal steroid treatment remained significant factors associated with MRI.

The disease-specific risk factors identified on multivariate modeling are listed in Table 5. The significant risk factors for each disease were as follows: male sex, RDS and PDA for NEC; outborn delivery for FIP; and twin pregnancy, PROM, and lack of maternal steroid treatment for MRI, showing that the risk factors of each disease were different.

#### Discussion

Although the macroscopic findings of NEC, FIP and MRI are clearly different from each other, it is often difficult to make a definitive diagnosis without laparotomy. For example, NEC without pneumatosis or portal vein gas may be diagnosed as a form of MRI with mild inflammation. Inaccurate diagnosis could nullify

**Table 2** Maternal and patient risk factors for surgical NEC

Conditional logistic regression models		Cases	Controls	<i>P</i>	OR (95%CI)	<i>P</i>
		n = 44 (%)	n = 86 (%)			
Twin pregnancy	Single	30 (68)	77 (90)		1.00	
	Twin	14 (32)	8 (9)		<b>5.21</b> (1.68–16.2)	0.004
Sex	Male	32 (73)	44 (51)	0.018	1.00	
	Female	12 (27)	42 (49)		<b>0.40</b> (0.18–0.89)	0.026
RDS	No	2 (5)	22 (26)	0.003	1.00	
	Yes	42 (95)	63 (74)		<b>16.3</b> (2.09–128)	0.008
Surfactant	No	1 (2)	16 (19)	0.008	1.00	
	Yes	43 (98)	69 (81)		<b>11.6</b> (1.46–92.0)	0.021
PDA	No	19 (43)	51 (59)	0.081	1.00	
	Yes	25 (57)	35 (41)		<b>2.33</b> (1.00–5.41)	0.050
Multivariate models		OR	95%CI			<i>P</i>
Twin pregnancy		2.33	0.41–13.1			0.340
Maternal steroid		0.40	0.09–1.71			0.215
Sex (female)		<b>0.23</b>	0.06–0.89			0.033
RDS		<b>35.7</b>	2.48–514			0.009
PDA		<b>10.9</b>	1.51–79.3			0.018

**Bold**, *P* < 0.05. NEC, necrotizing enterocolitis; PDA, patent ductus arteriosus; RDS, respiratory distress syndrome.

**Table 3** Maternal and patient risk factors for FIP

Conditional logistic regression models		Cases	Controls	<i>P</i>	OR (95%CI)	<i>P</i>
		n = 47 (%)	n = 92 (%)			
Outborn delivery	No	33 (70)	79 (86)	0.027	1.00	
	Yes	14 (30)	13 (14)		<b>5.47</b> (1.48–20.2)	0.011

**Bold**, *P* < 0.05. FIP, focal intestinal perforation.

**Table 4** Maternal and patient risk factors for MRI

Conditional logistic regression models		Cases	Controls	<i>P</i>	OR (95%CI)	<i>P</i>
		n = 42(%)	n = 81(%)			
PROM	No	24 (60)	59 (74)	0.124	1.00	
	Yes	16 (40)	21 (26)		<b>3.10</b> (1.05–9.13)	0.040
Maternal steroid	No	25 (66)	38 (48)	0.063	1.00	
	Yes	13 (27)	42 (49)		<b>0.38</b> (0.15–0.96)	0.042
Surfactant	No	6 (14)	21 (26)	0.139	1.00	
	Yes	36 (86)	60 (74)		<b>4.83</b> (1.01–23.1)	0.049
Multivariate models		OR	95%CI			<i>P</i>
Twin pregnancy		<b>4.25</b>	1.06–17.1			0.042
PROM		<b>6.85</b>	1.33–35.4			0.022
Maternal steroid		<b>0.23</b>	0.07–0.79			0.019
Surfactant		17.7	0.94–333			0.055

**Bold**, *P* < 0.05. MRI, meconium-related ileus; PROM, premature rupture of membrane.

**Table 5** Multivariate risk factors for NEC, MRI and FIP

	NEC	FIP	MRI
Outborn delivery		↑*	
Maternal steroid			↓*
PROM			↑*
Twin pregnancy			↑*
Sex (female)	↓*		
RDS	↑*		
PDA	↑*		

\**P* < 0.05. FIP, focal intestinal perforation; MRI, meconium-related ileus; NEC, necrotizing enterocolitis; PDA, patent ductus arteriosus; PROM, premature rupture of membranes; RDS, respiratory distress syndrome.

precise analysis of the etiology of each disease. In the present study, in order to avoid making an unclear or suspected diagnosis, we categorized the diseases based on the operative findings. Therefore, infants who did not undergo laparotomy were excluded. Consequently, the present incidence of surgical MRI was similar to that of surgical NEC and FIP. This suggests that MRI should be recognized as a major entity causing surgical intestinal disorder in VLBW infants, as well as NEC and FIP.

In the present study male sex, RDS and PDA were associated with the development of NEC. These results are consistent with those of a recent large cohort study showing that RDS and PDA are related to the onset of NEC.<sup>8</sup> An epidemiologic study in the USA has shown that the incidence of NEC increased after surfactant use became a standard of care in the 1990s.<sup>9</sup> This suggests that RDS requiring surfactant replacement therapy is associated with the development of NEC. In 2013, Wariki *et al.* showed that PDA was the most significant risk factor for NEC based on an international collaborative database of four Asian countries.<sup>10</sup> Furthermore, the Israel Neonatal Network reported that PDA is a significant risk factor associated with the development of NEC in VLBW infants.<sup>11</sup> The present results and these previous findings suggest that the reduced mesenteric blood perfusion due to PDA may subsequently result in the development of NEC. In contrast,

we did not identify any relationships between the development of NEC and previously reported risk factors, such as the use of red blood cell transfusions,<sup>12,13</sup> maternal cigarette smoking,<sup>14</sup> placental abruption<sup>15</sup> and pre-eclampsia.<sup>16</sup> The fact that the present study was limited to surgical NEC among VLBW infants may account for the differences between the present results and the findings of former reports. Given that preterm neonates who required surgery were smaller and sicker compared with those managed medically, surgical NEC should be evaluated as distinct from non-surgical NEC.<sup>17</sup> Regarding preventive factors of NEC, many previous studies have shown that the use of antenatal glucocorticoids decreases the incidence of NEC,<sup>18,19</sup> In contrast, Guthrie *et al.* reported that exposure to antenatal glucocorticoids was associated with an increased risk for NEC using a national database of the USA.<sup>17</sup> The effect of antenatal glucocorticoids on the development of NEC remains controversial. The present study did not show any associations between antenatal glucocorticoids use and NEC development. In order to determine whether antenatal glucocorticoids have a preventive effect on NEC, a prospective study based on a unified treatment protocol of antenatal glucocorticoids is necessary.

In contrast to that observed for NEC, we identified only one risk factor, “outborn delivery”, for FIP. The transfer of patients to participating hospitals after the development of intestinal perforation may account for this result. Although some studies have shown that the use of indomethacin for PDA closure is a significant independent risk factor for FIP,<sup>20,21</sup> other studies have identified postnatal use of dexamethasone, but not indomethacin, as a significant risk factor for FIP.<sup>22</sup> In the present study we found no associations of dexamethasone or indomethacin with FIP. Differing protocols for the dexamethasone and indomethacin therapy may account for the different results. Recently, there have been several reports indicating that congenital or acquired defects in the muscular layer of the intestine may be involved in the pathogenesis of FIP.<sup>6,23</sup> This suggests that FIP is due not only to acquired but also to congenital factors.

Regarding MRI, several risk factors including cesarean section, maternal MgSO<sub>4</sub> therapy and maternal diabetes have been identified, based on small cohorts.<sup>24,25</sup> Most of these studies, however, included both surgically and medically treated MRI, and both