30 Fukao T, Nakamura H, Nakamura K *et al.* Characterization of six mutations in five Spanish patients with mitochondrial acetoacetyl-CoA thiolase deficiency: Effects of amino acid substitutions on tertiary structure. *Mol. Genet. Metab.* 2002; **75**: 235–43.
31 Fukao T, Matsuo N, Zhang GX *et al.* Single base substitutions at the initiator codon in the mitochondrial acetoacetyl-CoA thiolase (ACAT1/T2) gene result in production of varying amounts of wild-type T2 polypeptide. *Hum. Mutat.* 2003; **21**: 587–92.
32 Fukao T, Zhang GX, Sakura N *et al.* The mitochondrial acetoacetyl-CoA thiolase (T2) deficiency in Japanese patients: Urinary organic acid and blood acylcarnitine profiles under stable conditions have subtle abnormalities in T2-deficient patients with some residual T2 activity. *J. Inherit. Metab. Dis.* 2003; **26**: 423–31.
33 Zhang GX, Fukao T, Rolland MO *et al.* Mitochondrial acetoacetyl-CoA thiolase (T2) deficiency: T2-deficient patients with "mild" mutation(s) were previously misinterpreted as normal by the coupled assay with tiglyl-CoA. *Pediatrics* 2004; **56**: 60–64.
34 Mrazova L, Fukao T, Halovd K *et al.* Two novel mutations in mitochondrial acetoacetyl-CoA thiolase deficiency. *J. Inherit. Metab. Dis.* 2005; **28**: 235–6.
35 Zhang G, Fukao T, Sakurai S, Yamada K, Michael Gibson K, Kondo N. Identification of Alu-mediated, large deletion-spanning exons 2-4 in a patient with mitochondrial acetoacetyl-CoA thiolase deficiency. *Mol. Genet. Metab.* 2006; **89**: 222–6.
36 Sakurai S, Fukao T, Haapalainen AM *et al.* Kinetic and expression analyses of seven novel mutations in mitochondrial acetoacetyl-CoA thiolase (T2): Identification of a Km mutant and an analysis of the mutational sites in the structure. *Mol. Genet. Metab.* 2007; **90**: 370–78.
37 Fukao T, Boneh A, Aoki Y, Kondo N. A novel single-base substitution (c.1124A>G) that activates a 5-base upstream cryptic splice donor site within exon 11 in the human mitochondrial acetoacetyl-CoA thiolase gene. *Mol. Genet. Metab.* 2008; **94**: 417–21.
38 Fukao T, Horikawa R, Naiki Y *et al.* A novel mutation (c.951C>T) in an exonic splicing enhancer results in exon 10 skipping in the human mitochondrial acetoacetyl-CoA thiolase gene. *Mol. Genet. Metab.* 2010; **100**: 339–44.
39 Fukao T, Nguyen HT, Nguyen NT *et al.* A common mutation, R208X, identified in Vietnamese patients with mitochondrial acetoacetyl-CoA thiolase (T2) deficiency. *Mol. Genet. Metab.* 2010; **100**: 37–41.
40 Thummler S, Dupont D, Acquaviva C, Fukao T, de Ricaud D. Different clinical presentation in siblings with mitochondrial acetoacetyl-CoA thiolase deficiency and identification of two novel mutations. *Tohoku J. Exp. Med.* 2010; **220**: 27–31.
41 Catanzano F, Ombrone D, Di Stefano C *et al.* The first case of mitochondrial acetoacetyl-CoA thiolase deficiency identified by expanded newborn metabolic screening in Italy: The importance of an integrated diagnostic approach. *J. Inherit. Metab. Dis.* 2010; **33**: 91–4.
42 Sarafoglou K, Matern D, Redlinger-Grosse K *et al.* Siblings with mitochondrial acetoacetyl-CoA thiolase deficiency not identified by newborn screening. *Pediatrics* 2011; **128**: 246–50.
43 Fukao T, Maruyama S, Ohura T *et al.* Three Japanese patients with beta-ketothiolase deficiency who share a mutation, c.431A>C (H144P) in ACAT1: Subtle abnormality in urinary organic acid analysis and blood acylcarnitine analysis using tandem mass spectrometry. *JIMD Rep.* 2012; **3**: 107–15.
44 Buhas D, Bernard G, Fukao T, Decarie JC, Chouinard S, Mitchell GA. A treatable new cause of chorea: Beta-ketothiolase deficiency. *Mov. Disord.* 2013; **28**: 1054–6.
45 Zschocke J, Ruiter JP, Brand J *et al.* Progressive infantile neurodegeneration caused by 2-methyl-3-hydroxybutyryl-CoA dehydrogenase deficiency: A novel inborn error of branched-chain fatty acid and isoleucine metabolism. *Pediatr. Res.* 2000; **48**: 852–5.
46 Ensenauer R, Niederhoff H, Ruiter JP *et al.* Clinical variability in 3-hydroxy-2-methylbutyryl-CoA dehydrogenase deficiency. *Ann. Neurol.* 2002; **51**: 656–9.
47 Olpin SE, Pollitt RJ, McMenamin J *et al.* 2-Methyl-3-hydroxybutyryl-CoA dehydrogenase deficiency in a 23-year-old man. *J. Inherit. Metab. Dis.* 2002; **25**: 477–82.
48 Ofman R, Ruiter JP, Feenstra M *et al.* 2-Methyl-3-hydroxybutyryl-CoA dehydrogenase deficiency is caused by mutations in the HADH2 gene. *Am. J. Hum. Genet.* 2003; **72**: 1300–7.
49 Sutton VR, O'Brien WE, Clark GD, Kim J, Wanders RJ. 3-Hydroxy-2-methylbutyryl-CoA dehydrogenase deficiency. *J. Inherit. Metab. Dis.* 2003; **26**: 69–71.
50 Poll-The BT, Wanders RJ, Ruiter JP *et al.* Spastic diplegia and periventricular white matter abnormalities in 2-methyl-3-hydroxybutyryl-CoA dehydrogenase deficiency, a defect of isoleucine metabolism: Differential diagnosis with hypoxic-ischemic brain diseases. *Mol. Genet. Metab.* 2004; **81**: 295–9.
51 Sass JO, Forstner R, Sperl W. 2-Methyl-3-hydroxybutyryl-CoA dehydrogenase deficiency: Impaired catabolism of isoleucine presenting as neurodegenerative disease. *Brain Dev.* 2004; **26**: 12–14.
52 Perez-Cerda C, Garcia-Villoria J, Ofman R *et al.* 2-Methyl-3-hydroxybutyryl-CoA dehydrogenase (MHBD) deficiency: An X-linked inborn error of isoleucine metabolism that may mimic a mitochondrial disease. *Pediatr. Res.* 2005; **58**: 488–91.
53 Cazorla MR, Verdu A, Perez-Cerda C, Ribes A. Neuroimage findings in 2-methyl-3-hydroxybutyryl-CoA dehydrogenase deficiency. *Pediatr. Neurol.* 2007; **36**: 264–7.
54 Lenski C, Kooy RF, Reyniers E *et al.* The reduced expression of the HADH2 protein causes X-linked mental retardation, choreoathetosis, and abnormal behavior. *Am. J. Hum. Genet.* 2007; **80**: 372–7.
55 Garcia-Villoria J, Navarro-Sastre A, Fons C *et al.* Study of patients and carriers with 2-methyl-3-hydroxybutyryl-CoA dehydrogenase (MHBD) deficiency: Difficulties in the diagnosis. *Clin. Biochem.* 2009; **42**: 27–33.
56 Yang SY, He XY, Olpin SE *et al.* Mental retardation linked to mutations in the HSD17B10 gene interfering with neurosteroid and isoleucine metabolism. *Proc. Natl Acad. Sci. USA* 2009; **106**: 14820–24.
57 Garcia-Villoria J, Gort L, Madrigal I *et al.* X-inactivation of HSD17B10 revealed by cDNA analysis in two female patients with 17beta-hydroxysteroid dehydrogenase 10 deficiency. *Eur. J. Hum. Genet.* 2010; **18**: 1353–5.
58 Rauschenberger K, Scholer K, Sass JO *et al.* A non-enzymatic function of 17beta-hydroxysteroid dehydrogenase type 10 is required for mitochondrial integrity and cell survival. *EMBO Mol. Med.* 2010; **2**: 51–62.
59 Seaver LH, He XY, Abe K *et al.* A novel mutation in the HSD17B10 gene of a 10-year-old boy with refractory epilepsy, choreoathetosis and learning disability. *PLoS ONE* 2011; **6**: e27348.
60 Zschocke J. HSD10 disease: Clinical consequences of mutations in the HSD17B10 gene. *J. Inherit. Metab. Dis.* 2012; **35**: 81–9.
61 Fukao T, Akiba K, Goto M *et al.* The first case in Asia of 2-methyl-3-hydroxybutyryl-CoA dehydrogenase deficiency (HSD10 disease) with atypical presentation. *J. Hum. Genet.* 2014; **59**: 609–14.

© 2015 Japan Pediatric Society