

How do you feel?

Date(MM/DD/YYYY) / /

Name :

ID :

Respondent-completed / Interview by Name:

How has your daily life been for the last month?

Please answer the following questions by circling the appropriate responses

(Exclude any difficulties caused by physical issues, e.g., pain).

Please ask for any help if needed.

You talk and ask about the same things repeatedly.	YES	NO	N/A Don't know
You have become unable to understand the context of facts.	YES	NO	N/A
You have become indifferent about clothing and other personal concerns.	YES	NO	N/A
You have begun to forget to turn off the faucet and/or close the door, and/or have become unable to clean up properly.	YES	NO	N/A
When doing two things at the same time, you forget one of them.	YES	NO	N/A
You have become unable to take medication under proper management.	YES	NO	N/A
You have begun to take a longer time to do work (e.g., household chores), which could be done quickly before.	YES	NO	N/A
You have become unable to make a plan.	YES	NO	N/A
You cannot understand complex topics.	YES	NO	N/A
You have become less interested and willing, and stopped hobbies, etc.	YES	NO	N/A
You have become more irritable and suspicious than before.	YES	NO	N/A
TOTAL SED-11Q SCORE			

Figure 1 *Continued*

For years, there has been a wide debate over whether or not community-wide population screening for dementia helps patients and saves healthcare costs. However, at present, there is insufficient evidence of the benefits to justify community-wide population screening from the perspective of clinical outcomes, emotional effects and

cost-effectiveness.^{8,37,38} The causes for concern are a lack of screening measures that are both sensitive and specific enough to detect dementia, and a shortage of local resources. As aforementioned, no tests or questionnaires are sensitive and specific enough to avoid false positives, and people would suffer from incidental findings that

were found to be false. Undergoing unnecessary procedures after screening alone could be a great mental burden, adding to anxiety and fear raised by screening results. Significant resources are required for population screening itself, and thus implementation would pose additional burdens to local communities who are already confronted with a shortage of social resources. Detection of dementia is the initial step in the medical and care process, and thus for effective implementation of population screening, medical and care follow-up systems should be required, including mental care and support.

Case findings appear to be a more appropriate method than population screening to avoid false positives.³⁹ In many cases, the first stage of diagnosis starts with suspicion of dementia by those who know the person well, such as family members, colleagues and neighbors.⁸ GPs and primary care physicians might also be keeping watchful eyes for signs of dementia. In a survey carried out in Canada, 26% of primary care physicians claimed that they routinely checked for signs of dementia in their patients.⁴⁰ Community nurses and staff at health-care centers may notice changes in community residents, and recommend them and/or their family members to consult with doctors.⁸ Screening tools could provide information to verify whether or not such suspicions should be regarded as signs of dementia, and further investigation should be required. It is recommended to use the screening tools when sufficient information has been accumulated to suspect dementia. In the case of population screening, the diagnosis process begins with screening using tests and/or questionnaires, often without any information, whereas in case findings, tests and/or questionnaires are carried out only after sufficient information is gathered to suspect dementia. In either case, ethical issues must be carefully considered, including protection of personal information, obtaining informed consent, relieving the mental burden of undergoing screening, and the problem of overcoming stigmatization and exclusion.

Issues to be addressed for effective implementation of early diagnosis

Accurate diagnosis, avoiding false positives and overdiagnosis

It is evident that effective therapy, intervention and support depend on an accurate diagnosis, and it should be noted that screening is absolutely different from diagnosis. Considering the mental burden and stigma that result from diagnosis, false positives should be carefully avoided.

One of the fundamental issues is a wide variation in GPs' abilities and confidence in diagnosing and managing dementia.⁴¹ At present, it is a practical resolution that the responsibility of diagnosis of dementia is shared

between generalists and specialists. The government launched a plan to designate Medical Centers for Dementia in 2008, and 171 hospitals had been designated up to May 2012. The functions of these centers are to establish medical-medical and medical-care cooperation in local communities. The GPs play an important gatekeeper role in referring people for diagnosis by a specialist, rather than taking on the burden of diagnosis themselves. Based on the referral, the centers diagnose patients, and once the diagnosis has been confirmed, the GPs are expected to provide both practical and emotional support for patients and their family members, and refer them for additional psychosocial support if required. If there is a need for a consultation with a specialist, the patients can revisit the center on referral by the GP. At present, the centers are overloaded because they are forced to provide daily practical and emotional support for patients as a result of the insufficient capabilities of GPs to deal with dementia. Thus, education of GPs is a pressing issue for treatment after detection of dementia. The Ministry of Health, Labor and Welfare in Japan certifies Dementia Support Doctors who lead and support the primary care doctors and other dementia care professionals, and holds the skill-up programs of dementia medicine for primary care doctors and GPs.

Human resource development under a community-based integrated care system

It is a challenge to ensure compatibility between cost-effectiveness and quality care, and quality care could be provided if there was effective cooperation and coordination between healthcare professionals and organizations.^{42,43} In Japan, the long-term care insurance system aims to promote the implementation of a community-based integrated care system with the goal of deinstitutionalization and promotion of at-home care. Under the system, allocation of social resources and capital was delegated to each local government for management and implementation, and basically, services, including medical treatment, care and community life support, are provided within local communities. The policy is based on the expectation that each community will create a "tailored" system to meet the health needs, health status and values of the community.^{44,45} Under the Five-Year Plan for Promotion of Measures against Dementia in Japan, the pilot project of the initial-phase intensive support teams has been launched to grope the community-based implementation methods of earlier detection and intervention; the support team consists of healthcare professionals including nurses, health nurses and occupational therapists.

Development of human resources is one of the urgent requirements. In addition to the education and training of medical and healthcare staff, community residents

are expected to act as participants, and not merely as beneficiaries of services.⁴⁶ It is vital to implement measures that involve all members of society, so that they can help and support each other towards the improvement of the living environment for the elderly. This can be achieved through cooperation and active involvement among all sectors of society, such as the national government, local governments, corporations, local communities, non-profit organizations, family members and individuals with dementia. Task shifting is one of the key concepts; that is, shifting some work carried out by healthcare professionals to volunteers in order to reduce the workload of professional workers. The policy encourages volunteers to assist in the lives of demented individuals, and good outcomes have been shown in a number of local communities. For implementation of beneficial community-based integrated care after diagnosis, promoting social participation and community involvement of the residents should be a key strategy in addressing the shortage of human resources.

Acknowledgments

Dr H Yamaguchi was supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, Sports, Culture and Technology, Japan (23300197), and a Grant-in-Aid for Scientific Research (H24-Ninchisho-Ippan-002, H25-Ninchisho-Ippan-008) from the Ministry of Health, Labor and Welfare, Japan. The authors are equally contributed to this article.

Disclosure statement

The authors declare no conflict of interest.

References

- Morris JC. Early-stage and preclinical Alzheimer disease. *Alzheimer Dis Assoc Disord* 2005; **19**: 163–165.
- Wolf DS, Gearing M, Snowdon DA, Mori H, Markesbery WR, Mirra SS. Progression of regional neuropathology in Alzheimer disease and normal elderly: findings from the Nun study. *Alzheimer Dis Assoc Disord* 1999; **13**: 226–231.
- Gauthier S, Reisberg B, Zaudig M *et al*. Mild cognitive impairment. *Lancet* 2006; **367**: 1262–1270.
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition*. Washington DC: American Psychiatric Association, 2013.
- Geslani DM, Tierney MC, Herrmann N, Szalai JP. Mild cognitive impairment: an operational definition and its conversion rate to Alzheimer's disease. *Dement Geriatr Cogn Disord* 2005; **19**: 383–389.
- Banerjee S, Willis R, Matthews D, Contell F, Chan J, Murray J. Improving the quality of care for mild to moderate dementia: an evaluation of the Croydon Memory Service Model. *Int J Geriatr Psychiatry* 2007; **22**: 782–788.
- Sheaff M, Sherriff I, McFarlane H, Giarch G. Early diagnosis of dementia in Plymouth. Plymouth; 2011. [Cited 2013 Sep 28]. Available from URL: <http://www1.plymouth.ac.uk/research/CRA/Documents/Early%20diagnosis%20of%20dementia%20in%20Plymouth%20Report.pdf>.
- Bamford C, Lamont S, Eccles M, Robinson L, May C, Bond J. Disclosing a diagnosis of dementia: a systematic review. *Int J Geriatr Psychiatry* 2004; **19**: 151–169.
- Husband HJ. Diagnostic disclosure in dementia: an opportunity for intervention? *Int J Geriatr Psychiatry* 2000; **15**: 544–547.
- Green KN, Billings LM, Roozendaal B, McLaugh JL, LaFerla FM. Glucocorticoids increase amyloid-beta and tau pathology in a mouse model of Alzheimer's disease. *J Neurosci* 2006; **26**: 9047–9056.
- Jolley DJ, Benbow SM. Stigma and Alzheimer's disease: causes, consequences and a constructive approach. *Int J Clin Pract* 2000; **54**: 117–119.
- Werner P, Heinik J. Stigma by association and Alzheimer's disease. *Aging Ment Health* 2008; **12**: 92–99.
- Blay SL, Toledo Pisa Peluso E. Public stigma: the community's tolerance of Alzheimer disease. *Am J Geriatr Psychiatry* 2010; **18**: 163–171.
- Werner P, Mittelman MS, Goldstein D, Heinik J. Family stigma and caregiver burden in Alzheimer's disease. *Gerontologist* 2012; **52**: 89–97.
- Brodsky H, Howarth GC, Mant A, Kurlle SE. General practice and dementia. A national survey of Australian GPs. *Med J Aust* 1994; **160**: 10–14.
- van Hout H, Vernooij-Dassen M, Bakker K, Blom M, Grol R. General practitioners on dementia: tasks, practices and obstacles. *Patient Educ Couns* 2000; **39**: 219–225.
- Holroyd S, Turnbull Q, Wolf AM. What are patients and their families told about the diagnosis of dementia? Results of a family survey. *Int J Geriatr Psychiatry* 2002; **17**: 218–221.
- Iliffe S, Manthorpe J, Eden A. Sooner or later? Issues in the early diagnosis of dementia in general practice: a qualitative study. *Fam Pract* 2003; **20**: 376–381.
- Banerjee S, Wittenberg R. Clinical and cost effectiveness of services for early diagnosis and intervention in dementia. *Int J Geriatr Psychiatry* 2009; **24**: 748–754.
- Weimer DL, Sager MA. Early identification and treatment of Alzheimer's disease: social and fiscal outcomes. *Alzheimers Dement* 2009; **5**: 215–226.
- Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975; **12**: 189–198.
- Nasreddine ZS, Phillips NA, Bedirian V *et al*. The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *J Am Geriatr Soc* 2005; **53**: 695–699.
- Meulen EF, Schmand B, van Campen JP *et al*. The seven minute screen: a neurocognitive screening test highly sensitive to various types of dementia. *J Neurol Neurosurg Psychiatry* 2004; **75**: 700–705.
- Cummings JL. Mini-Mental State Examination. Norms, normals, and numbers. *JAMA* 1993; **269**: 2420–2421.
- Crum RM, Anthony JC, Bassett SS, Folstein MF. Population-based norms for the Mini-Mental State Examination by age and educational level. *JAMA* 1993; **269**: 2386–2391.
- Lawrence J, Davidoff D, Katt-Lloyd D, Auerbach M, Hennen J. A pilot program of improved methods for community-based screening for dementia. *Am J Geriatr Psychiatry* 2001; **9**: 205–211.
- Espino DV, Lichtenstein MJ, Palmer RF, Hazuda HP. Evaluation of the mini-mental state examination's internal

- consistency in a community-based sample of Mexican-American and European-American elders: results from the San Antonio Longitudinal Study of Aging. *J Am Geriatr Soc* 2004; **52**: 822–827.
- 28 Lorentz WJ, Scanlan JM, Borson S. Brief screening tests for dementia. *Can J Psychiatry* 2002; **47**: 723–733.
 - 29 Holsinger T, Deveau J, Boustani M, Williams JW, Jr. Does this patient have dementia? *JAMA* 2007; **297**: 2391–2404.
 - 30 Jonker C, Geerlings MI, Schmand B. Are memory complaints predictive for dementia? A review of clinical and population-based studies. *Int J Geriatr Psychiatry* 2000; **15**: 983–991.
 - 31 Mograbi DC, Brown RG, Morris RG. Anosognosia in Alzheimer's disease—the petrified self. *Conscious Cogn* 2009; **18**: 989–1003.
 - 32 Maki Y, Amari M, Yamaguchi T, Nakaaki S, Yamaguchi H. Anosognosia: patients' distress and self-awareness of deficits in Alzheimer's disease. *Am J Alzheimers Dis Other Dement* 2012; **27**: 339–345.
 - 33 Morris JC. The Clinical Dementia Rating (CDR): current version and scoring rules. *Neurology* 1993; **43**: 2412–2414.
 - 34 Maki Y, Yamaguchi T, Yamaguchi H. Symptoms of early dementia-11 questionnaire (SED-11Q): a brief informant-operated screening for dementia. *Dement Geriatr Cogn Dis Extra* 2013; **3**: 131–142.
 - 35 Maki Y, Yamaguchi T, Yamaguchi H. Evaluation of anosognosia in Alzheimer's disease using the symptoms of early dementia-11 questionnaire (SED-11Q). *Dement Geriatr Cogn Dis Extra* 2013; **3**: 351–359.
 - 36 Galvin JE, Roe CM, Powlisha KK *et al*. The AD8: a brief informant interview to detect dementia. *Neurology* 2005; **65**: 559–564.
 - 37 Iliffe S, Robinson L, Brayne C *et al*. Primary care and dementia: 1. diagnosis, screening and disclosure. *Int J Geriatr Psychiatry* 2009; **24**: 895–901.
 - 38 Boustani M, Peterson B, Hanson L, Harris R, Lohr KN. Screening for dementia in primary care: a summary of the evidence for the U.S. Preventive Services Task Force. *Ann Intern Med* 2003; **138**: 927–937.
 - 39 Brayne C, Fox C, Boustani M. Dementia screening in primary care: is it time? *JAMA* 2007; **298**: 2409–2411.
 - 40 Valcour VG, Masaki KH, Curb JD, Blanchette PL. The detection of dementia in the primary care setting. *Arch Intern Med* 2000; **160**: 2964–2968.
 - 41 O'Connor DW, Pollitt PA, Hyde JB, Brook CP, Reiss BB, Roth M. Do general practitioners miss dementia in elderly patients? *BMJ* 1988; **297**: 1107–1110.
 - 42 Moss F, Garside P, Dawson S. Organisational change: the key to quality improvement. *Qual Health Care* 1998; **7** (Suppl): S1–S2.
 - 43 Plochg T, Klazinga NS. Community-based integrated care: myth or must? *Int J Qual Health Care* 2002; **14**: 91–101.
 - 44 Proenca EJ. Community orientation in health services organizations: the concept and its implementation. *Health Care Manage Rev* 1998; **23**: 28–38.
 - 45 Proenca EJ, Rosko MD, Zinn JS. Community orientation in hospitals: an institutional and resource dependence perspective. *Health Serv Res* 2000; **35**: 1011–1035.
 - 46 World Health Organization. The World Health Report 2000 – Health Systems: Improving Performance. Geneva; 2000 [Cited 2013 Sep 28]. Available from URL: <http://www.who.int/whr/2000/en/>.

Intensive rehabilitation for dementia improved cognitive function and reduced behavioral disturbance in geriatric health service facilities in Japan

Kenji Toba,¹ Yu Nakamura,² Hidetoshi Endo,¹ Jiro Okochi,³ Yukiko Tanaka,³ Chiyako Inaniwa,³ Akira Takahashi,³ Naoko Tsunoda,³ Kentaro Higashi,³ Motoharu Hirai,³ Hiroyuki Hirakawa,³ Shizuru Yamada,³ Yohko Maki,⁴ Tomoharu Yamaguchi⁴ and Haruyasu Yamaguchi⁴

¹National Center for Geriatrics and Gerontology, Ohbu, ²Department of Psychiatry, Kagawa University, Takamatsu, ³Japan Association of Geriatric Health Service Facilities, Tokyo, and ⁴Graduate School of Health Sciences, Gunma University, Maebashi, Japan

Aim: To examine the efficacy of rehabilitation for elderly individuals with dementia at intermediate facilities between hospitals and home, based on the policies for elderly individuals to promote community-based care at home and dehospitalization.

Methods: Participants were older adults with dementia newly admitted to intermediate facilities. A total of 158 in the intervention group who claimed Long-Term Care Insurance for three consecutive months, and 54 in the control group were included in the analysis. The interventions were carried out in a tailor-made manner to meet individual needs. The personal sessions were carried out three times a week for 3 months after admission by physical, occupational or speech therapists. Outcome measures were cognitive tests (Hasegawa Dementia Scale revised [HDS-R] and Mini-Mental State Examination), and observational assessments of dementia severity, activities of daily living (ADL), social activities, behavioral and psychological symptoms of dementia (BPSD) using a short version of the Dementia Disturbance Scale (DBD13), depressive mood, and vitality.

Results: Significant improvement in the intervention group was shown in cognitive function measured by HDS-R (interaction $F[1, 196] = 5.190, P = 0.024$), observational evaluation of dementia severity ($F[1, 198] = 9.550, P = 0.002$) and BPSD (DBD13; $F[1, 197] = 4.506, P = 0.035$). Vitality, social activities, depressive mood and ADL were significantly improved only in the intervention group, although interaction was not significant.

Conclusions: Significant improvement by intervention was shown in multiple domains including cognitive function and BPSD. Cognitive decline and worsening of BPSD are predictors of care burden and hospitalization, thus intensive rehabilitation for dementia was beneficial for both individuals with dementia and their caregivers. *Geriatr Gerontol Int* 2014; 14: 206–211.

Keywords: behavioral and psychological symptoms of dementia, clinical medicine, Dementia Disturbance Scale short version, dementia, geriatric medicine, rehabilitation, tailor-made.

Introduction

Promoting community-based care at home and dehospitalization is one of the main policies for elderly individuals. In order to reduce the length of hospital stay, it is recommended to establish a rehabilitation and care system for the elderly just after leaving hospital. Thus, the Japanese government established the “Geriatric

Health Service Facility” in 1986 (Long-Term Care Health Facility after 2000; Roken), which is a transitional facility between hospital and home or nursing home to provide medical treatment, nursing care, and rehabilitation. Elderly individuals are admitted to Roken after their condition has become stable in hospital, and stay until they are ready to return home. After returning home, Roken offers community-based rehabilitation and various care services to support home-based care, and facilitates networks for intraregional exchanges among municipalities, local healthcare and social welfare services.

Since Roken was launched, the number of inpatients with dementia has markedly increased. Hospitalization

Accepted for publication 20 March 2013.

Correspondence: Dr Kenji Toba MD PhD, National Center for Geriatrics and Gerontology, 35 Gengo, Morioka-cho, Ohbu-city, Aichi Japan 474-8511. Email: toba@ncgg.go.jp




Figure 1 Flow of participants in the intervention and control groups. HDS-R, Hasegawa Dementia Scale revised. MMSE, Mini-Mental State Examination.

itself can cause cognitive deterioration, even during a hospital stay for diseases other than dementia, and patients are often not expected to recover to their pre-hospitalization level.¹ Other predictors of hospitalization are caregivers' burden and the interrelationship with caregivers.² Behavioral and psychological symptoms of dementia (BPSD) are a source of distress for caregivers and a major reason for hospitalization.^{3,4} Additionally, disuse syndrome is triggered by psychological factors associated with dementia, such as a depressive and apathetic mood.⁵⁻⁹ Disuse syndrome can lead to deterioration of cognitive and physical function, which can result in repeated hospitalization.

To break the vicious cycle of repeated hospitalization, effective rehabilitation just after discharge from hospital is required, and Roken was singled out as the appropriate facility for the rehabilitation. Thus, in 2006, the Japanese Long-term Care Insurance system introduced intensive rehabilitation for individuals with dementia who were newly admitted to Roken, consisting of personal rehabilitation three times a week for 3 months. This rehabilitation has become widely practiced since its introduction. However, the efficacy has not been examined, although the rehabilitation is payable under long-term insurance. Thus, a model project was organized to examine the efficacy of the rehabilitation for dementia in Roken throughout Japan.

Methods

Study members

Study committee members were researchers excluding stakeholders of any Roken, and committee observers were staff of the Health and Welfare Bureau for the

Elderly, Ministry of Health, Labour and Welfare. The committee designed the research, selected 36 Rokens, and interpreted the data. Data were collected by rehabilitation staff in the 36 Rokens.

Participants

The study was carried out between July 2007 and February 2008. The flow of participants is shown in Figure 1. Survey slips were sent to the facilities in July 2007. The facilities were required to send them back after the pre-intervention and post-intervention assessment, respectively. Inclusion criteria of the intervention group were: (i) newly admitted patients with dementia diagnosed by *The Diagnostic and Statistical Manual of Mental Disorders IV*; (ii) with Mini-Mental State Examination (MMSE) or Hasegawa Dementia Scale revised (HDS-R) score between 5 and 25 at pre-intervention assessment; and (iii) who claimed Long-Term Care Insurance for three consecutive months. Inclusion criteria of the control group were: (i) and (ii), and (iii) who did not receive interventions. The participants were not randomized. We received 271 responses, and among them, 212 individuals met the inclusion criteria (158 in intervention group and 54 in control group; Table 1). Informed consent was given from all participants or their responsible care giver. The research plan was approved by the Ethics Board of the Japan Association of Geriatric Health Services Facilities.

Assessment

The assessment was minimized to reduce the burden of facilities staff. As the interventions were carried out by therapists during working time, it would have been

Table 1 Demographic data

		Intervention	Control	
<i>n</i>		158	54	
Male/female (%)		30.2/69.8	39.6/60.4	NS
Age		84.1 ± 7.1	87.3 ± 7.1	P = 0.005 [†]
Dementia	AD	22	7	NS
	VD	52	15	NS
	DLB	3	0	NS
	FTD	2	0	NS
	Others/unknown	79	32	NS

[†]Significant difference by two-sample *t*-test. AD, Alzheimer's disease; DLB, dementia with Lewy bodies; FTD, front-temporal dementia; M/F, male/female; NS, no significant difference by χ^2 test; VD, vascular dementia.

difficult to collect many data if the assessment were complicated. The assessment scales were chosen based on preliminary studies, which were carried out in the last 2 years.

Cognitive tests

The MMSE and HDS-R were carried out. HDS-R is similar to MMSE, but lays more weight on memory than does MMSE.

Questionnaires

For the assessment of subjective mood, the participants were required to answer the interview of a short version of the Geriatric Depression Scale (GDS;¹⁰ scores are between 0–5, high scores indicate more depressive mood). Facility care staff assessed activities of daily living (ADL), BPSD, N-Memory Scale (NM),¹¹ vitality index¹² and the Social Activity Scale. ADL was assessed using the Barthel Index (scoring was changed: total assistance of 0 to independence of 3 for each item, and full score of 15).¹³ In addition to ADL, the capacity for social interaction was measured using the Social Activity Scale, whose sub-items were conversation with facility staff members, conversation with other residents, organizing own belongings, participation in recreational activities, and outings (total assistance of 0 to independence of 3 for each item, and full score of 15). BPSD was evaluated using a short version of the Dementia Behavior Disturbance Scale (DBD;¹³ “never” of 0 to “usually” of 3 for each item and full score of 48).¹⁴ The NM Scale is an observational scale, which evaluates the stages of dementia in five domains: housework, social interaction and interest, communication, memory, and orientation (“impossible” of 0 to “normal” of 10 and full score of 50). The Vitality Index evaluates motivation in daily living, with sub-items of waking up, greetings, having meals, elimination, and participation in rehabilitation and/or recreation (“indifferent” of 0 to “voluntarily” of 2 and full score of 10).

Intervention

Before commencement of the study, a training workshop was held to introduce the intervention methods, whose efficacy was suggested by previous studies: such as reminiscence, reality orientation, memory rehabilitation, music therapy, physical exercise, occupational therapy, speech communication therapy and learning sessions.

The intervention was carried out in an individualized tailor-made manner.¹⁵ First, the individual functional profiles were assessed with regard to both abilities and disabilities to evaluate how to enhance the abilities and compensate for disabilities. Second, training activities were selected; the decision was shared between therapists and participants. Each personal session was took place three times a week for 3 months after admission by physical, occupational or speech therapists. Individuals in the control group took usual group therapies including exercise, singing songs and games.

Analysis of data

The data were analyzed using the Japanese version of SPSS for Windows version 19.0 (IBM Corporation, Armonk, NY, USA). For an initial baseline comparison between the intervention and control groups, two-sample *t*-tests were carried out; there was no significant difference between the two groups for any outcome measure. Participants who underwent the initial baseline and post-intervention assessments were included in the final analysis; dropout participants were excluded from the analysis. Repeated measures analysis of covariance (ANCOVA) with the covariate of age was used to analyze the completed cases. Age was used as a covariate, because the ages were significantly different between the two groups (Table 1). The interaction was examined to assess the differential effect between the intervention and control groups, and post-hoc “within subjects” analysis was carried out with Bonferroni correction. Regarding the measures where significant

interaction was shown, intention-to-treat analysis was also carried out; the participants who received the intervention but did not claim Long-Term Care Insurance for three consecutive months were included in the intention-to-treat analysis. A significant difference was set as $P < 0.05$.

Results

Demographic data of the participants are shown in Table 1. Analysis of 158 participants in the intervention group and 54 in the control group was carried out (Fig. 1). The number of participants who took donepezil during the intervention/observation period was two in both groups ($P = 0.269$, χ^2 -test).

Cognitive tests

Participants in the intervention group showed significant improvement in HDS-R score compared with those in the control group (interaction $F[1, 196] = 5.190$, $P = 0.024$; post-hoc intra-subject analysis: intervention group, $P = 0.001$, control group $P = 0.480$). There were no significant differences observed in MMSE (Table 2).

Questionnaire

The intervention group showed significant improvement compared with the control group in DBD¹³ ($F[1,197] = 4.506$, $P = 0.035$; post-hoc intra-subject analysis: intervention group, $P = 0.004$, control group $P = 0.413$) and NM Scale ($F[1,198] = 9.550$, $P = 0.002$; post-hoc intra-subject analysis: intervention group, $P < 0.001$, control group $P = 0.380$). Regarding the sub-items of the NM Scale, significant differences in interaction were observed for social interaction ($F[1,198] = 15.736$, $P < 0.001$), memory ($F[1,198] = 7.635$, $P = 0.006$) and orientation ($F[1,198] = 4.220$, $P = 0.041$).

Although the interaction was not significant, comparison between pre- and post-intervention showed significant improvement in ADL (Barthel Index), Social Activity Scale, motivation (Vitality Index) and mood (GDS) only in the intervention group after multiple correction (Table 2).

Intention-to-treat analysis

Significant differences remained in the intention-to-treat analysis in the HDS-R and NM Scale; HDS-R, interaction ($F[1, 230] = 4.466$, $P = 0.036$), post-hoc analysis within subjects: intervention group $P < 0.001$, control group $P = 0.585$; NM Scale, interaction ($F[1, 236] = 8.113$, $P = 0.005$), post-hoc analysis: intervention

Table 2 Outcome of intensive cognitive rehabilitation

	Intervention group		Control group		Interaction F (DF)	P	Intra-subject [†]	
	Pre mean \pm SD	Post mean \pm SD	Pre mean \pm SD	Post mean \pm SD			Intervention	Control
Cognitive test								
MMSE	19.1 \pm 4.5	19.4 \pm 5.5	19.5 \pm 4.9	18.2 \pm 7.4	1.780 (1,110)	0.185	0.542	0.234
HDS-R	16.9 \pm 5.7	17.9 \pm 6.5	17.0 \pm 5.9	16.7 \pm 6.3	5.190 (1,196)	0.024*	0.001**	0.480
Questionnaire								
NM	30.4 \pm 9.1	32.1 \pm 9.5	31.4 \pm 9.8	30.7 \pm 10.9	9.550 (1,198)	0.002**	$P < 0.001$ ***	0.380
ADL	16.4 \pm 7.1	17.3 \pm 7.1	15.7 \pm 7.0	15.9 \pm 6.9	1.448 (1,202)	0.230	0.001**	0.621
Activity	8.6 \pm 3.3	8.8 \pm 3.4	8.5 \pm 3.1	8.6 \pm 3.2	1.169 (1,200)	0.281	0.038*	0.972
Vitality	8.0 \pm 1.7	8.2 \pm 1.6	8.1 \pm 1.8	8.2 \pm 1.8	1.792 (1,199)	0.182	0.004**	0.864
DBD	4.5 \pm 5.1	4.0 \pm 4.1	4.5 \pm 4.2	4.8 \pm 4.7	4.506 (1,197)	0.035*	0.004**	0.413
GDS	2.5 \pm 1.8	2.4 \pm 1.9	2.3 \pm 1.5	2.4 \pm 1.5	2.048 (1,196)	0.154	0.042*	0.634

[†]Intra-subject: post-hoc analysis of intra-subject (comparison between pre- and post-intervention analysis). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. Activity, Original Activity Scale; ADL, Activities of daily living; DBD, Dementia Behavior Disturbance Scale; DF, degree of freedom; GDS, Geriatric Depression Scale; HDS-R, Hasegawa Dementia Scale revised; MMSE, Mini-Mental State Examination; NM, N-Memory Scale; Post, post-intervention assessment; Pre, pre-intervention assessment; Vitality, Vitality Index.

group $P < 0.001$, control group $P = 0.410$. The interaction of DBD was marginal; interaction ($F[1, 232] = 3.717, P = 0.055$), post-hoc analysis: intervention group $P = 0.007$, control group $P = 0.439$.

Discussion

Significant improvement by the intervention was shown in multiple domains; therefore, the intensive rehabilitation for dementia was beneficial for the individuals with dementia and also their caregivers. Pharmacological effects were thought to be negligible, as just two participants in both groups took donepezil during the intervention/observation period.

Regarding cognitive function, the effects of intensive rehabilitation for dementia were shown in both a cognitive test and observational evaluation of memory and orientation measured by NM Scale. In the symptomatic treatment of dementia, amelioration in daily living rather than in neuropsychological factors should be the therapeutic objectives, and thus the emphasis would be laid on improving performance in everyday life rather than on scores of cognitive tests.¹⁶ Besides, it is often pointed out that scores of cognitive tests cannot always be generalized to daily living, although cognitive tests are moderately predictive of functional status in everyday life.¹⁷ Therefore, mere enhancement of cognitive test scores is not sufficient, and beneficial changes in daily living are required. In the present study, cognitive improvement was shown in observational evaluation, in addition to a cognitive test. Cognitive enhancement is also beneficial for caregivers, because the severity of cognitive impairment could be a predictor of burden, in addition to BPSD.^{18,19} The effects of non-pharmacological approaches on cognitive function have not yet been established,^{16,19} and the present study could provide additional evidence for their benefit.

Amelioration of BPSD was also attained in the present study. Care for demented individuals requires allocation of longer times than for care of the elderly suffering from physical diseases. In particular, the presence of BPSD might induce more stress than do medical problems,^{4,20-23} and could result in depression or strain in caregivers.²⁴ Consequently, caregivers' burden is associated with an increased risk of institutionalization.²⁵ However, institutionalization could not solve caregivers' distress; a year after institutionalization, distress still persisted in caregivers.²⁶ In contrast, treatment of BPSD could help diminish caregiver burden.²⁷ Thus, it is beneficial both for individuals with dementia and their caregivers to reduce BPSD by rehabilitation in intermediate facilities between hospital and home.

In addition to enhancement of cognitive function and reduction of BPSD, improvement of social functioning and quality of life (QOL) should be the main outcomes of rehabilitation for dementia.¹⁶

Social isolation is associated with increased risk of mental decline,²⁸ whereas a rich social network and interaction might protect against mental decline.^{29,30} In demented individuals, symptoms of depression were a consistent predictor of QOL.³¹ In the present study, the intervention group showed improvement of social functioning measured by the Social Activity Scale, and amelioration of depressive mood measured by GDS.

Regarding the intervention, individualized tailor-made therapies were carried out, because the aim of the present study was to enhance each participant's ability to meet their individual needs, and not to show the efficacy of any specific method. Personally-relevant goals were identified, and the therapist worked with the individuals with dementia to devise strategies to cope with difficulties in their everyday lives by building on the person's strengths and developing ways of compensating for impairment.¹⁵ Personal selection was considered an essential therapeutic element to enhance the motivation and optimize the emotional impact of the training. Changing and combining methods were allowed during the intervention period.

The present study showed that intensive rehabilitation should be beneficial for both individuals with dementia and caregivers. To promote community-based care and dehospitalization, continuity of rehabilitation is desirable to maintain function after returning home; another mission of Roken is to offer community-based rehabilitation and various care services to support home-based care.

As a limitation, the participants were not randomized. By data cleaning, data including missing values were excluded so that the numbers of valid data were different among assessments. Finally, for evaluation of the effects on dehospitalization, a longitudinal follow-up study is required.

Acknowledgment

This study is partially supported by a Grant-in-Aid from the Ministry of Health Labor and Welfare (#19 Health Promotion Projects for Elderly-169 2-Na).

Disclosure statement

The authors declare no conflict of interest.

References

- 1 Zekry D, Herrmann FR, Grandjean R *et al.* Does dementia predict adverse hospitalization outcomes? A prospective study in aged inpatients. *Int J Geriatr Psychiatry* 2009; **24**: 283-291.
- 2 Andrieu S, Reynish E, Nourhashemi F *et al.* Predictive factors of acute hospitalization in 134 patients with Alzheimer's disease: a one year prospective study. *Int J Geriatr Psychiatry* 2002; **17**: 422-426.

- 3 Cerejeira J, Lagarto L, Mukaetova-Ladinska EB. Behavioral and psychological symptoms of dementia. *Front Neurol* 2012; **3**: 73.
- 4 Matsumoto N, Ikeda M, Fukuhara R *et al*. Caregiver burden associated with behavioral and psychological symptoms of dementia in elderly people in the local community. *Dement Geriatr Cogn Disord* 2007; **23**: 219–224.
- 5 Kobayashi T, Kato S. Depression-dementia medius: between depression and the manifestation of dementia symptoms. *Psychogeriatrics* 2011; **11**: 177–182.
- 6 Vilalta-Franch J, Calvo-Pexas L, Garre-Olmo J, Turro-Garriga O, Lopez-Pousa S. Apathy syndrome in Alzheimer's disease epidemiology: prevalence, incidence, persistence, and risk and mortality factors. *J Alzheimers Dis* 2012; **33**: 535–543.
- 7 Bjoerke-Bertheussen J, Ehrt U, Rongve A, Ballard C, Aarsland D. Neuropsychiatric symptoms in mild dementia with lewy bodies and Alzheimer's disease. *Dement Geriatr Cogn Disord* 2012; **34**: 1–6.
- 8 Quaranta D, Marra C, Rossi C, Gainotti G, Masullo C. Different apathy profile in behavioral variant of frontotemporal dementia and Alzheimer's disease: a preliminary investigation. *Curr Gerontol Geriatr Res* 2012; **2012**: 719250.
- 9 Xing Y, Wei C, Chu C *et al*. Stage-specific gender differences in cognitive and neuropsychiatric manifestations of vascular dementia. *Am J Alzheimers Dis Other Demen* 2012; **27**: 433–438.
- 10 Yesavage JA, Brink TL, Rose TL. Development and validation of a geriatric depression screening scale: a preliminary report. *J Psychiatr Res* 1982; **17**: 37–49.
- 11 Kobayashi T, Hariguchi S, Nishimura K *et al*. A new clinical scale for rating of mental states and activities of daily living of the elderly (NM scale and N-ADL). *Jpn J Clin Psychiatry* 1988; **17**: 1653–1668. (In Japanese.)
- 12 Toba K, Nakai R, Akishita M *et al*. Vitality Index as a useful tool to assess elderly with dementia. *Geriatr Gerontol Int* 2002; **2**: 23–29.
- 13 Mahoney FI, Barthel DW. Functional evaluation: the Barthel index. *Md State Med J* 1965; **14**: 61–65.
- 14 Machida A. Estimation of the reliability and validity of the short form version of the 28-item Dementia Behavior Disturbance scale. *Nippon Ronen Igakkai Zasshi* 2012; **49**: 463–467. (In Japanese, abstract English.)
- 15 Lam LC, Lui VW, Luk DN *et al*. Effectiveness of an individualized functional training program on affective disturbances and functional skills in mild and moderate dementia—a randomized control trial. *Int J Geriatr Psychiatry* 2010; **25**: 133–141.
- 16 Woods B, Aguirre E, Spector AE, Orrell M. Cognitive stimulation to improve cognitive functioning in people with dementia. *Cochrane Database Syst Rev* 2012; (2)CD005562.
- 17 Farias ST, Harrell E, Neumann C, Houtz A. The relationship between neuropsychological performance and daily functioning in individuals with Alzheimer's disease: ecological validity of neuropsychological tests. *Arch Clin Neuropsychol* 2003; **18**: 655–672.
- 18 Germain S, Adam S, Olivier C *et al*. Does cognitive impairment influence burden in caregivers of patients with Alzheimer's disease? *J Alzheimers Dis* 2009; **17**: 105–114.
- 19 Aguirre E, Woods RT, Spector A, Orrell M. Cognitive stimulation for dementia: a systematic review of the evidence of effectiveness from randomised controlled trials. *Ageing Res Rev* 2012; **12**: 253–262.
- 20 Black W, Almeida OP. A systematic review of the association between the Behavioral and Psychological Symptoms of Dementia and burden of care. *Int Psychogeriatr* 2004; **16**: 295–315.
- 21 Miyamoto Y, Tachimori H, Ito H. Formal caregiver burden in dementia: impact of behavioral and psychological symptoms of dementia and activities of daily living. *Geriatr Nurs* 2010; **31**: 246–253.
- 22 Marvardi M, Mattioli P, Spazzafumo L *et al*. The Caregiver Burden Inventory in evaluating the burden of caregivers of elderly demented patients: results from a multicenter study. *Ageing Clin Exp Res* 2005; **17**: 46–53.
- 23 Campbell P, Wright J, Oyeboode J *et al*. Determinants of burden in those who care for someone with dementia. *Int J Geriatr Psychiatry* 2008; **23**: 1078–1085.
- 24 Molyneux GJ, McCarthy GM, McEniff S, Cryan M, Conroy RM. Prevalence and predictors of carer burden and depression in carers of patients referred to an old age psychiatric service. *Int Psychogeriatr* 2008; **20**: 1193–1202.
- 25 Yaffe K, Fox P, Newcomer R *et al*. Patient and caregiver characteristics and nursing home placement in patients with dementia. *JAMA* 2002; **287**: 2090–2097.
- 26 Elmstahl S, Ingvad B, Annerstedt L. Family caregiving in dementia: prediction of caregiver burden 12 months after relocation to group-living care. *Int Psychogeriatr* 1998; **10**: 127–146.
- 27 Gitlin LN, Winter L, Dennis MP *et al*. Targeting and managing behavioral symptoms in individuals with dementia: a randomized trial of a nonpharmacological intervention. *J Am Geriatr Soc* 2010; **58**: 1465–1474.
- 28 Bassuk SS, Glass TA, Berkman LF. Social disengagement and incident cognitive decline in community-dwelling elderly persons. *Ann Intern Med* 1999; **131**: 165–173.
- 29 Fratiglioni L, Wang HX, Ericsson K, Maytan M, Winblad B. Influence of social network on occurrence of dementia: a community-based longitudinal study. *Lancet* 2000; **355**: 1315–1319.
- 30 Wang HX, Karp A, Winblad B, Fratiglioni L. Late-life engagement in social and leisure activities is associated with a decreased risk of dementia: a longitudinal study from the Kungsholmen project. *Am J Epidemiol* 2002; **155**: 1081–1087.
- 31 Naglie G, Hogan DB, Krahn M *et al*. Predictors of patient self-ratings of quality of life in Alzheimer disease: cross-sectional results from the Canadian Alzheimer's Disease Quality of Life Study. *Am J Geriatr Psychiatry* 2011; **19**: 881–890.

ORIGINAL ARTICLE

Twelve-week physical and leisure activity programme improved cognitive function in community-dwelling elderly subjects: a randomized controlled trial

Tadahiko KAMEGAYA,¹ Yumi ARAKI,² Hanami KIGURE,² Long-Term-Care Prevention Team of Maebashi City² and Haruyasu YAMAGUCHI¹

¹Department of Rehabilitation Sciences, Gunma University Graduate School of Health Science, and ²Division of Long-Term Care, Maebashi City Office, Maebashi, Japan

Correspondence: Dr Tadahiko Kamegaya PhD, 3-39-22 Showa-machi, Maebashi, Gunma 371-8514, Japan. Email: kamegaya@health.gunma-u.ac.jp

Received 6 September 2013; revision received 28 October 2013; accepted 10 November 2013.

Abstract

Background: Japan is one of the most rapidly ageing societies in the world. A number of municipalities have started services for the prevention of cognitive decline for community-dwelling elderly individuals, but the effectiveness of these services is currently insufficient. Our study explored the efficacy of a comprehensive intervention programme consisting of physical and leisure activities to prevent cognitive decline in community-dwelling elderly subjects.

Method: We administered a 12-week intervention programme consisting of physical and leisure activities aimed at enhancing participants' motivation to participate and support one another by providing a pleasant atmosphere, empathetic communication, praise, and errorless support. This programme for the prevention of cognitive decline was conducted as a service by the city of Maebashi. All participants underwent the Five-Cog test, which evaluated the cognitive domains of attention, memory, visuospatial function, language, and reasoning. Executive function was evaluated by the Wechsler Digit Symbol Substitution Test and Yamaguchi Kanji-Symbol Substitution Test. Subjective health status, level of social support, functional capacity, subjective quality of life, and depressive symptoms were assessed with a questionnaire. Grip strength test, timed up-and-go test, 5-m maximum walking times test, and functional reach test were performed to evaluate physical function. Fifty-two participants were randomly allocated to intervention ($n = 26$) and control ($n = 26$) groups. Twenty-six participants, aged between 65–87 years, received intervention once a week at a community centre. The programme was conducted by health-care professionals, with the help of senior citizen volunteers.

Results: The intervention group ($n = 19$) showed significant improvement on the analogy task of the Five-Cog test ($F_{1,38} = 4.242$, $P = 0.046$) and improved quality of life ($F_{1,38} = 4.773$, $P = 0.035$) as compared to the control group ($n = 24$).

Conclusion: A community-based 12-week intervention programme that aimed to enhance motivation to participate in activities resulted in improvements in some aspects of cognitive function and quality of life. Senior citizens who volunteered in the present intervention enabled the smooth implementation of the programme and alleviated the burden on professional staff.

Key words: community-dwelling elderly, senior citizen volunteer, service for prevention of cognitive decline.

INTRODUCTION

Japan is one of the most rapidly ageing societies in the world. The number of demented elderly people who need nursing care is predicted to be 3.5 million by 2015 and 4.7 million by 2025.¹ Since Japan's long-term care insurance system was revised in 2008 to emphasize the importance of preventing the need for long-term care, a number of municipalities have started services to help prevent cognitive decline in community-dwelling elderly individuals. The services focus on maintaining and improving the cognitive functions of those who do not need care at present. However, the effectiveness of these services is currently insufficient, and such public services still need to demonstrate their effectiveness in preventing cognitive decline.

Physical exercise intervention has been suggested as an efficient strategy to reduce the risk of cognitive decline. A meta-analysis that focused on elderly subjects with dementia and related cognitive impairments suggested that physical exercise increases fitness, physical function, cognitive function, and positive behaviour.² The results of a more recent meta-analysis showed that non-demented subjects who performed physical activity had a significantly reduced risk of cognitive decline.³ Also, recent randomized intervention studies reported that physical activity helps improve cognitive function of elderly patients.⁴⁻⁶

Participation in leisure activities is associated with a reduced risk of dementia by increasing cognitive reserve.⁷⁻⁹ Frequent participation in cognitively stimulating activities, such as playing games, is associated with a reduced risk of Alzheimer's disease.¹⁰ Previous randomized intervention studies have suggested that cognitively stimulating training or activity improves some aspects of cognitive ability in elderly subjects.^{11,12}

Physical and leisure activities in a pleasant atmosphere can be effective for the prevention of cognitive decline. It has been shown in an animal study that exercise in enriched environments has a suppressive effect on the accumulation of amyloid- β protein.¹³ We have proposed the efficacy of intervention conducted in a pleasant atmosphere with an emphasis on communication.¹⁴ Therefore, it could be meaningful to facilitate a pleasant atmosphere between participants and form a group in which participants enjoy mutual communication.

We performed a randomized controlled trial of a comprehensive intervention programme consisting of physical and leisure activities for prevention of cognitive decline in community-dwelling elderly subjects. The programme was conducted as a service of the city of Maebashi, Japan, and it was administered by the city's staff of health-care professionals along with senior citizen volunteers. In the intervention, a pleasant atmosphere, empathetic communication, praising each other, and errorless support were emphasized to enhance the participants' motivation to participate and support one another according to five principles of brain-activating rehabilitation.¹⁴

Our study explored the efficacy of this service for preventing cognitive decline in elderly residents.

METHOD

Participants

The intervention programme was carried out as a service of the municipality of Maebashi in 2012. The service for the prevention of cognitive decline targeted elderly subjects aged 65 years and older residing in Maebashi. Participants were recruited through a lecture on the prevention of cognitive decline for community residents, leaflets that went to 2986 households, and door-to-door visits by public health nurses, local welfare commissioners, and senior citizen volunteers to invite elderly residents to the programme.

The Medical Ethics Committee of Gunma University (Maebashi, Japan) approved this study (21-47), and written informed consent was obtained from all participants.

Initial screening

Participants ($n = 58$) were screened by a questionnaire and medical interview (Fig. 1). They were examined by a clinician who specialized in dementia. Those who were diagnosed as having dementia according to the criteria of International Statistical Classification of Diseases and Related Health Problems 10th Revision or who had a medical condition that made them unable to engage in physical activity were excluded. After screening and exclusion, 52 participants remained.

Participants who were excluded from the study were still able to participate in the intervention programme, as it was conducted as a community service available to all community dwellers.

Amnesic mild cognitive impairment was diagnosed according to the following criteria:¹⁵ reported memory




Figure 1 Flow of participants through the study.

complaint; objective memory impairment for age; essentially preserved general cognitive function; largely intact functional activities; and not demented. Reported memory complaint was examined based on information provided in the questionnaire administered at the baseline evaluation. Objective memory impairment for age was evaluated based on the score of the three-word delayed recall task of the Mini-Mental State Examination and questions about recent events or news by a clinician. Essentially preserved general cognitive function was evaluated based on the Mini-Mental State Examination score. Whether functional activities were largely intact was evaluated based on the score of Tokyo Metropolitan Institute of Gerontology Index of Competence.

Evaluation

Change in cognitive function was evaluated using the Five-Cog test, which evaluates attention, memory, visuospatial function, language, and reasoning. The Five-Cog test consists of five items: character position referencing task to evaluate attention, category

cued recall task to evaluate memory, clock drawing task to evaluate visuospatial function, animal name listing task to evaluate language ability, and analogy task to evaluate abstract reasoning ability.^{16,17} Participants were also evaluated using the Wechsler Digit Symbol Substitution Test and Yamaguchi Kanji-Symbol Substitution Test,¹⁸ which evaluate executive function.

To evaluate the physical function of each participant, a grip strength test, timed up-and-go test, 5-m maximum walking times test, and functional reach test were performed.

Participants were required to complete a questionnaire consisting of questions regarding age, sex, education, and previous/current medical history. Their subjective health status was evaluated with the question 'How is your health in general?' and scored using a rating scale from 1 (excellent) to 4 (poor). The level of social support was evaluated with the Lubben Social Network Scale Revised, which gauges social isolation in older adults by measuring the perceived social support from family, friends, and neighbours.¹⁹

Functional capacity was determined using the Tokyo Metropolitan Institute of Gerontology Index of Competence,²⁰ which is a multidimensional 13-item index of competence comprising three dimensions: instrumental self-maintenance, intellectual activity, and social role. The index was designed to measure higher level competence in community-dwelling elderly subjects.^{20,21} The Satisfaction in Daily Life, a simple measurement of subjective quality of life (QOL), was used to evaluate the life satisfaction of participants.^{22,23} The Satisfaction in Daily Life consists of 11 items: physical health, mental health, self-care, gait, housework, house facilities, partner and family relationships, hobby and leisure activities, social intercourse, economic state and social security, and having a job. Each item was rated from 1 (dissatisfied) to 5 (satisfied). The 15-item short version of the Geriatric Depression Scale was used to evaluate depressive symptoms.²⁴

Randomization

After receiving the initial screening and baseline evaluation, 52 eligible participants were randomly allocated to the intervention group or control group.

Intervention

A comprehensive intervention programme consisting of physical and leisure activities was designed. The physical activity programme was the primary content of the programme. The exercise programme included muscle-stretching exercise in a sitting position (17 items), muscle-strengthening exercise in a sitting position (3 items), muscle-strengthening exercise in a standing position (7 items), and aerobic exercise (3 items). These exercises require comprehensive abilities in physical activity involving body flexibility, muscle-strength, balance and endurance. The mean duration of the exercise programme was 45 min. Participants were encouraged to perform exercises based on the programme at home. Walking was recommended to participants as a regular exercise. Leisure activities, such as cooking, handcrafts, and competitive games, were included in the programme to stimulate participant's cognitive function. All participants allocated to the intervention group attended a weekly 2-h programme at a community centre located in the district where they resided for 12 weeks.

The programme was presented by health-care professionals, including a physical therapist, occupational therapist, and public health nurse. Each session was conducted by three health-care professionals and three to five senior citizen volunteers. Maebashi has promoted elderly participation in volunteer activities, particularly those focusing on long-term care prevention in the community. In the present intervention, 32 senior citizens participated as volunteers. They received training on brain-activating rehabilitation. In each programme, three to seven volunteers participated in assisting the professional staff. Their roles were to assist the professional staff in conducting the physical and leisure activities, support participants who needed assistance engaging in the programmes, facilitate communication among the participants and help maintain a pleasant atmosphere. Participants in the control group did not attend a programme during this period.

Statistical analysis

Statistical analysis was performed using the Japanese version of SPSS v. 17.0 (IBM, Armonk, NY, USA). For baseline comparison between the intervention and control groups, a χ^2 test was conducted in categorical data, and two-sample *t*-tests were conducted in continuous variables. No significant differences were observed between the two groups at baseline. Changes from baseline were analyzed by repeated measures ANCOVA, with covariance of age, sex, and years of education. Post-hoc analysis was conducted using Bonferroni correction. $P < 0.05$ indicated significance.

RESULTS

Flow of participants through the study

Figure 1 shows the flow of participants through the study. Fifty-two participants were randomized to the intervention ($n = 26$) and control ($n = 26$) groups. The attendance rate during the intervention was 77.6%. Nine were excluded from the analysis: seven in the intervention group attended the programme less than eight times and two in the control group were absent from the second evaluation. Finally, 43 subjects were included in analysis. Table 1 shows the demographic and clinical characteristics of the participants. Changes from baseline were investigated

Table 1 Demographic data of the participants at baseline

Characteristics	Intervention (<i>n</i> = 26)	Control (<i>n</i> = 26)	Total (<i>n</i> = 52)
Age [†] (years)	73.6 ± 5.6	76.2 ± 6.1	74.9 ± 5.9
Female, <i>n</i> (%)	24 (92.3)	23 (88.5)	47 (90.4)
Years of education [†]	11.6 ± 2.0	11.2 ± 2.4	11.2 ± 2.2
MMSE score [†]	27.6 ± 2.0	27.9 ± 1.6	27.7 ± 1.8
aMCI, <i>n</i> (%)	8 (30.8)	8 (30.8)	16 (30.8)

[†]Results are expressed as mean ± SD. aMCI, amnesic mild cognitive impairment; MMSE, Mini-Mental State Examination.

Table 2 Results of the test scores (*n* = 43)

Scale	Intervention group		Control group		Interaction		Post-hoc analysis,	
	Before [†]	After [†]	Before [†]	After [†]	F value	<i>P</i> -value	Intervention	Control
Cognition								
Character position referencing task	18.8 ± 7.9	21.7 ± 8.1	20.0 ± 9.0	22.4 ± 8.4	0.001	0.970	0.029	0.017
Cued recall task	13.6 ± 4.6	16.5 ± 5.5	12.7 ± 4.3	15.8 ± 4.4	0.370	0.547	<0.001	<0.001
Clock drawing task	6.9 ± 0.3	6.7 ± 0.7	6.8 ± 0.5	6.7 ± 0.6	1.102	0.300	0.114	0.833
Animal name listing task	14.7 ± 3.2	16.1 ± 3.5	14.9 ± 3.6	14.8 ± 3.3	2.999	0.91	0.047	0.735
Analogy task	9.4 ± 3.0	11.0 ± 3.1	9.7 ± 3.7	10.3 ± 2.5	4.242	0.046	<0.001	0.147
WDSST	45.0 ± 13.5	52.7 ± 15.0	46.6 ± 14.4	51.5 ± 18.5	1.165	0.287	<0.001	0.005
YKSST	43.9 ± 10.6	45.2 ± 15.0	42.5 ± 11.7	44.2 ± 13.7	0.096	0.759	0.542	0.249
Questionnaire								
Subjective health status	2.1 ± 0.5	2.0 ± 0.4	2.0 ± 3.0	2.1 ± 0.4	2.142	0.152	0.374	0.225
TMIG-IC	12.3 ± 1.0	12.2 ± 1.0	12.5 ± 0.9	12.3 ± 1.4	0.040	0.842	0.567	0.343
LSNS-R	20.1 ± 4.9	20.7 ± 5.4	17.9 ± 4.4	16.7 ± 4.9	1.824	0.185	0.572	0.160
SDL	43.5 ± 6.3	45.4 ± 5.3	45.5 ± 5.5	44.9 ± 6.0	4.773	0.035	0.029	0.428
GDS	3.2 ± 3.5	2.6 ± 3.0	2.0 ± 2.3	2.1 ± 1.8	0.840	0.365	0.289	0.848
Motor								
Grip strength	24.4 ± 6.3	24.8 ± 5.8	22.7 ± 5.3	24.2 ± 4.5	2.564	0.118	0.626	0.005
Timed up-and-go test	7.4 ± 1.0	7.0 ± 0.7	7.8 ± 2.0	8.0 ± 2.2	1.632	0.210	0.256	0.518
5-m maximum walking time	2.6 ± 0.4	2.5 ± 0.3	2.8 ± 0.8	3.1 ± 1.2	1.778	0.191	0.817	0.083
Functional reach test	34.9 ± 5.2	35.6 ± 4.4	35.0 ± 5.5	35.5 ± 4.6	0.001	0.982	0.681	0.675

[†]Results are expressed as mean ± SD. GDS, 15-item short version of the Geriatric Depression Scale; LSNS-R, Lubben Social Network Scale Revised; SDL, Satisfaction in Daily Life; TMIG-IC, Tokyo Metropolitan Institute of Gerontology Index of Competence; WDSST, Wechsler Digit Symbol Substitution Test; YKSST, Yamaguchi Kanji-Symbol Substitution Test.

with 19 participants in the intervention group and 24 in the control group (Fig. 1).

Analysis of all the participants

The intervention group had a significant increase in the score on the Five-Cog test's analogy task relative to the control group ($F_{1,38} = 4.242$, $P = 0.046$). Post-hoc analysis among the subjects showed a significant improvement in the intervention group ($P < 0.01$), but not in the control group ($P = 0.147$) (Table 2).

The score on the Five-Cog test's animal name listing task was significantly increased in the intervention group ($P = 0.047$), but not in the control group ($P = 0.735$). However, the interaction between the two groups was marginal ($F_{1,38} = 2.999$, $P = 0.091$).

With regard to the scores on the character position referencing task, category cued recall task and

Wechsler Digit Symbol Substitution Test, the results of the post-hoc analysis indicated a significant increase in both the intervention and control groups, but the interaction was not significant. There were no significant differences in the scores on the other cognitive tests.

Significant interaction was shown with regard to QOL in the intervention group ($F_{1,38} = 4.773$, $P = 0.035$). Post-hoc analysis showed significant improvement in the intervention group ($P = 0.029$), but not in the control group ($P = 0.428$) (Table 2). No significant differences were observed in other items investigated by the questionnaires, including subjective health status, social support, functional capacity and depressive symptoms.

None of the items on the physical function tests showed significant changes in the intervention group.

DISCUSSION

This study explored the efficacy of a comprehensive intervention programme for the prevention of cognitive decline in community-dwelling elderly subjects. The programme consisted of physical and leisure activities aimed at enhancing motivation to participate in activities, and it was run as a municipal service. Effects were investigated using cognitive, physical, functional, social, and behavioural outcome measures. Through the 12-week intervention, participants showed improvement in some aspects of cognitive function and QOL. The Five-Cog test's analogy task evaluated abstract reasoning ability, which tends to decline with incipient dementia, and showed significant improvement in the intervention group.²⁵ The benefit observed in this study could be the result of physical activity and leisure activities associated with the five principles of brain-activating rehabilitation and the support of senior citizen volunteers.¹⁴ Furthermore, the benefit could have resulted from the pleasant atmosphere that emphasized interactive communication. Throughout the programme, the staff and the volunteers were expected to facilitate communication among the participants and maintain a pleasant atmosphere. It is possible that a comprehensive intervention programme conducted in a pleasant atmosphere with interactive communication enhanced motivation, leading to improved cognitive function.^{26,27} The benefit shown in QOL may represent a sense of satisfaction and participation in the community among the participants. With regard to the scores on the character position referencing task, category cued recall task, and Wechsler Digit Symbol Substitution Test, the results of the post-hoc analysis indicated significant improvement in both the intervention and control groups. Participants in both groups were tested twice on these items, and the score of the second trial may have increased based on experience from the first test. This tendency might have reduced the interaction effect between the two groups.

Municipal programmes to prevent cognitive decline among community-dwelling elderly have not been widely established yet. Providing appropriate and effective programmes to prevent cognitive decline is a critical issue. We administered a 12-week comprehensive intervention programme for prevention of cognitive decline. Evidence suggests that physical

activity reduces the risk of cognitive decline among non-demented elderly subjects.³ Therefore, physical activity should be an aspect of any programme for preventing cognitive decline. Furthermore, it may be useful to offer physical activity programmes as a community service, as they are labour-saving and cost-effective. Leisure activities, such as handcrafts or competitive games, are effective in facilitating communication among participants and maintaining a pleasant atmosphere. These activities are also easily administered, similar to those within physical activity programmes.

This study's period of the intervention was relatively short. Previous randomized controlled trials of physical activity or cognitive training for prevention of cognitive decline examined longer periods of 6–12 months.^{6,11,12} The 12-week duration of the preventive programme was determined by the long-term care insurance system in Japan, and it was a feasible period as a service of a local government. The present study showed that a short-term 12-week intervention improved some aspects of cognitive function and QOL.

Use of senior citizen volunteers was emphasized in our intervention. Involvement of citizen volunteers could be effective for a community-based intervention programme for the prevention of cognitive decline.²⁶ It is important to develop human resources who can promote preventive care programmes throughout the community, as a shortage of professional staff can occur in an ageing society. Senior citizen volunteers who joined in the present intervention played important roles; they enabled smooth implementation of the programme and alleviated the burden on professional staff. In addition, they facilitated a pleasant atmosphere and easy communication among participants.

The present study has several limitations. The number of participants was small; the present study targeted 52 participants, but only 43 participants were included in analysis due to absence from the intervention and evaluation. The preponderance of female patients (90.4%) is a limitation of our study. The participants of this study were self-selected; thus, they may have been more health-conscious than the general elderly population, and greater efficacy on cognition and other functions could be expected. The relatively short intervention period was established in accordance with the long-term care insurance programme, and the intervention was carried out for just

3 months. A longer-term follow-up of the participants should be conducted to ensure lasting positive effects. These factors may limit the ability to generalize the results of the study.

In conclusion, to prevent cognitive decline, participants took part in a comprehensive 12-week intervention programme consisting of physical and leisure activities. The programme was associated with the five principles of brain-activating rehabilitation that enhanced motivation. Participants showed improvement in some aspects of cognitive function and QOL. Senior citizen volunteers enabled smooth implementation of the programme and alleviated the burden on professional staff. Thus, the present study demonstrates the potential of a community-based intervention programme to prevent cognitive decline.

ACKNOWLEDGMENT

The authors thank Rumi Shinohara and Yuko Tsunoda (Gunma University, Maebashi, Japan) for technical assistance. We would like to express our deep appreciation to all of the participants and senior citizen volunteers.

This study was supported in part by a Grant-in-Aid for Scientific Research (H22-Ninchisho-Ippan-004, H25-Ninchisho-Ippan-008) from the Ministry of Health, Labour and Welfare of Japan. Haruyasu Yamaguchi received support from a Grant-in-Aid for Scientific Research from Japan's Ministry of Education, Science, Sports, Culture and Technology (23300197 and 22650123).

Tadahiko Kamegaya prepared the manuscript. The intervention programme described in the present study was designed by Yumi Araki. Hanami Kigure, a physical therapist and public health nurse of the Long-Term-Care Prevention Team of Maebashi City, and Haruyasu Yamaguchi had final approval of the manuscript.

REFERENCES

- Ministry of Health, Labour and Welfare, Japan. The number of demented elderly people in Japan. [Cited 24 Aug 2012.] Available from URL: <http://www.mhlw.go.jp/stf/houdou/2r9852000002iau1.html>
- Heyn P, Abreu BC, Ottenbacher KJ. The effects of exercise training on elderly persons with cognitive impairment and dementia: a meta-analysis. *Arch Phys Med Rehabil* 2004; **85**: 1694–1704.
- Sofi F, Valecchi D, Bacci D *et al*. Physical activity and risk of cognitive decline: a meta-analysis of prospective studies. *J Intern Med* 2010; **269**: 107–117.
- Maki Y, Ura C, Yamaguchi T *et al*. Effects of intervention using a community-based walking program for prevention of mental decline: a randomized controlled trial. *J Am Geriatr Soc* 2012; **60**: 505–510.
- Erickson KI, Voss MW, Prakash RS *et al*. Exercise training increases size of hippocampus and improves memory. *Proc Natl Acad Sci U S A* 2011; **108**: 3017–3022.
- Lautenschlager NT, Cox KL, Flicker L *et al*. Effect of physical activity on cognitive function in older adults at risk for Alzheimer disease: a randomized trial. *JAMA* 2008; **300**: 1027–1037.
- Fabrigoule C, Letenneur L, Dartigues JF, Zarrouk M, Commenges D, Barberger-Gateau P. Social and leisure activities and risk of dementia: a prospective longitudinal study. *J Am Geriatr Soc* 1995; **43**: 485–490.
- Scarmeas N, Levy G, Tang MX, Manly J, Stern Y. Influence of leisure activity on the incidence of Alzheimer's disease. *Neurology* 2001; **57**: 2236–2242.
- Verghese J, Lipton RB, Katz MJ *et al*. Leisure activities and the risk of dementia in the elderly. *N Engl J Med* 2003; **348**: 2508–2516.
- Wilson RS, Mendes De Leon CF, Barnes LL *et al*. Participation in cognitively stimulating activities and risk of incident Alzheimer disease. *JAMA* 2002; **287**: 742–748.
- Ball K, Berch DB, Helmers KF *et al*. Effects of cognitive training interventions with older adults: a randomized controlled trial. *JAMA* 2002; **288**: 2271–2281.
- Olazaran J, Muniz R, Reisberg B *et al*. Benefits of cognitive-motor intervention in MCI and mild to moderate Alzheimer disease. *Neurology* 2004; **63**: 2348–2353.
- Lazarov O, Robinson J, Tang YP *et al*. Environmental enrichment reduces Abeta levels and amyloid deposition in transgenic mice. *Cell* 2005; **120**: 701–713.
- Yamaguchi H, Maki Y, Yamagami T. Overview of non-pharmacological intervention for dementia and principles of brain-activating rehabilitation. *Psychogeriatrics* 2010; **10**: 206–213.
- Petersen RC. Mild cognitive impairment as a diagnostic entity. *J Intern Med* 2004; **256**: 183–194.
- Sasaki M, Kodama C, Hidaka S *et al*. Prevalence of four subtypes of mild cognitive impairment and APOE in a Japanese community. *Int J Geriatr Psychiatry* 2009; **24**: 1119–1126.
- Miyamoto M, Kodama C, Kinoshita T *et al*. Dementia and mild cognitive impairment among non-responders to a community survey. *J Clin Neurosci* 2009; **16**: 270–276.
- Yamaguchi T, Maki Y, Kaiho A *et al*. Yamaguchi Kanji Symbol Substitution Test as a scale of executive function in old people for preventing dementia in community. *Jap J Geriatr Psychiatry* 2011; **22**: 587–594 (in Japanese with English abstract).
- Lubben J, Blozik E, Gillmann G *et al*. Performance of an abbreviated version of the Lubben Social Network Scale among three European community-dwelling older adult populations. *Gerontologist* 2006; **46**: 503–513.
- Koyano W, Shibata H, Nakazato K, Haga H, Suyama Y. Measurement of competence: reliability and validity of the TMIG Index of Competence. *Arch Gerontol Geriatr* 1991; **13**: 103–116.
- Fujiwara Y, Shinkai S, Kumagai S. Changes in TMIG-Index of Competence by subscale in Japanese urban and rural community older populations: six years prospective study. *Geriatr Gerontol Int* 2003; **3**: 63–68.

- 22 Hachisuka K, Nagayoshi M, Iwata N. The examination on similarity and difference of in daily life satisfaction SDL and SF-36 measurement concept. *Sumon ni kansuru Chosa Kenkyuhan Heisei 14 Nendo Sokatsu, Buntan Kenkyu Hokokusho* 2003; 133–135 (in Japanese with English abstract).
- 23 Nagayoshi M, Iwata N, Hachisuka K. Factors associated with life satisfaction in Japanese stroke outpatients. *Disabil Rehabil* 2008; **30**: 222–230.
- 24 Yesavage JA, Brink TL, Rose TL *et al.* Development and validation of a geriatric depression screening scale: a preliminary report. *J Psychiatr Res* 1982; **17**: 37–49.
- 25 Ritchie K, Artero S, Touchon J. Classification criteria for mild cognitive impairment: a population-based validation study. *Neurology* 2001; **56**: 37–42.
- 26 Kamegaya T, Long-Term-Care Prevention Team of Maebashi City, Maki Y *et al.* Pleasant physical exercise program for prevention of cognitive decline in community-dwelling elderly with subjective memory complaints. *Geriatr Gerontol Int* 2012; **12**: 673–679.
- 27 Yamaguchi H, Maki Y, Takahashi K. Rehabilitation for dementia using enjoyable video-sports games. *Int Psychogeriatr* 2011; **23**: 674–676.

Biochemical stages of amyloid- β peptide aggregation and accumulation in the human brain and their association with symptomatic and pathologically preclinical Alzheimer's disease

Ajeet Rijal Upadhaya,¹ Irina Kosterin,¹ Sathish Kumar,² Christine A. F. von Arnim,³ Haruyasu Yamaguchi,⁴ Marcus Fändrich,⁵ Jochen Walter² and Dietmar Rudolf Thal¹

1 Laboratory of Neuropathology, Institute of Pathology, Centre for Clinical Research at the University of Ulm, Ulm, Germany

2 Department of Neurology, University of Bonn, Bonn, Germany

3 Department of Neurology, University of Ulm, Ulm, Germany

4 Gunma University School of Health Sciences, Maebashi, Gunma, Japan

5 Institute for Pharmaceutical Biotechnology, University of Ulm, Ulm, Germany

Correspondence to: Dietmar R. Thal,
Laboratory of Neuropathology,
Institute of Pathology,
Centre for Clinical Research at the University of Ulm,
Helmholtzstrasse 8/1,
D-89081 Ulm
E-mail: dietmar.thal@uni-ulm.de

Correspondence may also be addressed to: Jochen Walter, Department of Neurology, University of Bonn, Sigmund Freud Strasse 25, D-53105 Bonn, Germany. E-mail: Jochen.Walter@ukb.uni-bonn.de

Alzheimer's disease is characterized by the deposition of amyloid- β peptide in the brain. N-terminal truncation resulting in the formation of $A\beta_{N3pE}$ and phosphorylation at serine 8 have been reported to modify aggregation properties of amyloid- β . Biochemically, soluble, dispersible, membrane-associated, and insoluble, plaque-associated amyloid- β aggregates have been distinguished. Soluble and dispersible amyloid- β aggregates are both in mixture with the extracellular or intracellular fluid but dispersible aggregates can be cleared from proteins in solution by ultracentrifugation. To clarify the role of phosphorylated amyloid- β and $A\beta_{N3pE}$ in soluble, dispersible, membrane-associated, and plaque-associated amyloid- β aggregates in the pathogenesis of Alzheimer's disease we studied brains from 21 cases with symptomatic Alzheimer's disease, 33 pathologically preclinical Alzheimer's disease cases, and 20 control cases. Western blot analysis showed that soluble, dispersible, membrane-associated and plaque-associated amyloid- β aggregates in the earliest preclinical stage of Alzheimer's disease did not exhibit detectable amounts of $A\beta_{N3pE}$ and phosphorylated amyloid- β . This stage was referred to as biochemical stage 1 of amyloid- β aggregation and accumulation. In biochemical amyloid- β stage 2, $A\beta_{N3pE}$ was additionally found whereas phosphorylated amyloid- β was restricted to biochemical amyloid- β stage 3, the last stage of amyloid- β aggregation. Phosphorylated amyloid- β was seen in the dispersible, membrane-associated, and plaque-associated fraction. All cases with symptomatic Alzheimer's disease in our sample fulfilled biochemical amyloid- β stage 3 criteria, i.e. detection of phosphorylated amyloid- β . Most, but not all, cases with pathologically preclinical Alzheimer's disease had biochemical amyloid- β stages 1 or 2. Immunohistochemistry confirmed the hierarchical occurrence of amyloid- β , $A\beta_{N3pE}$, and phosphorylated amyloid- β in amyloid plaques. Phosphorylated amyloid- β containing plaques were, thereby, seen in all symptomatic cases with Alzheimer's disease but only in a few non-demented control subjects. The biochemical amyloid- β stages correlated with the expansion of amyloid- β

Received July 9, 2013. Revised October 16, 2013. Accepted November 2, 2013.

© The Author (2014). Published by Oxford University Press on behalf of the Guarantors of Brain. All rights reserved.

For Permissions, please email: journals.permissions@oup.com

plaque deposition and with that of neurofibrillary tangle pathology. Taken together, we demonstrate that $A\beta_{N3pE}$ and phosphorylated amyloid- β are not only detectable in plaques, but also in soluble and dispersible amyloid- β aggregates outside of plaques. They occur in a hierarchical sequence that allows the distinction of three stages. In light of our findings, it is tempting to speculate that this hierarchical, biochemical sequence of amyloid- β aggregation and accumulation is related to disease progression and may be relevant for an increasing toxicity of amyloid- β aggregates.

Keywords: amyloid- β protein; phosphorylation; N-terminal truncation; pyroglutamate formation; soluble fraction; dispersible fraction

Abbreviations: $A\beta$ = amyloid- β ; CERAD = Consortium to Establish a Registry for Alzheimer's Disease; CSF = Cerebrospinal fluid; SDS = Sodium dodecyl sulfate

Introduction

Alzheimer's disease is characterized by the formation of amyloid- β ($A\beta$) protein aggregates in the brain (Alzheimer, 1907; Masters *et al.*, 1985; Hyman *et al.*, 2012). Various types of $A\beta$ aggregates have been described: soluble $A\beta$ oligomers, dispersible $A\beta$ oligomers, protofibrils and fibrils, membrane-associated (SDS soluble) $A\beta$ aggregates, and solid, plaque-associated (formic acid soluble) $A\beta$ fibrils (Harper *et al.*, 1997; Walsh *et al.*, 1997; Kaye *et al.*, 2003; Lesne *et al.*, 2006; Habicht *et al.*, 2007; Shankar *et al.*, 2008; Rijal Upadhaya *et al.*, 2012a). Recently, we showed that dispersible $A\beta$ oligomers, protofibrils, and fibrils are pathologically relevant forms of $A\beta$ aggregates that cause neurotoxic effects in a concentration-dependent manner as demonstrated in APP transgenic mouse models (Rijal Upadhaya *et al.*, 2012a). Dispersible $A\beta$ aggregates represent diffusible but non-soluble $A\beta$ aggregates that differ from insoluble membrane-associated and plaque-associated $A\beta$ aggregates. In contrast to plaque-associated and membrane-associated $A\beta$, dispersible $A\beta$ is thought to represent not fully dissolved $A\beta$ aggregates mixed with the extra- or intracellular fluid whereas membrane-associated and plaque-associated $A\beta$ are not mixed with the extracellular fluid and become detectable only after SDS or formic acid treatment. Dispersible $A\beta$ can be separated from soluble $A\beta$ by ultracentrifugation (Rijal Upadhaya *et al.*, 2012a). Currently, it is not clear whether dispersible $A\beta$ aggregates play a role in the human Alzheimer's disease brain.

Post-translational N-terminal truncation of the first two amino acids of the $A\beta$ peptide and subsequent pyroglutamate formation at the N-terminal glutamate resulting in $A\beta_{N3pE}$ (Saido *et al.*, 1995) as well as phosphorylation of serine 8 of $A\beta$ (Kumar *et al.*, 2011) have been described. Both $A\beta_{N3pE}$ and phosphorylated $A\beta$ occur in Alzheimer's disease plaques (Saido *et al.*, 1995; Kumar *et al.*, 2011). $A\beta_{N3pE}$ increases the aggregation propensity of $A\beta$ by changing the biophysical properties of $A\beta$ fibrils (Schlenzig *et al.*, 2009). Phosphorylated $A\beta$, on the other hand, promotes the formation of $A\beta$ oligomers that serve as nucleation sites for fibril formation (Kumar *et al.*, 2011). Here, it is suggested that these modifications of $A\beta$ play a role in Alzheimer's disease pathogenesis and the development of dementia, but it is not clear at which stage of the disease and in which types of aggregates these modifications may become evident.

Alzheimer's disease begins many years before the cognitive deficits become evident that allow its clinical diagnosis. The recently revised criteria for the clinical diagnosis of Alzheimer's disease introduced

preclinical Alzheimer's disease as a diagnosis for non-demented individuals with positive biomarkers for Alzheimer's disease, e.g. Alzheimer's disease-like Pittsburgh compound B retention in the brain (Dubois *et al.*, 2007; Sperling *et al.*, 2011). Current neuropathological guidelines for the assessment of Alzheimer's disease pathology recommend the description of the level of Alzheimer's disease pathology in a given brain regardless of the ante-mortem cognitive status (Hyman *et al.*, 2012). As such, by employing the neuropathological diagnosis of Alzheimer's disease pathology as a biomarker for Alzheimer's disease, pathologically preclinical Alzheimer's disease cases are those that were non-demented before death, but exhibit at least low levels of Alzheimer's disease pathology at autopsy, whereas symptomatic Alzheimer's disease cases exhibit significant Alzheimer's disease pathology and cognitive impairment (Monsell *et al.*, 2013). Non-Alzheimer's disease cases are those without any $A\beta$ plaques (Hyman *et al.*, 2012). The term 'pathologically preclinical Alzheimer's disease' for non-demented cases with amyloid plaques does not necessarily mean that these cases must convert into symptomatic Alzheimer's disease. It cannot be excluded that progression of pathology may cease in some of these cases although they fulfil the recommended criteria for low levels of Alzheimer's disease pathology. Whether the biochemical composition of cortical $A\beta$ aggregates has impact on the course of the disease and the development of dementia is not yet clear. Clinical criteria for preclinical Alzheimer's disease based on CSF- $A\beta$ and CSF-tau protein levels have been published (Vos *et al.*, 2013) that differ from the neuropathological definition of pathologically preclinical Alzheimer's disease that is used here, synonymous with the term 'asymptomatic Alzheimer's disease' (Monsell *et al.*, 2013).

To clarify the role of phosphorylation of $A\beta$ and $A\beta_{N3pE}$ formation for the clinicopathological stage of the disease and for the pattern of $A\beta$ aggregate types, we studied brains from 21 patients with Alzheimer's disease, 33 with pathologically preclinical Alzheimer's disease, and 20 non-demented control cases immunohistochemically and biochemically.

Materials and methods

Neuropathology

For morphological analysis, autopsy brains from 21 patients with Alzheimer's disease, 20 control cases without any $A\beta$ -pathology and 33 cases with pathologically preclinical Alzheimer's disease were used