

いる。Naaman ら³⁾が行った、全病期の乳癌患者を対象とした心理的介入の効果に関するシステマティック・レビューでは、不安の軽減(効果量: -0.40 , 95%CI: -0.72 – -0.08), 抑うつ軽減(効果量: -1.01 , 95%CI: -1.48 – -0.54), QOLの改善(効果量: 0.74 , 95%CI: 0.12 – 1.37)においていずれも中程度の効果が認められたが、さらなる臨床試験での追試が推奨されている。Fors ら⁴⁾は、1999~2008年に報告された18件(心理教育6件, 認知行動療法7件, 心理社会的サポート5件)のランダム化比較試験をレビューしたが、QOLや感情状態の改善において、いずれも短期的な効果であったと報告している。Taylor ら⁵⁾らは、心理社会的介入の性的問題に対する介入効果を報告した15件のランダム化比較試験をレビューし、心理教育の限定的な効果を報告しているが、介入効果を明らかにするためには、方法論的により洗練された研究が必要であると結論付けている。Anderson ら⁶⁾が行った、227人の術後乳癌を対象とした心理的介入(週1回4カ月間+月1回8カ月の小グループによる心理・行動学的介入)の効果に関するランダム化比較試験によると、介入群では再発(HR: 0.55 , 95%CI: 0.32 – 0.95 , $p=0.034$), 乳癌死亡(HR: 0.44 , 95%CI: 0.22 – 0.86 , $p=0.016$), 全死亡(HR: 0.51 , 95%CI: 0.28 – 0.93 , $p=0.028$)ともそのリスクが減少していた。本研究は、術後の早期乳癌患者を対象に心理社会的介入の生存期間に対する効果をみた最初の報告であり、今後の再現性の検証が必要である。今回採用した2件のシステマティック・レビューは、マインドフルネスに基づいたストレス軽減法とヨガに関する効果をみたものである。前者⁷⁾は9件(ランダム化比較試験は2件のみ)の研究をレビューし、ストレスに対する有効性(effect size: 0.71 , 95%CI: 0.51 – 0.90)を示しているものの、さらなる良質な研究の必要性を示唆している。後者⁸⁾は6件のランダム化比較試験をレビューしたが、不安や抑うつといった心理的な効果について有意な結果は得られなかった。わが国からの優れた報告もある。Fukui ら⁹⁾による、46人の日本の原発乳癌患者を対象とした心理社会的介入(6週の構造化された健康教育, 対処技術訓練, ストレス管理, 心理的サポート)のランダム化比較試験によると、介入群では6週目に活力と癌との前向きな取り組み(コーピング)において、対照群との比較で有意に高いスコアを示し、6カ月目まで効果は持続した。しかしこれ以降、わが国から心理社会的介入の有効性を評価したランダム化比較試験は報告されていない。

以上より、転移乳癌患者に対する心理社会的介入の効果は、心理的アウトカムの一部でしかなく、また持続期間も短期にとどまると結論付けられるため、推奨するための根拠は不十分である。また、原発乳癌患者対象の研究においても、効果は一部の指標で認められるものの、再現性に乏しく、評価指標の多重性の問題や対象症例数を考慮すると、エビデンスレベルの高いものは少数にとどまる。

心理社会的介入論文の限界は大きく分けて3つある。1つは、介入プログラムの信頼性・妥当性に関する問題であり、信頼性・妥当性に関する記述が不明確なものが多い。2つ目は評価指標の選択基準である。患者の負担を考慮すると、最小限の、確立された尺度を用いることが望ましいが、その点の記載が不十分な文献が多い。3つ目は症例数設定根拠と解析手法、結果の多重性の扱いである。症例数の設定根拠に触れている論文は極めて少ない。また、複数の評価指標を同時に用いる研究が多いが、プライマリー・エンドポイントの設定が明確でなく、どれかで有意差が出ればよしとする姿勢が多々見受けられる。そのために、エビデンスレベルを1段階下げざるを得なかった論文が多かった。

検索式

検索はPubMedにて、Breast Neoplasms/therapy, Psychotherapy, Social Support, psychosocial interventionのキーワードを用いて行った。検索期間は2012年9月～2014年9月とした。最終的に選択された海外文献該当102件より、タイトルと抄録により、CQに関連がないものや、明らかに抄録や総説とわかる論文、対象が乳癌に限らないものを除き44件選択、原論文査読により、エビデンスレベルが2c以下のもの、特殊性の強いもの、評価指標に心理的側面や生存が含まれていないものなどの文献15件を除外し、ランダム化比較試験26件とシステマティック・レビュー3件を抽出した。エビデンスの評価にはこの中からシステマティック・レビュー3件のみを採用し、結果、2013年版と併せた9件を本文の解説に引用した。国内文献に採択すべき論文はなかった。

参考文献

- 1) Mustafa M, Carson-Stevens A, Gillespie D, Edwards AG. Psychological interventions for women with metastatic breast cancer. *Cochrane Database Syst Rev.* 2013; 6: CD004253.
- 2) Chow E, Tsao MN, Harth T. Does psychosocial intervention improve survival in cancer? A meta-analysis. *Palliat Med.* 2004; 18(1): 25-31.
- 3) Naaman SC, Radwan K, Fergusson D, Johnson S. Status of psychological trials in breast cancer patients: a report of three meta-analyses. *Psychiatry.* 2009; 72(1): 50-69.
- 4) Fors EA, Bertheussen GF, Thune I, Juvet LK, Elvsaaas IK, Oldervoll L, et al. Psychosocial interventions as part of breast cancer rehabilitation programs? Results from a systematic review. *Psychooncology.* 2011; 20(9): 909-18.
- 5) Taylor S, Harley C, Ziegler L, Brown J, Velikova G. Interventions for sexual problems following treatment for breast cancer: a systematic review. *Breast Cancer Res Treat.* 2011; 130(3): 711-24.
- 6) Andersen BL, Yang HC, Farrar WB, Golden-Kreutz DM, Emery CF, Thornton LM, et al. Psychologic intervention improves survival for breast cancer patients: a randomized clinical trial. *Cancer.* 2008; 113(12): 3450-8.
- 7) Zainal NZ, Booth S, Huppert FA. The efficacy of mindfulness-based stress reduction on mental health of breast cancer patients: a meta-analysis. *Psychooncology.* 2013; 22(7): 1457-65.
- 8) Zhang J, Yang KH, Tian JH, Wang CM. Effects of yoga on psychologic function and quality of life in women with breast cancer: a meta-analysis of randomized controlled trials. *J Altern Complement Med.* 2012; 18(11): 994-1002.
- 9) Fukui S, Kugaya A, Okamura H, Kamiya M, Koike M, Nakanishi T, et al. A psychosocial group intervention for Japanese women with primary breast carcinoma. *Cancer.* 2000; 89(5): 1026-36.

Effectiveness of collage activity based on a life review in elderly cancer patients: A preliminary study

SATOKO SAKAGUCHI M.S., O.T.R.,¹ AND HITOSHI OKAMURA, PH.D., M.D.²

¹Graduate School of Health Sciences, Hiroshima University, Hiroshima, Japan

²Institute of Biomedical and Health Sciences, Hiroshima University, Hiroshima, Japan

(RECEIVED August 27, 2013; ACCEPTED January 12, 2014)

ABSTRACT

Objective: Much of the cancer rehabilitation research that has been conducted has consisted of relatively early recovery-of-function rehabilitation, and little attention has been paid to the psychosocial aspects of palliative rehabilitation. The aim of the present preliminary study was to examine the “narratives” of elderly cancer patients that emerged as a result of a life review performed in association with collage activity and to assess the effectiveness of this intervention.

Method: We conducted a collage activity based on a life review in two sessions. Some 11 cancer patients who were 65 years of age or older and receiving palliative care participated.

Evaluations using the Functional Assessment of Chronic Illness Therapy–Spiritual (FACIT–Sp) Scale, the Hospital Anxiety and Depression Scale (HADS), and the Self-Efficacy Scale for Terminal Cancer (SESTC) were administered before and immediately after the intervention.

Results: The mean scores for the FACIT–Sp and affect regulation efficacy on the SESTC significantly increased, while the mean HADS score significantly decreased. Regarding the impressions after completion of the intervention, generally favorable evaluations were heard from families and medical staff members as well as from the subjects.

Significance of results: A collage activity based on a life review may be effective for improving spiritual well-being, mitigating anxiety and depression, and improving self-efficacy. The collage itself was also useful in facilitating interactions with others, including family members, and the activity provided psychological support for families.

KEYWORDS: Collage activity, Elderly, Life review, Palliative care, Occupational therapy

INTRODUCTION

The use of rehabilitation to improve the quality of life (QoL) of cancer patients has attracted interest in recent years. However, much of the cancer rehabilitation research that has been conducted has consisted of relatively early recovery-of-function rehabilitation, and little research on maintenance or palliative rehabilitation has been performed. Hamaguchi and coworkers (2008) conducted a fact-finding survey examining cancer rehabilitation at 1045 institutions in Japan and reported that physical function training

accounted for a high proportion of cancer rehabilitation efforts, while little attention was directed toward spiritual or psychological suffering. While some research has certainly shown improvement in activities of daily living (ADL) in response to rehabilitation, decreases in function and reduced ADL are inevitable as cancer progresses. Moreover, the prevalence of depression appears to be especially elevated in patients with advanced cancer (Caplette-Gingras & Savard, 2008). Thus, attending to the psychological aspects of cancer is an important task.

In light of this situation, the importance of rehabilitation approaches based on psychosocial aspects aimed at maintaining and improving cancer patients' QoL is now attracting attention (Ronson, 2002). One of these approaches, a life review, is a type of

Address correspondence and reprint requests to: Hitoshi Okamura, Graduate School of Health Sciences, Hiroshima University, 1-2-3 Kasumi, Minami-ku, Hiroshima 734-8551, Japan. E-mail: hokamura@hiroshima-u.ac.jp.

reminiscence method, which was developed by Butler (1963) as a psychological support technique. This activity has been shown to be effective in relation to a patient's degree of life satisfaction, psychological well-being, depressed mood, and self-esteem (Hanaoka & Okamura, 2004). The reminiscence method has been shown to be a promising intervention method in cancer patients. However, there are relatively few studies concerning the reminiscence method for cancer patients. Wholihan (1992) demonstrated the effectiveness of utilizing a patient's photo albums or treasured objects. In Japan, Ando et al. (2007a,b) conducted several studies examining topics and programs for application of the reminiscence method in cancer patients. Out of consideration for the physical burden on patients, Ando et al. (2008) also developed a brief reminiscence method that can be completed in two interviews, and showed that the spiritual well-being, anxiety, and depression of terminal-stage cancer patients improved as a result of its application (Ando et al., 2010a,b). However, because the therapists produced the albums in this brief reminiscence method, no benefits were gained from enabling patients to leave behind a product that they themselves had created or from the experience of working through an actual activity. Regarding the benefits of having the subjects themselves participate in the activities, Chochinov et al. (2002a,b; 2005; 2011) demonstrated the effect of dignity therapy on distress and quality of life in terminally ill patients. Dignity therapy involves two or three shorter sessions. Patients review their lives with the aid of routine questions, and the session is recorded, edited, and transcribed. Hall and colleagues (2013) indicated that the most frequently perceived benefits of dignity therapy related to reminiscence, and that dignity therapy gave patients the opportunity to recall happy memories they could share with their families.

"Collage," on the other hand, affords the experience of an activity that has been employed widely in occupational therapy. "Collage" is a French term that means "pasting." It is produced by cutting existing images out of magazines, pamphlets, and the like, using scissors, rearranging the images on a posterboard, and then pasting the images onto the board. Collage came into being as an artistic technique in the early 1900s. It has the special characteristic of stimulating self-expression, and it is said to be capable of mitigating anxiety and conflicts experienced by patients, eliciting a positive mood and emotions (Aida et al., 2010). In the area of cancer care, Williams (2002) mentioned the use of collage in end-of-life care, and Forzoni (2010) recently investigated and reported on the use of collage in cancer patients. Another study reported the use of a combi-

nation of collage and the reminiscence method in elderly subjects. However, that study did not focus on the effectiveness of the activity, but rather centered on the reminiscence method itself.

Life review usually consists mainly of "talks" generated by the review. The subjects may check albums and other works prepared by therapists, but they do not create works using their own hands and bodies, and their intentions are not reflected in their works. However, it has been pointed out that patients' "feelings" are often sublimated by expressing them in "things (works and activities)." Therefore, we thought we might be able to expect changes in the subjects by having them create works using a collage along with life review, which could increase one's sense of self-affirmation and is effective for spiritual pain, and creative activities, which are said to promote self-expression, thereby leading to affirmative feelings and emotions, and by preserving the works.

The aim of the present study was to examine the "narrative" of elderly cancer patients that emerged as a result of a life review involving the production of a collage and to assess the effectiveness of this activity in relation to the psychosocial and spiritual aspects of their lives, as well as its impact on their experiences.

MATERIALS AND METHODS

Our study was conducted with the approval of the ethics committee of the Graduate School of Health Sciences of Hiroshima University.

Participants

The subjects of our study included patients receiving home-visit nursing care, patients in the palliative care unit of a rehabilitation hospital who were receiving palliative care, and patients visiting a clinic who were receiving palliative care at home. All the subjects met the following eligibility criteria: (1) elderly cancer patient aged 65 years or over, (2) capable of understanding the nature of the study and the contents of the question sheets (no cognitive problems), and (3) consent of the attending physician allowing the patient's participation in the study.

Intervention Method

Ando and colleagues (2008) previously developed a brief reminiscence method that can be completed in two sessions and suggested that the method was effective for improving both the spiritual and psychosocial aspects of a patient's condition. The content of the interviews is mainly based on a structured life review but does not parallel developmental stages. Out of consideration for the physical burden that the




Fig. 1. Example of a finished collage product. (Upper left)

[Memories of my husband]

Many times I invited my husband for a trip although he did not like traveling, and I and my husband had many trips.

I was rather quiet.

But if I wanted to do something, I did it immediately.

I wanted to go and see places I had never been before.

I also had strong curiosity.

It was fun visiting Keelung, Taiwan, glaciers, New Zealand, etc.

I felt more lonely because of my husband's death. It was painful beyond description.

(Upper right)

[When I look back]

We experience various things in our lives.

But I have done what I wanted to do.

I am the happiest person.

And children, grandchildren, great grandchildren...

A family made by my husband and myself is expanding and expanding.

I have been really happy.

(Lower left)

I have boarded fishing boats since I was in my teens, and I have worked hard.

It was a matter of course to send money to my parents.

I did all the sewing and washing with my hands.

I prepared my things by myself.

The flow of time we had to choose.

I went south to New Zealand and north to the Bering Sea.

(Lower right)

My first job was toilet cleaning and cooking.

An era when there was no water and little electricity.

Rice was washed with seawater, and rice with a lot of debris was filtered with gauze and boiled.

The rice had no taste at all. Embattled by the weather.

And, finally, we got a pile of king crab.

intervention method would impose on the elderly subjects in our study, we configured the sessions based on the following brief reminiscence method. Before the intervention, we conducted an orientation in which we introduced ourselves, explained the nature of the study, and confirmed the patient's consent to participate in the study. We then conducted the first life review session as follows.

(Content of the Interview)

- What do you think is most important in life?
- What memory in life has left the greatest impression?
- What was the turning point in your life? If there was an event or person that influenced you. Would you mind telling me what or who that was?
- What sort of role do you feel you have played in life?
- Is there something in your life that you feel proud of?
- If you were to express life in a single word, what word would express it best?
- Were any words or advice ever said to you by a person who was important to you?

During the week after the first interview, the authors summarized the spoken content using a word-for-word record and divided it into paragraphs, and then collected images from magazines, the Internet, and the library that reflected the spoken content. During the second session, while matching photographs that the subject had to the images that had been collected, an independent collage activity was performed with the help of the therapist, and the selected images and photographs were arranged in a single booklet (see Figure 1).

Measures

Sociodemographic and Medical Variables

Age, sex, family makeup, current residence, disease site, performance status (PS), and whether or not the subject was experiencing pain were assessed as sociodemographic and medical information.

ADL: Barthel Index

The Barthel Index was created in 1965 by Mahoney and Barthel (1965). Ratings are made based on 10 items (Ito & Kamakura, 1994), and the maximum score is 100. A special feature of this index is that the scores for each item are weighted according to the time and amount of nursing care required.

QoL: Functional Assessment of Chronic Illness Therapy–Spiritual (FACIT–Sp)

The FACIT–Sp is part of a self-report that was developed by Cella and colleagues (1993). It consists of 12 items and is composed of 2 main factors: Meaning of Life/Peace and Faith. The Japanese-language version was developed by Shimotsuma (2002), and its reliability and validity have been confirmed (Noguchi et al., 2004).

Anxiety and Depression: Hospital Anxiety and Depression Scale (HADS)

The HADS is a self-rated scale developed by Zigmond and Snaith (1983) to measure anxiety and depression. It consists of 14 items, and possible total scores for anxiety and depression each range from 0 to 21. The reliability and validity of the Japanese-language version (Kitamura, 1993) were confirmed by Kugaya et al. (1998).

Self-Efficacy: Self-Efficacy Scale for Terminal Cancer

The Self-Efficacy Scale for Terminal Cancer (SESTC) was devised by Hirai and coworkers (2002) and is composed of three subscales: “Affect Regulation Efficacy” (ARE), “Symptom Coping Efficacy” (SCE), and “ADL Efficacy” (ADE). There are 18 items in the full scale. Higher scores are considered to indicate greater self-efficacy.

Procedure

Physicians and nurses gave an oral explanation of the nature of the activity in advance to patients who were receiving palliative care between July 11 and December 28, 2011, and who met the eligibility criteria. The first author then visited the home or room of the patients who consented to participate and provided a written explanation of the purpose and methods of the study to each patient.

Scheduling of interviews was selected so as not to interfere with the conduct of patients' treatment or care, and the place selected was the bedside, day room, or rehabilitation room. If the patient had difficulty filling out the self-reported question sheets because of a physical problem, the author read the questions aloud to the subject and then the author and subject filled out the sheets together.

The evaluation forms were collected before the first session, and the first life review interview was then conducted. The content of the interview was recorded using an audio recorder. Whenever necessary, information was also collected from the patient's chart. The second session was held one week later.

At the end of the second session, the subject was asked to fill out the evaluation sheets again, and the evaluations were again collected.

Analysis

Statistical Analysis

After confirming the normality of the data, the changes in FACIT–Sp, HADS, and SESTC scores before and after the collage activity were analyzed using a paired *t* test.

Summaries of Impressions

After completing the collage, the impressions of the subjects, families, and medical staff members regarding the intervention were obtained and summarized separately. The *p* values for all the tests were two-tailed, and a *p* value less than 0.05 was considered significant. The Statistical Package for Social Science (SPSS, version 17.0 J for Windows) was employed to perform all statistical analyses.

RESULTS

Subjects' Participation and Background of the Participants

Some 15 patients met the eligibility criteria during the study period, and consent was obtained from 13 patients. The reasons why consent was not obtained from the other two patients included “I’m not good at details,” “It’s a lot of trouble,” “I feel reluctant to talk to a stranger,” and “I don’t want to think about it.” The reasons given by one patient who initially consented but later refused prior to the actual performance of the intervention were “I don’t want to talk about the terrible past” and “Doing this in the form of research would be too restricting for me.” During the collage activity, one patient read the life story that the therapist had summarized from the narrative and then reminisced extensively. Upon trying to give them the memories form, however, the patient looked at the album and gave up on the collage activity, saying, “I could never finish in just one day.” Thus, the intervention was ultimately completed in 11 subjects. The time allotted to conduct each session of the intervention in this study was about one hour. However, because in many cases it took much longer, the mean length of the first session was 84.5 ± 20.0 minutes, and the mean length of the second session was 93.6 ± 38.6 minutes. None of the subjects subsequently complained of not feeling well.

Table 1. Background data for the 11 subjects who completed the intervention

	Case 1	Case 2	Case 3	Case 4	Case 5	Case 6	Case 7	Case 8	Case 9	Case 10	Case 11
Age	73	72	70	82	85	67	76	74	80	65	87
Gender	Female	Male	Female	Male	Male	Male	Male	Male	Male	Female	Female
Current residence	Home	Hospital	Hospital	Home	Hospital	Hospital	Home	Home	Hospital	Hospital	Hospital
Marital status	Married	Married	Unmarried	Married	Married	Married	Married	Married	Married	Married	Married
Cancer site	Thyroid	Lung	Ovary	Lung	Colon	Lung	Liver	Stomach	Gallbladder	Lung	Lymphoma
Performance status	2	1	1	3	2	4	2	4	2	3	2
Pain	Absence	Absence	Presence	Presence	Presence	Presence	Presence	Presence	Absence	Presence	Absence
Barthel Index	100	100	100	85	95	45	85	45	100	85	80
FACIT–Sp	36	25	27	21	33	28	20	8	26	24	37
HADS	7	18	7	10	4	16	11	23	19	7	6
SESTC	54	101	139	113	150	91	133	87	107	89	113
Length of first session (min)	105	93	125	55	88	86	80	91	80	68	59
Length of second session (min)	113	100	133	65	96	33	140	125	128	44	53

FACIT–Sp: Functional Assessment of Chronic Illness Therapy–Spiritual; HADS: Hospital Anxiety and Depression Scale; SESTC: Self-Efficacy Scale for Terminal Cancer.

The background data on the 11 subjects who completed the intervention (7 males, 4 females; mean age 75.54 ± 7.2 years), their baseline FACIT–Sp, HADS, and SESTC scores, and the time required to complete sessions are presented in Table 1.

Changes in Evaluation Scale Scores

An examination of the changes in FACIT–Sp scores before and after the intervention showed that the mean score for Meaning of Life/Peace changed from 17.0 ± 5.3 to 22.8 ± 5.3 ($p = 0.004$), the mean score for Faith changed from 8.6 ± 3.2 to 12.0 ± 3.0 ($p = 0.001$), and the mean total score changed from 25.9 ± 8.1 to 34.9 ± 17.5 ($p = 0.002$).

An examination of the changes in HADS scores before and after the intervention showed that the mean score for anxiety changed from 5.6 ± 4.6 to 2.8 ± 2.0 ($p = 0.034$), the mean score for depression changed from 6.0 ± 2.4 to 3.6 ± 2.5 ($p = 0.042$), and the mean total score changed from 11.6 ± 6.3 to 6.4 ± 3.7 ($p = 0.026$).

Table 2. Impressions of subjects

In relation to the intervention:

- Now, I feel that doing this was really worthwhile.
- My gloomy feelings have lifted.
- For the first time in a long time I became completely absorbed, and I put all my energy into making it.
- I feel that this activity serves as a link between those who are living and those who will die.
- While I was making this, I completely forgot about my illness.

In relation to the life review:

- I nostalgically recalled things in the past.
- There has never been an opportunity to think so deeply and to try to recall the past.
- I was able to look back at how I had lived my life.
- I can't bring myself to say so to my son, but I came to feel grateful to him.

In relation to the product:

- I'm glad that I was able to make something that contains so much of my feelings.
- I think it will be a treasure at home.
- This is the best present we could give to our family in the condition we are in now.
- Making it as the two of us consulted with each other was fun.
- Making things is fun.

Table 3. Impressions of family members and medical staff

Impressions of family members:

- I think this is what mental rehabilitation is.
- I was moved.
- The patient was pleased, saying, "I'm glad I made it."
- It's something that's full of the patient.
- There were so many thoughts in this home!

Impressions of medical staff:

- It was good to see the patient speaking happily.
- The family was happy, too.
- I realized that there was another side of the patient.
- I really understood what it was that the patient cared about.
- It is difficult to explain collages to patients.

An examination of the changes in SESTC scores before and after the intervention showed that, while the mean score for ARE increased significantly from 30.6 ± 16.0 to 39.0 ± 11.5 ($p = 0.040$), no significant differences in mean scores for SCE ($p = 0.967$) or ADE ($p = 0.208$) or total mean scores ($p = 0.284$) were observed.

Impressions of the Intervention

Some of the summations of the impressions of the subjects, families, and medical staff members after completion of the intervention are given in Tables 2 and 3, where one can see that generally favorable evaluations were heard from family members and medical staff as well as from subjects. However, several participants had not grasped the full picture before the start of the collage activity, and this limitation is also listed as an impression.

DISCUSSION

As a result of performing the collage activity based on a life review, the FACIT–Sp scores of the 11 subjects in this study increased significantly, while their HADS scores decreased significantly. These results suggest that the intervention can be effective in increasing the spiritual well-being of patients, similar to the findings of previous studies (Ando et al., 2007a,b; 2008; 2010a,b). Our results also suggest that the intervention may be simultaneously effective for mitigating anxiety and depression. Moreover, the score for the ARE subscale of the SESTC also

increased significantly. Hirai and colleagues (2002) regarded improvements in the scores for this subscale to be a primary factor with direct effects on anxiety and depression. The level of self-efficacy has also been widely reported (Ewart, 1995; Beckham et al., 1997) to affect psychological adjustment, as corroborated by the results of our present study. Bandura (1977) explained self-efficacy as a subjective determination that "I am capable of doing this," and the "successful experience" of "even I can do that" was remarked upon among the impressions expressed after the intervention. It is suggested that this boost in confidence felt by subjects was linked to the increase in scores for this scale.

By contrast, no significant changes in feelings of efficacy in relation to physical symptoms or ADL were seen. This finding regarding feelings of efficacy as they relate to physical symptoms may have been attributable to the fact that none of the patients in this study complained of cancer pain severe enough to require the intervention to be stopped before completion, and symptom control in these patients was relatively good. Many participants had sufficient stamina to concentrate and spend time performing the life review and collage activity beyond the time that had been scheduled. The mean Barthel Index score before the intervention was 83.6 ± 20.5 , and the fact that subjects' original ADL capacity was relatively high might have been related to the results for efficacy in relation to the ADL. In particular, after excluding the subjects with a PS of four, the ADL of the nine remaining subjects was high, and they were able to perform basic body movements and their own body care movements independently.

According to the impressions of the subjects after completion of the intervention, their memories and past experiences were recalled, and most of the subjects spoke animatedly, saying that the intervention created feelings of nostalgia. Moreover, while looking at the finished product, some subjects evaluated the product highly, saying that it was a "treasure" or their "best present," and some said that by becoming engrossed in the collage activity they had forgotten about their illness, if only for a short time, and that they had been able to spend time in a meaningful way. These impressions were thought to have been affected by an improvement in the subjects' state of mind as a result of the self-affirmation and self-esteem created by the collage activity, during which the subjects enjoyed expressing themselves. Moreover, since the impression that "this activity serves as a link between those who are alive with those who will die" was also expressed, discussions of the continuity of relationships and one's own impermanence might have contributed to the spiritual care that is involved in relationships.

Among the impressions of the families who participated in the collage activity together with the subject, there were descriptions of the intervention as a "spiritual rehabilitation," and the family members thought favorably of the intervention, saying that the subjects themselves had enjoyed the activity. In addition, subjects' family members stated that they discovered new aspects of the subjects through this activity and that this activity was an opportunity for them to renew their knowledge of the subjects' feelings. It is said that people do not often express their feelings to others in Japan as compared to the situation in other countries (Benedict, 2006). Therefore, this activity seems to be more deeply significant as a tool with which subjects and their families communicate each other's feelings. As an approach to the family, Tajiri and colleagues (2006) pointed out that creating the product by working with and drawing closer to the patient is linked to a sense of achievement, and that in many cases the feeling of satisfaction of having been able to do something for the patient had an effect with regard to their psychological care. Thus, this activity may have also provided psychological care for the family as well. The results may be a possible avenue for future research. Furthermore, impressions were heard from the medical staff members that they had discovered new aspects of the subjects through the finished product, and that they had reconfirmed that it was something the subjects considered important. Moreover, the product was shown to not only link the subject and family as a means of communication, but also to have served as a topic for interaction between the staff and the subject.

This study has some limitations. The first is that, although only self-reported question sheets were used to evaluate the results of an intervention, whenever it was difficult for the subject to fill out the sheets because of physical problems the author read the questions aloud to the subject, and sometimes they read through the questions and filled out the sheets together. As a result, the evaluation methods sometimes varied according to the evaluator, and several biases may have occurred as a result. Second, because a control group was not included, the results may not be attributable to only the collage activity. The use of a control group and a controlled study design is needed for future studies. Third, no psychosocial changes over time were demonstrated after the product was completed. A detailed follow-up in addition to evaluations before and after performing the collage activity are needed. Finally, we configured the sessions based on the brief reminiscence method to avoid taxing patients. However, the possibility of adequately reviewing a life in such a short period of time is not clear. It should be better

to have a greater number of smaller interviews where more adequate information could be gathered without overtaxing patients.

CONCLUSIONS

The possibility of objective and subjective effectiveness of a collage activity based on a life review was demonstrated in elderly cancer patients. Our results suggest that collage activity can be effective in improving spiritual well-being, mitigating anxiety and depression, and improving self-efficacy. In addition, in terms of the impact of the work activity on subjects and their family members, the product was shown to be useful as a tool for interacting with others, including family members, and that the activity was capable of providing psychological care to family members who performed the activity together with the subject. Future studies are needed where the activity is actually put into practice so that the link between a collage activity based on a previous life review and an improvement in the quality of life of patients and their families can be clarified as a useful psychosocial approach to treatment of elderly cancer patients.

REFERENCES

- Aida, Y., Okamura, T., Takenouchi, S., et al. (2010). The effects of intervention using collage in long-stay psychiatric inpatient groups [in Japanese]. *Journal of Japanese Occupational Therapy Association*, 29, 60–72a.
- Ando, M., Tsuda, A. & Morita, T. (2007a). Life review interviews on the spiritual well-being of terminally ill cancer patients. *Supportive Care in Cancer*, 15, 225–231.
- Ando, M., Morita, T. & O'Connor, S.J. (2007b). Primary concerns of advanced cancer patients identified through the structured life review process: A qualitative study using a text mining technique. *Palliative & Supportive Care*, 5, 265–271.
- Ando, M., Morita, T., Okamoto, T., et al. (2008). One-week short-term life review interview can improve spiritual well-being of terminally ill cancer patients. *Psycho-Oncology*, 17, 885–890.
- Ando, M., Morita, T., Akechi, T., et al. (2010a). Japanese Task Force for Spiritual Care: Efficacy of short-term life-review interviews on the spiritual well-being of terminally ill cancer patients. *Journal of Pain and Symptom Management*, 39, 993–1002.
- Ando, M., Morita, T., Miyashita, M., et al. (2010b). Effects of bereavement life review on spiritual well-being and depression. *Journal of Pain and Symptom Management*, 40, 453–459.
- Bandura, A. (1977). Self-efficacy: Toward a unifying theory of behavior change. *Psychological Review*, 84, 195–215.
- Beckham, J.C., Burker, E.J. & Lytle, B.L. (1997). Self-efficacy and adjustment in cancer patients: A preliminary report. *Behavioral Medicine*, 23, 138–142.
- Benedict, R. (2006). *Chrysanthemum and the sword: Patterns of Japanese culture*. Boston: Mariner Books.
- Butler, R.N. (1963). The life review: An interpretation of reminiscence in the aged. *Psychiatry*, 26, 65–76.
- Caplette-Gingras, A. & Savard, J. (2008). Depression in women with metastatic breast cancer: A review of the literature. *Palliative & Supportive Care*, 6, 377–387.
- Cella, D., Tulsky, D.S. & Gray, G. (1993). The Functional Assessment of Cancer Therapy (FACT) Scale: Development and validation of the general measure. *Journal of Clinical Oncology*, 11, 1407–1418.
- Chochinov, H.M., Hack, T., Hassard, T., et al. (2002a). Dignity in the terminally ill: A cross-sectional, cohort study. *Lancet*, 360, 2026–2030.
- Chochinov, H.M., Hack, T., McClement, S., et al. (2002b). Dignity in the terminally ill: A developing empirical model. *Social Science & Medicine*, 54, 433–443.
- Chochinov, H.M., Hack, T., Hassard, T., et al. (2005). Dignity therapy: A novel psychotherapeutic intervention for patients near the end of life. *Journal of Clinical Oncology*, 23, 5520–5525.
- Chochinov, H.M., Kristjanson, L.J., Breitbart, W., et al. (2011). Effect of dignity therapy on distress and end-of-life experience in terminally ill patients: A randomised controlled trial. *Lancet Oncology*, 12, 753–762.
- Ewart, C.K. (1995). Self-efficacy and recovery from heart attack: Implication for a social cognitive analysis of exercise and emotion. In *Self-efficacy, adaptation, and adjustment: Theory, research, and application*. J.E. Maddux (ed.), pp. 203–226. New York: Plenum Press.
- Forzoni, S. (2010). Art therapy with cancer patients during chemotherapy sessions: An analysis of the patients' perception of helpfulness. *Palliative & Supportive Care*, 8, 41–48.
- Hall, S., Goddard, C., Speck, P.W., et al. (2013). "It makes you feel that somebody is out there caring": A qualitative study of intervention and control participants' perceptions of the benefits of taking part in an evaluation of dignity therapy for people with advanced cancer. *Journal of Pain and Symptom Management*, 45, 712–725.
- Hamaguchi, T., Okamura, H., Nakaya, N., et al. (2008). Survey of the current status of cancer rehabilitation in Japan. *Disability and Rehabilitation*, 21, 1–6.
- Hanaoka, H. & Okamura, H. (2004). Study on effects of life review activities on the quality of life of the elderly: A randomized controlled trial. *Psychotherapy and Psychosomatics*, 73, 302–311.
- Hirai, K., Suzuki, Y. & Tsuneto, A. (2002). Development of the self-efficacy scale for terminally ill cancer patients [in Japanese]. *Japanese Journal of Psychosomatic Medicine*, 41, 19–27.
- Ito, T. & Kamakura, N. (1994). *ADL and its related fields* [in Japanese]. Tokyo: Igaku-shoin.
- Kitamura, T. (1993). Hospital Anxiety and Depression Scale [in Japanese]. *Archives of Psychiatric Diagnosis and Clinical Evaluation*, 4, 371–372.
- Kugaya, A., Akechi, T., Okuyama, T., et al. (1998). Screening for psychological distress in Japanese cancer patients. *Japanese Journal of Clinical Oncology*, 28, 333–338.
- Mahoney, F.I. & Barthel, D.W. (1965). Functional evaluation: The Barthel Index. *Maryland State Medical Journal*, 14, 61–65.
- Noguchi, W., Ohno, T., Morita, S., et al. (2004). Reliability and validity of the Functional Assessment of Chronic Illness Therapy–Spiritual (FACIT–Sp) Japanese Version in Cancer Patients [in Japanese]. *Japanese Journal of General Hospital Psychiatry*, 16, 42–47.