

表1 機種の違いによる抗ウイルス剤の回収率

単位 %

農薬名	試験部位	メーカー名				
		Thermo	waters	Agilent	AB	島津
アマンタジン	A	12.0	194.8	96.0	104.0	94.4
	D	26.0	104.8	84.0	96.0	114.6
	E	294.0	99.6	78.0	72.0	130.0
リマンタジン	A	48.8	220.0	71.2	100.5	47.2
	D	32.5	105.6	72.8	37.4	80.4
	E	25.7	110.3	75.4	51.6	94.9
オセルタミビル	A	200.0	221.6	76.0	184.0	200.0
	D	98.0	106.0	76.0	74.0	100.0
	E	100.0	105.6	84.0	88.0	42.0
ザナミビル	A	117.8	212.4	85.0	87.6	83.8
	D	104.4	107.5	90.6	87.4	63.5
	E	105.5	107.4	75.1	82.1	57.0
アシクロビル	A	182.4	237.0	69.6	94.4	104.0
	D	99.2	105.2	79.6	104.0	96.8
	E	99.2	106.7	70.8	100.0	96.0
アルビドール	A	86.6	80.9	69.6	104.0	49.7
	D	105.0	79.8	73.4	70.6	76.3
	E	111.8	75.1	77.7	76.6	76.3
イミキモド	A	-	95.6	112.0	32.0	44.0
	D	114.0	88.0	142.0	20.0	78.0
	E	118.0	66.4	142.0	16.0	104.0
ベラミビル	A	32.0	53.2	76.0	104.0	48.0
	D	32.0	62.4	52.0	46.0	92.0
	E	18.0	50.0	46.0	52.0	122.0
ラニナミビル	A	-	-	13.9	31.7	56.4
	D	-	105.5	4.4	3.1	95.9
	E	-	144.8	5.5	3.2	95.6
リバビリン	A	176.0	205.6	112.0	-	100.0
	D	99.0	102.4	122.0	-	132.0
	E	104.0	83.8	128.0	-	180.0

添加量： A:鶏筋肉(0.01 mg/kg) B:鶏筋肉(0.1 mg/kg) C:鶏卵(0.1 mg/kg)