

The environment itself can be a source of unintentional skin exposure to nanoparticles. Environmental nanoparticles in urban areas arise primarily from combustion sources such as burning coal, fuel oil, and biomass; and waste and motor vehicle emissions typically account for the greatest proportion of environmental nanoparticles [27, 28]. Comprising only a small proportion of ambient air particles by mass but a large proportion in terms of number [29], airborne nanoparticles are usually present as aggregates (diameter, ~ 100 nm) of very small (diameter, ≤ 10 nm) primary nanoparticles [28]. Although information regarding the health effects of environmental exposure is limited to effects after inhalation [30], skin exposure to environmental nanoparticles occurs in the same contexts in which they are inhaled.

Recent reports acknowledge that naturally occurring nanoparticles represent a previously unrecognized opportunity for exposure [31]. For example, various metal objects, including earrings and wire, spontaneously generate metal nanoparticles [32], and laundering silver-embedded textiles releases silver nanoparticles [33]. These naturally occurring metal nanoparticles are thought to arise through the nucleation of metal ions released due to chemical or photochemical reduction [32, 34].

Together, these findings suggest that we are exposed to nanoparticles via the skin merely as a consequence of our daily lives—for example, whenever we wear metal accessories or dress in clothes containing or washed with metal-embellished fabric. In addition, nanoparticles abound in nature. Various soils are naturally rich in nanoparticles, and earthquakes generate massive quantities of new soil-derived nanoparticles via mechanical grinding [31]. Although unavoidable, exposure to nanoparticles via the airways and skin is not a new problem but a long-standing facet of everyday life, at least in terms of naturally occurring nanoparticles.

6.2.2 *Skin Penetration by Nanoparticles*

We know little regarding how naturally occurring nanoparticles or those emitted during industrial processes enter and cross the skin. However, the intense research done to confirm the safety of nanomaterials has taught us about the skin-penetrating characteristics of nanoparticles.

6.2.2.1 **Titanium Dioxide and Zinc Oxide**

Primary nanoparticles of titanium dioxide and zinc oxide are 10–20 nm in diameter, but they typically exist as 30- to 150-nm aggregates and frequently are used in cosmetics [35]. In a minipig model of human skin, inductively coupled plasma mass spectroscopy failed to detect any significant increase in the titanium concentration in the dermis or at draining lymph nodes after 22 days of sunscreen application [36]; in the same study, transmission electron microscopy (TEM) of the dermis revealed only a few titanium particles, equivalent to 10^{-6} to 10^{-4} % of the total amount applied. In another study, multiphoton microscopy of human skin *in vivo* 4 h after the application

of zinc oxide nanoparticles revealed zinc oxide nanoparticles in the stratum corneum but none in viable epidermis [37]; a similar quantitative analysis was unable to document any noteworthy penetration of the skin by titanium nanoparticles [38]. Therefore, even though aggregates of titanium dioxide nanoparticles and zinc oxide nanoparticles penetrate the skin in some situations, the rate is extremely low or below the limit of detection of most modern methods of quantitation.


6.2.2.2 Silica Nanoparticles

Manufactured synthetic silica nanoparticles are amorphous silica and are divided into two main types: pyrogenic silica and wet-process silica (i.e., precipitated silica and silica gels) [20, 39]. All pyrogenic silica nanoparticles exist as aggregates and agglomerates; some of the wet-process types also are well dispersed. Pyrogenic silica nanoparticles are widely used in cosmetics, but available data regarding the skin penetration of silica nanoparticles largely derive from the wet-process type, which is used for coatings and ink-receptive papers and as a filter aid in food production. In a mouse model involving epicutaneous application (“skin painting”) for 3 days or 28 days, TEM analysis revealed well-dispersed silica nanoparticles (diameter, 70 nm) not only in epidermal Langerhans cells but also in the dermis and draining lymph nodes [40, 41], but the images obtained suggested that the penetration rate was minimal, at most. Fluorescence microscopy and flow cytometry showed that fluorescently labeled silica nanoparticles (diameter, 42 nm) enter keratinocytes and Langerhans cells in human tape-stripped epidermis; the authors stated that tape stripping removed part of the stratum corneum and contents of the follicular infundibulum [42]. Although quantitative data regarding skin penetration by silica nanoparticles is currently unavailable, well-dispersed silica nanoparticles likely will enter viable skin at some point during our daily lives.

6.2.2.3 Quantum Dot Nanoparticles

Quantum dot nanoparticles are intrinsically fluorescent, and their dispersion characteristics are easily manipulated by modifying the particle surface; consequently, they are widely used as models to investigate the biodistribution of nanoparticles. In an early study, confocal scanning microscopy suggested that two types of quantum dot nanoparticles with different coatings (PEG [polyethylene glycol]-carboxylic acid and PEG-amine; diameter, 15 to 45 nm) penetrated the epidermis of pigs [43]; however in a follow-up study using TEM and inductively coupled plasma-optical emission spectroscopy, this same group was unable to detect the entry of a similar nanoparticle preparation (40 nm with PEG modification) [44]. In contrast, our TEM analysis revealed that 40-nm quantum dot nanoparticles penetrated slightly into viable skin in mice [40], and confocal microscopy and TEM both demonstrated very low, but qualitatively higher, penetration of quantum dot nanoparticles in UVR-exposed skin compared with intact skin in mice [45]. The findings of another group suggested that, among the preparations and pH conditions they evaluated,

Fig. 6.1 Skin exposure to nanoparticles



only pegylated quantum dot nanoparticles at pH 8.3 penetrated intact human skin [46]. These interesting, but contradictory, observations and quantitative data suggest that, like other nanoparticles, well-dispersed quantum dot nanoparticles can enter our bodies through skin, but the amount that enters is minimal.

The mechanism underlying the penetration of skin by nanoparticles remains unclear. Skin presents two physical barriers: the stratum corneum and tight junctions (TJs). To be absorbed through the skin, a medicine must be less than 500 Da in molecular weight [47]; that is, a compound exceeding 500 Da cannot cross the TJ barrier. Therefore all nanoparticles are thought to be too large to penetrate the TJ barrier, at least when it is fully functional. However, nanoparticles may be able to elude the TJs in hair follicles [48], whose heterogeneously differentiated epithelial cells and their various functions may impart some flexibility in regard to penetration [49]. In addition, Langerhans cells can uptake external antigens despite the presence of an intact TJ barrier [50], thus representing another possible mechanism for the skin penetration of nanoparticles.

The skin is the body's primary defense against the environment and thus is under constant assault from mechanical irritation, microbial pathogens, and chemical insults. Therefore, although the skin barrier is quite strong, it is not always completely protective. For example, tape stripping [51], mechanical flexion [52], and UV exposure [45] all increase nanoparticle penetration in skin. Therefore, in reality, nanoparticles likely gain access into our body through skin that is damaged during the activities of daily living. However, we want to reiterate that the rate of any possible penetration is extremely low, given that state-of-the-art quantitative techniques are unable to detect significant numbers of penetrated particles (Fig. 6.1).

6.3 Potential Hazards of Skin Exposure to Nanoparticles

The potential hazards associated with nanoparticles are divided into two main categories according to the context in which the exposure occurs.

6.3.1 Potential Direct Health Effects of Nanoparticles

Whether nanoparticles are directly inflammatory and cytotoxic currently is the main topic of debate regarding their health risk [53–56]. However, because the rate at which nanoparticles enter through skin is extremely low, we think that they are unlikely to directly cause significant inflammation and cytotoxicity in most situations in skin. Therefore, we now focus on the potential for nanoparticles to lead to mutation and sensitization, adverse effects that are sometimes induced even at relatively low-level skin exposure to a chemical substance.

Many reports suggest that nanoparticles are mutagenic [57–60], but almost all of these studies indicate that the mechanisms underlying the observed mutagenicity were ROS dependent [57–60]. ROS are not “evil” in and of themselves; instead the duration and intensity of exposure determine whether their health effects are harmful or beneficial [61, 62]. For example, high doses of the herbicides, paraquat and juglone, shorten the life span of *Caenorhabditis elegans* by inducing ROS, whereas low doses extend the organism’s life span [63, 64]. However, because only very low numbers of nanoparticles enter the body via the skin, the mutagenicity of nanoparticles is unlikely to occur through ROS-dependent mechanisms. Additional research is warranted to address potential mechanisms for the mutagenicity of nanoparticles and the conditions that enhance their entry or accumulation.

Whether nanoparticles act as directly as sensitizing agents has not been determined. A local lymph node assay, which identifies chemical sensitizers by their capacity to induce the proliferation of cells from draining lymph nodes after dermal exposure [65], was unable to detect any sensitizing ability associated with amine-modified polystyrene nanoparticles (diameter, 50 nm) or titanium dioxide primary nanoparticles (diameter, <25 nm) [66]. Another study evaluated the sensitizing ability of two types of wet-process silica nanoparticles, mesoporous silica and colloidal silica, both of which were about 100 nm in diameter [67]. After three consecutive days of skin painting, mesoporous silica but not colloidal silica increased the ear thickness of mice, prompting the authors to conclude that mesoporous silica nanoparticles acted as a chemical sensitizer. However, the changes in ear thickness were slight, and unlike the positive control (2,4-dinitrofluorobenzene, a strong chemical sensitizer), the mesoporous silica nanoparticles did not elicit any cell proliferation in the local lymph node assay [67].

To date, the only nanoparticle that is confirmed to be antigenic is fullerene: the immunization of mice with complete Freund’s adjuvant and a C₆₀ fullerene derivative conjugated to bovine thyroglobulin successfully induced C₆₀ fullerene-

specific antibodies [68, 69]. However, evaluating the sensitizing ability based on OECD Guideline 406, Ema et al. however could not detect any sensitizability of C₆₀ fullerene [69]. These results suggest that C₆₀ fullerene must be haptenized to induce specific antibody. However, such haptenization seems unlikely under natural conditions, implying that C₆₀ fullerene would be safe in terms of sensitizing ability, according to our current knowledge [70]. Considering all of these findings together, we conclude that the notion that nanoparticles act as chemical sensitizers after skin exposure has not been confirmed.

In contrast, several epidemiologic studies indicate that airborne particulates containing sensitizing metals contribute to the onset of metal allergy [71–73]. Considering metal nanoparticles release metal ions, a supposed cause of metal allergy, metal nanoparticles may act as sensitizers, not because of their own antigenicity, but because of the antigenicity of their released metal ions. Future safety evaluations of nanoparticles need to address this potential function of nanoparticles as indirect sensitizing agents.

6.3.2 Combined Exposure to Nanoparticles and Other Substances

Exposure to nanoparticles often occurs simultaneously with exposure to other chemical compounds and environmental allergens [30]. However, little is known about the hazards of combined skin exposure to nanoparticles and other substances. As mentioned earlier, exposure to environmental nanoparticles facilitates the onset and severity of allergic diseases. Because the most important exposure route for these effects is unknown, the possibility that concurrent skin exposure to nanoparticles and allergen contributes to the onset of allergy needs to be considered.

We first focus on allergic contact dermatitis, which typically is induced by a chemical sensitizer. In one study, injection of titanium dioxide nanoparticles (diameter, 15–25 nm) was done before 2,4-dinitrochlorobenzene-induced expansion of cells from the draining lymph nodes of mice [74]. Furthermore, 3 consecutive days of skin painting with a mixture of mesoporous silica nanoparticles and 2,4-dinitrofluorobenzene induced more severe ear swelling in mice than 2,4-dinitrofluorobenzene alone did [67]. Furthermore, the fragrances in cosmetics, which frequently contain nanoparticles as well, are a leading cause of allergic contact dermatitis [75]. Additional studies to reveal the mechanisms of these nanoparticle-associated effects and to identify the threshold amount for an adverse response are needed urgently.

Atopic allergies are IgE-related allergic conditions, such as atopic dermatitis. As mentioned earlier, exposure to environmental nanoparticles is one of the most important factors in the induction or aggravation of atopic allergy. Yanagisawa et al. found that intradermal injection of mixture of mite allergen, which is the

major cause of atopic dermatitis [76], and titanium dioxide nanoparticles or polystyrene nanoparticles aggravated atopic dermatitis-like skin lesions in the NC/Nga mouse model [77, 78]. We confirmed that intradermal injections of silica nanoparticles caused similar adverse effects, and smaller nanoparticles caused more severe reactions [79]. In contrast, skin painting of zinc oxide nanoparticles with ovalbumin and staphylococcal enterotoxin B did not exacerbate atopic dermatitis-like skin lesions [80]. These findings together suggest that although some nanoparticles might aggravate atopic dermatitis directly, this potential might be weaker under non-laboratory exposure conditions. One potential mechanism for the role of nanoparticles in atopic allergy is that the interaction of nanoparticles and allergen changes the skin penetration kinetics of the allergen and induces IgE-biased immune responses, which are a characteristic feature of human atopic allergies [81, 82]. For this effect, skin penetration by nanoparticles is unnecessary. Therefore studies designed to obviate nanoparticles' possible health risks should focus on not only the direct immunomodulating effects of nanoparticles but also on their potential to interfere with a healthy response to coexisting chemical substances or allergens.

6.4 Conclusion

Existing data are inconclusive regarding the health risk of skin exposure to nanoparticles. Although qualitative data suggest that nanoparticles enter the body via the skin in some situations, almost every quantitative study has concluded that the penetrating amount of nanoparticles is less than the detection limit of present technology, at least in the model used. To continue the discussion regarding the adverse effects of skin-acquired nanoparticles, we propose that further quantitative analyses are needed to determine the rate at which nanoparticles penetrate the skin and any potential for their accumulation. We also believe that additional research should be focused on determining whether nanoparticles cause immunologic sensitization, either directly or by promoting the sensitizing effects of co-exposed substances. These future studies will not only reveal ways through which we can accommodate an environment rich in nanoparticles; they also will provide insight into means to improve the safety and efficacy of nanomedicines for skin care.

References

1. Auffan M, Rose J, Bottero JY, Lowry GV, Jolivet JP, Wiesner MR. Towards a definition of inorganic nanoparticles from an environmental, health and safety perspective. *Nat Nanotechnol.* 2009;4(10):634–41. doi:10.1038/nnano.2009.242.
2. Cheng Z, Al Zaki A, Hui JZ, Muzykantov VR, Tsourkas A. Multifunctional nanoparticles: cost versus benefit of adding targeting and imaging capabilities. *Science.* 2012;338(6109):903–10. doi:10.1126/science.1226338.

3. Bowman DM, van Calster G, Friedrichs S. Nanomaterials and regulation of cosmetics. *Nat Nanotechnol.* 2010;5(2):92. doi:10.1038/nnano.2010.12.
4. Jain RK, Stylianopoulos T. Delivering nanomedicine to solid tumors. *Nat Rev Clin Oncol.* 2010;7(11):653–64. doi:10.1038/nrclinonc.2010.139.
5. Peters R, Kramer E, Oomen AG, Rivera ZE, Oegema G, Tromp PC, et al. Presence of nano-sized silica during *in vitro* digestion of foods containing silica as a food additive. *ACS Nano.* 2012;6(3):2441–51. doi:10.1021/nm204728k.
6. Nel A, Xia T, Madler L, Li N. Toxic potential of materials at the nanolevel. *Science.* 2006;311(5761):622–7. doi:10.1126/science.1114397.
7. Setyawati MI, Tay CY, Chia SL, Goh SL, Fang W, Neo MJ, et al. Titanium dioxide nanomaterials cause endothelial cell leakiness by disrupting the homophilic interaction of VE-cadherin. *Nat Commun.* 2013;4:1673. doi:10.1038/ncomms2655.
8. Yamashita K, Yoshioka Y, Higashisaka K, Mimura K, Morishita Y, Nozaki M, et al. Silica and titanium dioxide nanoparticles cause pregnancy complications in mice. *Nat Nanotechnol.* 2011;6(5):321–8. doi:10.1038/nnano.2011.41.
9. Janssen NA, Brunekreef B, van Vliet P, Aarts F, Meliefste K, Harssema H, et al. The relationship between air pollution from heavy traffic and allergic sensitization, bronchial hyperresponsiveness, and respiratory symptoms in Dutch schoolchildren. *Environ Health Perspect.* 2003;111(12):1512–18.
10. Kulbok PA, Baldwin JH. From preventive health behavior to health promotion: advancing a positive construct of health. *ANS Adv Nurs Sci.* 1992;14(4):50–64.
11. Bartra J, Mullol J, del Cuavillo A, Davila I, Ferrer M, Jauregui I, et al. Air pollution and allergens. *J Invest Allergol Clin Immunol.* 2007;17 Suppl 2:3–8.
12. McCreanor J, Cullinan P, Nieuwenhuijsen MJ, Stewart-Evans J, Malliarou E, Jarup L, et al. Respiratory effects of exposure to diesel traffic in persons with asthma. *N Engl J Med.* 2007;357(23):2348–58. doi:10.1056/NEJMoa071535.
13. Peters A, Wichmann HE, Tuch T, Heinrich J, Heyder J. Respiratory effects are associated with the number of ultrafine particles. *Am J Respir Crit Care Med.* 1997;155(4):1376–83. doi:10.1164/ajrccm.155.4.9105082.
14. Delfino RJ, Sioutas C, Malik S. Potential role of ultrafine particles in associations between airborne particle mass and cardiovascular health. *Environ Health Perspect.* 2005;113(8):934–46.
15. Maynard AD, Kuempel ED. Airborne nanostructured particles and occupational health. *J Nanoparticle Res.* 2005;7:587–614.
16. Tsuji JS, Maynard AD, Howard PC, James JT, Lam CW, Warheit DB, et al. Research strategies for safety evaluation of nanomaterials, part IV: risk assessment of nanoparticles. *Toxicol Sci.* 2006;89(1):42–50. doi:10.1093/toxsci/kfi339.
17. Popov AP, Priezzhev AV, Lademann J, Myllylä R. TiO₂ nanoparticles as an effective UV-B radiation skin-protective compound in sunscreens. *J Phys D Appl Phys.* 2005;38(15):2564–70.
18. Tyner KM, Wokovich AM, Godar DE, Doub WH, Sadrieh N. The state of nano-sized titanium dioxide (TiO₂) may affect sunscreen performance. *Int J Cosmet Sci.* 2011;33(3):234–44. doi:10.1111/j.1468-2494.2010.00622.x.
19. Weir A, Westerhoff P, Fabricius L, Hristovski K, von Goetz N. Titanium dioxide nanoparticles in food and personal care products. *Environ Sci Technol.* 2012;46(4):2242–50. doi:10.1021/es204168d.
20. Merget R, Bauer T, Kupper HU, Philippou S, Bauer HD, Breitstadt R, et al. Health hazards due to the inhalation of amorphous silica. *Arch Toxicol.* 2002;75(11–12):625–34.
21. Yin JJ, Lao F, Fu PP, Wamer WG, Zhao Y, Wang PC, et al. The scavenging of reactive oxygen species and the potential for cell protection by functionalized fullerene materials. *Biomaterials.* 2009;30(4):611–21. doi:10.1016/j.biomaterials.2008.09.061.
22. Lens M. Use of fullerenes in cosmetics. *Recent Patents Biotechnol.* 2009;3(2):118–23.
23. Lens M. Recent progresses in application of fullerenes in cosmetics. *Recent Patents Biotechnol.* 2011;5(2):67–73.

24. Marambio-Jones C, Hoek EMV. A review of the antibacterial effects of silver nanomaterials and potential implications for human health and the environment. *J Nanoparticle Res.* 2010;12:1531–51.
25. Quadros ME, Marr LC. Silver nanoparticles and total aerosols emitted by nanotechnology-related consumer spray products. *Environ Sci Technol.* 2011;45(24):10713–19. doi:10.1021/es202770m.
26. Kuhlbusch TA, Asbach C, Fissan H, Gohler D, Stintz M. Nanoparticle exposure at nanotechnology workplaces: a review. *Part Fibre Toxicol.* 2011;8:22. doi:10.1186/1743-8977-8-22.
27. Lighty JS, Veranth JM, Sarofim AF. Combustion aerosols: factors governing their size and composition and implications to human health. *J Air Waste Manage Assoc.* 2000;50(9):1565–618; discussion 619–22.
28. Zhu Y, Hinds WC, Kim S, Sioutas C. Concentration and size distribution of ultrafine particles near a major highway. *J Air Waste Manage Assoc.* 2002;52(9):1032–42.
29. Donaldson K, Stone V, Seaton A, MacNee W. Ambient particle inhalation and the cardiovascular system: potential mechanisms. *Environ Health Perspect.* 2001;109 Suppl 4:523–7.
30. Li N, Xia T, Nel AE. The role of oxidative stress in ambient particulate matter-induced lung diseases and its implications in the toxicity of engineered nanoparticles. *Free Radic Biol Med.* 2008;44(9):1689–99. doi:10.1016/j.freeradbiomed.2008.01.028.
31. Wiesner MR, Lowry GV, Casman E, Bertsch PM, Matson CW, Di Giulio RT, et al. Meditations on the ubiquity and mutability of nano-sized materials in the environment. *ACS Nano.* 2011;5(11):8466–70. doi:10.1021/nn204118p.
32. Glover RD, Miller JM, Hutchison JE. Generation of metal nanoparticles from silver and copper objects: nanoparticle dynamics on surfaces and potential sources of nanoparticles in the environment. *ACS Nano.* 2011;5(11):8950–7. doi:10.1021/nn2031319.
33. Mitrano DM, Rimmel E, Wichser A, Erni R, Height M, Nowack B. Presence of nanoparticles in wash water from conventional silver and nano-silver textiles. *ACS Nano.* 2014;8(7):7208–19. doi:10.1021/nn502228w.
34. Yin Y, Liu J, Jiang G. Sunlight-induced reduction of ionic Ag and Au to metallic nanoparticles by dissolved organic matter. *ACS Nano.* 2012;6(9):7910–19. doi:10.1021/nn302293r.
35. Schilling K, Bradford B, Castelli D, Dufour E, Nash JF, Pape W, et al. Human safety review of “nano” titanium dioxide and zinc oxide. *Photochem Photobiol Sci.* 2010;9(4):495–509. doi:10.1039/b9pp00180h.
36. Sadrieh N, Wokovich AM, Gopee NV, Zheng J, Haines D, Parmiter D, et al. Lack of significant dermal penetration of titanium dioxide from sunscreen formulations containing nano- and submicron-size TiO₂ particles. *Toxicol Sci.* 2010;115(1):156–66. doi:10.1093/toxsci/kfq041.
37. Zvyagin AV, Zhao X, Gierden A, Sanchez W, Ross JA, Roberts MS. Imaging of zinc oxide nanoparticle penetration in human skin *in vitro* and *in vivo*. *J Biomed Opt.* 2008;13(6):064031. doi:10.1117/1.3041492.
38. Senzui M, Tamura T, Miura K, Ikarashi Y, Watanabe Y, Fujii M. Study on penetration of titanium dioxide (TiO₂) nanoparticles into intact and damaged skin *in vitro*. *J Toxicol Sci.* 2010;35(1):107–13.
39. Fruijtier-Polloth C. The toxicological mode of action and the safety of synthetic amorphous silica—a nanostructured material. *Toxicology.* 2012;294(2–3):61–79. doi:10.1016/j.tox.2012.02.001.
40. Nabeshi H, Yoshikawa T, Matsuyama K, Nakazato Y, Matsuo K, Arimori A, et al. Systemic distribution, nuclear entry and cytotoxicity of amorphous nanosilica following topical application. *Biomaterials.* 2011;32(11):2713–24. doi:10.1016/j.biomaterials.2010.12.042.
41. Hirai T, Yoshikawa T, Nabeshi H, Yoshida T, Akase T, Yoshioka Y, et al. Dermal absorption of amorphous nanosilica particles after topical exposure for three days. *Die Pharmazie.* 2012;67(8):742–3.

42. Rancan F, Gao Q, Graf C, Troppens S, Hadam S, Hackbarth S, et al. Skin penetration and cellular uptake of amorphous silica nanoparticles with variable size, surface functionalization, and colloidal stability. *ACS Nano*. 2012;6(8):6829–42. doi:10.1021/nn301622h.
43. Ryman-Rasmussen JP, Riviere JE, Monteiro-Riviere NA. Penetration of intact skin by quantum dots with diverse physicochemical properties. *Toxicol Sci*. 2006;91(1):159–65. doi:10.1093/toxsci/kfj122.
44. Zhang LW, Yu WW, Colvin VL, Monteiro-Riviere NA. Biological interactions of quantum dot nanoparticles in skin and in human epidermal keratinocytes. *Toxicol Appl Pharmacol*. 2008;228(2):200–11. doi:10.1016/j.taap.2007.12.022.
45. Mortensen LJ, Oberdorster G, Pentland AP, Delouise LA. *In vivo* skin penetration of quantum dot nanoparticles in the murine model: the effect of UVR. *Nano Lett*. 2008;8(9):2779–87. doi:10.1021/nl801323y.
46. Prow TW, Monteiro-Riviere NA, Inman AO, Grice JE, Chen X, Zhao X, et al. Quantum dot penetration into viable human skin. *Nanotoxicology*. 2012;6(2):173–85. doi:10.3109/17435390.2011.569092.
47. Bos JD, Meinardi MM. The 500 Dalton rule for the skin penetration of chemical compounds and drugs. *Exp Dermatol*. 2000;9(3):165–9.
48. Hansen S, Lehr CM. Transfollicular delivery takes root: the future for vaccine design? *Expert Rev Vaccines*. 2014;13(1):5–7. doi:10.1586/14760584.2014.862500.
49. Brandner JM, McIntyre M, Kief S, Wladykowski E, Moll I. Expression and localization of tight junction-associated proteins in human hair follicles. *Arch Dermatol Res*. 2003;295(5):211–21. doi:10.1007/s00403-003-0418-3.
50. Kubo A, Nagao K, Yokouchi M, Sasaki H, Amagai M. External antigen uptake by Langerhans cells with reorganization of epidermal tight junction barriers. *J Exp Med*. 2009;206(13):2937–46. doi:10.1084/jem.20091527.
51. Kuchler S, Abdel-Mottaleb M, Lamprecht A, Radowski MR, Haag R, Schafer-Korting M. Influence of nanocarrier type and size on skin delivery of hydrophilic agents. *Int J Pharm*. 2009;377(1–2):169–72. doi:10.1016/j.ijpharm.2009.04.046.
52. Rouse JG, Yang J, Ryman-Rasmussen JP, Barron AR, Monteiro-Riviere NA. Effects of mechanical flexion on the penetration of fullerene amino acid-derivatized peptide nanoparticles through skin. *Nano Lett*. 2007;7(1):155–60. doi:10.1021/nl062464m.
53. Winter M, Beer HD, Hornung V, Kramer U, Schins RP, Forster I. Activation of the inflammasome by amorphous silica and TiO₂ nanoparticles in murine dendritic cells. *Nanotoxicology*. 2011;5(3):326–40. doi:10.3109/17435390.2010.506957.
54. Kumar K, Saini S, Mehta RD, Bumb RA. Contact leucoderma caused by lemon. *Indian J Dermatol Venereol Leprol*. 1996;62(1):61.
55. Lunov O, Syrovets T, Loos C, Nienhaus GU, Mailander V, Landfester K, et al. Amino-functionalized polystyrene nanoparticles activate the NLRP3 inflammasome in human macrophages. *ACS Nano*. 2011;5(12):9648–57. doi:10.1021/nn203596e.
56. Stern ST, Adiseshaiah PP, Crist RM. Autophagy and lysosomal dysfunction as emerging mechanisms of nanomaterial toxicity. *Part Fibre Toxicol*. 2012;9:20. doi:10.1186/1743-8977-9-20.
57. Kang SJ, Kim BM, Lee YJ, Chung HW. Titanium dioxide nanoparticles trigger p53-mediated damage response in peripheral blood lymphocytes. *Environ Mol Mutagen*. 2008;49(5):399–405. doi:10.1002/em.20399.
58. Trouiller B, Reliene R, Westbrook A, Solaimani P, Schiestl RH. Titanium dioxide nanoparticles induce DNA damage and genetic instability *in vivo* in mice. *Cancer Res*. 2009;69(22):8784–9. doi:10.1158/0008-5472.CAN-09-2496.
59. Ye Y, Liu J, Xu J, Sun L, Chen M, Lan M. Nano-SiO₂ induces apoptosis via activation of p53 and Bax mediated by oxidative stress in human hepatic cell line. *Toxicol In Vitro*. 2010;24(3):751–8. doi:10.1016/j.tiv.2010.01.001.

60. Nabeshi H, Yoshikawa T, Matsuyama K, Nakazato Y, Tochigi S, Kondoh S, et al. Amorphous nanosilica induce endocytosis-dependent ROS generation and DNA damage in human keratinocytes. *Part Fibre Toxicol.* 2011;8:1. doi:10.1186/1743-8977-8-1.
61. Ristow M. Unraveling the truth about antioxidants: mitohormesis explains ROS-induced health benefits. *Nat Med.* 2014;20(7):709–11. doi:10.1038/nm.3624.
62. Kawagishi H, Finkel T. Unraveling the truth about antioxidants: ROS and disease: finding the right balance. *Nat Med.* 2014;20(7):711–13. doi:10.1038/nm.3625.
63. Heidler T, Hartwig K, Daniel H, Wenzel U. *Caenorhabditis elegans* lifespan extension caused by treatment with an orally active ROS-generator is dependent on DAF-16 and SIR-2.1. *Biogerontology.* 2010;11(2):183–95. doi:10.1007/s10522-009-9239-x.
64. Yang W, Hekimi S. A mitochondrial superoxide signal triggers increased longevity in *Caenorhabditis elegans*. *PLoS Biol.* 2010;8(12), e1000556. doi:10.1371/journal.pbio.1000556.
65. Kimber I, Hilton J, Dearman RJ, Gerberick GF, Ryan CA, Basketter DA, et al. An international evaluation of the murine local lymph node assay and comparison of modified procedures. *Toxicology.* 1995;103(1):63–73.
66. Park YH, Jeong SH, Yi SM, Choi BH, Kim YR, Kim IK, et al. Analysis for the potential of polystyrene and TiO₂ nanoparticles to induce skin irritation, phototoxicity, and sensitization. *Toxicol In Vitro.* 2011;25(8):1863–9. doi:10.1016/j.tiv.2011.05.022.
67. Lee S, Yun HS, Kim SH. The comparative effects of mesoporous silica nanoparticles and colloidal silica on inflammation and apoptosis. *Biomaterials.* 2011;32(35):9434–43. doi:10.1016/j.biomaterials.2011.08.042.
68. Chen BX, Wilson SR, Das M, Coughlin DJ, Erlanger BF. Antigenicity of fullerenes: antibodies specific for fullerenes and their characteristics. *Proc Natl Acad Sci U S A.* 1998;95(18):10809–13.
69. Braden BC, Goldbaum FA, Chen BX, Kirschner AN, Wilson SR, Erlanger BF. X-ray crystal structure of an anti-Buckminsterfullerene antibody fab fragment: biomolecular recognition of C(60). *Proc Natl Acad Sci U S A.* 2000;97(22):12193–7. doi:10.1073/pnas.210396197.
70. Ema M, Matsuda A, Kobayashi N, Naya M, Nakanishi J. Dermal and ocular irritation and skin sensitization studies of fullerene C60 nanoparticles. *Cutan Ocul Toxicol.* 2013;32(2):128–34. doi:10.3109/15569527.2012.727937.
71. Mann E, Ranft U, Eberwein G, Gladtko D, Sugiri D, Behrendt H, et al. Does airborne nickel exposure induce nickel sensitization? *Contact Dermatitis.* 2010;62(6):355–62. doi:10.1111/j.1600-0536.2010.01725.x.
72. Otani S, Onishi K, Mu H, Yokoyama Y, Hosoda T, Okamoto M, et al. The relationship between skin symptoms and allergic reactions to Asian dust. *Int J Environ Res Pub Health.* 2012;9(12):4606–14. doi:10.3390/ijerph9124606.
73. Swinnen I, Goossens A. An update on airborne contact dermatitis: 2007–2011. *Contact Dermatitis.* 2013;68(4):232–8. doi:10.1111/cod.12022.
74. Hussain S, Smulders S, De Vooght V, Ectors B, Boland S, Marano F, et al. Nano-titanium dioxide modulates the dermal sensitization potency of DNCB. *Part Fibre ogy.* 2012;9:15. doi:10.1186/1743-8977-9-15.
75. Fonacier LS, Dreskin SC, Leung DY. Allergic skin diseases. *J Allergy Clin Immunol.* 2010;125(2 Suppl 2):S138–49. doi:10.1016/j.jaci.2009.05.039.
76. Sanda T, Yasue T, Oohashi M, Yasue A. Effectiveness of house dust-mite allergen avoidance through clean room therapy in patients with atopic dermatitis. *J Allergy Clin Immunol.* 1992;89(3):653–7.
77. Yanagisawa R, Takano H, Inoue K, Koike E, Kamachi T, Sadakane K, et al. Titanium dioxide nanoparticles aggravate atopic dermatitis-like skin lesions in NC/Nga mice. *Exp Biol Med.* 2009;234(3):314–22. doi:10.3181/0810-RM-304.
78. Yanagisawa R, Takano H, Inoue KI, Koike E, Sadakane K, Ichinose T. Size effects of polystyrene nanoparticles on atopic dermatitislike skin lesions in NC/NGA mice. *Int J Immunopathol Pharmacol.* 2010;23(1):131–41.