

serum α 1,3fucosyltransferase activity [38]. In fact, the patient was determined to possess the *pf* gene homozygously (*pf/pf*) indicating occurrence of lethal mutation in the *FUT6* genes which resulted in the absence of fucosylated glycans in AGP [37]. However, a couple of fucosylated glycans were still detected as less than several % of the total glycans in AGP (Fig 4b). Accordingly, it remains to be examined a significance of poorly expressed fucosylated glycans in such a cancer patient with *FUT6*-deficiency (details of molecular and enzymatic studies on fucosylation of AGP in patient N with *FUT6*-deficiency will be published elsewhere).

Discussion

Proteins are frequently modified post-translationally and more than half of all proteins in nature are glycosylated [39] resulting in being present as glycoproteins with covalently attached glycans having highly heterogeneous and diverse structures. It must therefore be plausible that these glycans have significant implication and responsibility for the structure and function of such glycoproteins. It has been widely recognized that glycans play important roles in not only protein folding and clearance but also cell to cell interaction, recognition and binding together with frequent occurrences of modifications in glycans themselves [40–42]. With the aid of detailed analyses of glycan structures in well characterized glycoproteins, it has been demonstrated that aberrant and novel glycans are synthesized associated with various biological processes, in particular, malignancies [2,43]. Hence, cancer-associated changes of glycan molecules in glycoproteins as well as in other glycoconjugates such as glycolipids [44,45] have been widely demonstrated. Further, such changes are recognized as one of the most promising targets to improve existing tumor markers by means of predicting prognosis of cancer patients more precisely [46].

Whereas, since all these glycans are synthesized enzymatically through catalytic actions of a series of glycosyltransferases, such enzymes could be an important target to detect glycan-based changes in malignancies involving possible establishment of a novel biomarker for diagnosis of cancer. In fact, our previous studies demonstrated the usefulness of analyzing α 1,2- [47–49], α 1,3- [20,21,37,38,50–54], α 1,4- [55–58] and α 1,6fucosyltransferase [59] activities using specific sugar-acceptors, which emphasized the specific role of the fucosylated antigens appeared to play on carcinogenesis [1,47–49]. In particular, significantly elevated activities of α 1,3fucosyltransferase in plasma and/or serum have been observed in patients with diverse cancers [20,21,51–54]. It has, therefore, been suggested that measurement of serum α 1,3fucosyltransferase activity has some superiority over diagnosis with tumor-associated antigen levels in the same samples as a novel biomarker for patients in earlier clinical stages and with smaller size of tumors. Although the possibility cannot be excluded that the enzyme with elevated activities is originated from tumor itself as observed in patients with colorectal carcinoma [60], the origin of α 1,3fucosyltransferase in serum has been considered to be the liver. It has also been determined that α 1,3fucosyltransferase in serum is encoded by the *FUT6* gene and the missense mutation that occurs in the *FUT6* gene results in the deficiency of serum α 1,3fucosyltransferase activity [37,38,61]. Therefore, tumor-associated elevation of serum α 1,3fucosyltransferase activities is assumed to occur in the liver but not originated from tumors formed.

Human AGP is generally recognized as the major serum protein with highly glycosylated glycans and to be originated from the liver. Glycans in AGP consist of five complex type glycans including di-, tri- and tetraantennary structures [25,33]. It has also been determined by our recent mass spectrometric analysis that almost all the terminal galactose residues in every glycan chains are sialylated and the α 1,3fucosyl residues are present on the lactosamine structures in both tri- and tetraantennary glycans and elongated tetraantennary ones with repeating lactosamine structure. Hence, our previous investigations convincingly demonstrated both

presence of a significant correlation between activities of α 1,3fucosyltransferase and total amounts of fucosylated glycans in AGP using the same serum samples [34] and absence of fucosylated glycans in AGP obtained from FUT6-deficient individuals [37]. Furthermore, such a-fuco-glycans of AGP obtained from FUT6-deficient individuals could be fucosylated on their tri- and tetraantennary structures using the recombinant FUT6 enzyme or purified plasma α 1,3fucosyltransferase under the presence of GDP-fucose as a sugar-donor [S. Yazawa, unpublished observation]. It could, therefore, be strongly suggested that tumor-dependent, increased α 1,3fucosylated glycans in AGP are synthesized by an action of the hepatic FUT6 gene encoded α 1,3fucosyltransferase with highly elevated activities.

There have been many reports indicating that analyses of changes occurred in both glycan structures and concentrations of AGP in serum could be useful for the diagnosis and successful management of various diseases including diverse cancers [31,62–74]. Our previous studies suggested that fucosylation of AGP implied significant potential as a novel biomarker for diagnosis and prognosis of cancer patients, which must be strongly associated with tumor-dependent changes of serum α 1,3fucosyltransferase activity. Indeed, a strong correlation of levels between fucosylated AGP and α 1,3fucosyltransferase activity in the same serum samples was demonstrated previously. Further, the fucosylation index of AGP glycoforms and the relative abundance of fucosylated glycans in AGP (FUCAGP) determined by means of CAIE method [34] and mass spectrometric analysis [35], respectively, were found, for the first time, to be useful for predicting postoperative cancer patients.

In the present study, we first focused on levels of FUCAGP and serum AGP concentrations in patients with various cancers in different clinical stages. Levels of FUCAGP and serum AGP were analyzed in relation to their clinicopathological features in patients ($n = 30$) who received surgical interventions and/or various chemotherapy treatments. As demonstrated previously [34,35], significant elevated levels of both FUCAGP and serum AGP concentrations were observed in cancer patients at preoperative periods when compared with those in healthy controls. No significant correlation was found between levels of FUCAGP and AGP in both cancer patients and healthy controls, respectively. Interestingly, a very low level of FUCAGP was found in a patient who underwent neoadjuvant chemotherapy. Whereas, as demonstrated previously [34,35], increases of relative amount of diantennary glycans rather than those of tri- or tetraantennary glycans in AGP occurred primarily along with a new synthesis of AGP shortly after operation. This rapid response seemed to occur commonly in patients at the time of operation, and to involve as a consequence of the acute phase reaction toward surgical treatment. During the periods, fucosylated glycans were scarcely detected in AGP, since as described above, only tri- and tetraantennary glycans in AGP were fucosylated through α 1,3linkages.

Previous studies on glycoforms and corresponding glycan structures also indicated that, secondarily, AGP with highly fucosylated and branched glycan chains increased specifically in patients with poor prognosis but not in patients with good prognosis. Further, levels of FUCAGP were suggested to become a clinically relevant biomarker over existing diagnostic tumor markers. However, levels of FUCAGP and AGP followed in our previous studies [34, 35] were determined for a limited duration up to one and a half year after operation, and in particular, validation of the fucosylation index of AGP glycoforms and levels of FUCAGP was scarcely performed for predicting treatment outcome in therapeutic settings. Accordingly, we then focused on analyzing variability of serum FUCAGP in patients who had been followed for several years after operation and had undergone various chemotherapies during the period in connection with patients' responses to medication and various chemotherapy treatments. Seventeen patients including ten patients who died due to the recurrence and/or metastasis of tumors were followed up by analyzing levels of FUCAGP and AGP concentrations in their serum samples together with general surveillance with CT scans and tumor marker diagnosis.

Results from follow-up studies of FUCAGP levels conducted depending on patients' prognoses indicated that elevated levels of FUCAGP were retained and no striking drop of the level was observed in patients who seemed not to respond to any repeated treatments of chemotherapy and died with a short survival after operation. Whereas, patients who survived for a longer period (up to 1639 POD) under the therapeutic settings with repeated chemotherapies showed partly different changes of FUCAGP levels. First, patients whose FUCAGP levels were found to extremely elevate shortly after operation and survived for a longer period with repeated chemotherapy seemed to respond to the first line of adjuvant chemotherapy resulting that the level fell and in some cases dropped below the cut-off level. However, either recurrence and/or metastasis or no experience of complete remission occurred, and subsequently they died due to the progress of disease accompanying constantly elevated levels of FUCAGP. Second, patients whose FUCAGP levels had been retained at relatively low and around the cut-off level under receiving repeated chemotherapy treatments showed an incomplete response to the latest line treatment of chemotherapy resulting in progress of disease and constant elevation of FUCAGP levels.

Whereas, patients who had been survived during the observation period with and without chemotherapy treatment showed either clear response to repeated chemotherapy treatments or no clinical recurrence during the period of observation. It was of particular interest that in these patients with good prognosis, elevated levels of FUCAGP once after receiving the previous line treatment of chemotherapy fell around the cut-off level by receiving the latest line treatment of chemotherapy and low levels of FUCAGP after operation were retained around the normal range without adjuvant chemotherapy treatment. Therefore, these follow-up studies of FUCAGP levels in patients with different prognosis suggested that patients with poor prognosis possessed extremely elevated levels in the latest period but patient with good prognosis seemed to retain relatively low levels around the normal range. Further, in patients who seemed to respond well to chemotherapy treatment and survived, elevated levels found in association with the recurrence and/or metastasis of tumors dropped to the normal range after treatment with the latest line treatment of chemotherapy against such clinical events. It is more noteworthy that all such changes in FUCAGP levels seemed to occur frequently in advance before detection of recurrence and/or metastasis of tumors and respective tumor markers coming up positive.

One of the obvious evidences to provide that fucosylation of the AGP molecule is accompanied solely by the *FUT6* gene-encoded hepatic α 1,3fucosyltransferase has been obtained from molecular and enzymatic studies on *FUT6* deficient individuals. Lack of serum α 1,3fucosyltransferase due to the *FUT6*-deficiency has scarcely been found so far and only restricted numbers of individuals whose AGP molecule has no fucosylated glycan have been reported previously [37,38,75]. The patient N was supposed to possess the lethal mutation in the *FUT6* genes resulting in possessing the mutant *pf* gene homozygously as demonstrated previously [37]. While, mass spectrometric analysis of AGP glycans in this patient showed that up to several % of glycans in AGP were still fucosylated and a couple of fucosylated glycans were constantly detected during the period of observation (Fig 4b). One of the seven individuals with *FUT6*-deficiency in our previous study showed to possess a small amount of fucosylated glycoforms in AGP [37]. These fucosylated glycans might be synthesized by the action of a certain α 1,3fucosyltransferase other than *FUT6*-encoded α 1,3fucosyltransferase. CEA levels in the patient N were continuously above the normal range even though CT scans revealed no significant remarks during the period. Accordingly, CEA levels in this patient might be subjected to be false-positive range as demonstrated recently [76]. It has been widely recognized that CA 19-9 could not be used as tumor marker in Lewis-negative cancer patients because individuals with Lewis negative genotypes lack the key enzyme to synthesize Fuc α 1,4GlcNAc linkage, part

of its antigenic determinant [57,77]. Hence, the results suggest that FUCAGP level could not be useful in patients with FUT6 deficiency as a clinically relevant biomarker.

The function of plasma AGP and its potential physiologic significance as an acute phase protein have generated profound interest [33,78–81]. Generally, AGP has highly glycosylated *N*-glycan chains on the molecule with more than 45% content and with abundant heterogeneity. It has therefore been demonstrated that despite expression of different genomic variants and mRNA levels of AGP, drastic changes occur frequently in not only serum concentrations but also their glycan structures including branching, sialylation and, in particular, fucosylation degrees as posttranslational modification involving in rapid responses to various biological phenomena [25–35,78–84]. It was also shown that each AGP glycoform was renewed at a rate of 15% per day of the plasma pool [85]. A wide variety of changes and a large amount of replacements were also observed in AGP glycans from various cancer patients who received surgical interventions and chemotherapies [34].

Glycoproteomic approach to investigate tumor-associated alteration of glycans in glycoproteins has been developed recently. As one of the most powerful tools, mass spectrometric analyses through MALDI TOF or LC-MS/MS together with specific enrichment methods such as the use of lectin-affinity toward glycans targeted have been applied for a comprehensive analysis of large amounts of glycans. Indeed, findings of usefulness of *Aleuria aurantia* lectin (AAL) for isolation of fucosylated antigens [86,87] and characterization of its detailed binding specificity [88] have been carried on into further applications of this lectin to recent glycoproteomic studies using human plasma and/or serum samples [34,71,73,89–92]. Whereas, AGP is a major serum protein and due to its original aspect as an acute phase protein, changes of AGP concentrations in serum waver depending on the degree of inflammation caused by a series of diseases and biological events. As expected from our previous studies [34,35], serum AGP levels were found to change differently from FUCAGP ones during the follow-up period in most of the patients. It was also suggested that AGP level changed indistinguishably in patients with poor and good prognosis after operation. Further, it was not unusual observation to see aberrantly elevated AGP levels in serum samples from cancer patients up to several times higher than those in healthy controls. It should therefore be important for targeting serum AGP molecule to focus on not only the fucosylated AGP glycans but also relative abundance of fucosylated AGP glycans after exclusion of large amounts of background noise.

In conclusion, in the present study, FUCAGP could be a clinically relevant biomarker of cancer progression as well as cancer prognosis. The FUCAGP level was found to closely relate to patients' response to various chemotherapies. Further, instead of our previous CAIE technique for measuring AGP glycoforms which was not simple or convenient for assaying large numbers of samples at attempt, the evaluation of AGP glycans in the present study, by means of an AGP-prep-DOCK and mass spectrometric analyses, was aimed for a quick determination of serum AGP glycans. Additionally, recently we developed and improved AGPAS software for very rapid determination of FUCAGP levels in serum samples. These reflect our long standing view that rapid and accurate measurement of FUCAGP in serum samples might advance diagnosis of better treatment outcomes of cancer patients. We believe that changes in FUCAGP level in a cohort of patients followed up while undergoing various chemotherapeutic regimens warrant further investigation.

Acknowledgments

We gratefully acknowledge the valuable comments on this work by late Prof. Naohisa Kochibe, Gunma University and also to Dr. Touko Hirano and Ms. Mariko Nakamura for their excellent technical assistance.

Author Contributions

Conceived and designed the experiments: SY TA. Performed the experiments: SY RS. Analyzed the data: SY RT TY RS AM ARS TA. Contributed reagents/materials/analysis tools: SY RT TY RS AM ARS HK TA. Wrote the paper: SY.

References

1. Hakomori S, Kannagi R. Glycosphingolipids as tumor-associated and differentiation markers. *J Natl Cancer Inst.* 1983; 71:231–51. PMID: [6576183](#)
2. Varki A, Kannagi R, Toole BP. Glycosylation changes in cancer. In: Varki A, Cummings RD, Esko JD, Freeze HH, Stanley P, Bortozzi CR, et al., editors. *Essentials of Glycobiology*. 2nd ed. New York: Cold Spring Harbor Lab. Press; 2009. p. 617–32.
3. Gold P, Freedman SO. Demonstration of tumor-specific antigens in human colonic carcinomata by immunological tolerance and absorption techniques. *J Exp Med.* 1965; 121:439–62. PMID: [14270243](#)
4. Koprowski H, Stepewski Z, Mitchell K, Herlyn M, Herlyn D, Fuhrer P. Colorectal carcinoma antigens detected by hybridoma antibodies. *Somatic Cell Genet.* 1979; 5:957–71. PMID: [94699](#)
5. Koprowski H, Brockhaus M, Blaszyk M, Magnani JL, Stepewski Z, Ginsburg V. Lewis blood-type may affect the incidence of gastrointestinal cancer. *Lancet* 1982; 1:1332–3. PMID: [6177980](#)
6. Magnani JL, Nilsson B, Brockhaus M, Zopf D, Stepewski Z, Koprowski H, et al. A monoclonal antibody-defined antigen associated with gastrointestinal cancer is a ganglioside containing sialylated lacto-N-fucopentaose II. *J Biol Chem.* 1982; 257:14365–9. PMID: [7142214](#)
7. Abelev GI. Alpha-fetoprotein in ontogenesis and its association with malignant tumors. *Adv Cancer Res.* 1971; 14:295–358. PMID: [4107670](#)
8. Stenman UH, Sutinen ML, Selander RK, Tontti K, Schröder J. Characterization of a monoclonal antibody to human alpha-fetoprotein and its use in affinity chromatography. *J Immunol Methods.* 1981; 46:337–45. PMID: [6171597](#)
9. Bast RC Jr, Feeney M, Lazarus H, Nadler LM, Colvin RB, Knopp RC. Reactivity of a monoclonal antibody with human ovarian carcinoma. *J Clin Invest.* 1981; 68:1331–7. PMID: [7028788](#)
10. Klug TL, Bast RC Jr, Niloff JM, Knapp RC, Zurawski VR Jr. Monoclonal antibody immunoradiometric assay for an antigenic determinant (CA 125) associated with human epithelial ovarian carcinomas. *Cancer Res.* 1981; 44:1048–53.
11. Fukushi Y, Kannagi R, Hakomori S, Shepard T, Kulander BG, Singer JW. Location and distribution of difucoganglioside (VI3NeuAcV3III3Fuc₂NL₆) in normal and tumor tissues defined by its monoclonal antibody FH6. *Cancer Res.* 1985; 45:3711–7. PMID: [4016748](#)
12. Kannagi R, Fukushi Y, Tachikawa T, Noda A, Shin S, Shigeta K, et al. Quantitative and qualitative characterization of human cancer-associated serum glycoprotein antigens expressing fucosyl or sialyl-fucosyl type 2 chain polylectosamine. *Cancer Res.* 1986; 46:2619–26. PMID: [3008996](#)
13. Itzkowitz SH, Bloom EJ, Kokal WA, Modin G, Hakomori S, Kim YS. Sialosyl-Tn. A novel mucin antigen associated with prognosis in colorectal cancer patients. *Cancer* 1990; 66: 1960–6. PMID: [2224793](#)
14. Inoue M, Ogawa H, Nakanishi K, Tanigawa O, Karino K, Endo J. Clinical value of sialyl Tn antigen in patients with gynecologic tumors. *Obstet Gynecol.* 1991; 75:1032–6.
15. Ni XG, Bai XF, Mao YL, Shao YF, Wu JX, Shan Y, et al. The clinical value of serum CEA, CA19-9, and CA242 in the diagnosis and prognosis of pancreatic cancer. *Eur J Surg Oncol.* 2005; 31:164–9. PMID: [15698733](#)
16. Schutter EM, Davelaar EM, van Kamp GJ, Verstraeten RA, Kenemans P, Verheijen RH. The differential diagnostic potential of a panel of tumor markers (CA 125, CA 15–3, and CA 72–4 antigens) in patients with a pelvic mass. *Am J Obstet Gynecol.* 2002; 187:385–92. PMID: [12193930](#)
17. Gaspar MJ, De Miguel J, García Díaz JD, Diez M. Clinical utility of a combination of tumour markers in the diagnosis of malignant pleural effusions. *Anticancer Res.* 2008; 28:2947–52. PMID: [19031938](#)
18. Yang AP, Liu J, Lei HY, Zhang QW, Zhao L, Yang GH. CA72-4 combined with CEA, CA125 and CA19-9 improves the sensitivity for the early diagnosis of gastric cancer. *Clin Chim Acta.* 2014; 437:183–6. doi: [10.1016/j.cca.2014.07.034](#) PMID: [25086284](#)
19. Lai H, Jin Q, Liu Y, Mo X, Li B, He K, et al. Combined use of lysyl oxidase, carcino-embryonic antigen, and carbohydrate antigens improves the sensitivity of biomarkers in predicting lymph node metastasis and peritoneal metastasis in gastric cancer. *Tumour Biol.* 2014; 35:10547–54. doi: [10.1007/s13277-014-2355-5](#) PMID: [25060181](#)

20. Yazawa S, Asao T, Nagamachi Y, Abbas SA, Matta KL. Tumor-related elevation of serum $\alpha(1\text{--}3)\text{-L}$ -fucosyltransferase activity in gastric cancer. *J Cancer Res.* 1989; 115:451–5.
21. Asao T, Yazawa S, Nagamachi Y, Abbas SA, Matta KL. Serum $\alpha(1\text{--}3)\text{-L}$ -fucosyltransferase, carcinoembryonic antigen, and sialyl Lewis X-i antigen levels in lung cancer. *Cancer.* 1989; 64:2541–5. PMID: [2819663](#)
22. Hakomori S, Nudelman E, Levery SB, Kannagi R. Novel fucolipids accumulating in human adenocarcinoma. I. Glycolipids with di- or trifucosylated type 2 chain. *J Biol Chem.* 1984; 259:4672–80. PMID: [6707026](#)
23. Fukushi Y, Hakomori S, Nudelman E, Cochran N. Novel fucolipids accumulating in human adenocarcinoma. II. Selective isolation of hybridoma antibodies that differentially recognize mono-, di-, and trifucosylated type 2 chain. *J Biol Chem.* 1984; 259:4681–5. PMID: [6200484](#)
24. Nichols EJ, Kannagi R, Hakomori S, Krantz MJ, Fuks A. Carbohydrate determinants associated with carcinoembryonic antigen (CEA). *J Immunol.* 1985; 135:1911–3. PMID: [2410505](#)
25. van Dijk W, Havenaar EC, Brinkman-van der Linden EC. Alpha1-acid glycoprotein (orosomucoid): pathophysiological changes in glycosylation in relation to its function. *Glycoconjugate J.* 1995; 12:227–33.
26. Hansen JE, Larsen VA, Bøg-Hansen TC. The microheterogeneity of alpha 1-acid glycoprotein in inflammatory lung disease, cancer of the lung and normal health. *Clin Chim Acta.* 1984; 138:41–7. PMID: [6713687](#)
27. Bories PN, Feger J, Benbernou N, Rouzeau JD, Agneray J, Durand G. Prevalence of tri- and tetraantennary glycans of human alpha 1-acid glycoprotein in release of macrophage inhibitor of interleukin-1 activity. *Inflammation.* 1990; 14:315–23. PMID: [2361735](#)
28. Hrycaj P, Sobieska M, Mackiewicz S, Müller W. Microheterogeneity of alpha 1-acid glycoprotein in early and established rheumatoid arthritis. *J Rheumatol.* 1993; 20:2020–4. PMID: [8014928](#)
29. Mackiewicz A, Mackiewicz K. Glycoforms of serum alpha 1-acid glycoprotein as markers of inflammation and cancer. *Glycoconjugate J.* 1995; 12:241–7.
30. Brinkman-Van der Linden CM, Havenaar EC, Van Ommen CR, Van Kamp GJ, Gooren LJ, Van Dijk W. Oral estrogen treatment induces a decrease in expression of sialyl LewisX on α_1 -acid glycoprotein in females and male-to-female transsexuals. *Glycobiology.* 1996; 6:407–12. PMID: [8842704](#)
31. Shiyan SD, Bovin NV. Carbohydrate composition and immunomodulatory activity of different glycoforms of alpha 1-acid glycoprotein. *Glycoconjugate J.* 1997; 14:631–8.
32. Zimmermann-Belsing T, Feldt-Rasmussen U, From G, Perrild H, Bøg-Hansen TC. Long-term pathologic changes of α_1 -acid glycoprotein (orosomucoid) glycoforms in autoimmune thyroid disease. *Autoimmunity.* 2002; 35:441–7. PMID: [12685872](#)
33. Fournier T, Medjoubi-N N, Porquet D. Alpha-1-acid glycoprotein. *Biochim Biophys Acta.* 2000; 1482:157–71. PMID: [11058758](#)
34. Hashimoto S, Asao T, Takahashi J, Yagihashi Y, Nishimura T, Saniabadi AR, et al. α_1 -Acid glycoprotein fucosylation as a marker of carcinoma progression and prognosis. *Cancer* 2004; 15:2825–36.
35. Asao T, Yazawa S, Nishimura T, Hayashi T, Shimaoka H, Saniabadi AR, et al. Development of a novel system for mass spectrometric analysis of cancer-associated fucosylation in plasma α_1 -acid glycoprotein. *BoiMed Res Internat.* 2013; (Supplement published in *World Biomed Frontiers* ISSN:2328-0166, 2014).
36. Sobin LH, Gospodarowicz MK, Wittekind C. *TNM Classification of Malignant Tumours.* 7th ed. NY: Wiley-Blackwell; 2009.
37. Tanaka S, Yazawa S, Noguchi K, Nishimura T, Miyanaga K, Kochibe N, et al. Molecular analysis of plasma $\alpha_1,3$ -fucosyltransferase deficiency and development of the methods for its genotyping. *Exp Clin Immunogenet.* 2001; 18:1–12. PMID: [11150848](#)
38. Yazawa S, Tanaka S, Nishimura T, Miyanaga K, Kochibe N. Plasma $\alpha_1,3$ -fucosyltransferase deficiency in schizophrania. *Exp Clin Immunogenet.* 1999; 16:125–30. PMID: [10394050](#)
39. Apweiler R, Hermjakob H, Sharon N. On the frequency of protein glycosylation, as deduced from analysis of the SWISS-PROT database. *Biochim Biophys Acta.* 1999; 1473:4–8. PMID: [10580125](#)
40. Shental-Bechor D, Levy Y. Folding of glycoproteins: toward understanding the biophysics of the glycosylation code. *Curr Opin Struct Biol.* 2009; 19:524–33. doi: [10.1016/j.sbi.2009.07.002](#) PMID: [19647993](#)
41. Fukuda MN, Sasaki H, Lopez L, Fukuda M. Survival of recombinant erythropoietin in the circulation: the role of carbohydrates. *Blood.* 1989; 73:84–9. PMID: [2910371](#)
42. Gu J, Isaji T, Xu Q, Kariya Y, Gu W, Fukuda T, et al. Potential roles of N-glycosylation in cell adhesion. *Glycoconjugate J.* 2012; 29:599–607.

43. Fukuda M. Possible roles of tumor-associated carbohydrates antigens. *Cancer Res.* 1996; 56:2237–44. PMID: [8625291](#)
44. Hakomori S. Tumor associated glycolipid antigens, their metabolism and organization. *Chem Phys Lipids.* 1986; 42:209–33. PMID: [2435423](#)
45. Hakomori S. Tumor malignancy defined by aberrant glycosylation and sphingo(glyco)lipid metabolism. *Cancer Res.* 1996; 56:5309–18. PMID: [8968075](#)
46. Dube DH, Bertozzi CR. Glycans in cancer and inflammation—potential for therapeutics and diagnostics. *Nat Rev Drug Discov.* 2005; 4:477–88. PMID: [15931257](#)
47. Yazawa S, Nakamura J, Asao T, Nagamachi Y, Sagi M, Matta KL, et al. Aberrant α 1 \rightarrow 2fucosyltransferase found in human colorectal carcinoma involved in the accumulation of Leb and Y antigens in colorectal tumors. *Jpn J Cancer Res.* 1993; 84:989–95. PMID: [8407568](#)
48. Yazawa S, Nishimura T, Ide M, Asao T, Okamura A, Tanaka S, et al. Tumor-related expression of α 1,2fucosylated antigens on colorectal carcinoma cells and its suppression by cell-mediated priming using sugar acceptors for α 1,2fucosyl-transferase. *Glycobiology.* 2002; 12:545–53. PMID: [12213787](#)
49. Tsuboi K, Asao T, Ide M, Hashimoto S, Noguchi K, Kominato Y, et al. α 1,2Fucosylation is a superior predictor of postoperative prognosis for colorectal cancer compared with blood group A, B, or sialyl Lewis X antigen generated within colorectal tumor tissues. *Ann Surg Oncol.* 2007; 14:1880–9. PMID: [17375356](#)
50. Madiyalakan R, Yazawa S, Abbas SA, Barlow JJ, Matta KL. Use of *N*-acetyl-2'-*O*-methylactosamine as a specific acceptor for the determination of α -L-(1 \rightarrow 3)-fucosyltransferase in human serum. *Anal Biochem.* 1986; 52:2–8.
51. Yazawa S, Madiyalakan R, Izawa H, Asao T, Furukawa K, Matta KL. Cancer-associated elevation of α (1 \rightarrow 3)-L-fucosyltransferase activity in human serum. *Cancer* 1988; 62:516–20. PMID: [3390792](#)
52. Yazawa S, Asao T, Nagamachi Y, Matta KL. Elevated activities of serum α (1 \rightarrow 3) -L-fucosyltransferase in human cancer. *J Tumor Oncol.* 1989; 4:355–62.
53. Tachikawa T, Yazawa S, Asao T, Shin S, Yanaihara N. Novel method for quantifying α (1 \rightarrow 3)-L-fucosyltransferase activity in serum. *Clin Chem.* 1991; 37:2081–6. PMID: [1764783](#)
54. Hada T, Fukui K, Ohno M, Akamatsu S, Yazawa S, Enomoto K, et al. Increased plasma α (1 \rightarrow 3)-L-fucosyltransferase activities in patients with hepatocellular carcinoma. *Glycoconjugate J.* 1995; 12:627–31.
55. Yazawa S, Asao T, Izawa H, Miyamoto Y, Matta KL. The presence of CA19-9 in serum and saliva from Lewis blood-group negative cancer patients. *Jpn J Cancer Res.* 1988; 79:538–43. PMID: [3133342](#)
56. Yazawa S, Madiyalakan R, Jain RK, Shimoda N, Matta KL. Use of benzyl 2-acetamido-2-deoxy-3-*O*-(2-*O*-methyl- β -D-galactosyl)- β -D-glucopyranoside [2'-*O*-methylacto-*N*-biose I β Bn] as a specific acceptor for GDP-fucose: *N*-acetyl-glucosaminide α (1 \rightarrow 4)-L-fucosyltransferase. *Anal Biochem.* 1990; 187:374–8. PMID: [2382836](#)
57. Yazawa S, Nishihara S, Iwasaki H, Asao T, Nagamachi Y, Matta KL, et al. Genetic and enzymatic evidence for Lewis enzyme expression in Lewis-negative cancer patients. *Cancer Res.* 1995; 55:1473–8. PMID: [7882355](#)
58. Kubishiro K, Tsukazaki K, Sakuma Y, Sakayori M, Yazawa S, Nozawa S. Enzymatic basis for the accumulation of Lewisb antigen in uterine endometrial cancer. *Jpn J Cancer Res.* 1995; 86:361–367. PMID: [7775258](#)
59. Yazawa S, Kochibe N, Nishimura T, Shima C, Takai I, Adachi M, et al. A novel method for determination of α 1,6fucosyltransferase activity using a reducing oligosaccharide from egg yolk as a specific acceptor. *Glycoconjugate J.* 1998; 15:863–71.
60. Asao T, Kuwano H, Nakamura J, Okamura A, Berger EG, Matta KL, et al. Tumor cells as the origin of elevated serum α 1,3fucosyltransferase in association with malignancy. *Clin Exp Metastasis.* 2001; 18:605–10.
61. Mollicone R, Reguigne I, Fletcher A, Aziz A, Rustam M, Weston BW, et al. Molecular basis for plasma α (1,3)-fucosyltransferase gene deficiency (FUT6). *J Biol Chem.* 1994; 269:12662–71. PMID: [8175676](#)
62. Hansen JE, Larson VA, Bøg-Hansen TC. The microheterogeneity of alpha 1-acid glycoprotein in inflammatory lung disease, cancer of the lung and normal health. *Clin Chim Acta.* 1984; 27:41–7.
63. Hansen JE, Iversen J, Lihme A, Bøg-Hansen TC. Acute phase reaction, heterogeneity, and microheterogeneity of serum proteins as nonspecific tumor markers in lung cancer. *Cancer* 1987; 60:1630–5. PMID: [2441848](#)
64. Dobryszcka W, Katnik I. Interaction of haptoglobin with concanavalin A and wheat germ agglutinin: basis research and clinical application. In: Breborowicz J, Mackiewicz A, editors. *Affinity electrophoresis: principles and application.* Boca Raton: CRC Press; 1992. p. 211–5.

65. Kushner I, Mackiewicz A. Acute phase proteins. In: Mackiewicz A, Kushmer I, Baumann H, editors. *Molecular biology, biochemistry and clinical applications*. Boca Raton: CRC Press; 1993. p. 4–19.
66. Mackiewicz A, Mackiewicz K. Glycoforms of serum alpha 1-acid glycoprotein as markers of inflammation and cancer. *Glycoconjugate J*. 1995; 12:241–7.
67. Van Dijk W, Brinkmann-Vander Linden ECM, Navenaar EC. Glycosylation of α 1- acid glycoprotein (orosomucoid) in healthy and diseases: occurrence, regulation and possible functional implications. *TIGG*. 1998; 10:235–45.
68. Kratz E, Poland DCW, Van Dijk W, Katnik-Prastowska I. Alterations of branching and differential expression of sialic acid on alpha-1-aid glycoprotein in human seminal plasma. *Clin Chim Acta*. 2003; 331:87–95. PMID: [12691868](#)
69. Olewicz-Gawlik A, Korczowska-Lacka I, Lacki JK, Klama K, Hrycaj P. Fucosylation of serum α 1-acid glycoprotein in rheumatoid arthritis patients treated with infliximab. *Clin Rheumatol*. 2007; 26:1679–84. PMID: [17310270](#)
70. Mondal G, Chatterjee U, Das HR, Chatterjee BP. Enhanced expression of α 1-acid glycoprotein and fucosylation in hepatitis B patients provides an insight into pathogenesis. *Glycoconjugate J*. 2009; 26:1225–34.
71. Ongay S, Martin-Alvarez PJ, Neustüss C, de Frutos M. Statistical evaluation of CZE-UV and CZE-ESI-MS data of intact α -1-acid glycoprotein isoforms for their use as potential biomarkers in bladder cancer. *Electrophoresis*. 2010; 31:3314–25. PMID: [22216449](#)
72. Ahn YH, Shin PM, Oh NR, Park GW, Kim H, Yoo JS. A lectin-coupled, targeted proteomic mass spectrometry (MRM MS) platform for identification of multiple liver cancer biomarkers in human plasma. *J Proteomics*. 2012; 75:5507–5515. doi: [10.1016/j.jprot.2012.06.027](#) PMID: [22789673](#)
73. Ferens-Sieczkowska M, Kratz EM, Kossowska B, Passowicz-Muszyńska E, Jankowska R. Comparison of haptoglobin and alpha₁-acid glycoprotein glycosylation in the sera of small cell and non-small cell lung cancer patients. *Postepy Hig Med Dosw*. 2013; 67:828–36.
74. Ahn YH, Ji ES, Oh NR, Kim YS, Ko JH, Yoo JS. Differential proteomic approach for identification and verification of aberrantly glycosylated proteins in adenocarcinoma lung cancer (ADLC) plasmas by lectin-capturing and targeted mass spectrometry. *J Proteomics*. 2014; 106:221–9. doi: [10.1016/j.jprot.2014.04.031](#) PMID: [24780727](#)
75. Brinkman-Van der Linden EC, Millicone R, Oriol R, Larson G, Van den Eijnden DH, Van Dijk W. A missense mutation in the *FUT6* gene results in total absence of α 3-fucosylation of human α -1-acid glycoprotein. *J Biol Chem*. 1996; 271:14492–5. PMID: [8662894](#)
76. Litvak A, Cercek A, Segal N, Reidy-Lagunes D, Stadler ZK, Yaeger RD, et al. False-positive elevations of carcinoembryonic antigen in patients with a history of resected colorectal cancer. *J Natl Compr Cancer Netw*. 2014; 12:907–13.
77. Yazawa S, Asao T, Izawa H, Miyamoto Y, Matta KL. The presence of CA19-9 in serum and saliva from Lewis blood-group negative cancer patients. *Jpn J Cancer Res*. 1988; 79:538–43. PMID: [3133342](#)
78. Chiu KM, Mortensen RF, Osmand AP, Gewurz H. Interaction of alpha₁-acid glycoprotein with the immune system. I. Purification and effects upon lymphocytes responsiveness. *Immunology*. 1977; 32:997–1005. PMID: [142068](#)
79. Kremer JMH, Wiling J, Janssen LHM. Drug binding to human alpha-1-acid glycoprotein in health and disease. *Pharmacol Rev*. 1988; 40:1–47. PMID: [3064105](#)
80. Libert C, Brouckaert P, Fiers W. Protection by α -1-acid glycoprotein against tumor necrosis factor-induced lethality. *J Exp Med*. 1994; 180:1571–5. PMID: [7931089](#)
81. Haston JL, FitzGerald O, Kane D, Smith KD. The influence of α -1-acid glycoprotein on collagenase-3 activity in early rheumatoid arthritis. *Biomed Chromatogr*. 2003; 17:361–4. PMID: [13680845](#)
82. Poland DCW, Garcia Vallejo JJ, Niessen HWM, Nijmeyer R, Calafat J, Hack CE, et al. Activated human PMN synthesize and release a strongly fucosylated glycoform of α -1-acid glycoprotein, which is transiently deposited in human myocardial infarction. *J Leukoc Biol*. 2005; 78:453–61. PMID: [15647324](#)
83. Chavan MM, Kawle PD, Mehta NG. Increased sialylation and defucosylation of plasma proteins are early events in the acute phase response. *Glycobiology*. 2005; 15:838–48. PMID: [15858075](#)
84. Levander L, Gunnarsson P, Grenegård M, Rydén I, Pålsson P. Effects of α 1-acid glycoprotein fucosylation on its Ca²⁺ mobilizing capacity in neutrophils. *Scand J Immunol*. 2009; 69:412–20. doi: [10.1111/j.1365-3083.2009.02240.x](#) PMID: [19508372](#)
85. Poland DCW, Kulik W, van Dijk W, Hallemeesch MM, Jakobs C, de Meer K. Distinct glycoforms of human α -1-acid glycoprotein have comparable synthesis rates: a [¹³C]valine-labelling study in healthy humans. *Glycoconjugate J*. 2004; 20:99–105.

86. Yazawa S, Furukawa K, Kochibe N. Isolation of fucosyl glycoproteins from human erythrocyte membranes by affinity chromatography using *Aleuria aurantia* lectin. *J Biochem.* 1984; 96:1737–42. PMID: [6530394](#)
87. Yazawa S, Kochibe N, Asao T. A simple procedure for isolation of tumor-associated antigens by affinity chromatography using fucose-specific *Aleuria aurantia* lectin. *Immunol Invest.* 1990; 19:319–27. PMID: [2210805](#)
88. Yamashita K, Kochibe N, Ohkura T, Ueda I, Kobata A. Fractionation of L-fucose-containing oligosaccharides on immobilized *Aleuria aurantia* lectin. *J Biol Chem.* 1985; 260:4688–93. PMID: [3988732](#)
89. Comunale MA, Wang M, Hafner J, Krakover J, Rodemich L, Kopenhaver B, et al. Identification and development of fucosylated glycoproteins as biomarker of primary hepatocellular carcinoma. *J Proteome Res.* 2009; 8:595–602. doi: [10.1021/pr800752c](#) PMID: [19099421](#)
90. Wu J, Xie X, Liu Y, He J, Benitez R, Buckanovich RJ, et al. Identification and confirmation of differentially expressed fucosylated glycoproteins in the serum of ovarian cancer patients using a lectin array and LC-MS/MS. *J Proteome Res.* 2012; 11:4541–52. doi: [10.1021/pr300330z](#) PMID: [22827608](#)
91. Selvaraju S, Rassi ZE. Targeting human serum fucose by an integrated liquid-phase multicolumn platform operating in "cascade" to facilitate comprehensive mass spectrometric analysis of disease-free and breast cancer sera. *Proteomics.* 2013; 13:1701–13. doi: [10.1002/pmic.201200524](#) PMID: [23533108](#)
92. Liu L, Yan B, Huang J, Gu Q, Wang L, Fang M, et al. The identification and characterization of novel N-glycan-based biomarkers in gastric cancer. *PLoS ONE* 2013; 8:e77821, 1–11.