

研究成果の刊行に関する一覧表

雑誌

	発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
1	Asayama K, Satoh M, Murakami Y, Ohkubo T, Nagasawa SY, Tsuji I, Nakayama T, Okayama A, Miura K, Imai Y, Ueshima H, Okamura T.	Cardiovascular risk with and without antihypertensive drug treatment in the Japanese general population: participant-level meta-analysis.	Hypertension.	63	1189-97	2014
2	Satoh M, Ohkubo T, Asayama K, Murakami Y, Sakurai M, Nakagawa H, Iso H, Okayama A, Miura K, Imai Y, Ueshima H, Okamura T.	Combined effect of blood pressure and total cholesterol levels on long-term risks of subtypes of cardiovascular death: evidence for cardiovascular prevention from observational cohorts in Japan.	Hypertension.	65	517-24	2015
3	Nakamura K, Nakagawa H, Murakami Y, Kitamura A, Kiyama M, Sakata K, Tsuji I, Miura K, Ueshima H, Okamura T.	Smoking increases the risk of all-cause and cardiovascular mortality in patients with chronic kidney disease.	Kidney Int.	88	1144-52	2015
4	Zhang W, Iso H, Murakami Y, Miura K, Nagai M, Sugiyama D, Ueshima H, Okamura T; EPOCH-JAPAN GROUP.	Serum Uric Acid and Mortality Form Cardiovascular Disease: EPOCH-JAPAN Study.	J Atheroscler Thromb.	23	692-703	2016
5	Li Y, Iso H, Cui R, Murakami Y, Yatsuya H, Miura K, Nagasawa S, Ueshima H, Okamura T; EPOCH- JAPAN Research Group.	Serum γ -glutamyltransferase and Mortality due to Cardiovascular Disease in Japanese Men and Women.	J Atheroscler Thromb.	23(7)	792-99	2016
6	Nakai M, Miyamoto Y, Higashiyama A, Murakami Y, Nishimura K, Yatsuya H, Saitoh S, Sakata K, Iso H, Miura K, Ueshima H, Okamura T; EPOCH-JAPAN Research Group.	Calibration between the Estimated Probability of the Risk Assessment Chart of Japan Atherosclerosis Society and Actual Mortality Using External Population: Evidence for Cardiovascular Prevention from Observational Cohorts in Japan (EPOCH-JAPAN).	J Atheroscler Thromb.	23(2)	176-95	2016
7	Hirakawa Y, Ninomiya T, Kiyohara Y, Murakami Y, Saitoh S, Nakagawa H, Okayama A, Tamakoshi A, Sakata K, Miura K, Ueshima H, Okamura T; EPOCH-JAPAN Research Group.	Age-specific impact of diabetes mellitus on the risk of cardiovascular mortality: An overview from the Evidence for Cardiovascular Prevention from Observational Cohorts in the Japan Research Group (EPOCH-JAPAN).	J Epidemiol.	27(3)	123-29	2017
8	Hirata T, Sugiyama D, Nagasawa SY, Murakami Y, Saitoh S, Okayama A, Iso H, Irie F, Sairenchi T, Miyamoto Y, Yamada M, Ishikawa S, Miura K, Ueshima H, Okamura T.	A pooled analysis of the association of isolated low levels of high-density lipoprotein cholesterol with cardiovascular mortality in Japan.	Eur J Epidemiol.		in press	2016
9	Nagai M, Murakami Y, Tamakoshi A, Kiyohara Y, Yamada M, Ukawa S, Hirata T, Tanaka S, Miura K, Ueshima H, Okamura T; EPOCH-JAPAN Research Group.	Fasting but not casual blood glucose is associated with pancreatic cancer mortality in Japanese:EPOCH-JAPAN.	Cancer Causes Control		in press	2017

10	村上義孝	特集 わが国における脳・心血管疾患予防のための研究 EPOCH-JAPAN	呼吸と循環	64(1)	57-63	2016
11	Kubota Y, Higashiyama A, Imano H, Sugiyama D, Kawamura K, Kadota A, Nishimura K, Miyamatsu N, Miyamoto Y, Okamura T.	Serum Polyunsaturated Fatty Acid Composition and Serum High-Sensitivity C-Reactive Protein Levels in Healthy Japanese Residents: The KOBE Study.	J Nutr Health Aging.	19	719-28	2015
12	Hirata T, Higashiyama A, Kubota Y, Nishimura K, Sugiyama D, Kadota A, Nishida Y, Imano H, Nishikawa T, Miyamatsu N, Miyamoto Y, Okamura T.	HOMA-IR values are associated with glycemic control in Japanese subjects without diabetes or obesity: the KOBE Study.	J Epidemiol.	25	407-14	2015
13	Tatsumi Y, Higashiyama A, Kubota Y, Sugiyama D, Nishida Y, Hirata T, Kadota A, Nishimura K, Imano H, Miyamatsu N, Miyamoto Y, Okamura T.	Underweight young women without later weight gain are at high risk for osteopenia after midlife: the KOBE study.	J Epidemiol.	26(11)	572-78	2016
14	西田陽子、原田 成、武林亨、岡村智教	新しいコホート研究の立ち上げと今後の展望：神戸研究と鶴岡メタボロームコホート研究	呼吸と循環	64	71-77	2016
15	Kuwabara K, Harada S, Sugiyama D, Kurihara A, Kubota Y, Higashiyama A, Hirata T, Nishida Y, Kawasaki M, Takebayashi T, Okamura T.	Relationship between non-high-density lipoprotein cholesterol and low-density lipoprotein cholesterol in the general population.	J Atheroscler Thromb.	23(4)	477-90	2016
16	Hirata T, Higashiyama A, Kubota Y, Sugiyama D, Kuwabara K, Tatsumi Y, Hirata A, Nishida Y, Kadota A, Imano H, Nishikawa T, Miyamatsu N, Miyamoto Y, Okamura T.	Impact of flushing response on the relationship between alcohol consumption and gamma-glutamyl transpeptidase: the KOBE study.	Nihon Arukoru Yakubutsu Igakkai Zasshi.	51	173-83	2016
17	Fujihara K, Sugawara A, Heianza Y, Sairenchi T, Irie F, Iso H, Doi M, Shimano H, Watanabe H, Sone H, Ota H.	Utility of the triglyceride level for predicting incident diabetes mellitus according to the fasting status and body mass index category: the Ibaraki Prefectural Health Study.	J Atheroscler Thromb.	21(11)	1152-69	2014
18	Tsujimoto T, Sairenchi T, Iso H, Irie F, Yamagishi K, Watanabe H, Tanaka K, Muto T, Ota H.	The dose-response relationship between body mass index and the risk of incident stage ≥ 3 chronic kidney disease in a general Japanese population: the Ibaraki prefectural health study (IPHS).	J Epidemiol.	24(6)	444-51	2014
19	Xu D, Murakoshi N, Sairenchi T, Irie F, Igarashi M, Nogami A, Tomizawa T, Yamaguchi I, Yamagishi K, Iso H, Ota H, Aonuma K.	Anemia and Reduced Kidney Function as Risk Factors for New Onset of Atrial Fibrillation (from the Ibaraki Prefectural Health Study).	Am J Cardiol.	115(3)	328-33	2015
20	Murakoshi N, Xu D, Sairenchi T, Igarashi M, Irie F, Tomizawa T, Tada H, Sekiguchi Y, Yamagishi K, Iso H, Yamaguchi I, Ota H, Aonuma K.	Prognostic impact of supraventricular premature complexes in community-based health checkups: The Ibaraki Prefectural Health Study.	Eur Heart J.	36(3)	170-78	2015

21	Nagai K, Sairenchi T, Irie F, Watanabe H, Ota H, Yamagata K.	Relationship between Estimated Glomerular Filtration Rate and Cardiovascular Mortality in a Japanese Cohort with Long-Term Follow-Up.	PLoS One	11(6)	e0156792	2016
22	Tsukinoki R, Okamura T, Watanabe M, Kokubo Y, Higashiyama A, Nishimura K, Takegami M, Murakami Y, Okayama A, Miyamoto Y.	Blood pressure, low-density lipoprotein cholesterol, and incidences of coronary artery disease and ischemic stroke in Japanese: the Suita study.	Am J Hypertens.	27(11)	1362-69	2014
23	Nishimura K, Okamura T, Watanabe M, Nakai M, Takegami M, Higashiyama A, Kokubo Y, Okayama A, Miyamoto Y.	Predicting coronary heart disease using risk factor categories for a Japanese urban population, and comparison with the framingham risk score: the suita study.	J Atheroscler Thromb.	21(8)	784-98	2014
24	Kosaka T, Kokubo Y, Ono T, Sekine S, Kida M, Kikui M, Yamamoto M, Watanabe M, Amano A, Maeda Y, Miyamoto Y.	Salivary inflammatory cytokines may be novel markers of carotid atherosclerosis in a Japanese general population: the Suita study.	Atherosclerosis.	237(1)	123-28	2014
25	Sakamoto Y, Kokubo Y, Toyoda K, Watanabe M, Tanigawa T, Miyamoto Y.	Sleep-disordered breathing is associated with elevated human atrial natriuretic peptide levels in a Japanese urban population: the Suita study.	Int J Cardiol.	173(2)	334-35	2014
26	Kosaka T, Ono T, Yoshimuta Y, Kida M, Kikui M, Nokubi T, Maeda Y, Kokubo Y, Watanabe M, Miyamoto Y	The effect of periodontal status and occlusal support on masticatory performance: the Suita study.	J Clin Periodontol.	41(5)	497-503	2014
27	Kokubo Y, Watanabe M, Higashiyama A, Nakao YM, Kobayashi T, Watanabe T, Okamura T, Okayama A, Miyamoto Y.	Interaction of Blood Pressure and Body Mass Index With Risk of Incident Atrial Fibrillation in a Japanese Urban Cohort: The Suita Study.	Am J Hypertens.	28	1355-61	2015
28	Miyamoto K, Aiba T, Arihiro S, Watanabe M, Kokubo Y, Ishibashi K, Hirose S, Wada M, Nakajima I, Okamura H, Noda T, Nagatsuka K, Noguchi T, Anzai T, Yasuda S, Ogawa H, Kamakura S, Shimizu W, Miyamoto Y, Toyoda K, Kusano K.	Impact of renal function deterioration on adverse events during anticoagulation therapy using non-vitamin K anta-gonist oral anticoa-gulants in patients with atrial fibrillation.	Heart Vessels.	31(8)	1327-36	2016
29	Maruyama K, Kokubo Y, Yamanaka T, Watanabe M, Iso H, Okamura T, Miyamoto Y.	The reasonable reliability of a self-administered food frequency questionnaire for an urban, Japanese, middle-aged population: the Suita study.	Nutr Res.	35(1)	14-22	2015
30	Kawase Ishihara K, Kokubo Y, Yokota C, Hida E, Miyata T, Toyoda K, Matsumoto M, Minematsu K, Miyamoto Y.	Effect of Plasma Fibrinogen, High-Sensitive C-Reactive Protein, and Cigarette Smoking on Carotid Atherosclerosis: The Suita Study.	J Stroke Cerebrovasc Dis.	24	2385-89	2015
31	Tatsumi Y, Watanabe M, Nakai M, Kokubo Y, Higashiyama A, Nishimura K, Kobayashi T, Takegami M, Nakao YM, Watanabe T, Okayama A, Okamura T, Miyamoto Y.	Changes in Waist Circumference and the Incidence of Type 2 Diabetes in Community-Dwelling Men and Women: The Suita Study.	J Epidemiol.	25	489-95	2015

32	Turin TC, Okamura T, Raheen Afzal A, Rumana N, Watanabe M, Higashiyama A, Nakao YM, Nakai M, Takegami M, Nishimura K, Kokubo Y, Okayama A, Miyamoto Y.	Impact of hypertension on the lifetime risk of coronary heart disease.	Hypertens Res.	39(7)	548-51	2016
33	Turin TC, Okamura T, Afzal AR, Rumana N, Watanabe M, Higashiyama A, Nakao Y, Nakai M, Takegami M, Nishimura K, Kokubo Y, Okayama A, Miyamoto Y.	Hypertension and lifetime risk of stroke.	J Hypertens.	34(1)	116-22	2016
34	Yokota C, Sawamura T, Watanabe M, Kokubo Y, Fujita Y, Kakino A, Nakai M, Toyoda K, Miyamoto Y, Minematsu K.	High Levels of Soluble Lectin-Like Oxidized Low-Density Lipoprotein Receptor-1 in Acute Stroke: An Age- and Sex-Matched Cross-Sectional Study.	J Atheroscler Thromb.	23(10)	1222-26	2016
35	Kosaka T, Ono T, Kida M, Kikui M, Yamamoto M, Yasui S, Nokubi T, Maeda Y, Kokubo Y, Watanabe M, Miyamoto Y.	A multifactorial model of masticatory performance: the Suita study.	J Oral Rehabil.	43(5)	340-47	2016
36	Kikui M, Kokubo Y, Ono T, Kida M, Kosaka T, Yamamoto M, Watanabe M, Maeda Y, Miyamoto Y.	Relationship between Metabolic Syndrome Components and Periodontal Disease in a Japanese General Population: the Suita Study.	J Atheroscler Thromb.		in press	2016
37	Kikui M, Ono T, Kokubo Y, Kida M, Kosaka T, Yamamoto M, Nokubi T, Watanabe M, Maeda Y, Miyamoto Y.	Relationship between metabolic syndrome and objective masticatory performance in a Japanese general population: The Suita study.	J Dent.	56	53-57	2017
38	Gotoh S, et al.	Trends in the incidence and survival of intracerebral hemorrhage by its location in a Japanese community.	Circ J.	78	403-9	2014
39	Imamura T, et al.	Non-high-density lipoprotein cholesterol and the development of coronary heart disease and stroke subtypes in a general Japanese population: the Hisayama Study.	Atherosclerosis.	233	343-48	2014
40	Kojima I, et al.	A low ankle brachial index is associated with an increased risk of cardiovascular disease: the Hisayama Study.	J Atheroscler Thromb.	21	966-73	2014
41	Kishimoto H, et al.	Midlife and late-life handgrip strength and risk of cause-specific death in a general Japanese population: the Hisayama Study.	J Epidemiol Community Health.	68	663-68	2014
42	Mukai N, et al.	Trends in the prevalence of type 2 diabetes and prediabetes in community-dwelling Japanese subjects: the Hisayama Study.	J Diabetes Investig.	5	162-69	2014
43	Mukai N, et al.	Thresholds of various glycemic measures for diagnosing diabetes based on prevalence of retinopathy in community-dwelling Japanese subjects: the Hisayama Study.	Cardiovasc Diabetol.	13	45	2014

44	Izumaru K, et al.	Serum 1, 25-dihydroxyvitamin D and the development of kidney dysfunction in a Japanese community: the Hisayama Study.	Circ J.	78	732-37	2014
45	Mukai N, et al.	Association of hemoglobin A1c and glycated albumin with carotid atherosclerosis in community-dwelling Japanese subjects: the Hisayama Study.	Cardiovasc Diabetol.	14	84	2015
46	Gotoh S, et al.	Hematocrit and the risk of cardiovascular disease in a Japanese community: the Hisayama Study.	Atherosclerosis.	242	199-204	2015
47	Ohara T, et al.	Midlife and late-life smoking and risk of dementia in the community: the Hisayama Study.	J Am Geriatr Soc.	63	2332-39	2015
48	Kondo H, et al.	Angiotensin I-converting enzyme gene polymorphism enhances the effect of hypercholesterolemia on the risk of coronary heart disease in a general Japanese population: the Hisayama Study.	J Atheroscler Thromb.	22	390-403	2015
49	Takae K, et al.	Serum uric acid as a risk factor for chronic kidney disease in a Japanese community - the Hisayama Study -	Circ J.	80	1857-62	2016
50	Hata J, et al.	Serum angiopoietin-like protein 2 is a novel risk factor for cardiovascular disease in the community: the Hisayama Study.	Arterioscler Thromb Vasc Biol.	36	1686-91	2016
51	Hirabayashi N, et al.	Association between diabetes and hippocampal atrophy in elderly Japanese: the Hisayama Study.	Diabetes Care.	39	1543-49	2016
52	Kishimoto H, et al.	The long-term association between physical activity and risk of dementia in the community: the Hisayama Study.	Eur J Epidemiol.	31	267-74	2016
53	Ikeda F, et al.	Combination of Helicobacter pylori antibody and serum pepsinogen as a good predictive tool of gastric cancer incidence: 20-Year prospective data from the Hisayama Study.	J Epidemiol.	26	629-36	2016
54	立川佳美、Cologne JB、山田美智子、大石和佳、飛田あゆみ、古川恭治、高橋規郎、中村 典、小笹晃太郎、赤星正純、藤原佐枝子、Shore RE	親の放射線被ばくと多因子疾患有病率との関連：被爆二世健康診断調査（第2報）	広島医学	67(4)	375-78	2014
55	Yamada M, Landes RD, Mimori Y, Nagano Y, Sasaki H.	Trajectories of cognitive function in dementia-free subjects: Radiation Effects Research Foundation Adult Health Study.	J Neurol Sci.	351	115-19	2015
56	Imaizumi M, Ohishi W, Nakashima E, Sera, N, Neriishi, K, Yamada, M, Tatsukawa Y, Takahashi I, Fujiwara S, Sugino K, Ando T, Usa T, Kawakami A, Akahoshi M, Hida A.	Association of radiation dose with prevalence of thyroid nodules among atomic bomb survivors exposed in childhood (2007-2011).	JAMA Intern Med.	175	228-36	2015
57	Yamada M, Landes RD, Mimori Y, Nagano Y, Sasaki H.	Radiation effects on cognitive function among atomic bomb survivors exposed at or after adolescence.	Am J Med.	129	586-91	2016

58	Izumi S, Sakata R, Yamada M, Cologne J.	Interaction between a single exposure and age in cohort-based hazard rate models impacted the statistical distribution of age at onset.	J Clin Epidemiol.	71	43-50	2016
59	斎藤重幸	高血圧	最新肥満症学 日本臨床増刊号	72	583-85	2014
60	斎藤重幸	JNC8 ガイドライン	血圧	21	496-97	2014
61	斎藤重幸	肥満症・メタボリックシンドローム、 高血圧発症・進展の関わり	臨床高血圧学 日本臨床	72	269-72	2014
62	斎藤重幸	日本高血圧学会高血圧治療ガイドラ イン (JSH) の変遷	循環器内科	14	385-91	2014
63	斎藤重幸	糖尿病合併高血圧の降圧目標をめぐ って	血圧	21	926-31	2014
64	Yoshida Y, Akasaka H, Saitoh, S, Shimamoto K, Miura T.	Comparative effects of telmisartan and valsartan as add-on agents for hypertensive patients with morning blood pressure insufficiency controlled by amlodipine monotherapy.	Hypertens Res.	37	225-31	2014
65	Furuhashi M, Moniwa N, Mita T, Fuseya T, Ishimura S, Ohno K, Shibata S, Tanaka M, Watanabe Y, Akasaka H, Ohnishi H, Yoshida H, Takizawa H, Saitoh S, Ura N, Shimamoto K, T Miura.	Urinary angiotensin-converting enzyme 2 in hypertensive patients may be increased by Olmesartan, an angiotensin II Receptor Blocker.	Am J Hypertens.	28(1)	15-21	2015
66	Ihira H, Furuna T, Mizumoto A, Makino K, Saitoh S, Ohnishi H, Shimada H, Makizako H.	Subjective physical and cognitive age among community-dwelling older peoples aged 75 years and Older: differences with chronological age and its associated factors.	Aging Ment Health.	19(8)	756-61	2014
67	Fuseya T, Furuhashi M, Yuda S, Muranaka A, Kawamukai M, Mita T, Ishimura S, Watanabe Y, Hoshina K, Tanaka M, Ohno K, Akasaka H, Ohnishi H, Yoshida H, Saitoh S, Shimamoto K, Miura T.	Elevation of circulating fatty acid-binding protein 4 is independently associated with left ventricular diastolic dysfunction in a general population.	Cardiovasc Diabetol.	http://ww w.cardiab. com/cont ent/13/1/12 6	1-9	2014
68	Miki T, Tobisaw T, Sato T, Tanno M, Yano T, Akasaka H, Kuno A, Ogasawara M, Murase H, Saitoh S, Miura T.	Does glycemic control reverse dispersion of ventricular repolarization in type 2 diabetes?	Cardiovasc Diabetol.	http://ww w.cardiab. com/cont ent/13/1/12 5	1-10	2014
69	斎藤重幸	高血圧治療における RAS 抑制薬の 位置づけ - わが国のガイドラインの 立場から	Angiotensin Research.	12	17-21	2015
70	斎藤重幸	高血圧	Mordan Physician.	35(2)	213-15	2015
71	斎藤重幸	JSH2009 から JSH2014 への変更の概 要	プラクティス	32	155-61	2015
72	斎藤重幸、赤坂 憲、大西 浩文、島本和明、甲斐久史	JSH2014 理解と応用 実地医家への アンケート調査より	血圧	22	893-97	2015

73	齋藤重幸	どうして高血圧を治療しなければならないのか	循環器ナーシング	6	23-31	2016
74	Furuhashi M, Saitoh S, Shimamoto K, Miura T.	Fatty Acid-Binding Protein 4 (FABP4): Pathophysiological Insights and Potent Clinical Biomarker of Metabolic and Cardiovascular Diseases. Clinical Medicine Insights: Cardiology. Supplementary Issue: Inflammation, Atherosclerosis and Coronary Artery Disease.	Clin Med Insights Cardiol.	8(13)	23-33	2015
75	Midori Sato M, Ohnishi H, Ohnishi-Kameyama M, Matsunaga I, Naito S, Ippoushi K, Oike H, Nagata T, Akasaka H, Saitoh S, Shimamoto K, Kobori M.	Estimated Daily Intake and Seasonal Food Sources of Quercetin in Japan.	Nutrients.	7	2345-58	2015
76	Fujii M, Ohnishi H, Saitoh S, Akasaka H, Miura T, Mori M	The combination of abdominal obesity and high-sensitivity C-reactive protein predicts new-onset hypertension in the general Japanese population: the Tanno-Sobetsu study.	Hypertens Res.	38	426-32	2015
77	齋藤重幸	体質を考慮した高血圧の予防と治療	日本体質学会雑誌	78	32-36	2016
78	鷲尾昌一、田中裕士、猪股慎一郎、高橋弘毅、齋藤重幸、三浦哲嗣、島本和明、大西浩文、森 満	喫煙は肺がんのリスクを上昇させ、緑茶と緑黄色野菜はリスクを低下させる 北海道民を対象とした症対象研究の結果より	臨床と研究	93	93-96	2016
79	Ohnishi H, Saitoh S, Akasaka H, Furukawa T, Mori M, Miura T.	Combination of type 2 diabetes and smoking increases total cancer mortality in Japanese men using competing risk analysis: the Tanno-Sobetsu study.	Diabetol Int.	7	167-72	2016
80	Ohnishi H, Saitoh S, Akasaka H, Furukawa T, Mori M, Miura T.	Impact of longitudinal status change in metabolic syndrome defined by two different criteria on new onset of type 2 diabetes in a general Japanese population: the Tanno-Sobetsu Study.	Diabetol Metab Syndr.	8	1-9	2016
81	Wang C, Yatsuya H, Tamakoshi K, Iso H, Tamakoshi A.; JACC Study Group.	Milk drinking and mortality: findings from the Japan collaborative cohort study.	J Epidemiol.	25	66-73	2015
82	Ikehara S, Iso H, Wada Y, Tanabe N, Watanabe Y, Kikuchi S, Tamakoshi A; JACC Study Group.	Television Viewing Time and Mortality From Stroke and Coronary Artery Disease Among Japanese Men and Women - The Japan Collaborative Cohort Study.	Circ J	79	2389-95	2015
83	Shirakawa T, Iso H, Yamagishi K, Yatsuya H, Tanabe N, Ikehara S, Ukawa S, Tamakoshi A.	Watching television and risk of mortality from pulmonary embolism among Japanese men and women: the JACC study (Japan collaborative cohort).	Circulation.	134	355-57	2016
84	Hayashi R, Iso H, Cui R, Tamakoshi A; JACC Study Group.	Occupational physical activity in relation to risk of cardiovascular mortality: The Japan Collaborative Cohort Study for Evaluation for Cancer Risk (JACC Study).	Prev Med.	89	286-91	2016

85	Umesawa M, Iso H, Fujino Y, Kikuchi S, Tamakoshi A; JACC Study Group.	Salty Food Preference and Intake and Risk of Gastric Cancer: The JACC Study.	J Epidemiol.	26	242-28	2016
86	Kumagai Y, Chou WT, Tomata Y, Sugawara Y, Kakizaki M, Nishino Y, Tsuji I.	Dietary Patterns and colorectal cancer risk in Japan: the Ohsaki Cohort Study.	Cancer Causes Control.	25	727-36	2014
87	Sugiyama K, Tomata Y, Kaiho Y, Honkura K, Sugawara Y, Tsuji I.	Association between Coffee Consumption and Incident Risk of Disabling Dementia in Elderly Japanese: The Ohsaki Cohort 2006 Study.	J Alzheimers Dis.	50(2)	491-500	2016
88	Honkura K, Tomata Y, Sugiyama K, Kaiho Y, Watanabe T, Zhang S, Sugawara Y, Tsuji I.	Defecation frequency and cardiovascular disease mortality in Japan: the Ohsaki cohort study.	Atherosclerosis.	246	251-56	2016
89	Zhang S, Tomata Y, Sugiyama K, Kaiho Y, Honkura K, Watanabe T, Tanji F, Sugawara Y, Tsuji I.	Body mass index and the risk of incident functional disability in elderly Japanese: The OHSAKI Cohort 2006 Study.	Medicine.	95(31)	e4452	2016
90	Tomata Y, Sugiyama K, Kaiho Y, Honkura K, Watanabe T, Zhang S, Sugawara Y, Tsuji I.	Dietary Patterns and Incident Dementia in Elderly Japanese: The Ohsaki Cohort 2006 Study.	J Gerontol A Biol Sci Med Sci.	71(10)	1322-28	2016
91	Tomata Y, Sugiyama K, Kaiho Y, Honkura K, Watanabe T, Zhang S, Sugawara Y, Tsuji I.	Green Tea Consumption and the Risk of Incident Dementia in Elderly Japanese: The Ohsaki Cohort 2006 Study.	Am J Geriatr Psychiatry.	24(10)	881-89	2016
92	Tomata Y, Sugiyama K, Kaiho Y, Sugawara Y, Hozawa A, Tsuji I.	Predictive ability of a simple subjective memory complaints scale for incident dementia: Evaluation of Japan's national checklist, the "Kihon Checklist".	Geriatr Gerontol Int.		in press	2016
93	Sugiyama K, Sugawara Y, Tomata Y, Nishino Y, Fukao A, Tsuji I.	The association between coffee consumption and bladder cancer incidence in a pooled analysis of the Miyagi Cohort Study and Ohsaki Cohort Study.	Eur J Cancer Prev.	26(2)	125-30	2017
94	Zhang S, Tomata Y, Sugiyama K, Sugawara Y, Tsuji I.	Mushroom consumption and incident dementia in elderly Japanese: The Ohsaki Cohort 2006 Study.	J Am Geriatr Soc.		in press	2017
95	Kaiho Y, Sugawara Y, Sugiyama K, Tomata Y, Endo Y, Toyama H, Yamauchi M, Tsuji I.	Impact of Pain on Incident Risk of Disability in Elderly Japanese: Cause-specific Analysis.	Anesthesiology.		in press	2017
96	Matsumoto A, et al.	Day-to-Day variability in home blood pressure is associated with cognitive decline: the Ohasama study.	Hypertension.	63	1333-38	2014
97	Imai E, et al.	Animal protein intake is associated with higher-level functional capacity in elderly adults: the Ohasama study.	J Am Geriatr Soc.	62	426-34	2014
98	Satoh M, et al.	Aldosterone-to-renin ratio and nocturnal blood pressure decline assessed by self-measurement of blood pressure at home: the Ohasama Study.	Clin Exp Hypertens.	36	108-14	2014

99	Tsubota-Utsugi M, et al.	Personality traits as predictors of decline in higher-level functional capacity over a 7-year follow-up in older adults: the Ohasama Study.	Tohoku J Exp Med.	234	197-207	2014
100	Satoh M, et al.	A survey of self-medication practices and related factors in the general population: the Ohasama study.	YAKUGAKU ZASSHI.	134	1347-55	2014
101	Tsubota-Utsugi M, et al.	Living situations associated with poor dietary intake among healthy Japanese elderly: the Ohasama Study.	J Nutr Health Aging.	19	375-82	2015
102	Satoh M, et al.	Association between N-terminal pro B-type natriuretic peptide and day-to-day blood pressure and heart rate variability in a general population: the Ohasama study.	J Hypertens.	33	1536-41	2015
103	Satoh M, et al.	Long-Term Stroke Risk Due to Partial White-Coat or Masked Hypertension Based on Home and Ambulatory Blood Pressure Measurements: The Ohasama Study.	Hypertension.	67	48-55	2016
104	Murakami K, et al.	Impaired higher-level functional capacity as a predictor of stroke in community-dwelling older adults: the Ohasama study.	Stroke.	47	323-28	2016
105	Murakami K, et al.	Menstrual factors and stroke incidence in Japanese postmenopausal women: the Ohasama study.	Neuroepidemiology.	47	109-16	2016
106	Tsubota-Utsugi M, et al.	Lacunar infarcts rather than white matter hyperintensity as a predictor of future higher-level functional decline: The Ohasama Study.	J Stroke Cerebrovasc Dis.	26	376-84	2017
107	Sakurai M, Nakamura K, Miura K, Takamura T, Yoshita K, Nagasawa SY, Morikawa Y, Ishizaki M, Kido T, Naruse Y, Suwazono Y, Sasaki S, Nakagawa H.	Sugar-sweetened beverage and diet soda consumption and the 7-year risk for type 2 diabetes mellitus in middle-aged Japanese men.	Eur J Nutr.	53	251-58	2014
108	Sakurai M, Nakamura K, Miura K, Yoshita K, Takamura T, Nagasawa SY, Morikawa Y, Ishizaki M, Kido T, Naruse Y, Nakashima M, Nogawa N, Suwazono Y, Nakagawa H.	Association between serum thyroid stimulating hormone within the normal range and indices of obesity in Japanese men and women.	Intern Med.	53	669-74	2014
109	Yoneyama S, Sakurai M, Nakamura K, Morikawa Y, Miura K, Nakashima M, Yoshita K, Ishizaki M, Kido T, Naruse Y, Nogawa K, Suwazono Y, Sasaki S, Nakagawa H.	Associations between rice, noodle, and bread intake and sleep quality in Japanese men and women.	PLoS One.	9	e105198	2014
110	Morikawa Y, Nakamura K, Sakurai M, Nagasawa SY, Ishizaki M, Nakashima M, Kido T, Naruse Y, Nakagawa H.	The effect of age on the relationships between work-related factors and heavy drinking.	J Occup Health.	56	141-49	2014

111	Nakamura K, Sakurai M, Miura K, Morikawa Y, Nagasawa SY, Ishizaki M, Kido T, Naruse Y, Nakashima M, Nogawa K, Suwazono Y, Nakagawa H.	HOMA-IR and the risk of hyperuricemia: A prospective study in non-diabetic Japanese men.	Diabetes Res Clin Pract.	106	154-60	2014
112	Nakamura K, Sakurai M, Miura K, Morikawa Y, Nagasawa SY, Ishizaki M, Kido T, Naruse Y, Nogawa K, Nakashikma M, Suwazono Y, Nakagawa H.	Overall sleep status and high sensitivity C-reactive protein: a prospective study in a Japanese population.	J Sleep Res.	23	717-27	2014
113	Sakurai M, Nakamura K, Miura K, Takamura T, Yoshita K, Nagasawa SY, Morikawa Y, Ishizaki M, Kido T, Naruse Y, Nakashima M, Nogawa K, Suwazono Y, Sasaki S, Nakagawa H.	Dietary carbohydrate intake, presence of obesity, and the incident risk of type 2 diabetes in Japanese men.	J Diabetes Invest.	7	343-51	2016
114	Nakamura K, Sakurai M, Morikawa Y, Nagasawa SY, Miura K, Ishizaki M, Kido T, Naruse Y, Nakashikma M, Nogawa K, Suwazono Y, Nakagawa H.	Serum ferritin, insulin resistance, and β -cell dysfunction: a prospective study in normoglycemic Japanese men.	Exp Clin Endocrinol Diabet .	125	12-20	2017
115	Makita S, Onoda T, Ohsawa M, Tanno K, Tanaka F, Omama S, Yoshida Y, Ishibashi Y, Itai K, Sakata K, Ohta M, Kuribayashi T, Ogasawara K, Ogawa A, Okayama A, Nakamura M.	Bradycardia is associated with future cardiovascular diseases and death in men from the general population.	Atherosclerosis.	236	116-20	2014
116	Koeda Y, Tanaka F, Segawa T, Ohta M, Ohsawa M, Tanno K, Makita S, Ishibashi Y, Omama S, Onoda T, Nakamura M.	Usefulness of risk grading system using albuminuria for predicting cardiovascular events and all-cause death in chronic kidney disease: a population-based prospective cohort study in Japan.	Int J Cardiol.	175	576-77	2014
117	Ando A, Tanno K, Ohsawa M, Onoda T, Sakata K, Tanaka F, Makita S, Nakamura M, Omama S, Ogasawara K, Ishibashi Y, Kuribayashi T, Koyama T, Itai K, Ogawa A, Okayama A.	Associations of number of teeth with risks for all-cause mortality and cause-specific mortality in middle-aged and elderly men in the northern part of Japan: the Iwate-KENCO study.	Community Dent Oral Epidemiol.	42	358-65	2014
118	Sato K, Segawa T, Tanaka F, Takahashi T, Tanno K, Ohsawa M, Onoda T, Itai K, Sakata K, Omama S, Ogasawara K, Ishibashi Y, Makita S, Okayama A, Nakamura M.	Cardiovascular risk stratification with plasma B-type natriuretic peptide levels in a community-based hypertensive cohort.	Am J Cardiol.	113	682-86	2014
119	Makita S, Tanaka F, Onoda T, Ohsawa M, Tanno K, Omama S, Yoshida Y, Ishibashi Y, Segawa T, Takahashi T, Satoh K, Itai K, Sakata K, Ohta M, Kuribayashi T, Ogasawara K, Ogawa A, Okayama A, Nakamura M.	Elevated blood pressure at the first measurement predicts cardiovascular disease independently from the subsequent second reading in men, but not in women.	Clin Exp Hypertens.	37	39-44	2015

120	Ohsawa M, Okamura T, Ogasawara K, Ogawa A, Fujioka T, Tanno K, Yonekura Y, Omama S, Turin TC, Itai K, Ishibashi Y, Morino Y, Itoh T, Miyamatsu N, Onoda T, Kuribayashi T, Makita S, Yoshida Y, Nakamura M, Tanaka F, Ohta M, Sakata K, Okayama A.	Relative and absolute risks of all-cause and cause-specific deaths attributable to atrial fibrillation in middle-aged and elderly community dwellers.	Int J Cardiol.	184	692-98	2015
121	Koeda Y, Tanaka F, Segawa T, Ohta M, Ohsawa M, Tanno K, Makita S, Ishibashi Y, Itai K, Omama S, Onoda T, Sakata K, Ogasawara K, Okayama A, Nakamura M.	Comparison between urine albumin-to-creatinine ratio and urine protein dipstick testing for prevalence and ability to predict the risk for chronic kidney disease in the general population (Iwate-KENCO study): a prospective community-based cohort study.	BMC Nephrol.	17	46	2016
122	Tanaka F, Komi R, Makita S, Onoda T, Tanno K, Ohsawa M, Itai K, Sakata K, Omama S, Yoshida Y, Ogasawara K, Ishibashi Y, Kuribayashi T, Okayama A, Nakamura M; Iwate-Kenco Study Group.	Low-grade albuminuria and incidence of cardiovascular disease and all-cause mortality in nondiabetic and normotensive individuals.	J Hypertens.	34	506-12	2016
123	Ohsawa M, Tanno K, Okamura T, Yonekura Y, Kato K, Fujishima Y, Obara W, Abe T, Itai K, Ogasawara K, Omama S, Turin TC, Miyamatsu N, Ishibashi Y, Morino Y, Itoh T, Onoda T, Kuribayashi T, Makita S, Yoshida Y, Nakamura M, Tanaka F, Ohta M, Sakata K, Okayama A.	Standardized Prevalence Ratios for Atrial Fibrillation in Adult Dialysis Patients in Japan.	J Epidemiol.	26	272-76	2016
124	大澤 正樹, 岡山 明	【わが国における脳・心血管疾患予防のための疫学研究】 岩手県北コホート	呼吸と循環	64	37-48	2016
125	Iso H, Imano H, Yamagishi K, Ohira T, Cui R, Noda H, Sato S, Kiyama M, Okada T, Hitsumoto S, Tanigawa T, Kitamura A; CIRCS Investigators.	Fasting and non-fasting triglycerides and risk of ischemic cardiovascular disease in Japanese men and women: the Circulatory Risk in Communities Study (CIRCS).	Atherosclerosis.	237	361-68	2014
126	Chei CL, Yamagishi K, Ikeda A, Noda H, Maruyama M, Cui , Imano H, Kiyama M, Kitamura A, Asada T, Iso H; CIRCS Investigators.	C-reactive protein levels and risk of disabling dementia with and without stroke in Japanese: the Circulatory Risk in Communities Study (CIRCS).	Atherosclerosis.	236	438-43	2014
127	Cui R, Yamagishi K, Imano H, Ohira T, Tanigawa T, Hitsumoto S, Kiyama M, Okada T, Kitamura A, Iso H; CIRCS investigators.	Relationship between the ankle-brachial index and the risk of coronary heart disease and stroke: the circulatory risk in communities study.	J Atheroscler Thromb.	21	1283-89	2014

128	Tanaka A, Cui R, Kitamura A, Liu K, Imano H, Yamagishi K, Kiyama M, Okada T, Iso H; CIRCS Investigators.	Heavy Alcohol Consumption is Associated with Impaired Endothelial Function.	J Diabetes Complications.	30(5)	834-38	2016
129	Hori M, Kitamura A, Kiyama M, Imano H, Yamagishi K, Cui R, Umesawa M, Muraki I, Okada T, Sankai T, Ohira T, Saito I, Tanigawa T, Iso H; CIRCS Investigators.	Fifty-year Time Trends in Blood Pressures, Body Mass Index and their Relations in a Japanese Community: The Circulatory Risk in Communities Study (CIRCS).	J Atheroscler Thromb.		in press	2016
130	Hayama-Terada M, Muraki I, Imano H, Kiyama M, Okada T, Yamagishi K, Cui R, Ohira T, Umesawa M, Sankai T, Sato S, Kitamura A, Iso H; CIRCS Investigators.	Diabetes Trend and Impact on Risk of Cardiovascular Disease in Middle-Aged Japanese People - The CIRCS Study.	Circ J.	80(11)	2343-48	2016
131	Ishikawa J, Ishikawa S, Kario K.	Levels of cornell voltage and cornell product for predicting cardiovascular and stroke mortality and morbidity in the general Japanese population.	Circ J.	78	465-75	2014
132	Takeuchi E, Kotani K, Ishikawa S, Gotoh T, Kayaba K, Nakamura Y, Kajii E.	Relationship between screening plasma glucose concentrations and cancer- and all-cause mortality: the Jichi Medical School (JMS) cohort study.	J Public Health.	22	505-11	2014
133	Yamasaki K, Kayaba K, Ishikawa S.	Soy and Soy Products Intake, All-Cause Mortality, and Cause-Specific Mortality in Japan: The Jichi Medical School Cohort Study.	Asia Pac J Public Health.		doi: 10.1177/1010539514539545.	2014
134	Ishikawa J, Ishikawa S, Kario K.	Prolonged corrected QT interval is predictive of future stroke events even in subjects without ECG-diagnosed left ventricular hypertrophy.	Hypertension.	65	554-60	2015
135	Ishikawa J, Ishikawa S, Kario K.	Relationships between the QTc interval and cardiovascular, stroke, or sudden cardiac mortality in the general Japanese population.	J Cardiol.	65	237-42	2015
136	Sakamaki T, Hara M, Kayaba K, Kotani K, Ishikawa S.	Coffee Consumption and Incidence of Subarachnoid Hemorrhage: The Jichi Medical School Cohort Study.	J Epidemiol.	26	71-75	2016
137	Mieno M. N, Tanaka N, Arai T, Kawahara T, Kuchiba A, Ishikawa S, Sawabe M.	Accuracy of Death Certificates and Assessment of Factors for Misclassification of Underlying Cause of Death.	J Epidemiol.	26	191-98	2016
138	Tadenuma S, Kanda H, Ishikawa S, Kayaba K, Gotoh T, Nakamura Y, Kajii E.	Salt Preference and the Incidence of Cardiovascular Disease in a Japanese General Population: The Jichi Medical School Cohort Study.	Health.	8	105-15	2016
139	Sugiyama T, Ishikawa S, Kotani K, Gotoh T, Itoh Y, Kayaba K, Kajii E.	Relationship Between Serum High-Sensitivity C-Reactive Protein and Myocardial Infarction in a General Japanese Population.	J Clin Lab Anal.	30	999-1002	2016

140	Kawate N, Kayaba K, Hara M, Kotani K, Ishikawa S.	Body mass index and stroke incidence in Japanese community residents: The Jichi Medical School (JMS) Cohort Study.	J Epidemiol.		in press	2016
141	Shiraishi T, Ishikawa S, Kario K, Kayaba K, Kajii E.	Factor VII and incidence of myocardial infarction in a Japanese population: The Jichi Medical School Cohort Study.	J Clin Lab Anal.		in press	2017
142	Toyoshima H, Otsuka R, Hashimoto S, Tamakoshi K, Yatsuya H.	Body mass index-modified relationship of chronic mental stress with resting blood pressure during 5 years in Japanese middle-aged male workers.	Circ J.	78(6)	1379-86	2014
143	Yatsuya H, Nihashi T, Li Y, Hotta Y, Matsushita K, Muramatsu T, Otsuka R, Matsunaga M, Yamashita K, Wang C, Uemura M, Harada A, Fukatsu H, Toyoshima H, Aoyama A, Tamakoshi K.	Independent association of liver fat accumulation with insulin resistance.	Obes Res Clin Pract.	8(4)	e350-55	2014
144	Uemura M, Yatsuya H, Hilawe EH, Li Y, Wang C, Chiang C, Otsuka R, Toyoshima, Tamakoshi K, Aoyama A.	Breakfast Skipping is Positively Associated with Incidence of Type 2 Diabetes Mellitus - Evidence from Aichi Workers' Cohort Study.	J Epidemiol.	25(5)	351-58	2015
145	Hilawe EH, Yatsuya H, Li Y, Uemura M, Wang C, Chiang C, Toyoshima H, Tamakoshi K, Zhang Y, Kawazoe N, Aoyama A.	Smoking and Diabetes: Is the Association Mediated by Adiponectin, Leptin, or C-reactive Protein?	J Epidemiol.	25(2)	99-109	2015
146	Wang C, Yatsuya H, Tamakoshi K, Toyoshima H, Wada K, Li Y, Hilawe EH, Uemura M, Chiang C, Zhang Y, Otsuka R, Ota A, Hirakawa Y, Aoyama A.	Association between parental history of diabetes and the incidence of type 2 diabetes mellitus differs according to the sex of the parent and offspring's body weight: A finding from a Japanese worksite-based cohort study.	Prev Med.	81	49-53	2015
147	Zhang Y, Yatsuya H, Li Y, Chiang C, Hirakawa Y, Kawazoe N, Tamakoshi K, Toyoshima H, Aoyama A.	Long-term weight change slope, weight fluctuation and risk of type 2 diabetes mellitus in middle-aged Japanese men and women: findings of Aichi Workers' Cohort Study.	Nutr Diabetes.		in press	2017

書籍

著者氏名	論文タイトル名	書籍全体の 編集者名	書籍名	出版社名	出版地	出版年	ページ
斎藤重幸	血管疾患	日本高血圧学会 編	高血圧専門医ガイ ドブック	診断と治療社	東京	2014	144-47
斎藤重幸	糖尿病	日本循環器病予 防学会編	第7版循環器病予 防ハンドブック	保健同人社	東京	2014	273-89
斎藤重幸	疫学からみた高齢者 高血圧治療の意義	楽木宏美	高齢者高血圧の 治療と管理 - JSH2014改訂を ふまえて	先端医学社	東京	2014	32-38
Kokubo Y, Iwashima Y, Kamide K.	Introduction to Hypertension: Types, Causes and Complications.	Jagadeesh G, Balakumar P, Khin Maung U, eds.	Pathophysiology and Pharmaco- therapy of Cardiovascular Disease	Spriner	U.S.A.	2015	635-53
斎藤重幸	高血圧の疫学	浦信行編	高血圧診療の Q&A	中外医学社	東京	2015	1-15
斎藤重幸	メタボリックシンド ローム	山口徹、北原光男 編	今日の治療指針 2015	医学書院	東京	2015	715-17