

雑誌（英文）

| 発表者氏名 | 論文タイトル名 | 発表誌名 | 巻号 | ページ | 出版年 |
|---|--|--------------------------------|-----------------|-----------|------|
| Mori K, Yoshii T, Hirai T, Iwanami A, Takeuchi K, Yamada T, Seki S, Tsuji T, Fujiyoshi K, Furukawa M, Nishimura S, Wada K, Koda M, Furuya T, Matsuyama Y, Hasegawa T, Takeshita K, Kimura A, Abematsu M, Haro H, Ohba T, Watanabe M, Katoh H, Watanabe K, Ozawa H, Kanno H, Imagama S, Ito Z, Fujibayashi S, Yamazaki M, Matsumoto M, Nakamura M, Okawa A, Kawaguchi Y. | Prevalence and distribution of ossification of the supraspinous/interspinous ligaments in symptomatic patients with cervical ossification of the posterior longitudinal ligament of the spine: a CT-based multicenter cross-sectional study. | BMC Musculoskeletal Disorders. | 17(1) | 492 | 2016 |
| Inose H, Yamada T, Mulati M, Hirai T, Ushio S, Yoshii T, Kato T, Kawabata S, Okawa A. | Bone Turnover Markers as a New Predicting Factor for Non-union After Spinal Fusion Surgery. | Spine | <i>In press</i> | | 2016 |
| Yoshii T, Sakai K, Hirai T, Yamada T, Inose H, Kato T, Enomoto M, Tomizawa S, Kawabata S, Arai Y, Okawa A | Anterior decompression with fusion versus posterior decompression with fusion for massive cervical ossification of the posterior longitudinal ligament with a $\geq 50\%$ canal occupying ratio: a multicenter retrospective study. | Spine J. | 16(11) | 1351-1357 | 2016 |

| | | | | | |
|---|---|--------------------|-----------|----------|------|
| Hirai T, Yoshii T, Iwanami A, Takeuchi K, Mori K, Yamada T, Wada K, Koda M, Matsuyama Y, Takeshita K, Abe matsu M, Haro H, Watanabe M, Watanabe K, Oza wa H, Kanno H, Imagama S, Fuji bayashi S, Yamazaki M, Matsumoto M, Nakamura M, Okawa A, Kawaguchi Y. | Prevalence and Distribution of Ossified Lesions in the Whole Spine of Patients with Cervical Ossification of the Posterior Longitudinal Ligament A Multicenter Study (JO SL CT study). | PLoS One. | 11(8) | e0160117 | 2016 |
| Yoshii T, Hirai T, Sakai K, Inose H, Kato T, Okawa A. | Cervical pedicle screw placement using intraoperative computed tomography imaging with a mobile scanner gantry. | Eur Spine J | 25(6) | 1690-7 | 2016 |
| Sakai K, Yoshii T, Hirai T, Arai Y, Torigoe I, Tomori M, Sato H, Okawa A. | Cervical Sagittal Imbalance is a Predictor of Kyphotic Deformity After Laminoplasty in Cervical Spondylotic Myelopathy Patients Without Preoperative Kyphotic Alignment. | Spine | Feb;41(4) | 299-305 | 2016 |
| Kashii M, Matsuo Y, Sugiura T, Fujimori T, Nagamoto Y, Makino T, Kaito T, Ebina K, Iwasaki M, Yoshikawa H. | Circulating sclerostin and dickkopf-1 levels in ossification of the posterior longitudinal ligament of the spine | J Bone Miner Metab | 34(3) | 315-324 | 2016 |
| Kawaguchi Y, Nakano M, Yasuda T, Seki S, Hori T, Suzuki K, Makino H, Kimura T | Characteristics of ossification of the spinal ligament; incidence of ossification of the ligamentum flavum in patients with cervical ossification of the posterior longitudinal ligament - Analysis of the whole spine using multidetector CT - | J Orthop Sci | 21(4) | 439-45 | 2016 |

| | | | | | |
|--|---|-------------------|-----------------|---------|------|
| Takahashi H, Koda M, Hashimoto M, Furuya T, Sakuma T, Kato K, Okawa A, Inada T, Kamiya K, Ota M, Maki S, Takahashi K, Yamazaki M, Mannoji C | Transplanted peripheral blood stem cells mobilized by granulocyte colony-stimulating factor promoted hindlimb functional recovery after spinal cord injury in mice | Cell Transplant | 25 | 283-292 | 2016 |
| Maki S, Aramomi M, Matsuura Y, Furuya T, Ota M, Iijima Y, Saito J, Suzuki T, Mannoji C, Takahashi K, Yamazaki M, Koda M | Paravertebral foramen screw fixation for posterior cervical spine fusion: biomechanical study and description of a novel technique | J Neurosurg Spine | <i>in press</i> | | 2016 |
| Maki S, Koda M, Ota M, et al. | Reduced field-of-view diffusion tensor imaging of the spinal cord shows motor dysfunction of the lower extremities in patients with cervical compression myelopathy. | Spine | <i>in press</i> | | 2016 |
| Fujii K, Abe T, Kubota S, Marushima A, Kawamoto H, Ueno T, Matsushita A, Nakai K, Saotome K, Kadone H, Endo A, Haginoya A, Hada Y, Matsumura A, Sankai Y, Yamazaki M | The voluntary driven exoskeleton Hybrid Assistive Limb (HAL) for postoperative training of thoracic ossification of the posterior longitudinal ligament: a case report | J Spinal Cord Med | 9 | 1-7 | 2016 |
| Kubota S, Fujii K, Marushima A, Ueno T, Haginoya A, Endo A, Kadone H, Kawamoto H, Shimizu Y, Hada Y, Matsumura A, Sankai Y, Yamazaki M | Improvement of walking ability using Hybrid Assistive Limb training in a patient with severe thoracic myelopathy caused by ossification of the posterior longitudinal ligament. A case report | J Spine | S7 | 003 | 2016 |

| | | | | | |
|---|--|-------------------|-----------------------|---------|------|
| Ikumi A, Kubota S, Shimizu Y, Kadone H, Marushima A, Ueno T, Kawamoto H, Hada Y, Matsumura A, Sankai Y, Yamazaki M | Decrease of spasticity after hybrid assistive limb(®) training for a patient with C4 quadriplegia due to chronic SCI | J Spinal Cord Med | 20 | 1-6 | 2016 |
| Koda M, Furuya T, Rokkaku T, Murakami M, Iijima Y, Saito J, Kitamura M, Ohtori S, Orita S, Inage K, Yamazaki M, Mannoji C | Drop finger caused by 8th cervical nerve root impairment: a clinical case series | Eur Spine J | [Epub ahead of print] | | 2016 |
| Maki S, Koda M, Saito J, Takahashi S, Inada T, Kamiya K, Ota M, Iijima Y, Masuda Y, Matsumoto K, Kojima M, Takahashi K, Obata T, Yamazaki M, Furuya T | Tract-specific diffusion tensor imaging reveals laterality of neurological symptoms in patients with cervical compression myelopathy | World Neurosurg | 96 | 184-190 | 2016 |
| Saito J, Maki S, Kamiya K, Furuya T, Inada T, Ota M, Iijima Y, Takahashi K, Yamazaki M, Aramomi M, Mannoji C, Koda M | Outcome of posterior decompression with instrumented fusion surgery for K-line (-) cervical ossification of the longitudinal ligament | J Clin Neurosci | 32 | 57-60 | 2016 |
| Ota M, Furuya T, Maki S, Inada T, Kamiya K, Iijima Y, Saito J, Takahashi K, Yamazaki M, Aramomi M, Mannoji C, Koda M | Addition of instrumented fusion after posterior decompression surgery suppresses thickening of ossification of the posterior longitudinal ligament of the cervical spine | J Clin Neurosci | 34 | 162-165 | 2016 |

| | | | | | |
|--|--|-----------------|---------|-------------|------|
| Koda M, Furuya T, Kinoshita T, Miyashita T, Ota M, Maki S, Ijima Y, Saito J, Takahashi K, Yamazaki M, Aramomi M, Mannoji C | Dropped head syndrome after cervical laminoplasty: A case-control study | J Clin Neurosci | 32 | 88-90 | 2016 |
| Koda M, Mochizuki M, Konishi H, Aiba A, Kadota R, Inada T, Kamiya K, Ota M, Maki S, Takahashi K, Yamazaki M, Mannoji C, Furuya T | Comparison of clinical outcomes between laminoplasty, posterior decompression with instrumented fusion, and anterior decompression with fusion for K-line (-) cervical ossification of the posterior longitudinal ligament | Eur Spine J | 25 | 2294-2301 | 2016 |
| Maki S, Koda M, Iijima Y, Furuya T, Inada T, Kamiya K, Ota M, Saito J, Okawa A, Takahashi K, Yamazaki M | Medially-shifted rather than high-riding vertebral arteries preclude safe pedicle screw insertion | J Clin Neurosci | 29 | 169-172 | 2016 |
| Koda M, Furuya T, Okawa A, Inada T, Kamiya K, Ota M, Maki S, Takahashi K, Yamazaki M, Aramomi M, Ikeda O, Mannoji C | Mid- to long-term outcomes of posterior decompression with instrumented fusion for thoracic ossification of the posterior longitudinal ligament | J Clin Neurosci | 27 | 87-90 | 2016 |
| Ito Z, Matsuyama Y, Ando M, Kawabata S, Kanchiku T, Kida K, Fujiwara Y, Yamada K, Yamamoto N, Kobayashi S, Saito T, Wada K, Tadokoro N, Takahashi M, Satomi K, Shinomiya K, Tani T | Postoperative Paralysis from Thoracic Ossification of Posterior Longitudinal Ligament (OPLL) Surgery-Risk factor of Neurologic Injury: Nationwide Multi-Institution Survey | Spine | 41(19): | E1159-E1163 | 2016 |

| | | | | | |
|---|---|----------------------------|-------|-----------|------|
| Katsumi K, Izumi T, Ito T, Hirano T, Watanabe K, Ohashi M. | Posterior instrumented fusion suppresses the progression of ossification of the posterior longitudinal ligament: A comparison of laminoplasty with and without instrumented fusion by 3-dimensional analysis. | European Spine Journal | 25(5) | 1634-1640 | 2016 |
| Ito Z, Matsuyama Y, Ando M, Kawabata S, Kanchiku T, Kida K, Fujiwara Y, Yamada K, Yamamoto N, Kobayashi S, Saito T, Wada K, Satomi K, Shinomiya K, Tani T | A Multicenter Study by the Monitoring Committee of the Japanese Society for Spine Surgery and Related Research. | Global Spine J | 6(3) | 234-241 | 2016 |
| Aizawa T et al. | Increasing Incidence of Degenerative Spinal Diseases in Japan during 25 Years: The Registration System of Spinal Surgery in Tohoku University Spine Society. | Tohoku J Exp Med | 238 | 153-63 | 2016 |
| Yahata K et al. | Low-energy extracorporeal shock wave therapy for promotion of vascular endothelial growth factor expression and angiogenesis and improvement of locomotor and sensory functions after spinal cord injury. | J Neurosurg Spine | 25 | 745-55 | 2016 |
| Shibata A, Machida J, Yamaguchi S, Kimura M, Tatematsu T, Miyachi H, Matsushita M, Kitoh H, Ishiguro N, Nakayama A, Higashi Y, Shimoza to K, Tokita Y | Characterization of novel Runx2 mutation with alanine tract expansion from Japanese cleidocranial dysplasia patient | Mutagenesis | 31 | 61-67 | 2016 |
| Hasegawa S, Kitoh H, Ohkawara B, Mishima K, Matsushita M, Masuda A, Ishiguro N, Ohno K | Tranilast stimulates endochondral ossification by upregulating SOX9 and RUNX2 promoters | Biochem Biophys Res Commun | 470 | 356-361 | 2016 |

| | | | | | |
|--|--|------------------------------|--------|-----------|------|
| Sugiura K, Ohno A, Kono M, Kitoh H, Itomi K, Akiyama M | Hyperpigmentation over the metacarpophalangeal joints the malleoli in a case of hyaline fibromatosis syndrome with ANTXR2 mutations | J Eur Acad Dermatol Venereol | 30 | 44-46 | 2016 |
| Matsushita M, Kitoh H, Mishima K, Kadono I, Sugiura H, Hasegawa S, Nishida Y, Ishiguro N | Low bone mineral density in achondroplasia and hypochondroplasi | Pediatr In | 58 | 705-708 | 2016 |
| Hasegawa S, Victoria T, Kayserili H, Zackai E, Nishimura G, Haga N, Nakashima Y, Miyazaki O, Kitoh H | Characteristic calcaneal ossification: an additional early radiographic finding in infants with fibrodysplasia ossificans progressiva | Ped Radiol | 46(11) | 1568-1572 | 2016 |
| Melovitz-Vasan C, Kitoh H, Vasan N | A rare case of femoral hypoplasia (proximal focal femoral deficiency) with overlapping phenotype of mullerian duct syndrome (Mayer-Rokitansky-Hauser syndrome) | Int J Anat Res | 4 | 3312-3316 | 2016 |
| Ohya J, Oshima Y, Oka H, Saiki F, Taniguchi Y, Matsubayashi Y, Tanaka S, Chikuda H, Takeshita K | Patient Satisfaction with Posterior Decompression Surgery for Cervical Ossification of the Posterior Longitudinal Ligament: Prognostic Radiographic Factors and Patient-Reported Outcomes for the Effectiveness of Surgical Treatment. | World Neurosurgery | 96 | 272-279 | 2016 |
| Imajo Y, Kanchiku T, Suzuki H, Yoshida Y, Nishida N, Goto H, Suzuki M, Taguchi T | Intracranial epidural hemorrhage during lumbar spinal surgery | Spinal Cord Ser Cases | 7(2) | 15040 | 2016 |
| Funaba M, Kanchiku T, Imajo Y, Yamamoto H, Hirura Y, Fujimoto K, Taguchi T, Kido K | Relation between distal motor latency delay and sensory action potential absence of the median nerve in carpal tunnel syndrome | Clin Neurolog Neurosurg | 148 | 45-48 | 2016 |
| Kanchiku T, Imajo Y, Suzuki H, Yoshida Y, Akashi K, Taguchi T | Correlation Between Spinal Cord Function Assessed by Intraoperative SCEPs and Morphology of the Compressed Spinal Cord on MRI | Clin Spine Surg | 29(10) | E496-E501 | 2016 |

| | | | | | |
|---|--|--------------------|---------|--------------|------|
| Suetomi Y, Kanchiku T, Nishijima S, Imajo Y, Suzuki H, Yoshida Y, Nishida N, Taguchi T | Application of diffusion tensor imaging for the diagnosis of segmental level of dysfunction in cervical spondylotic myelopathy | Spinal Cord | 54(5) | 390-5 | 2016 |
| Ito Z, Matsuyama Y, Ando M, Kawabata S, Kanchiku T, Kida K, Fujiwara Y, Yamada K, Yamamoto N, Kobayashi S, Saito T, Wada K, Tadokoro N, Takahashi M, Satomi K, Shinomiya K, Tani T. | Postoperative paralysis from thoracic ossification of posterior longitudinal ligament surgery risk factor of neurologic injury: Nationwide multi-institutional survey. | Spine | 41 | E1159-E1163. | 2016 |
| Fujiwara H, Makino T, Yonenobu K, Honda H, Kaito T | Correlations and time-dependent changes of upper arm performance tests, the Japanese Orthopaedic Association score, and a newly developed patient-based outcome measure: the Japanese Orthopaedic Cervical Myelopathy Questionnaire. | Bone Joint J. | 98-B(7) | 990-996 | 2016 |
| Yoshii T, Hirai T, Yamada T, Inose H, Kato T, Sakai K, Enomoto M, Kawabata S, Arai Y, Okawa A | Intraoperative evaluation using mobile computed tomography in anterior cervical decompression with floating method for massive ossification of the posterior longitudinal ligament. | J Orthop Surg Res. | 12(1) | 12 | 2017 |
| Kawaguchi Y, Nakano M, Yasuda T, Seki S, Hori T, Suzuki K, Makino H, Kitazima I, Kimura T | Serum biomarkers in patients with ossification of the posterior longitudinal ligament (OPLL):-inflammation in OPLL- | PLoS One | 12(5) | e0174881 | 2017 |

| | | | | | |
|--|--|---------------------|-----------------|------------|------|
| Shimizu Y, Kadone H, Kubota S, Ikumi A, Abe T, Marushima A, Ueno T, Endo A, Kawamoto H, Saotome K, Matsushita A, Matsumura A, Sankai Y, Hada Y, Yamazaki M | Active elbow flexion is possible in C4 quadriplegia using hybrid assistive limb (HAL®) technology: A case study | J Spinal Cord Med | 29 | 1-7 | 2017 |
| Kubota S, Abe T, Kadone H, Fujii K, Shimizu Y, Marushima A, Ueno T, Kawamoto H, Hada Y, Matsumura A, Sankai Y, Yamazaki M | Walking ability following Hybrid Assistive Limb treatment for a patient with chronic myelopathy after surgery for cervical ossification of the posterior longitudinal ligament | J Spinal Cord Med | <i>in press</i> | | 2017 |
| Kimura A et al | "Fall-related deterioration of subjective symptoms in patients with cervical myelopathy." | Spine | 42 | E398-E403. | 2017 |
| Mori K, Kasahara T, Mimura T, Nishizawa K, Nakamura A, Imai S. | Prevalence of thoracic diffuse idiopathic skeletal hyperostosis (DISH) in Japanese: Results of chest CT-based cross-sectional study. | J Orthop Sci | 22(1) | 38-42 | 2017 |
| Matsushita M, Mishiima K, Esaki R, Ishiguro N, Ohno K, Kitoh H | Maternal administration of mifepristone for the treatment of foramen magnum stenosis in transgenic mice with achondroplasia | J Neurosurg Pediatr | 19 | 91-95 | 2017 |
| Imajo Y, Kanchiku T, Suzuki H, Yoshida Y, Funaba M, Nishida N, Fujimoto K, Taguchi T | Effects of differences in age and body height on normal values of central motor conduction time determined by F-waves | J Spinal Cord Med | 40(2) | 181-187 | 2017 |
| Nishida N, Kanchiku T, Kato Y, Imajo Y, Suzuki H, Yoshida Y, Ohgi J, Chen X, Taguchi T | Cervical ossification of the posterior longitudinal ligament: factors affecting the effect of posterior decompression | J Spinal Cord Med | 40(1) | 93-99 | 2017 |

| | | | | | |
|---|--|-------------------|--------------|------------------|-------------|
| <p>Kimura A, Seichi A, Takeshita K, Inoue H, Kato T, Yoshii T, Furuya T, Kodama M, Takeuchi K, Matsunaga S, Seki S, Ishikawa Y, Imagama S, Yamazaki M, Mori K, Kawasaki Y, Fujita K, Endo K, Sato K, Okawa A.</p> | <p>Fall-related Deterioration of Subjective Symptoms in Patients with Cervical Myelopathy.</p> | <p>Spine</p> | <p>42(7)</p> | <p>E398-E403</p> | <p>2017</p> |
| <p>Yoshii T, Hirai T, Sakai K, Sotome S, Enomoto M, Yamada T, Inose H, Kato T, Kawabata S, Okawa A.</p> | <p>Anterior Cervical Corpectomy and Fusion Using a Synthetic Hydroxyapatite Graft for Ossification of the Posterior Longitudinal Ligament.</p> | <p>Orthopedic</p> | <p>40(2)</p> | <p>E334-E339</p> | <p>2017</p> |