

厚生労働科学研究費補助金（化学物質リスク研究事業）
分担研究報告書

**Predictors of folate status among pregnant Japanese women: the Hokkaido Study
on Environment and Children's Health, 2002 -2012**

研究代表者 岸 玲子 北海道大学環境健康科学研究教育センター 特別招へい教授
研究分担者 荒木 敦子 北海道大学環境健康科学研究教育センター 准教授
研究分担者 宮下ちひろ 北海道大学環境健康科学研究教育センター 特任講師
研究分担者 池野多美子 北海道大学環境健康科学研究教育センター 特任講師
研究分担者 水上 尚典 北海道大学大学院医学研究科生殖・発達医学講座 教授
研究分担者 遠藤 俊明 札幌医科大学医学部産婦人科学講座 准教授
研究分担者 千石 一雄 旭川医科大学医学部産婦人科学講座 教授
研究分担者 佐々木成子 北海道大学大学院医学研究科 助教
研究分担者 花岡 知之 北海道大学環境健康科学研究教育センター 客員教授

研究要旨

The International Clearinghouse for Birth Defects, Surveillance and Research (ICBDSR) reports a rise in the prevalence rate of spina bifida in Japan. We determined first trimester folate status of Hokkaido women and identified potential predictors. Participants were 15266 pregnant women of The Hokkaido Study on Environment and Children's Health Cohort. Data was extracted from self-reported questionnaires and biochemical assay results. Demographic determinants of low folate status were younger maternal age (Adjusted odds ratio [AOR]: 1.48, 95% CI: 1.32, 1.66); lower educational level (AOR: 1.27, 95% CI: 1.17, 1.39); and lower annual income (AOR: 1.11, 95% CI: 1.01, 1.22). Plasma cotinine concentrations of 1.19 – 65.21nmol/L increased the risk of low folate status (AOR: 1.20, 95% CI: 1.10, 1.31); and concentration > 65.21nmol/L further increased the risk (AOR: 1.91, 95% CI: 1.70, 2.14). Most favorable predictor was folic acid (FA) supplements use (AOR: 0.19, 95% CI: 0.17, 0.22). Certain socio-demographic factors influence folate status among pregnant Japanese women. Modifiable negative and positive predictors are active and passive tobacco smoking, and folic acid supplements use. Avoiding both active and passive tobacco smoking, and using folic acid supplements could improve folate status of Japanese women.

研究協力者

Yila Thamar Ayo、伊藤久美子、小林澄貴、馬場俊明、Braithmoh Tiilola
(北海道大学環境健康科学研究教育センター)
Houman Goudaruzi (外国人特別研究員)

A. 研究目的

Folate as a cofactor in one-carbon metabolism is very essential in all cellular processes, especially in conditions of

rapid cell replications and tissue growth like pregnancy. The role of synthetic folic acid (FA) supplements in the prevention of neural tube defects (NTDs) has been well documented^{1,2}. Several countries with policies of FA food fortification, do report a 30.00% to 70.00% reduction in the incidence of NTDs³. As reported by the International Clearinghouse for Birth Defects, Surveillance and Research

(ICBDSR), Japan has experienced an increase in the prevalence of spina bifida over the past few decades. While in countries like the United States and England the prevalence of NTDs is about 3 per 10,000 births, Japan has a prevalence of 5.2⁴; ⁵. Efforts have been made to determine folate status of pregnant Japanese women in other regions of Japan but not Hokkaido⁶; ⁷; ⁸; ⁹. However, probable small sample sizes, different folate assay techniques and varied folate status definitions/cut-off levels, might have yielded inconsistent results. For instance, three previous studies have reported a wide range of folate status among pregnant Japanese women. A study from Aichi Prefecture which defined normal versus inadequate folate status as $\geq 6.80\text{nmol/L}$ and $< 6.80\text{nmol/L}$ respectively, reported folate inadequacy in 1.00% of 41 pregnant and 154 non pregnant participants⁶. Another study from Tokyo metropolis among pregnant women in all trimesters reported low folate status in 67.00% of 118 women in their first trimester and 79.30% overall. The study defined normal, low and deficient folate statuses as having $> 13.60\text{nmol/L}$, $6.80 - 13.60\text{nmol/L}$, and $< 6.80\text{nmol/L}$, respectively⁸. The third study from Tokyo metropolis was conducted among 58 young university women. Normal folate status was categorized as having $\geq 13.60\text{nmol/L}$; low status as having $6.80 - 13.59\text{nmol/L}$; while folate deficient group had $< 6.80\text{nmol/L}$ of folate concentration. Results showed 12.10% had folate deficiency, and 58.60% had low status⁹. In Hokkaido Prefecture, such reports are scarce. We recently conducted a genetic study using the first

part of this cohort's data. Low folate status was reported among 28.70% of the study population ($N = 1784$), but the scope of the study excluded a detailed exploration of the demographic and lifestyle predictors of folate status¹⁰. In contrast to previous smaller studies in Japan, this study uses data from a large cohort to explore the various demographic and lifestyle factors that may influence first trimester folate status of Japanese women in Hokkaido.

B . 研究方法

Study location and population

Participants of this study were pregnant women recruited during their first trimester (< 13 weeks of gestation) from thirty-seven health facilities across Hokkaido Prefecture. They are participants of the ongoing large-scale birth cohort of the Hokkaido Study on Environment and Children's Health. The study is broadly aimed at observing the health effects of intrauterine exposures to various environmental and genetic factors on fetal development, outcomes of pregnancy, and subsequent childhood health. Details of the study have been described elsewhere¹¹; ¹². Briefly, the ongoing large-scale cohort started in 2002, with a full blown large scale version from February 2003. A total of 20816 women were recruited between 2002 and March 2012. All pregnant Japanese women who presented at any of the participating healthcare facilities for prenatal care during their first trimester, were considered eligible for the study. However, only those who agreed to participate in the study were contacted and recruited. Data was generated from these

participants by means of baseline questionnaires, biochemical assays, hospital birth records, and four months post-partum health records. We finally used a total of 15266 participants, details of selection is shown in the flow chart (**Fig. 1**).

Certain repeated self-reported information obtained from birth records and postpartum questionnaires were used to compare with the baseline questionnaires in order to improve quality and missing information in the whole data. Otherwise, these data were not used in the analysis of this report.

Biochemical assays

Non-fasting whole blood samples were collected from participants, pre-treated and sera obtained. Sera were stored promptly at 4°C until they were transported on ice to a commercial laboratory (SRL Corporation Inc, Tokyo, Japan), for folate assay. The ADVIA Centaur Folate Assay Protocol is one of the automated Competitive Protein Binding (CPB) Immunoassay Technology. Folate is quantified by direct chemiluminescent acridinium ester technology¹³. This technique has an acceptable imprecision of less than 10.00%, with an advanced Quality Control (QC) package. It has an analytical sensitivity of 0.91nmol/L. It can detect from small volumes of as low as 10uL of biological specimen, thus, making it a method of choice in large epidemiological studies¹⁴. Specimen preparations, shipping, and assays, were done in batches, depending on new recruitments. All laboratory analysts were blinded to participants' information. Because there is no standard classification of folate status

from automated immunoassay techniques, we adopted the World Health Organization's (WHO) classification guidelines¹⁵. Nicotine is the toxic chemical in tobacco products and its predominant metabolite is cotinine. Cotinine can be detected in biological specimens as biomarker of exposure to tobacco. In this study, we used plasma cotinine concentrations to quantitatively classify active and passive smoking status. The details of measurements of plasma cotinine are described in our previous report¹⁶.

Definition of variables

The dependent variable was folate status. Folate status was classified as: folate deficiency, defined as (<6.80nmol/L) of serum folate; suboptimal status (6.80 -13.59nmol/L) and optimal folate status (\geq 13.60nmol/L)¹⁵. Folate deficiency was reported in 0.52% of the study population. To improve study power, and because non fasting serum was used for folate assay, we merged this group with the suboptimal category. Active and passive exposure to tobacco smoking statuses were classified based on plasma cotinine cut-off points established in a previous report¹⁶. A non-smoker was defined as having plasma cotinine concentration of less than 1.19nmol/L. A person exposed to environmental tobacco smoke (ETS) as having 1.19 – 65.21nmol/L; and active smoker as having greater than 65.21nmol/L of plasma cotinine concentration. Prenatal folic acid (FA) supplement use was defined as “a report on the use of FA supplements before or after conception”. Other nutritional

厚生労働科学研究費補助金（化学物質リスク研究事業）
分担研究報告書

supplements use was defined as “any report on intake of nutritional supplements other than FA, before or after conception”. Ingestion of alcoholic beverages was categorized based on frequency of intake: “monthly, weekly or daily”. Self-reported active tobacco smoking was categorized based on the number of cigarette sticks smoked per day. Light smokers (<10 cigarette sticks per day); moderate smokers (10-19 cigarette sticks per day); and heavy smokers (≥ 20 cigarette sticks per day). ETS exposure at home was defined as “living with one or more active smokers”. ETS at work place referred to “working with one or more active smokers at work place”. In this study, lifestyle habits considered were alcoholic beverage consumption, nutritional supplements use, and tobacco use. Potential predictors of folate status were identified based on previous reports. In this study, year of enrolment, maternal age, parity, BMI, educational level, household income, occupation, use of nutritional supplements, active and passive cigarette smoking, alcohol intake, season of the year, and geographical location, were identified as putative predictors.

Statistical analyses

Statistical tests of associations included Pearson's chi squared tests, and Fischer's exact tests for categorical variables. Skewed serum folate and plasma cotinine concentrations were log-transformed during the preliminary descriptive analyses, thereafter back-transformed. Differences in mean folate levels were explored using ANOVA with post-hoc analyses to correct for multiple

comparisons. However, the main regression analyses were performed using qualitative folate status. We imputed the missing values present in the data via Multivariate Imputation by Chained Equations (MICE), as implemented in the R package *mice*, obtaining $m = 10$ imputed data sets. MICE is a Markov Chain Monte Carlo method that uses the correlation structure of the data and imputes missing data values for each incomplete variable m times by regression of incomplete variables on the other available variables iteratively. We used Bayesian logistic regression and fitted the model to the $m = 10$ imputed data set, with dichotomized folate status as the outcome variable, and the followings as potential predictor variables: age, BMI, parity, educational level, income, occupation, region, year of enrollment, season of the year at enrollment, folic acid supplements use, other nutritional supplements use, alcohol intake, active cigarette smoking, and exposure to environmental tobacco smoke (ETS) both at home and work place. We used results of plasma cotinine concentration to quantitatively classify active smoking and passive exposure to tobacco products, and regressed against folate status, with adjustment for all other potential predictors. We reported pooled estimates for the main effects of the predictor variables in the model. P-values for testing for the presence of a linear trend are also reported for predictor variables with more than two categories. Reported effects, confidence intervals and p-values are pooled over the $m = 10$ imputed data sets. Additionally, we reported the value of the McFadden's

pseudo- R^2 pooled over these data sets. All statistical analyses were performed using JMP 11 Pro Statistical Software Package (SAS, Cary, NC, USA), except for the binary logistic regression model which required multiple imputation of missing data and was performed using R version 3.2.2 (Vienna, Austria). An alpha level of significance was set at <0.05 .

（倫理面への配慮）

The Institutional Ethical Board for Human Gene and Genome studies at Hokkaido University Graduate School of Medicine approved the study protocol.

C. 研究結果

Overall, geometric mean (standard deviation) of serum folate concentration was 17.77 (3.58)nmol/L. Among women with optimal folate status, the geometric mean (standard deviation) was 20.67 (3.26) nmol/L, and 10.83 (2.65) nmol/L among participants with suboptimal folate status. One sided lower limit tolerance interval at 95% of the population was 8.47nmol/L. Prevalence of folate deficiency was 0.52%. Suboptimal folate status constituted 25.65%, while optimal folate status was reported in 73.83% of the population (**Table 1**). Initial descriptive analyses using folate as a continuous variable revealed mean serum folate concentrations increased with increasing maternal age ($p<0.001$), educational status ($p<0.001$), annual income ($p<0.001$), FA supplements use ($p<0.001$), and other nutritional supplements use ($p<0.001$). Mean serum folate concentrations decreased with increasing number of cigarette sticks smoked per day ($p<0.001$), ETS exposure at home ($p<0.001$), and increasing plasma

cotinine concentrations ($p<0.001$). Exposure to ETS at both home and at work was associated with low folate status, $p <0.001$. About 60.00% of those with folate deficiency were exposed to both ETS at home and at work place. Other associations were geographical region, year of enrolment into the study, and season of the year (data not shown). Serum folate inversely correlated with plasma cotinine concentration ($r = -0.2000$, $p <0.001$, data not shown). Significant differences were observed in mean plasma cotinine concentrations among nonusers of FA supplements and users, with geometric mean (SD) of 46.41 (23.23)nmol/L, and 25.27 (15.32)nmol/L, $p <0.001$, respectively. Also, geometric mean (SD) between nonusers and users of other nutritional supplements was 42.49 (21.91)nmol/L, and 34.99 (20.17)nmol/L, $p = 0.028$, respectively (**Fig. 2**). Users of FA supplements were likely to be those with chronic inter-current medical conditions, those who had fertility treatments, and those who were also users of other nutritional supplements. 7.00% of folic acid users started intake more than 3months before conception. Another 8.00% started 1 month before conception, while majority (more than 60.00%) started use following confirmation of pregnancy. The average frequency of use per week was 3 times. Multivitamins reported were found to contain various doses of folic acid in the range of 100 μ g to 200 μ g per tablet (data not shown).

In the regression model, the value of the McFadden's pseudo- R^2 pooled over the $m = 10$ imputed data sets was -8.69% . The demographic determinants of low

厚生労働科学研究費補助金（化学物質リスク研究事業）
分担研究報告書

folate status identified were lower maternal age (AOR: 1.48, 95% CI: 1.32, 1.66, $p < 0.001$); lower educational level (AOR: 1.27, 95% CI: 1.17, 1.39, $p < 0.001$); lower annual income (AOR: 1.11, 95% CI: 1.01, 1.22, $p = 0.024$); residing in the south and eastern regions (AOR: 1.25, 95% CI: 1.14, 1.38, $p < 0.001$), and (AOR: 1.15, 95% CI: 1.05, 1.25, $p = 0.003$), respectively. Being enrolled into the study between 2005 and 2007 was associated with an increase in the risk of low folate status (AOR: 1.23, 95% CI: 1.12, 1.35, $p < 0.001$); while recruitment between 2008 and 2010 reduced the likelihood of having low folate status (AOR: 0.81, 95% CI: 0.73, 0.90, $p < 0.001$), respectively. Being enrolled during summer, autumn, and winter were associated with higher likelihood of low folate status (AOR: 1.12, 95% CI: 1.02, 1.24, $p = 0.023$); (AOR: 1.13, 95% CI: 1.02, 1.25, $p = 0.015$), and (AOR: 1.13, 95% CI: 1.01, 1.27, $p = 0.037$), respectively. Lower BMI (AOR: 0.84, 95% CI: 0.74, 0.94, $p = 0.006$; and unemployment were associated with risk reduction (AOR: 0.87, 95% CI: 0.80, 0.94, $p = 0.001$), (**Table 2**).

Lifestyle factors that reduced the odds of low folate status were the use of FA supplements (AOR: 0.19, 95% CI: 0.17, 0.22, $p < 0.001$); other nutritional supplements (AOR: 0.55, 95% CI: 0.48, 0.64, $p < 0.001$); and weekly alcohol consumption (AOR: 0.75, 95% CI: 0.62, 0.90, $p = 0.003$), respectively. Lifestyle factors that increased the odds of low folate status were active cigarette smoking and ETS exposure. Smoking < 10 cigarette sticks per day was associated with increased odds (AOR: 1.42, 95% CI:

1.23, 1.64, $p < 0.001$); while smoking between 10 to 19 cigarette sticks per day was associated with an increased risk (AOR: 2.28, 95% CI: 1.92, 2.71, $p < 0.001$). However, smoking ≥ 20 cigarette sticks per day was not statistically significant, but $p_{\text{trend}} < 0.001$. Exposure to ETS at home increased the odds of low folate status (AOR: 1.23, 95% CI: 1.13, 1.34, $p < 0.001$), and exposure to ETS at workplace also increased the odds of low folate status (AOR: 1.16, 95% CI: 1.02, 1.31, $p = 0.02565$), (**Tables 2**).

Using plasma cotinine concentrations to classify active and passive exposure to tobacco products, **Tables 3** shows that participants with plasma cotinine levels between 1.19 – 65.21nmol/L were 1.20 times more likely to have low folate status (AOR: 1.20, 95% CI: 1.10, 1.31, $p < 0.001$); while those with levels > 65.21 nmol/L had a twofold increase in risk (AOR: 1.91, 95% CI: 1.70, 2.14, $p < 0.001$); $p_{\text{trend}} < 0.001$.

D. 考察

To our knowledge, this report presents robust information on demographic and lifestyle predictors of folate status in a relatively large cohort of pregnant Japanese women. Majority (73.83%) of participants had optimal first trimester folate status. Only 0.52% had serum folate concentrations below 6.80nmol/L, a level clinically considered a negative folate balance, while 25.65% of the population had marginal folate status. Lower tolerance limit of 8.47nmol/L implies a negative folate balance for this population. Our findings contrast those from Tokyo where more than 50.00% of the study population of pregnant women

had low folate status.

Demographic predictors of folate status

Low folate status was associated with younger maternal age, higher BMI, educational level and annual income. Cigarette smoking rate is on the increase among young Japanese women, and a quest to achieve a lower BMI via dieting is in vogue among women of reproductive age. These factors may invariably compromise nutritional status including folate among younger women^{4; 17}. Micronutrients deficiencies including folate in overweight/obese people have been reported by some previous studies¹⁸. Possible mechanisms postulated have been: decrease dietary intake, current cigarette smoking, and possible low serum/plasma concentrations as a result of increased intravascular volume¹⁹. Consistent with our findings, socio-economic status has been reported to influence folate intake among Japanese workers²⁰. Also, educational attainment was reported in Belgium²¹, and Australia²². In USA, older maternal age, higher education, and higher income status, have been reported to predict the use of FA supplements²³. In this study, these factors might have favored higher folate status. Other demographic factors associated with suboptimal folate status have been reported from other countries, and these include household size²⁴, season of the year²⁵, rural residence²⁶, and region²⁷. We observed that residing in the southern and eastern regions; and seasons of the year were associated with the risk of low folate status.

Traditionally, most Japanese women are full-time house wives. This may explain

why the unemployed had lower risk. Working women are likely to skip their meals and may prefer fast foods as reported among children of working women²⁸. Of note here is that employment status was broadly classified. Further exploration based on job types may shade more insight on this observation.

Unfavorable lifestyle predictors of folate status

We report self-reported active cigarette smoking and ETS exposure as the major modifiable unfavorable predictors of folate status. Although we could not demonstrate a dose-response pattern in the odds, especially among heavy smokers during pregnancy; this may probably be related to a small subgroup size. Using plasma cotinine biomarker, the risk of low folate status increased in a dose-response pattern. Contrary to this result, another study in Tokyo found no lifestyle habits as risk factors for suboptimal folate status⁸. However, our result is consistent with reports from other developed countries, where lifestyle factors are commonly observed as predictors of folate status. Folate depleting effects of active smoking and ETS exposure have been reported^{29; 30; 31; 32; 33; 34; 35}. Possible biologic mechanisms of folate depletion in active and passive smokers include decreased intake^{29; 33}, inactivating effects of organic nitrites, cyanates, and nitrous oxide on circulating folates^{34; 36}, and direct effects of oxidative stress or increased folate turnover^{31; 37}. We observed lower mean plasma cotinine concentrations among nutritional supplements users. Nutritional

supplements users are more likely to practice healthy lifestyles.

Favorable lifestyle predictors of folate status

FA supplements use is the major modifiable predictor of optimal folate status. This report further confirms the well documented role of FA supplements in improving folate status. Other nutritional supplements used also correlated positively with folate status, probably because most multivitamins also contain FA. Other nutritional supplements used included multivitamins, trace elements, herbs, proteins, ginseng and energy drinks. Over-the-counter (OTC) multivitamins used contained various doses of folic acid in the range of 100 μ g to 200 μ g per tablet according to the brand names reported by study participants, majority of whom were recruited between 2002 and 2010. However, lately, folic acid content seems to have been increased by drug makers (up to 480 μ g/tablet). This may reflect in our findings of increase in mean folate concentrations of participants enrolled from 2010 and beyond, and a reduction in the risk of having low folate status. In this study, majority of folic acid supplements users did not use it because of pregnancy. Those who used it for prenatal purpose started only after confirming they were pregnant. This information may impact on the crucial periconceptional period for prevention of NTDs. Within Japan, some smaller studies outside Hokkaido did report that using FA supplements increased blood folate concentrations more than using dietary sources of folate only. They also observed that Japanese women in their

reproductive age do not meet the daily Recommended Dietary Allowance (RDA) of 440 μ g for folate^{5; 7; 8; 38; 39}. Although the Japanese Government has recommended that women of reproductive age or those who plan to become pregnant should take 400 μ g/day of FA supplements, scholars have reported that the level of awareness and compliance with the recommendations are still low^{5; 40}. Furthermore, across the Asian sub-region, prenatal FA supplements use is not a routine prenatal care practice⁴¹. Our findings are similar to other reports emerging from China, Malaysia and Indonesia. Of these three, mandatory fortification is legislated only in Indonesia^{42; 43; 44}. Internationally, studies from other developed countries without food fortification policies are reporting increasing incidence of suboptimal folate concentrations^{21; 45; 46}. Our result on the role of alcoholic beverage consumption on folate is consistent with a previous study in Czech Republic, where moderate beer consumption correlated with higher plasma folate⁴⁷. Conversely, chronic heavy alcohol consumption is associated with folate deficiency via numerous mechanisms⁴⁸. We stand with the universal recommendation that pregnant women should abstain from consuming alcoholic beverages, because of adverse fetal effects⁴⁹.

Strengths and limitations

This study is the first to utilize a large population of pregnant Japanese women who were recruited early enough within the stage of embryonic neurulation and organogenesis. Epidemiologically, the study identified demographic and lifestyle

determinants of folate status at this critical stage of neural tube formation. Identifying modifiable lifestyle factors as favourable and unfavourable determinants can lay a sound foundation for Public Health intervention policies. All information about the type or brand name of nutritional supplements used, the timing and duration of use were self-reported, hence the risk of bias. However, nutritional supplements use and smoking status were validated by biomarkers to avoid misclassification bias. For instance, the difference observed in folate biomarker concentrations among FA users and non-users was an indication of valid self-reported use. Also, comparable results were obtained with plasma cotinine and self-reported cigarette smoking or ETS exposure. Serum folate was used as an indicator of folate status. Erythrocyte folate signifies tissue folate reserves and is not subject to dietary fluctuations exhibited by serum/plasma folate concentrations, thus making it a more reliable choice. However, because erythrocyte folate assay is more complex, serum folate assay was preferred to conduct this large epidemiological study. Two previous studies have justified its use in epidemiologic studies^{50;51}. This study involved only women who presented at the designated health facilities and consented to participate, therefore may not be representative of the general population. Finally, our findings are more of statistical correlations and not in any way signifying causality. Future randomized controlled trials employing erythrocyte folate and known dosages of folic acid supplements may be more informative.

Implications

The implication of active and passive tobacco smoking in the determination of folate status is of public health importance because an increasing prevalence of tobacco smoking among younger Japanese women is being reported⁵². Optimal first trimester folate status is central in this subpopulation. It may be helpful to consider policies that could improve folate status in this group. Mandatory food fortification with FA might be a great precautionary measure. Although, there are emerging controversies about prenatal FA exposure and epigenetic effects⁵³, however, the folate depleting effects of tobacco smoke may constitute a huge public health challenge in the prevention of NTDs and other birth defects in Japan. Although this Hokkaido cohort data recorded only eight (0.04%) cases of isolated NTDs, the national rate is the second highest in developed countries after Germany.

E . 結論

In conclusion, demographic and lifestyle factors likely predict folate status of Hokkaido women. Active cigarette smoking and ETS exposure are the major modifiable unfavourable predictors of folate status; while the use of FA supplement and FA containing multivitamins are the major favourable predictors. FA supplementation may correct the folate deficits associated with tobacco smoking.

F. 研究発表

1. 論文発表

厚生労働科学研究費補助金（化学物質リスク研究事業）
分担研究報告書

Under minor review with British Journal of Nutrition (BJN).

2. 学会発表

I.Manokhina, T.A. Yila, W.P. Robinson. Towards Accurate Quantification of miRNAs for Clinical Use: Evaluation of Technical and Biological Confounders. 4th Annual Canadian Human and Statistical Genetics Meeting, 18 – 21 April, 2015 Vancouver, BC, Canada.

G. 知的財産権の出願・登録状況（予定を含む。）

1. 特許取得

なし

2. 実用新案登録

なし

3. その他

なし

引用文献

1. (1991) Vitamins to prevent neural tube defects. *Lancet* **338**, 894-895.
2. Cragan JD, Roberts HE, Edmonds LD *et al.* (1995) Surveillance for anencephaly and spina bifida and the impact of prenatal diagnosis--United States, 1985-1994. *MMWR CDC Surveill Summ* **44**, 1-13.
3. Initiative FF Fifteen Years of Fortifying With FA Reduces Birth Defects; Averts Healthcare Expenses. Flour Fortification Initiative. http://www.ffinetwork.org/about/stay_informed/publications/documents/FolicAcidBackground.pdf (accessed April 6 2015)
4. Takimoto H, Tamura T (2006) Increasing trend of spina bifida and decreasing birth weight in relation to declining body-mass index of young

women in Japan. *Med Hypotheses* **67**, 1023-1026.

5. Kondo A, Kamihira O, Ozawa H (2009) Neural tube defects: prevalence, etiology and prevention. *Int J Urol* **16**, 49-57.

6. Kondo A, Kamihira O, Shimosuka Y *et al.* (2005) Awareness of the role of folic acid, dietary folate intake and plasma folate concentration in Japan. *J Obstet Gynaecol Res* **31**, 172-177.

7. Mito N, Takimoto H, Umegaki K *et al.* (2007) Folate intakes and folate biomarker profiles of pregnant Japanese women in the first trimester. *Eur J Clin Nutr* **61**, 83-90.

8. Matsuzaki M, Haruna M, Ota E *et al.* (2008) Dietary folate intake, use of folate supplements, lifestyle factors, and serum folate levels among pregnant women in Tokyo, Japan. *J Obstet Gynaecol Res* **34**, 971-979.

9. Ihara H, Watanabe T, Aoki Y, *et al.* (2009) Dietary folate intake and serum folate status in Japanese women of childbearing age. *J Anal Bio-Sci* **32**, 5.

10. Yila TA, Sasaki S, Miyashita C *et al.* (2012) Effects of maternal 5,10-methylenetetrahydrofolate reductase C677T and A1298C Polymorphisms and tobacco smoking on infant birth weight in a Japanese population. *J Epidemiol* **22**, 91-102.

11. Kishi R, Sasaki S, Yoshioka E *et al.* (2011) Cohort profile: the Hokkaido study on environment and children's health in Japan. *Int J Epidemiol* **40**, 611-618.

12. Kishi R, Kobayashi S, Ikeno T *et al.* (2013) Ten years of progress in the Hokkaido birth cohort study on environment and children's health: cohort profile--updated 2013. *Environ Health*

厚生労働科学研究費補助金（化学物質リスク研究事業）
分担研究報告書

- Prev Med* **18**, 429-450.
13. Diagnostics B Package insert: ADVIA Centaur assay manual. Bayer Diagnostics, Revision AT.
14. Sujaku K, Ogiwara T, Y K (2006) Determination of vitamin B12 and folate by ADVIA Centaur. *J Anal Bio-Sci*. 2006; 29:235-42. *J Anal Bio-Sci* **29**, 235-242.
15. Sauberlich HE (July 27, 1999) Folate (FA, pteroylmonoglutamic acid, folacin). In *Laboratory Tests for the Assessment of Nutritional Status*, Second Edition ed., pp. Pages 103-134. Washington D.C: CRC Press
16. Sasaki S, Braimoh TS, Yila TA *et al.* (2011) Self-reported tobacco smoke exposure and plasma cotinine levels during pregnancy--a validation study in Northern Japan. *Sci Total Environ* **412-413**, 114-118.
17. Takimoto H, Yoshiike N, Kaneda F *et al.* (2004) Thinness among young Japanese women. *Am J Public Health* **94**, 1592-1595.
18. Damms-Machado A, Weser G, Bischoff SC (2012) Micronutrient deficiency in obese subjects undergoing low calorie diet. *Nutr J* **11**, 34.
19. Aasheim ET, Hofso D, Hjelmestaeth J *et al.* (2008) Vitamin status in morbidly obese patients: a cross-sectional study. *Am J Clin Nutr* **87**, 362-369.
20. Miyaki K, Song Y, Taneichi S *et al.* (2013) Socioeconomic status is significantly associated with the dietary intakes of folate and depression scales in Japanese workers (J-HOPE Study). *Nutrients* **5**, 565-578.
21. Vandevijvere S, Amsalkhir S, Van Oyen H *et al.* (2012) Determinants of folate status in pregnant women: results from a national cross-sectional survey in Belgium. *Eur J Clin Nutr* **66**, 1172-1177.
22. Gall S, Seal J, Taylor R *et al.* (2012) Folate status and socio-demographic predictors of folate status, among a national cohort of women aged 26-36 in Australia, 2004-2006. *Aust N Z J Public Health* **36**, 421-426.
23. Branum AM, Bailey R, Singer BJ (2013) Dietary supplement use and folate status during pregnancy in the United States. *J Nutr* **143**, 486-492.
24. Thoradeniya T, Wickremasinghe R, Ramanayake R *et al.* (2006) Low folic acid status and its association with anaemia in urban adolescent girls and women of childbearing age in Sri Lanka. *Br J Nutr* **95**, 511-516.
25. Hao L, Ma J, Stampfer MJ *et al.* (2003) Geographical, seasonal and gender differences in folate status among Chinese adults. *J Nutr* **133**, 3630-3635.
26. Garcia-Casal MN, Osorio C, Landaeta M *et al.* (2005) High prevalence of folic acid and vitamin B12 deficiencies in infants, children, adolescents and pregnant women in Venezuela. *Eur J Clin Nutr* **59**, 1064-1070.
27. Zhao Y, Hao L, Zhang L *et al.* (2009) Plasma folate status and dietary folate intake among Chinese women of childbearing age. *Matern Child Nutr* **5**, 104-116.
28. Gaina A, Sekine M, Chandola T *et al.* (2009) Mother employment status and nutritional patterns in Japanese junior high schoolchildren. *Int J Obes (Lond)* **33**, 753-757.
29. Mathews F, Yudkin P, Smith RF *et al.* (2000) Nutrient intakes during pregnancy: the influence of smoking status and age. *J Epidemiol Community Health* **54**, 17-23.
30. Trobs M, Renner T, Scherer G *et al.*

厚生労働科学研究費補助金（化学物質リスク研究事業）
分担研究報告書

- (2002) Nutrition, antioxidants, and risk factor profile of nonsmokers, passive smokers and smokers of the Prevention Education Program (PEP) in Nuremberg, Germany. *Prev Med* **34**, 600-607.
31. Ulvik A, Ebbing M, Hustad S *et al.* (2010) Long- and short-term effects of tobacco smoking on circulating concentrations of B vitamins. *Clin Chem* **56**, 755-763.
32. Mannino DM, Mulinare J, Ford ES *et al.* (2003) Tobacco smoke exposure and decreased serum and red blood cell folate levels: data from the Third National Health and Nutrition Examination Survey. *Nicotine Tob Res* **5**, 357-362.
33. Ortega RM, Requejo AM, Lopez-Sobaler AM *et al.* (2004) Smoking and passive smoking as conditioners of folate status in young women. *J Am Coll Nutr* **23**, 365-371.
34. Erdemir EO, Bergstrom J (2007) Effect of smoking on folic acid and vitamin B12 after nonsurgical periodontal intervention. *J Clin Periodontol* **34**, 1074-1081.
35. Vardavas CI, Linardakis MK, Hatzis CM *et al.* (2008) Smoking status in relation to serum folate and dietary vitamin intake. *Tob Induc Dis* **4**, 8.
36. Ozerol E, Ozerol I, Gokdeniz R *et al.* (2004) Effect of smoking on serum concentrations of total homocysteine, folate, vitamin B12, and nitric oxide in pregnancy: a preliminary study. *Fetal Diagn Ther* **19**, 145-148.
37. Yanbaeva DG, Dentener MA, Creutzberg EC *et al.* (2007) Systemic effects of smoking. *Chest* **131**, 1557-1566.
38. Kondo A, Shimosuga Y, Oguchi H *et al.* (2008) [Folic acid reduces the risk of neural tube defects: awareness and folate intake among pregnant women in 2006]. *Hinyokika Kyo* **54**, 537-542.
39. Kondo A, Asada Y, Shibata K *et al.* (2011) Dietary folate intakes and effects of folic acid supplementation on folate concentrations among Japanese pregnant women. *J Obstet Gynaecol Res* **37**, 331-336.
40. Japan MoH (Dec 28, 2000) Ministry of Health: Department of Maternal and Child Health, Bureau of Children and Families, Tokyo: Information on promoting intake of FA in order to reduce children afflicted with neural tube defects among young women who are capable of becoming pregnant [BoCaF Department of Maternal and Child Health, editor]. Tokyo.
41. Oi S (2003) Current status of prenatal management of fetal spina bifida in the world: worldwide cooperative survey on the medico-ethical issue. *Childs Nerv Syst* **19**, 596-599.
42. Ren A, Zhang L, Li Z *et al.* (2006) Awareness and use of folic acid, and blood folate concentrations among pregnant women in northern China--an area with a high prevalence of neural tube defects. *Reprod Toxicol* **22**, 431-436.
43. Khor GL, Duraisamy G, Loh SP *et al.* (2006) Dietary and blood folate status of Malaysian women of childbearing age. *Asia Pac J Clin Nutr* **15**, 341-349.
44. Green TJ, Skeaff CM, Venn BJ *et al.* (2007) Red cell folate and predicted neural tube defect rate in three Asian cities. *Asia Pac J Clin Nutr* **16**, 269-273.
45. Brough L, Rees GA, Crawford MA *et al.* (2009) Social and ethnic differences in folic acid use preconception and during early pregnancy in the UK: effect on

厚生労働科学研究費補助金（化学物質リスク研究事業）
分担研究報告書

maternal folate status. *J Hum Nutr Diet* **22**, 100-107.

46. McNulty B, Pentieva K, Marshall B *et al.* (2011) Women's compliance with current folic acid recommendations and achievement of optimal vitamin status for preventing neural tube defects. *Hum Reprod* **26**, 1530-1536.

47. Mayer O, Jr., Simon J, Rosolova H (2001) A population study of the influence of beer consumption on folate and homocysteine concentrations. *Eur J Clin Nutr* **55**, 605-609.

48. Halsted CH, Villanueva JA, Devlin AM *et al.* (2002) Metabolic interactions of alcohol and folate. *J Nutr* **132**, 2367S-2372S.

49.

<http://www.icap.org/PolicyTools/ICAPBlueBook/BlueBookModules/10DrinkingandPregnancy/tabid/171/Default.aspx> ICAP Blue Book 10. Drinking and Pregnancy

50. Galloway M, Rushworth L (2003)

Red cell or serum folate? Results from the National Pathology Alliance benchmarking review. *J Clin Pathol* **56**, 924-926.

51. Drogan D, Klipstein-Grobusch K, Wans S *et al.* (2004) Plasma folate as marker of folate status in epidemiological studies: the European Investigation into Cancer and Nutrition (EPIC)-Potsdam study. *Br J Nutr* **92**, 489-496.

52. Takimoto H, Yokoyama T, Yoshiike N *et al.* (2005) Increase in low-birth-weight infants in Japan and associated risk factors, 1980-2000. *J Obstet Gynaecol Res* **31**, 314-322.

53. Dolinoy DC (2007) Epigenetic gene regulation: early environmental exposures. *Pharmacogenomics* **8**, 5-10.

厚生労働科学研究費補助金（化学物質リスク研究事業）
分担研究報告書

Table1: Distributions of maternal characteristics by folate status: The Hokkaido Study on Environment and Children's Health 2002-2012, Japan (N = 15266).

Variables	Categories	n	Folate status (nmol/L)					
			Deficient (<6.80)		Suboptimal (6.80 – 13.59)		Optimal (≥13.60)	
			n	(%)	n	(%)	n	(%)
			79	(0.52)	3916	(25.65)	11271	(73.83)
Age (years)	<20	110	2	(1.82)	53	(48.18)	55	(50.00)
	20-24	1685	16	(0.95)	614	(36.44)	1055	(62.61)
	25-29	4564	25	(0.55)	1249	(27.37)	3290	(72.09)
	30-34	5393	24	(0.45)	1258	(23.33)	4111	(76.23)
	≥35	2827	9	(0.32)	572	(20.23)	2246	(79.45)***
Parity	Nulliparous	5983	27	(0.45)	1549	(25.89)	4407	(73.66)
	Parous	8035	38	(0.47)	2125	(26.45)	5872	(73.08)
BMI (Kg/m ²)	<18.50	2532	8	(0.32)	660	(26.07)	1864	(73.62)
	18.50 -24.99	10576	45	(0.43)	2649	(25.05)	7882	(74.53)
	25.00 -29.00	1225	17	(1.39)	347	(28.33)	861	(70.29)
	≥30.00	313	4	(1.28)	92	(29.39)	217	(69.33)***
Educational level	Junior high school	768	4	(0.52)	283	(36.85)	481	(62.63)
	High school	6573	49	(0.75)	1946	(29.61)	4578	(69.65)
	College	5948	17	(0.29)	1301	(21.87)	4630	(77.84)
	University	1580	7	(0.44)	283	(17.91)	1290	(81.65)***
Annual income (million JPY)	<3	2914	21	(0.72)	915	(31.40)	1978	(67.88)
	3-4,999	5709	23	(0.40)	1462	(25.61)	4224	(73.99)
	5-7,999	3215	13	(0.40)	716	(22.27)	2486	(77.33)
	≥8	889	4	(0.45)	164	(18.45)	721	(81.10)***
Occupation	Unemployed	6464	31	(0.48)	1568	(24.26)	4865	(75.26)
	Employed	8802	48	(0.55)	2348	(26.68)	6406	(72.78)**
Tobacco smoking (cigarette sticks/day)	No	13599	59	(0.44)	3249	(23.96)	10251	(75.60)
	<10	975	8	(0.82)	343	(35.18)	624	(64.00)
	10-19	630	11	(1.75)	290	(46.03)	329	(52.22)
	≥20	102	1	(0.98)	34	(33.33)	67	(65.69)***
eETS at home	No	5763	25	(0.43)	1178	(20.44)	4560	(79.13)
	Yes	9503	54	(0.57)	2738	(28.81)	6711	(70.62)***
eETS at work place	No	1530	11	(0.72)	383	(25.03)	1136	(74.25)
	Yes	13736	68	(0.50)	3533	(25.72)	10135	(73.78)
Combined ETS exposure at home and work place	None	724	4	(0.55)	149	(20.58)	571	(78.87)
	Work place	5039	21	(0.42)	1029	(20.42)	3989	(79.16)
	Home only	806	7	(0.87)	234	(29.03)	565	(70.10)
	Home and work place	8697	47	(0.54)	2504	(28.79)	6146	(70.67)***
Plasma cotinine status (nmol/L)	<1.19	5874	22	(0.37)	1142	(19.44)	4710	(80.18)
	1.19-65.21	7113	35	(0.49)	1905	(26.78)	5173	(72.73)
	>65.21	2279	22	(0.97)	869	(38.13)	1388	(60.90)***

厚生労働科学研究費補助金（化学物質リスク研究事業）
分担研究報告書

Table1 (continued)

Variables	Categories	n	Folate status (nmol/L)					
			Deficient (<6.80)		Suboptimal (6.80 – 13.59)		Optimal (≥13.60)	
			n	(%)	n	(%)	n	(%)
			79	(0.52)	3916	(25.65)	11271	(73.83)
Alcohol intake (frequency)	No	8084	37	(0.46)	1965	(24.31)	6082	(75.24)
	Monthly	5590	34	(0.61)	1586	(28.37)	3970	(71.02)
	Weekly	723	3	(0.41)	160	(22.13)	560	(77.46)
	Daily	869	5	(0.58)	205	(23.59)	659	(75.83)***
Folic Acid supplements use	No	13559	74	(0.62)	3672	(30.68)	8224	(68.71)
	Yes	1707	5	(0.15)	244	(7.40)	3047	(92.45)***
Other nutritional supplements use	No	13956	74	(0.53)	3660	(26.23)	10222	(73.24)
	Yes	1310	5	(0.38)	256	(19.54)	1049	(80.08)***
Region	Central	6718	27	(0.40)	1522	(22.66)	5169	(76.94)
	South	3589	18	(0.50)	1076	(29.98)	2495	(69.52)
	East	4765	33	(0.69)	1271	(26.67)	3461	(72.63)
	Other regions	194	1	(0.52)	47	(24.23)	146	(75.26)***
	Year of enrolment	2002 -2004	4623	15	(0.32)	1290	(27.90)	3318
	2005 - 2007	5651	35	(0.62)	1675	(29.64)	3941	(69.74)
	2008 -2010	4063	22	(0.54)	782	(19.25)	3259	(80.21)
	2011-2012	929	7	(0.75)	169	(18.19)	735	(81.05)***
Season of the year at enrolment	Spring	3850	31	(0.81)	1010	(26.23)	2809	(72.96)
	Summer	3720	17	(0.46)	987	(26.53)	2716	(73.01)
	Autum	2424	13	(0.54)	629	(25.95)	1782	(73.51)
	Winter	5272	18	(0.34)	1290	(24.47)	3964	(75.19)*

SD, Standard deviation ; n, number of participants; BMI, Body mass index; JPY, Japanese Yen; ETS, Environmental Tobacco smoke.

P values were derived from Pearson's chi squared tests and Fisher's exact tests. Means and standard deviations generated from Students t-tests and ANOVA with post-hoc analysis. All percentages are row percentages.

Values may not add up to 100% due to missing values. Levels of significance: *p <0.050; **p<0.010; ***p<0.001.

厚生労働科学研究費補助金（化学物質リスク研究事業）
分担研究報告書

Table 2: Estimated effects of demographic characteristics and lifestyle factors on folate status: The Hokkaido Study on Environment and Children's Health 2002-2012, Japan (N = 15266)

Variables	Categories	n	Folate status (nmol/L)					
			Deficient (<6.80)		Suboptimal (6.80 – 13.59)		Optimal (≥13.60)	
			n	(%)	n	(%)	n	(%)
			79	(0.52)	3916	(25.65)	11271	(73.83)
Age (years)	<20	110	2	(1.82)	53	(48.18)	55	(50.00)
	20-24	1685	16	(0.95)	614	(36.44)	1055	(62.61)
	25-29	4564	25	(0.55)	1249	(27.37)	3290	(72.09)
	30-34	5393	24	(0.45)	1258	(23.33)	4111	(76.23)
	≥35	2827	9	(0.32)	572	(20.23)	2246	(79.45)***
Parity	Nulliparous	5983	27	(0.45)	1549	(25.89)	4407	(73.66)
	Parous	8035	38	(0.47)	2125	(26.45)	5872	(73.08)
BMI (Kg/m ²)	<18.50	2532	8	(0.32)	660	(26.07)	1864	(73.62)
	18.50 -24.99	10576	45	(0.43)	2649	(25.05)	7882	(74.53)
	25.00 -29.00	1225	17	(1.39)	347	(28.33)	861	(70.29)
	≥30.00	313	4	(1.28)	92	(29.39)	217	(69.33)***
Educational level	Junior high school	768	4	(0.52)	283	(36.85)	481	(62.63)
	High school	6573	49	(0.75)	1946	(29.61)	4578	(69.65)
	College	5948	17	(0.29)	1301	(21.87)	4630	(77.84)
	University	1580	7	(0.44)	283	(17.91)	1290	(81.65)***
Annual income (million JPY)	<3	2914	21	(0.72)	915	(31.40)	1978	(67.88)
	3-4,999	5709	23	(0.40)	1462	(25.61)	4224	(73.99)
	5-7,999	3215	13	(0.40)	716	(22.27)	2486	(77.33)
	≥8	889	4	(0.45)	164	(18.45)	721	(81.10)***
Occupation	Unemployed	6464	31	(0.48)	1568	(24.26)	4865	(75.26)
	Employed	8802	48	(0.55)	2348	(26.68)	6406	(72.78)**
Tobacco smoking (cigarette sticks/day)	No	13599	59	(0.44)	3249	(23.96)	10251	(75.60)
	<10	975	8	(0.82)	343	(35.18)	624	(64.00)
	10-19	630	11	(1.75)	290	(46.03)	329	(52.22)
	≥20	102	1	(0.98)	34	(33.33)	67	(65.69)***
°ETS at home	No	5763	25	(0.43)	1178	(20.44)	4560	(79.13)
	Yes	9503	54	(0.57)	2738	(28.81)	6711	(70.62)***
°ETS at work place	No	1530	11	(0.72)	383	(25.03)	1136	(74.25)
	Yes	13736	68	(0.50)	3533	(25.72)	10135	(73.78)
Combined ETS exposure at home and work place	None	724	4	(0.55)	149	(20.58)	571	(78.87)
	Work place	5039	21	(0.42)	1029	(20.42)	3989	(79.16)
	Home only	806	7	(0.87)	234	(29.03)	565	(70.10)
	Home and work place	8697	47	(0.54)	2504	(28.79)	6146	(70.67)***
Plasma cotinine status (nmol/L)	<1.19	5874	22	(0.37)	1142	(19.44)	4710	(80.18)
	1.19-65.21	7113	35	(0.49)	1905	(26.78)	5173	(72.73)
	>65.21	2279	22	(0.97)	869	(38.13)	1388	(60.90)***

厚生労働科学研究費補助金（化学物質リスク研究事業）
分担研究報告書

Table2 (continued)

Variables	Categories	n	Folate status (nmol/L)					
			Deficient (<6.80)		Suboptimal (6.80 – 13.59)		Optimal (≥13.60)	
			n	(%)	n	(%)	n	(%)
			79	(0.52)	3916	(25.65)	11271	(73.83)
Alcohol intake (frequency)	No	8084	37	(0.46)	1965	(24.31)	6082	(75.24)
	Monthly	5590	34	(0.61)	1586	(28.37)	3970	(71.02)
	Weekly	723	3	(0.41)	160	(22.13)	560	(77.46)
	Daily	869	5	(0.58)	205	(23.59)	659	(75.83)***
Folic Acid supplements use	No	13559	74	(0.62)	3672	(30.68)	8224	(68.71)
	Yes	1707	5	(0.15)	244	(7.40)	3047	(92.45)***
Other nutritional supplements use	No	13956	74	(0.53)	3660	(26.23)	10222	(73.24)
	Yes	1310	5	(0.38)	256	(19.54)	1049	(80.08)***
Region	Central	6718	27	(0.40)	1522	(22.66)	5169	(76.94)
	South	3589	18	(0.50)	1076	(29.98)	2495	(69.52)
	East	4765	33	(0.69)	1271	(26.67)	3461	(72.63)
	Other regions	194	1	(0.52)	47	(24.23)	146	(75.26)***
	Year of enrolment	2002 -2004	4623	15	(0.32)	1290	(27.90)	3318
	2005 - 2007	5651	35	(0.62)	1675	(29.64)	3941	(69.74)
	2008 -2010	4063	22	(0.54)	782	(19.25)	3259	(80.21)
	2011-2012	929	7	(0.75)	169	(18.19)	735	(81.05)***
Season of the year at enrolment	Spring	3850	31	(0.81)	1010	(26.23)	2809	(72.96)
	Summer	3720	17	(0.46)	987	(26.53)	2716	(73.01)
	Autum	2424	13	(0.54)	629	(25.95)	1782	(73.51)
	Winter	5272	18	(0.34)	1290	(24.47)	3964	(75.19)*

n, number of participants; BMI, Body mass index; JPY, Japanese Yen; ETS, Environmental Tobacco smoke. AOR, Adjusted odds ratio; CI, Confidence interval.

Regression model adjusted for maternal age, parity, BMI, educational level, annual income, occupation, geographical region, year of enrolment into the study, season of the year at enrolment, nutritional supplements use, alcohol intake, and active and passive smoking.

Levels of significance: *p <0.050; **p<0.010; ***p<0.001. McFadden's pseudo- R^2 = 8.69%.

All percentages are row percentages. Values may not add up to 100% due to missing values. NS, not significant.

†Other nutritional supplements used included multivitamins, trace elements, herbs, proteins, ginseng and energy drinks.

厚生労働科学研究費補助金（化学物質リスク研究事業）
分担研究報告書

Table 3: Estimated effects of active and passive cigarette smoking based on plasma cotinine concentrations on folate status: The Hokkaido Study on Environment and Children's Health 2002-2012, Japan (N = 15266)

Smoking status	Plasma cotinine levels (nmol/L)	Folate status (nmol/L)		AOR	95%CI	P _{trend}
		Suboptimal (<13.60) n (%)	Optimal (≥13.60) n (%)			
		3995 (26.17)	11271 (73.83)			
Non smoker	<1.19	1164 (19.82)	4710 (80.18)	1.00	Reference	<0.001
ETS exposed	1.19 – 65.21	1940 (27.27)	5173 (72.73)	1.20	(1.10, 1.31)**	
Active smoker	>65.21	891 (39.10)	1388 (60.90)	1.91	(1.70, 2.14)***	

n, number of participants; BMI, Body mass index; JPY, Japanese Yen; ETS, Environmental Tobacco smoke. AOR, Adjusted odds ratio; CI, Confidence interval.

Regression model adjusted for maternal age, parity, BMI, educational level, annual income, occupation, geographical region, year of enrolment into the study, season of the year at enrolment, nutritional supplements use, alcohol intake, and active and passive smoking.

Levels of significance: *p <0.050; **p<0.010; ***p<0.001.

All percentages are row percentages. Values may not add up to 100% due to missing values. McFadden's pseudo- $R^2 = 8.53\%$.

厚生労働科学研究費補助金（化学物質リスク研究事業）
分担研究報告書




Fig. 1: Study selection chart: The Hokkaido Study on Environment and Children's Health, 2002 – 2012, Japan.

厚生労働科学研究費補助金（化学物質リスク研究事業）
分担研究報告書




Fig. 2: Mean plasma cotinine concentrations by nutritional supplements use among participants: The Hokkaido Study on Environment and Children's Health, 2002 – 2012, Japan.