

資料 13 兩側副腎皮質多結節性過形成文獻

1. Berthon A, et al. Age-dependent effects of Armc5 haploinsufficiency on adrenocortical function. *Hum Mol Genet.* 26(18):3495-507, 2017. PubMed: 28911199.
2. Hu Y, et al. Armc5 deletion causes developmental defects and compromises T-cell immune responses. *Nat Commun.*8:13834, 2017. PubMed: 28169274.
3. Albiger NM, et al. A multicenter experience on the prevalence of ARMC5 mutations in patients with primary bilateral macronodular adrenal hyperplasia: from genetic characterization to clinical phenotype. *Endocrine.* 55(3):959-68, 2017. PubMed: 27094308.
4. Lodish M, Stratakis CA. A genetic and molecular update on adrenocortical causes of Cushing syndrome. *Nat Rev Endocrinol.* 12(5):255-62, 2016. PubMed: 26965378.
5. Drougat L, et al. Genetics of primary bilateral macronodular adrenal hyperplasia: a model for early diagnosis of Cushing's syndrome? *Eur J Endocrinol.* 173(4):M121-31, 2015. PubMed: 26264719.
6. Correa R, et al. The ARMC5 gene shows extensive genetic variance in primary macronodular adrenocortical hyperplasia. *Eur J Endocrinol.* 173(4):435-40, 2015. PubMed: 26162405.
7. El Ghorayeb N, et al. Multiple aberrant hormone receptors in Cushing's syndrome. *Eur J Endocrinol.* 173(4):M45-60, 2015. PubMed: 25971648.
8. Espiard S, et al. ARMC5 Mutations in a Large Cohort of Primary Macronodular Adrenal Hyperplasia: Clinical and Functional Consequences. *J Clin Endocrinol Metab.* 100(6):E926-35, 2015. PubMed: 25853793.
9. Li J, Yang CH. Diagnosis and treatment of adrenocorticotrophic hormone-independent macronodular adrenocortical hyperplasia: A report of 23 cases in a single center. *Exp Ther Med.* 9(2):507-12, 2015. PubMed: 25574224.
10. Fragoso MC, et al. Genetics of primary macronodular adrenal hyperplasia. *J Endocrinol.* 224(1):R31-43, 015. PubMed: 25472909.
11. Elbelt U, et al. Molecular and clinical evidence for an ARMC5 tumor syndrome: concurrent inactivating germline and somatic mutations are associated with both primary macronodular adrenal hyperplasia and meningioma. *J Clin Endocrinol Metab.*

100(1):E119-28, 2015. PubMed: 25279498.

12. Gagliardi L, et al. ARMC5 mutations are common in familial bilateral macronodular adrenal hyperplasia. *J Clin Endocrinol Metab.* 99(9):E1784-92, 2014. PubMed: 24905064.
13. Alencar GA, et al. ARMC5 mutations are a frequent cause of primary macronodular adrenal Hyperplasia. *J Clin Endocrinol Metab.* 99(8):E1501-9, 2014. PubMed: 24708098.
14. Faucz FR, et al. Macronodular adrenal hyperplasia due to mutations in an armadillo repeat containing 5 (ARMC5) gene: a clinical and genetic investigation. *J Clin Endocrinol Metab.* 99(6):E1113-9, 2014. PubMed: 24601692.
15. Lacroix A. Heredity and cortisol regulation in bilateral macronodular adrenal hyperplasia. *N Engl J Med.* 369(22):2147-9, 2013. PubMed: 24283229.
16. Louiset E, et al. Intraadrenal corticotropin in bilateral macronodular adrenal hyperplasia. *N Engl J Med.* 369(22):2115-25, 2013. PubMed: 24283225.
17. Assie G, et al. ARMC5 mutations in macronodular adrenal hyperplasia with Cushing's syndrome. *N Engl J Med.* 369(22):2105-14, 2013. PubMed: 24283224.
18. Vezzosi D, et al. Phosphodiesterase 11A(PDE11A) gene defects in patients with ACTH-independent macronodular adrenal hyperplasia (AIMAH): functional variants may contribute to genetic susceptibility of bilateral adrenal tumors. *J Clin Endocrinol Metab.* 97(11):E2063-9, 2012. PubMed: 22996146.
19. Rauschecker M, Stratakis CA. Molecular genetics of adrenocortical tumor formation and potential pharmacologic targets. *Minerva Endocrinol.* 37(2):133-9, 2012. PubMed: 22691887.
20. Almeida MQ, et al. Integrated genomic analysis of nodular tissue in macronodular adrenocortical hyperplasia: progression of tumorigenesis in a disorder associated with multiple benign lesions. *J Clin Endocrinol Metab.* 96(4):E728-38, 2011. PubMed: 21252250.
21. Assie G, et al. Systematic analysis of G protein-coupled receptor gene expression in adrenocorticotropin-independent macronodular adrenocortical hyperplasia identifies novel targets for pharmacological control of adrenal Cushing's syndrome. *J Clin Endocrinol Metab.* 95(10):E253-62, 2010. PubMed: 20660048.

22. de Groot JW, et al. Aberrant expression of multiple hormone receptors in ACTH-independent macronodular adrenal hyperplasia causing Cushing's syndrome. *Eur J Endocrinol*. 163(2):293-9, 2010. PubMed: 20460422.
23. Lacroix A, et al. Aberrant G-protein coupled receptor expression in relation to adrenocortical overfunction. *Clin Endocrinol (Oxf)*. 73(1):1-15, 2010. PubMed: 19719763.
24. Gagliardi L, et al. Familial vasopressin-sensitive ACTH-independent macronodular adrenal hyperplasia (VPs-AIMAH): clinical studies of three kindreds. *Clin Endocrinol (Oxf)*. 70(6):883-91, 2009. PubMed: 19018784.
25. Louiset E, et al. Expression of vasopressin receptors in ACTH-independent macronodular bilateral adrenal hyperplasia causing Cushing's syndrome: molecular, immunohistochemical and pharmacological correlates. *J Endocrinol*. 196(1):1-9, 2008. PubMed: 18180312.
26. Albiger NM, et al. Food-dependent Cushing's syndrome: from molecular characterization to therapeutical results. *Eur J Endocrinol*. 157(6):771-8, 2007. PubMed: 18057385.
27. Bourdeau I, et al. 17q22-24 chromosomal losses and alterations of protein kinase a subunit expression and activity in adrenocorticotropin-independent macronodular adrenal hyperplasia. *J Clin Endocrinol Metab*. 91(9):3626-32, 2006. PubMed: 16772351.
28. Lee S, et al. Ectopic expression of vasopressin V1b and V2 receptors in the adrenal glands of familial ACTH-independent macronodular adrenal hyperplasia. *Clin Endocrinol (Oxf)*. 63(6):625-30, 2005. PubMed: 16343095.
29. Christopoulos S, Bourdeau I, Lacroix A. Clinical and subclinical ACTH-independent macronodular adrenal hyperplasia and aberrant hormone receptors. *Horm. Res*. 64: 119-131, 2005. PubMed: 16215323
30. Bertherat J, et al. In vivo and in vitro screening for illegitimate receptors in adrenocorticotropin-independent macronodular adrenal hyperplasia causing Cushing's syndrome: identification of two cases of gonadotropin/gastric inhibitory polypeptide-dependent hypercortisolism. *J Clin Endocrinol Metab*. 90(3):1302-10, 2005. PubMed: 15585558.
31. Fragoso MCBV, et al. Cushing's syndrome secondary to adrenocorticotropin-independent macronodular adrenocortical hyperplasia due to activating mutations of GNAS1 gene.

J Clin Endocrinol Metab. 88: 2147-2151, 2003. PubMed: 12727968.

32. Cartier D, et al. Overexpression of serotonin-4 receptors in cisapride-responsive adrenocorticotropin-independent bilateral macronodular adrenal hyperplasia causing Cushing's syndrome. J Clin Endocrinol Metab. 88: 248-254, 2003. PubMed: 12519861.
33. Mune T, et al. Eutopic overexpression of vasopressin V1a receptor in adrenocorticotropin-independent macronodular adrenal hyperplasia. J Clin Endocrinol Metab. 87: 5706-5713, 2002. PubMed: 12466375.
34. Nies C, et al. Familial ACTH-independent Cushing's syndrome with bilateral macronodular adrenal hyperplasia clinically affecting only female family members. Exp Clin Endocrinol Diabetes. 110(6):277-83, 2002. PubMed: 12373631.
35. Lacroix A, Hamet P, Boutin JM. Leuprolide acetate therapy in luteinizing hormone-dependent Cushing's syndrome. N Engl J Med. 341: 1577-1581, 1999. PubMed: 10564687.
36. Swain JM, et al. Corticotropin-independent macronodular adrenal hyperplasia: a clinicopathologic correlation. Arch Surg. 133(5):541-5, 1998; discussion 545-6. PubMed: 9605918.
37. Lebrethon MC, et al. Food-dependent Cushing's syndrome: characterization and functional role of gastric inhibitory polypeptide receptor in the adrenals of three patients. J Clin Endocrinol Metab. 83: 4514-4519, 1998. PubMed: 9851802.
38. Lacroix A, et al. Propranolol therapy for ectopic beta-adrenergic receptors in adrenal Cushing's syndrome. N Engl J Med. 337: 1429-1434, 1997. PubMed: 9358140.
39. Wada N, et al. Adrenocorticotropin-independent bilateral macronodular adrenocortical hyperplasia: immunohistochemical studies of steroidogenic enzymes and post-operative course in two men. Eur J Endocrinol. 134(5):583-7, 1996. PubMed: 8664979.
40. de Herder WW, et al. Food-dependent Cushing's syndrome resulting from abundant expression of gastric inhibitory polypeptide receptors in adrenal adenoma cells. J Clin Endocrinol Metab. 81: 3168-3172, 1996. PubMed: 8784063.
41. Horiba N, et al. Lysine vasopressin stimulation of cortisol secretion in patients with adrenocorticotropin-independent macronodular adrenal hyperplasia. J Clin Endocrinol Metab. 80(8):2336-41, 1995. PubMed: 7629226.

42. Findlay JC, et al. Familial adrenocorticotropin-independent Cushing's syndrome with bilateral macronodular adrenal hyperplasia. *J Clin Endocrinol Metab.* 76(1):189-91, 1993. PubMed: 8380604.
43. Sasano H, Suzuki T, Nagura H. ACTH-independent macronodular adrenocortical hyperplasia: immunohistochemical and in situ hybridization studies of steroidogenic enzymes. *Mod Pathol.* 7(2):215-9, 1994. PubMed: 8008746.
44. Reznik Y, et al. Food-dependent Cushing's syndrome mediated by aberrant adrenal sensitivity to gastric inhibitory polypeptide. *N Engl J Med.* 327: 981-986, 1992. PubMed: 1325609.
45. Lacroix A, et al. Gastric inhibitory polypeptide-dependent cortisol hypersecretion--a new cause of Cushing's syndrome. *N Engl J Med.* 327: 974-980, 1992. PubMed: 1325608.
46. Hamet P, et al. Cushing syndrome with food-dependent periodic hormonogenesis. *Clin Invest Med.* 10: 530-533, 1987. PubMed: 2831001.
47. Kirschner MA, et al. Cushing's syndrome: nodular cortical hyperplasia of adrenal glands with clinical and pathological features suggesting adrenocortical tumor. *J Clin Endocrinol.* 24: 947-955, 1964. PubMed: 14228534.
48. Aiba M, et al. Adrenocorticotropic hormone-independent bilateral adrenocortical macronodular hyperplasia as a distinct subtype of Cushing's syndrome: enzyme histochemical and ultrastructural study of four cases with a review of the literature. *Am J Clin Path.* 96: 334-340, 1991. PubMed: 1652202.