

研究成果の刊行に関する一覧表

書籍

| 著者氏名 | 論文タイトル名 | 書籍全体の編集者名 | 書籍名 | 出版社名 | 出版地 | 出版年 | ページ |
|------|---|--------------------------------|---------------------------|---------|-----|------|-----------|
| 片山一朗 | 第X章 皮膚疾患 | 松田暉 / 萩原俊男 / 難波光義 / 鈴木久美 / 林直子 | 疾病と治療 | 南江堂 | | 2017 | 編集 |
| 片山一朗 | Ⅲ皮膚癢痒症 1皮膚癢痒症. | 渡辺晋一・古川福実 | 皮膚疾患 最新の治療2017-2018 | 南江堂 | 東京 | 2017 | pp55-6 |
| 片山一朗 | 第X章 皮膚疾患 2. 皮膚疾患各論 E. 蕁麻疹 | 松田暉、萩原俊男、難波光義、鈴木久美、林直子 | 疾病と治療 | 南江堂 | 東京 | 2017 | pp81-2 |
| 片山一朗 | コラム2. | 秋山一男、大田健、近藤直実 | アトピー性皮膚炎とフィラグリン アレルギーのはなし | 朝倉書店 | 東京 | 2017 | p97 |
| 片山一朗 | 12リウマチ・アレルギー 12-34その他のアレルギー疾患 1)じんま診・血管浮腫 | 矢崎義雄 | 内科学 第11版 | 朝倉書店 | 東京 | 2017 | 1334-6 |
| 鈴木民夫 | 尋常性白斑 | 福井次矢ら | 今日の治療指針2017私はこう治療している | 医学書院 | 東京 | 2017 | 1226 |
| 鈴木民夫 | 母斑細胞母斑 | 渡辺晋一ら | 皮膚疾患最新の治療2017 2018 | 南江堂 | 東京 | 2017 | 241-242 |
| 鈴木民夫 | 尋常性白斑 | 猿田享男, 北村惣一郎 | 1336専門家による私の治療2017-18年度版 | 日本医事新報社 | 東京 | 2017 | 1023-1025 |

雑誌

| 発表者氏名 | 論文タイトル名 | 発表誌名 | 巻号 | ページ | 出版年 |
|--|--|---------------------------------------|----|-------|------|
| Ito S, Okura M, Wakamatsu K, Yamashita T | The potent pro-oxidant activity of rhododendrol-eumelanin induces cysteine depletion in B16 melanoma cells | <i>Pigment Cell Melanoma Research</i> | 30 | 63-67 | 2017 |

| | | | | | |
|--|---|--|------------|------------------|------|
| Ito S, Agata M, Okochi K, Wakamatsu K | The potent pro-oxidant activity of rhododendrol-eumelanin is enhanced by ultraviolet A radiation | <i>Pigment Cell Melanoma Research</i> | 31 | Published online | 2018 |
| Ito S, Wakamatsu, K | Biochemical mechanism of rhododendrol-induced leukoderma | <i>International Journal of Molecular Sciences</i> | 19 | 522 | 2018 |
| Goto N, Tsujimoto M, Masaki T, Ito S, Wakamatsu K, Nishigori C | 4-(4-Hydroxyphenyl)-2-butanol (rhododendrol)-induced melanocyte cytotoxicity is enhanced by UVB exposure through generation of oxidative stresses | <i>Experimental Dermatology</i> | in press | | 2018 |
| Takeuchi Y, et al | Clinical response to PD-1 blockade correlates with a sub-fraction of peripheral central memory CD4+ T cells in patients with malignant melanoma. | <i>Int Immunol.</i> | 30(1) | 13-22. | 2018 |
| Yamaga K, et al | Claudin-3 loss causes leakage of sweat from the sweat gland to contribute to the pathogenesis of atopic dermatitis. | <i>J Invest Dermatol.</i> | in press | | 2017 |
| Yang F, et al | Dysregulation of autophagy in melanocytes contributes to hypopigmented macules in tuberous sclerosis complex. | <i>J Dermatol Sci</i> | in press | | 2017 |
| Yang F, et al | Uncoupling of ER/mitochondrial oxidative stress in mTORC1 hyperactivation-associated skin hypopigmentation. | <i>J Invest Dermatol.</i> | in press | | 2017 |
| Murota H, et al | Periostin in the pathogenesis of skin diseases. | <i>Cell Mol Life Sci.</i> | 74 | 4321-4328 | 2017 |
| Murota H, et al | Impact of Jumihaidokuto (Shi-Wei-Bai-Du-Tang) on treatment of chronic spontaneous urticaria: A randomized controlled study. | <i>Chin J Integr Med.</i> | in press | | 2017 |
| Kurata R, et al | Three-dimensional cell shapes and arrangements in human sweat glands as revealed by whole-mount immunostaining. | <i>PLoS One.</i> | 12 | e0178709 | 2017 |
| Nakagami H, et al | Physician-initiated first-in-human clinical study using a novel angiogenic peptide, AG30/5C, for patients with severe limb ulcers. | <i>Geriatr Gerontol Int.</i> | Nov;17(11) | 2150-2156 | 2017 |
| Itoi-Ochi S, et al | Occult HIV infection in Japanese rupioid psoriasis. | <i>J Dermatol.</i> | Jul;44(7) | e172-e173 | 2017 |

| | | | | | |
|------------------------|---|-----------------------------------|------|---------|------|
| Arase N, et al | Novel autoantibody against the β 2-glycoprotein I/human leucocyte antigen-DR complex in patients with refractory cutaneous ulcers. | <i>Br J Dermatol.</i> | 178 | 272-275 | 2018 |
| Kato K, et al | Novel interferon- γ enzyme-linked immunoSpot assay using activated cells for identifying hypersensitivity-inducing drug culprits. | <i>J Dermatol Sci.</i> | 86 | 222-229 | 2017 |
| Kamei R, et al | Successful treatment of a refractory dysbiotic intestinal pseudo-obstruction in a patient with systemic sclerosis-polymyositis overlap syndrome by intravenous immunoglobulin administration possibly related to gut flora normalisation. 35 Suppl. | <i>Clin Exp Rheumatol.</i> | 106 | 214-215 | 2017 |
| Kaneko S, et al | Usefulness of Sweat Management for Patients with Adult Atopic Dermatitis, regardless of Sweat Allergy: A Pilot Study. | <i>Biomed Res Int.</i> | 2017 | 8746745 | 2017 |
| Katayama I, et al | Japanese Society of Allergology. Japanese guidelines for atopic dermatitis 2017. | <i>Allergol Int.</i> | 66 | 230-247 | 2017 |
| Katayama I, et al | Itch: Its perception and involvement in allergy. | <i>Allergol Int.</i> | 66 | 1-2 | 2017 |
| Yamaoka T, et al | Value of ultrasonography findings for nail psoriasis before and after adalimumab administration. | <i>Clin Exp Dermatol.</i> | 42 | 201-203 | 2017 |
| Gan EY, et al | VGICC Repigmentation in vitiligo: position paper of the Vitiligo Global Issues Consensus Conference. | <i>Pigment Cell Melanoma Res.</i> | 30 | 28-40 | 2017 |
| Murota H, et al | Exacerbating factors of itch in atopic dermatitis. | <i>Allergol Int.</i> | 66 | 8-13 | 2017 |
| Wataya-Kaneda M, et al | Efficacy and Safety of Topical Sirolimus Therapy for Facial Angiofibromas in the Tuberous Sclerosis Complex : A Randomized Clinical Trial. | <i>JAMA Dermatol.</i> | 153 | 39-48 | 2017 |
| Deguchi A, et al | Anti-RNA polymerase III antibody positive limited cutaneous systemic sclerosis with cryoglobulin-induced digital gangrene. | <i>Clin Exp Dermatol.</i> | 42 | 200-201 | 2017 |
| Tanaka A, et al | Critical renal adverse event induced by nivolumab therapy in a stage IV melanoma patient. | <i>J Dermatol.</i> | 44 | 727-728 | 2017 |

| | | | | | |
|------------------|--|-----------------------------------|-------------------|-----------------------------|------|
| Hemmi A, et al. | Waardenburg syndrome type IIE in a Japanese patient caused by a novel non-frame-shift duplication mutation in the SOX10 gene. | <i>J Dermatol</i> | On line published | doi: 10.1111/1346-8138.1415 | 2017 |
| Omura R, et al | Ultrastructural study of dyschromatosis symmetrica hereditaria with widespread pigmentary eruption. | <i>J Dermatol.</i> | 44 | 150-151 | 2017 |
| Okamura K, et al | Characterization of melanosomes and melanin in Japanese patients with Hermansky-Pudlak syndrome types 1, 4, 6, and 9. | <i>Pigment Cell Melanoma Res.</i> | 31 | 267-276 | 2018 |
| Okamura K, et al | Microsatellite polymorphism located immediately upstream of the phosphatidylinositol glycan, class K gene (PIGK) affects its expression, which correlates with tyrosinase activity in human melanocytes. | <i>J Dermatol Sci</i> | 85 | 131-134 | 2017 |
| Ozaki S, et al | Melanotic Malignant Melanoma in Oculocutaneous Albinism Type 4. | <i>Acta Derm Venereol.</i> | 97 | 287-288 | 2017 |
| Gan EY, et al | VGICC.: Repigmentation in vitiligo: position paper of the Vitiligo Global Issues Consensus Conference. | <i>Pigment Cell Melanoma Res.</i> | 30 | 28-40 | 2017 |
| Hayashi M, et al | Spectrophotometer is useful for assessing vitiligo and chemical leukoderma severity by quantifying color difference with surrounding normally pigmented skin. | <i>Skin Res Technol.</i> | On line published | doi: 10.1111/srt.12410. | 2017 |