

厚生労働行政推進調査事業費補助金（肝炎等克服政策研究事業）
分担研究報告書

肝がん・重度肝硬変の治療に係るガイドラインの作成等に資する研究

工藤正俊 近畿大学医学部 消化器内科 主任教授

研究要旨

データ収集プラットフォーム作成

A. 研究目的

肝癌重度肝硬変のうち特に肝癌の治療に係る各国の研究状況と診療の実態調査。

B. 研究方法

肝癌に関する各種国際学会に出席し、各国における各ステージにおける治療成績、治療の適応などにおいて我が国と比較を行った。

C. 研究結果

日本の肝癌のサーベイランス、診断、治療は他国と比べて優れているということ、および治療方針なども他国と大きく異なっていることが明らかとなった。

D. 考察

今後日本において肝癌や重度肝硬変において効率的に進めるためには、患者の経済的サポートも重要と考えられた。

E. 結論

日本の肝癌の診療レベルは世界的に優れているが、更に患者の経済的援助が必要と考えられた。

G. 研究発表

1) 論文発表

1. Harding JJ, Watanabe T, El-Dika I, Nishida N, Abou-Alfa GK, Kudo M: 2.5 “Gastrointestinal Malignancies”; 2.5.3 Hepatocellular carcinoma. In “ESMO Handbook of Immunology”, Haanen J, Lugowska I, Garassino MC, Califano R, ed., 2018;191-202.
2. Kudo M, Cheng AL, Park JW, Park JH, Liang PC, Hidaka H, Izumi N, Heo J, Lee YJ, Sheen IS, Chiu CF, Arioka H, Morita S, Arai Y: Orantinib versus placebo combined with transcatheter arterial chemoembolisation in patients with unresectable hepatocellular carcinoma (ORIENTAL): a randomised, double-blind, placebo-controlled, multicentre, phase 3 study. *Lancet Gastroenterol Hepatol* 2018;3:37-46.
3. akayasu K, Arai S, Sakamoto M, Matsuyama Y, Kudo M, Kaneko S, Nakashima O, Kadoya M, Izumi N, Takayama T, Ku Y, Kumada T, Kubo S, Kokudo T, Hagiwara Y, Kokudo N; Liver Cancer Study Group of Japan: Impact of resection and ablation for single hypovascular hepatocellular carcinoma ≤ 2 cm analysed with propensity score weighting. *Liver Int* 2018;38:484-493.

4. Kudo M: Lenvatinib may drastically change the treatment landscape of hepatocellular carcinoma. *Liver Cancer* 2018;7:1-19.
5. Kudo M: Combination cancer immunotherapy in hepatocellular carcinoma. *Liver Cancer* 2018;7:20-27.
6. Kudo M, Finn RS, Qin S, Han KH, Ikeda K, Piscaglia F, Baron A, Park JW, Han G, Jassem J, Blanc JF, Vogel A, Komov D, Evans TRJ, Lopez C, Dutcus C, Guo M, Saito K, Kraljevic S, Tamai T, Ren M, Cheng AL: Lenvatinib versus sorafenib in first-line treatment of patients with unresectable hepatocellular carcinoma: a randomised phase 3 non-inferiority trial. *Lancet* 2018;391:1163-1173.
7. Ikeda M, Kudo M, Aikata H, Nagamatsu H, Ishii H, Yokosuka O, Torimura T, Morimoto M, Ikeda K, Kumada H, Sato T, Kawai I, Yamashita T, Horio H, Okusaka T, Miriplatin TACE Study Group: Transarterial chemoembolization with miriplatin vs. epirubicin for unresectable hepatocellular carcinoma: a phase III randomized trial. *J Gastroenterol* 2018;53:281-290.
8. Minami Y, Minami T, Hagiwara S, Ida H, Ueshima K, Nishida N, Murakami T, Kudo M: Ultrasound-ultrasound image overlay fusion improves real-time control of radiofrequency ablation margin in the treatment of hepatocellular carcinoma. *Eur Radiol* 2018;28:1986-1993.
9. Kudo M, Kang YK, Park JW, Qin S, Inaba Y, Assenat E, Umeyama Y, Lechuga MJ, Valota O, Fujii Y, Martini JF, Williams JA, Obi S: Regional differences in efficacy, safety and biomarkers for second-line axitinib in patients with advanced hepatocellular carcinoma: from a randomized phase II study. *Liver Cancer*, 2018;7:148-164.
10. Kudo M, Ueshima K, Yokosuka O, Ogasawara S, Obi S, Izumi N, Aikata H, Nagano H, Hatano E, Sasaki Y, Hino K, Kumada T, Yamamoto K, Imai Y, Iwadou S, Ogawa C, Okusaka T, Kanai F, Akazawa K, Yoshimura KI, Johnson P, Arai Y; SILIUS study group: Sorafenib plus low-dose cisplatin and fluorouracil hepatic arterial infusion chemotherapy versus sorafenib alone in patients with advanced hepatocellular carcinoma (SILIUS): a randomised, open label, phase 3 trial. *Lancet Gastroenterol Hepatol* 2018;3:424-432.
11. Raoul JL, Kudo M, Finn RS, Edeline J, Reig M, Galle PR: Systemic therapy for intermediate and advanced hepatocellular carcinoma: Sorafenib and beyond. *Cancer Treat Rev* 2018;68:16-24.
12. Kudo M: Cabozantinib as a second-line agent in advanced hepatocellular carcinoma. *Liver Cancer* 2008;7:123-133.
13. Kudo M: Management of hepatocellular carcinoma in Japan as a world-leading model. *Liver Cancer* 2018;7:134-147.
14. Chau I, Park JO, Ryoo BY, Yen CJ, Poon R, Pastorelli D, Blanc JF, Kudo M, Pfiffer T, Hatano E, Chung HC, Kopeckova K, Phelip JM, Brandi G, Ohkawa S, Li CP, Okusaka T, Hsu Y, Abada PB, Zhu AX: Alpha-fetoprotein kinetics in patients with hepatocellular carcinoma receiving ramucirumab or placebo: an analysis of the phase 3 REACH study. *Brit J Cancer* 2018;119:19-26.
15. Yen CJ, Muro K, Kim TW, Kudo M, Shih JY, Lee KW, Chao Y, Kim SW, Yamazaki K, Sohn J, Cheng R, Zhang Y, Binder P, Mi G, Orlando M, Chung HC: Ramucirumab safety in East Asian patients: A meta-analysis of six global, randomized, double-blind, placebo-controlled,

- phase III clinical trials. *J Glob Oncol* 2018;1-12,2018.
16. Zhu AX, Finn RS, Edeline J, Cattan S, Ogasawara S, Palmer D, Verslype C, Zagonel V, Fartoux L, Vogel A, Sarker D, Verset G, Chan SL, Knox J, Daniele B, Webber AL, Ebbinghaus SW, Ma J, Siegel AB, Cheng AL, Kudo M, for the KEYNOTE-224 investigators: Pembrolizumab in patients with advanced hepatocellular carcinoma previously treated with sorafenib (KEYNOTE-224): a non-randomised, open-label phase 2 trial. *Lancet Oncol* 2018;19:940-952.
 17. Nishida N, Kudo M: Immune checkpoint blockade for the treatment of human hepatocellular carcinoma. *Hepatol Res* 48:622-634, 2018.
 18. Dietrich CF, Averkiou M, Nielsen MB, Barr RG, Burns PN, Calliada F, Cantisani V, Choi B, Chammas MC, Clevert DA, Claudon M, Correas JM, Cui XW, Cosgrove D, D'Onofrio M, Dong Y, Eisenbrey J, Fontanilla T, Gilja OH, Ignee A, Jenssen C, Kono Y, Kudo M, Lassau N, Lyshchik A, Franca Meloni M, Moriyasu F, Nolsøe C, Piscaglia F, Radzina M, Saftoiu A, Sidhu PS, Sporea I, Schreiber-Dietrich D, Sirlin CB, Stanczak M, Weskott HP, Wilson SR, Willmann JK, Kim TK, Jang HJ, Vezeridis A, Westerway S: How to perform contrast-enhanced ultrasound (CEUS). *Ultrasound Int Open* 2018;4:E2-15.
 19. Kudo M: Proposal of primary endpoints for TACE combination trials with systemic therapy: Lessons learned from 5 negative trials and the positive TACTICS trial. *Liver Cancer* 2018;7:225-234.
 20. Kudo M: Extremely high objective response rate of Lenvatinib: Its clinical relevance and changing the treatment paradigm in hepatocellular carcinoma. *Liver Cancer* 2018;7:215-224.
 21. Kudo M: Ramucirumab as second-line systemic therapy in hepatocellular carcinoma. *Liver Cancer* 2018;7:305-311.
 22. Kudo M: Systemic therapy for hepatocellular carcinoma: Latest advances. *Cancers* 2018;10:E412.
 23. Chan AWH, Zhong J, Berhane S, Toyoda H, Cucchetti A, Shi K, Tada T, Chong CCN, Xiang BD, Li LQ, Lai PBS, Mazzaferro V, García-Fiñana M, Kudo M, Kumada T, Roayaie S, Johnson PJ: Development of pre and post-operative models to predict early recurrence of hepatocellular carcinoma after surgical resection. *J Hepatol* 2018;69:1284-1293.
 24. Hagiwara S, Nishida N, Watanabe T, Ida H, Sakurai T, Ueshima K, Takita M, Komeda Y, Nishijima N, Osaki Y, Kudo M: Sustained antiviral effects and clearance of hepatitis surface antigen after combination therapy with entecavir and pegylated interferon in chronic hepatitis B. *Antivir Ther* 2018;23:513-521.
 25. Chau I, Peck-Radosavljevic M, Borg C, Malferttheiner P, Seitz JF, Park JO, Ryoo BY, Yen CJ, Kudo M, Poon R, Pastorelli D, Blanc JF, Chung HC, Baron AD, Okusaka T, Bowman L, Cui ZL, Girvan AC, Abada PB, Yang L, Zhu AX: Corrigendum to "Ramucirumab as second-line treatment in patients with advanced hepatocellular carcinoma following first-line therapy with sorafenib: Patient-focused outcome results from the randomised phase III REACH study". *Eur J Cancer* 2018;100:135-136.
 26. erraioli G, Wong VW, Castera L, Berzigotti A, Sporea I, Dietrich CF, Choi BI, Wilson SR, Kudo M, Barr RG: Liver ultrasound

- elastography: An update to the World Federation for Ultrasound in Medicine and Biology guidelines and recommendations. *Ultrasound Med Biol* 2018;44:2419-2440.
27. Tak WY, Ryoo BY, Lim HY, Kim DY, Okusaka T, Ikeda M, Hidaka H, Yeon JE, Mizukoshi E, Morimoto M, Lee MA, Yasui K, Kawaguchi Y, Heo J, Morita S, Kim TY, Furuse J, Katayama K, Aramaki T, Hara R, Kimura T, Nakamura O, Kudo M: Phase I/II study of first-line combination therapy with sorafenib plus resminostat, an oral HDAC inhibitor, versus sorafenib monotherapy for advanced hepatocellular carcinoma in east Asian patients. *Invest New Drugs* 2018;36:1072-1084.
 28. Nishida N, Nishimura T, Kaido T, Minaga K, Yamao K, Kamata K, Takenaka M, Ida H, Hagiwara S, Minami Y, Sakurai T, Watanabe T, Kudo M: Molecular scoring of hepatocellular carcinoma for predicting metastatic recurrence and requirements of systemic chemotherapy. *Cancers* 2018;10: E367.
- 2) 学会発表
1. Finn RS, Kudo M, Kang YK, Yen CJ, Galle PR, Llovet JM, Assenat E, Brandi G, Lim HY, Pracht M, Rau KM, Merle P, Motomura K, Ohno I, Daniele B, Shin D, Gerken G, Abada P, Hsu Y, Zhu AX: Ramucirumab (RAM) as second-line treatment in patients with advanced hepatocellular carcinoma (HCC) and elevated baseline α -fetoprotein (AFP): an analysis of AFP kinetics in the phase 3 REACH-2 study. American Society of Clinical Oncology, Gastrointestinal Cancers Symposium (ASCO-GI 2019), San Francisco, USA, January 17-19, 2019.
 2. Kudo M, Matilla A, Santoro A, Melero I, Gracian A, Acosta MR, Choo SP, El-Khoueiry AB, Kuromatsu R, El-Rayes B, Numata K, Itoh Y, Di Costanzo F, Crysler O, Reig M, Shen Y, Neely J, dela Cruz C, Baccan C, Sangro B: CheckMate-040: Nivolumab (NIVO) in patients (Pts) with advanced hepatocellular carcinoma (aHCC) and Child-Pugh B (CPB) status. American Society of Clinical Oncology, 2019 Gastrointestinal Cancers Symposium (ASCO-GI 2019), San Francisco, USA, January 17-19, 2019.
 3. Kudo M, Okusaka T, Motomura K, Ohno I, Morimoto M, Seo S, Wada Y, Sato S, Yamashita T, Furukawa M, Aramaki T, Nadano S, Ohkawa K, Fujii H, Kudo T, Furuse J, Takai H, Homma G, Yoshikawa R, Zhu AX: Ramucirumab as second-line treatment in patients with advanced hepatocellular carcinoma (HCC) and elevated alpha-fetoprotein (AFP) following first-line sorafenib: Pooled efficacy and safety in Japanese patients across two global randomized phase III studies (REACH-2 and REACH). American Society of Clinical Oncology, 2019 Gastrointestinal Cancers Symposium (ASCO-GI 2019), San Francisco, USA, January 17-19, 2019.
 4. Okusaka T, Ikeda K, Kudo M, Finn RS, Qin S, Han KH, Cheng AL, Piscaglia F, Kobayashi M, Sung M, Chen M, Wyrxicz L, Yoon JH, Ren Z, Stepan D, Dutcus C, Tamai T, Ren M, Hayato S, Kumada H: Safety and efficacy of lenvatinib by starting dose (8 mg or 12 mg) based on body weight in patients with unresectable hepatocellular carcinoma in REFLECT. American Society of Clinical Oncology, 2019 Gastrointestinal Cancers Symposium (ASCO-GI 2019), San Francisco, USA, January 17-19, 2019.
 5. Sung M, Finn RS, Qin S, Han KH, Ikeda K,

- Cheng AL, Kudo M, Tateishi R, Ikeda M, Breder V, Rau KM, Ma YT, Alsina A, Ryoo BY, Ren Z, Mody K, Dutcus C, Tamai T, Saito K, Piscaglia F: Association between overall survival and adverse events with lenvatinib treatment in patients with hepatocellular carcinoma (REFLECT). American Society of Clinical Oncology, 2019 Gastrointestinal Cancers Symposium (ASCO-GI 2019), San Francisco, USA, January 17-19, 2019.
6. Kudo M, Finn RS, Qin S, Han KH Ikeda K, Cheng AL, Piscaglia F, Ueshima K, Aikata H, Vogel A, Lopez C, Pracht M, Meng Z, Daniele B, Park JW, Palmer D, Dutcus C, Tamai T, Saito K, Lencioni R: Analysis of survival and objective response (OR) in patients with hepatocellular carcinoma in a phase 3 study of lenvatinib (REFLECT). American Society of Clinical Oncology, 2019 Gastrointestinal Cancers Symposium (ASCO-GI 2019), San Francisco, USA, January 17-19, 2019.
 7. Alsina A, Kudo M, Vogel A, Cheng AL, Tak WY, Ryoo BY, Evans TJ, Lopez CL, Daniele B, Misir S, Ren M, Izumi N, Qin S, Finn RS: Subsequent anticancer medication following first-line lenvatinib: a posthoc responder analysis from the phase 3 REFLECT study in unresectable hepatocellular carcinoma. American Society of Clinical Oncology, 2019 Gastrointestinal Cancers Symposium (ASCO-GI 2019), San Francisco, USA, January 17-19, 2019.
 8. Ogasawara S, Ueshima K, Ikeda M, Yasui Y, Terashima T, Yamashita T, Obi S, Sato S, Aikata H, Ohmura T, Kuroda H, Ohki T, Nagashima K, Kurosaki M, Chayama K, Kaneko S, Izumi N, Kato N, Kudo M, Omata M: Sorafenib versus hepatic arterial infusion chemotherapy in patients with advanced hepatocellular carcinoma: A Japanese multi-center large cohort study. American Society of Clinical Oncology, 2019 Gastrointestinal Cancers Symposium (ASCO-GI 2019), San Francisco, USA, January 17-19, 2019.
 9. Kudo M: Invited Lecture “Newer diagnostic algorithm for a solid tumor <3 cm in a cirrhotic”. Hot Topics at AASLD 2018, Hyderabad, India, December 16, 2018.
 10. Kudo M: Invited Lecture “HCC and cholangiocarcinoma”. Hot Topics at AASLD 2018, Hyderabad, India, December 16, 2018.
 11. Yoo C, Oh DY, Choi HJ, Kudo M, Ueno M, Kondo S, Chen LT, Osada M, Helwig C, Dussault I, Ikeda M: M7824 (MSB0011359C), a bifunctional fusion protein targeting transforming growth factor β (TGF- β) and PD-L1, in Asian patients with pretreated biliary tract cancer (BTC): Efficacy by BTC subtype. European Society for Medical Oncology Congress (ESMO-Asia 2018), Singapore, November 23-25, 2018.
 12. Kang YK, Kudo M, Lim HY, Hsu CH, Vogel A, Brandi G, Cheng R, Carton I, Abada P, Hsu Y, Zhu A, Yen CJ: Efficacy and safety of ramucirumab (RAM) in Asian and non-Asian patients with advanced hepatocellular carcinoma (HCC) and elevated alpha-fetoprotein (AFP): Subgroup analysis from two randomized studies. European Society for Medical Oncology Congress (ESMO-Asia 2018), Singapore, November 23-25, 2018.
 13. Heo J, Cheng AL, Raoul JL, Peck-Radosavljevic M, Kudo M, Nakajima K, Bayh I, Lin SM, Lee HC: Practice patterns, radiologic tumor response, and deterioration of liver function after transarterial chemoembolization (TACE): Final analysis of OPTIMIS in Korea and other regions.

- European Society for Medical Oncology Congress (ESMO-Asia 2018), Singapore, November 23-25, 2018.
14. Llovet JM, Kudo M, Finn R, Galle PR, Blanc JF, Okusaka T, Chau I, Abada PB, Hsu Y, Zhu AX: Ramucirumab as second-line treatment in patients with hepatocellular carcinoma (HCC) and elevated alpha-fetoprotein (AFP) following sorafenib: pooled results from two global phase 3 studies (REACH-2 and REACH). American Association for the study of liver diseases (AASLD 2018), San Francisco, USA, November 9-13, 2018.
 15. Peck-Radosavljevic, Kudo M, Raoul JL, Lee HC, Decaens T, Heo J, Lin SM, Shan H, Yang Y, Bayh I, Nakajima K, Cheng AL: Outcomes of patients (pts) with hepatocellular carcinoma (HCC) treated with transarterial chemoembolization (TACE): Global OPTIMIS final analysis. American Association for the study of liver diseases (AASLD 2018), San Francisco, USA, November 9-13, 2018.
 16. Hiraoka A, Kumada T, Tsuji K, Takaguchi K, Itobayashi E, Kariyama K, Ochi H, Tajiri K, Hirooka M, Shimada N, Ishikawa T, Tachi Y, Tada T, Toyoda H, Nouse K, Joko K, Hiasa Y, Michitaka K, Kudo M: Validation of modified ALBI grade for more detailed assessment of hepatic function in hepatocellular carcinoma patients: multicenter analysis. American Association for the study of liver diseases (AASLD 2018), San Francisco, USA, November 9-13, 2018.
 17. Nishida N, Kudo M: Stem cell feature and immune-suppressive microenvironment in human hepatocellular carcinoma. American Association for the study of liver diseases (AASLD 2018), San Francisco, USA, November 9-13, 2018.
 18. Koizumi Y, Hirooka M, Yada N, Tamaki N, Izumi N, Kudo M, Hiasa Y: New diagnostic method for hepatic steatosis using attenuation measurement by ultrasound B mode: comparison with controlled attenuation parameter. American Association for the study of liver diseases (AASLD 2018), San Francisco, USA, November 9-13, 2018.
 19. Kudo M, Matilla A, Santoro A, Melero I, Gracian AC, Acosta MR, Choo SP, El-Khoueiry AB, Kuromatsu R, El-Rayes B, Numata K, Itoh Y, Di Costanzo F, Crysler O, Reig M, Shen Y, Neely J, dela Cruz C, Baccan C, Sangro B: Nivolumab in patients with Child-Pugh B advanced hepatocellular carcinoma (aHCC) in the CheckMate-040 study. American Association for the study of liver diseases (AASLD 2018), San Francisco, USA, November 9-13, 2018.
 20. Cheng AL, Yen CJ, Okusaka T, Ikeda M, Hsu CH, Wu SY, Morizane C, Hashimoto Y, Ueshima K, Ohtomo T, Tanaka T, Kudo M: A phase I, open-label, multi-center, dose-escalation study of codrituzumab, an anti-glypican-3 monoclonal antibody, in combination with atezolizumab in patients with locally advanced or metastatic hepatocellular carcinoma. European Society for Medical Oncology (ESMO 2018), Munich, Germany, October 19-23, 2018.
 21. Raoul JL, Decaens T, Burak K, Koskinas J, Villadsen GE, Heurgue-Berlot A, Bayh I, Cheng AL, Kudo M, Lee HC, Nakajima K, Peck-Radosavljevic M: Practice patterns and deterioration of liver function after transarterial chemoembolization (TACE) in hepatocellular carcinoma (HCC): Final analysis of OPTIMIS in Europe and Canada.

- European Society for Medical Oncology (ESMO 2018), Munich, Germany, October 19-23, 2018.
22. Qin S, Finn R, Kudo M, Meyer T, Vogel A, Ducreux M, Macarulla T, Tomasello G, Boisserie F, Hou J, Li C, Song J, Zhu A: Global phase 3 study of tislelizumab versus sorafenib as first-line treatment in patients with advanced hepatocellular carcinoma (HCC): A trial-in-progress. European Society for Medical Oncology (ESMO 2018), Munich, Germany, October 19-23, 2018.
 23. Zhu AX, Finn R, Galle P, Llovet JM, Blanc JF, Okusaka T, Chau I, Cella D, Girvan A, Gable J, Bowman L, Hsu Y, Abada P, Kudo M: Ramucirumab as second-line treatment in patients with advanced hepatocellular carcinoma (HCC) and elevated alpha-fetoprotein (AFP) following first-line sorafenib: patient reported outcome results across two phase 3 studies (REACH-2 and REACH). European Society for Medical Oncology (ESMO 2018), Munich, Germany, October 19-23, 2018.
 24. Yoo C, Oh DY, Choi HJ, Kudo M, Ueno M, Kondo S, Chen LT, Osada M, Helwig C, Dussault I, Ikeda M: M7824 (MSB0011359C), a bifunctional fusion protein targeting PD-L1 and TGF- β , in Asian patients with pretreated biliary tract cancer: Preliminary results from a phase I trial. European Society for Medical Oncology (ESMO 2018), Munich, Germany, October 19-23, 2018.
 25. Qin S, Finn RS, Kudo M, Meyer T, Vogel A, Ducreux M, Macarulla TM, Tomasello G, Boisserie F, Hou J, Li C, Song J, Zhu AX: Efficacy and safety of Tislelizumab, an anti-PD-1 antibody, versus sorafenib as a potential first-line treatment in patients with advanced hepatocellular carcinoma in a phase 3, randomized, multicenter study: A Trial-in-Progress. 12th Annual Conference International Liver Cancer Association (ILCA), London, United Kingdom, September 14-16, 2018.
 26. Galle PR, Kudo M, Kang YK, Yen CJ, Finn R, Llovet JM, Assenat E, Brandi G, Lim HY, Pracht M, Rau KM, Merle P, Motomura K, Ohno I, Daniele B, Shin DB, Gerken G, Abada P, Hsu Y, Zhu AX: Ramucirumab versus placebo as second-line treatment in patients with advanced hepatocellular carcinoma and elevated baseline alpha-fetoprotein following first-line sorafenib (REACH-2): a phase 3, randomized, double-blind, placebo-controlled trial. 12th Annual Conference International Liver Cancer Association (ILCA), London, United Kingdom, September 14-16, 2018.
 27. Finn R, Frenette C, Granito A, Ikeda M, Lim HY, Merle P, Ozgurdal K, Kudo M: A prospective, observational study to assess the safety and effectiveness of regorafenib in patients with unresectable hepatocellular carcinoma (uHCC) in routine clinical practice (REFINE). 12th Annual Conference International Liver Cancer Association (ILCA), London, United Kingdom, September 14-16, 2018.
 28. Peck-Radosavljevic M, Kudo M, Raoul JL, Lee HC, Decaens T, Heo J, Lin SM, Shan H, Yang Y, Bayh I, Nakajima K, Cheng AL: Outcomes of patients (PTS) with hepatocellular carcinoma (HCC) treated with transarterial chemoembolization (TACE): Global optimis final analysis. 12th Annual Conference International Liver Cancer Association (ILCA), London, United Kingdom, September 14-16, 2018.

29. Ueshima K, Kudo M, Ikeda M, Torimura T, Tanabe N, Aikata H, Izumi N, Yamasaki T, Nojiri S, Hino K, Tsumura H, Kuzuya T, Isoda N, Yasui K, Yoshimura K, Okusaka T, Furuse J, Kokudo N, Okita K, Arai Y, and TACTICS study group: Randomized, open label, multicenter, phase II trial of transcatheter arterial chemoembolization (TACE) therapy in combination with sorafenib as compared with TACE alone in patients with hepatocellular carcinoma: TACTICS trial. 12th Annual Conference International Liver Cancer Association (ILCA), London, United Kingdom, September 14-16, 2018.
30. Kudo M, Izumi N, Kaneko S, Kobayashi M, Azuma MK, Copher R, Meier G, Ishii M, Ikeda S: A cost-effectiveness analysis of Lenvatinib compared with sorafenib in unresectable hepatocellular carcinoma allowing for AFP adjustment in overall survival in Japan from the REFLECT phase 3 clinical trial. 12th Annual Conference International Liver Cancer Association (ILCA), London, United Kingdom, September 14-16, 2018.
31. Bruix J, Merle P, Granito A, Huang YH, Bodoky G, Yokosuka O, Rosmorduc O, Breder V, Gerolami R, Masi G, Ross PJ, Qin S, Song T, Bronowicki JP, Ollivier-Houmand I, Kudo M, LeBerre MA, Baumhauer A, Meinhardt G, Han G and on behalf of the RESORCE Investigators: Overall survival (OS) update: 2-year follow-up from the phase 3 RESORCE trial of regorafenib for patients with hepatocellular carcinoma (HCC) progressing on sorafenib. 12th Annual Conference International Liver Cancer Association (ILCA), London, United Kingdom, September 14-16, 2018.
32. Lencioni R, Kudo M, Finn RS, Qin S, Han KH, Ikeda K, Cheng AL, Piscaglia F, Han G, Ikeda M, Simon K, Komov D, OuYang X, Evans TR, Sung M, Binder T, Damon A, Kraljevic S, Ren M, Ryoo BY: Independent imaging review (IIR) results in a phase 3 trial of lenvatinib (LEN) vs sorafenib (SOR) in first-line treatment of patients (PTS) with unresectable hepatocellular carcinoma (UHCC). 12th Annual Conference International Liver Cancer Association (ILCA), London, United Kingdom, September 14-16, 2018.
33. Izumoto H, Hiraoka A, Kumada T, Hirooka M, Tsuji K, Itobayashi E, Kariyama K, Ishikawa T, Tajiri K, Ochi H, Tada T, Toyoda H, Nouse K, Joko K, Hiasa Y, Ninomiya T, Michitaka K, Kudo M: Newly proposed tools for assessment of hepatic function for hepatocellular carcinoma staging and treatment planning-usefulness of modified ALBI grade. 12th Annual Conference International Liver Cancer Association (ILCA), London, United Kingdom, September 14-16, 2018.
34. Cheng AL, Raoul JL, Lee HC, Bayh I, Nakajima K, Peck-Radosavljevic M, Kudo M: Acute and chronic deterioration in liver function after transarterial chemoembolization (TACE) in patients with hepatocellular carcinoma (HCC): the final analysis of OPTIMIS. 12th Annual Conference International Liver Cancer Association (ILCA), London, United Kingdom, September 14-16, 2018.
35. Evans TR, Kudo M, Finn RS, Han KH, Cheng AL, Kraljevic S, Ren M, Dutcus CE, Piscaglia F, Sung MW: Urine protein: creatinine ratio (UPCR) vs 24-h urine protein for the management of proteinuria: results from a phase 3 study of lenvatinib (LEN) vs sorafenib

(SOR) in hepatocellular carcinoma (HCC).
12th Annual Conference International Liver
Cancer Association (ILCA), London, United
Kingdom, September 14-16, 2018.

36. Finn RS, Kudo M, Cheng AL, Wyrwicz L,
Ngan R, Blanc JF, Baron A, Vogel A, Ikeda M,
Piscaglia F, Han KH, Qin S, Minoshima Y,
Kanekiyo M, Ren M, Dairiki R, Tamai T,
Dutcus C, Funahashi Y, Evans TRJ: Final
analysis of serum biomarkers in patients (pts)
from the phase 3 study of lenvatinib (LEN) vs
sorafenib (SOR) in unresectable hepatocellular
carcinoma (uHCC) [REFLECT]. European
Society for Medical Oncology (ESMO 2018),
Munich, Germany, October 19-23, 2018.

H. 知的財産権の出願・登録状況(予定を含む)

- 1) 特許取得：なし
- 2) 実用新案登録：なし
- 3) その他：なし