

別紙 4

研究成果の刊行に関する一覧表

書籍

| 著者氏名 | 論文タイトル名 | 書籍全体の編集者名 | 書 籍 名 | 出版社名 | 出版地 | 出版年 | ページ |
|------|---------|-----------|-------|------|-----|-----|-----|
| なし | | | | | | | |
| | | | | | | | |

雑誌

| 発表者氏名 | 論文タイトル名 | 発表誌名 | 巻号 | ページ | 出版年 |
|---|---|----------------------------------|----------|--|------|
| Ando W, Takao M, Tani T, Uemura K, Hamada H, Sugano N. | Geographical distribution of the associated factors of osteonecrosis of the femoral head, using the designated intractable disease database in Japan. | Mod Rheumatol. | In press | doi:10.1093/mr/roab065. | 2021 |
| Sato R, Ando W, Fukushima W, Sakai T, Hamada H, Takao M, Ito K, Sugano N. | Epidemiological study of osteonecrosis of the femoral head using the national registry of designated intractable diseases in Japan. | Mod Rheumatol. | In press | doi:10.1093/mr/roab047. | 2021 |
| Uemura K, Otake Y, Takao M, Soufi M, Kawasaki A, Sugano N, Sato Y. | Automated segmentation of an intensity calibration phantom in clinical CT images using a convolutional neural network. | Int J Comput Assist Radiol Surg. | 16(11) | 1855-1864. doi:10.1007/s11548-021-02345-w.. | 2021 |
| Iwasa M, Ando W, Uemura K, Hamada H, Takao M, Sugano N. | Pelvic incidence is not associated with the development of hip osteoarthritis. | Bone Joint J. | 103(11) | 1656-1661. doi:10.1302/0301-620X.103B11.BJJ-2021-0472.R1. | 2021 |

| | | | | | |
|--|---|----------------------|----------|--|------|
| Suetsugu H, Kim K, Yamamoto T, Bang SY, Sakamoto Y, Shin JM, Sugano N, Kim JS, Mukai M, Lee YK, Ohmura K, Park DJ, Takahashi D, Ahn GY, Karino K, Kwon YC, Miyamura T, Kim J, Nakamura J, Motomura G, Kuroda T, Niino H, Miyamoto T, Takeuchi T, Ikari K, Amano K, Tada Y, Yamaji K, Shimizu M, Atsumi T, Seki T, Tanaka Y, Kubo T, Hisada R, Yoshioka T, Yamazaki M, Kabata T, Kajino T, Ohta Y, Okawa T, Naito Y, Kaneuji A, Yasunaga Y, Ohzono K, Tomizuka K, Koido M, Matsuda K, Okada Y, Suzuki A, Kim BJ, Kochi Y, Lee HS, Ikegawa S, Bae SC, Terao C. | Novel susceptibility loci for steroid-associated osteonecrosis of the femoral head in systemic lupus erythematosus. | Hum Mol Genet | In press | doi:10.1093/hmg/ddab306. | 2021 |
| Sato R, Ando W, Hamada H, Takao M, Saito M, Sugano N. | Differences in knee joint degeneration between primary hip osteoarthritis and hip osteoarthritis secondary to hip developmental dysplasia: A propensity score-based analysis. | Mod Rheumatol | 31(6) | 1221-1227. doi:10.1080/14397595.2021.1883261. | 2021 |
| Thiengwittayaporn S, Budhiparoma N, Tanavalee C, Tantavivut S, Sorial RM, Li C, Kim KI; Asia-Pacific (AP) Region Venous Thromboembolism (VTE) Consensus Group. | Asia-Pacific venous thromboembolism consensus in knee and hip arthroplasty and hip fracture surgery: Part 3. Pharmacological venous thromboembolism prophylaxis. | Knee Surg Relat Res. | 33(1) | 24. doi:10.1186/s43019-021-00100-8. | 2021 |
| Ngarmukos S, Kim KI, Wongsak S, Chotanaphuti T, Inaba Y, | Chen CF, Liu D; Asia-Pacific (AP) Region Venous Thromboembolism (VTE) Consensus Group. Asia-Pacific venous thromboembolism consensus in knee and hip arthroplasty and hip fracture surgery: Part 1. Diagnosis and risk factors. | Knee Surg Relat Res. | 33(1) | doi:10.1186/s43019-021-00099 | 2021 |
| Takashima K, Nakahara I, Hamada H, Ando W, Takao M, Uemura K, Sugano N. | A carbon fiber-reinforced polyetheretherketone intramedullary nail improves fracture site visibility on postoperative radiographic images. | Injury. | 52(8) | 2225-2232. doi:10.1016/j.injury.2021.03.048. | 2021 |

| | | | | | |
|---|--|----------------------|--------|--|------|
| Ando W, Takao M, Hamada H, Uemura K, Sugano N. | Comparison of the accuracy of the cup position and orientation in total hip arthroplasty for osteoarthritis secondary to developmental dysplasia of the hip between the Mako robotic arm-assisted system and computed tomography-based navigation. | Int Orthop. | 45(7) | 1719-1725. doi:10.1007/s00264-021-05015-3. Epub 2021 | 2021 |
| Amarase C, Tanavalee A, Larbpaiboonpong V, Lee MC, Crawford RW, Matsubara M, Zhou Y. | Asia-Pacific (AP) Region Venous Thromboembolism (VTE) Consensus Group. Asia-Pacific venous thromboembolism consensus in knee and hip arthroplasty and hip fracture surgery: Part 2. Mechanical venous thromboembolism prophylaxis. | Knee Surg Relat Res. | 33(1) | 20 doi:10.1186/s43019-021-00101-7. | 2021 |
| Hines JT, Jo WL, Cui Q, Mont MA, Koo KH, Cheng EY, Goodman SB, Ha YC, Hernigou P, Jones LC, Kim SY, Sakai T, Sugano N, Yamamoto T, Lee MS, Zhao D, Drescher W, Kim TY, Lee YK, Yoon BH, Baek SH, Ando W, Kim HS, Park JW. | Osteonecrosis of the Femoral Head: an Updated Review of ARCO on Pathogenesis, Staging and Treatment. | J Korean Med Sci. | 36(24) | 177. doi:10.3346/jkms.2021.36.e177. | 2021 |
| Asada R, Abe H, Hamada H, Fujimoto Y, Choe H, Takahashi D, Ueda S, Kuroda Y, Miyagawa T, Yamada K, Tanaka T, Ito J, Morita S, Takagi M, Tetsunaga T, Kaneuji A, Inaba Y, Tanaka S, Matsuda S, Sugano N, Akiyama H. | Femoral head collapse rate among Japanese patients with pre-collapse osteonecrosis of the femoral head. | J Int Med Res. | 49(6) | 00605211023336. doi:10.1177/03000605211023336. | 2021 |
| Kuroda Y, Tanaka T, Miyagawa T, Hamada H, Abe H, Ito T, Ihara T, Asada R, Fujimoto Y, Takahashi D, Tetsunaga T, Kaneuji A, Takagi M, Inaba Y, Morita S, Sugano N, Tanaka S, Matsuda S, Akiyama H; | TRION trial collaborators: Recombinant human FGF-2 for the treatment of early-stage osteonecrosis of the femoral head: TRION, a single-arm, multicenter, Phase II trial. | Regen Med. | 16(6) | 535-548. doi:10.2217/rme-2021-0049.. | 2021 |

| | | | | | |
|---|--|-------------------|--------|---|------|
| Hashimoto Y, Ando W, Sakai T, Takao M, Hamada H, Sugano N. | The Effects of Rasp Over size on the Clinical and Radiographic Outcomes of Total Hip Arthroplasty With a Collared Satin-Finished Composite Beam Cemented Stem. | J Arthroplasty. | 36(6) | 2055-2061. doi:10.1016/j.arth.2021.01.044. | 2021 |
| Cheng EY, Cui Q, Goodman SB, Ando W, Baek SH, Bakker C, Drescher W, Hernigou P, Jones LC, Kim H, Kim SY, Kim TY, Ha YC, Koo KH, Lee MS, Mont MA, Reichert I, Sakai T, Salem HS, Sierra RJ, Stronach B, Sugano N, Yamamoto T, Yoon BH, Zhao D. | Diagnosis and Treatment of Femoral Head Osteonecrosis: A Protocol for Development of Evidence-Based Clinical Practice Guidelines. | Surg Technol Int. | 38 | 371-378. doi:10.52198/21.STI.38.OS1437. | 2021 |
| Aoki K, Kurashige M, Ichii M, Higaki K, Sugiyama T, Kaito T, Ando W, Sugano N, Sakai T, Shibayama H; HANDAI Clinical Blood Club, Takaori-Kondo A, Morii E, Kanakura Y, Nagasawa T | Identification of CXCL12-abundant reticular cells in human adult bone marrow. | Br J Haematol. | 193(3) | 659-668. doi:10.1111/bjh.17396 | 2021 |
| Tani T, Ando W, Fukushima W, Hamada H, Takao M, Ito K, Sakai T, Sugano N. | Geographic distribution of the incidence of osteonecrosis of the femoral head in Japan and its relation to smoking prevalence. | Mod Rheumatol. | 30 | 1-7. doi:10.1080/14397595. | 2021 |
| Cui Q, Jo WL, Koo KH, Cheng EY, Drescher W, Goodman SB, Ha YC, Hernigou P, Jones LC, Kim SY, Lee KS, Lee MS, Lee YJ, Mont MA, Sugano N, Taliaferro J, Yamamoto T, Zhao D. | ARCO Consensus on the Pathogenesis of Non-traumatic Osteonecrosis of the Femoral Head | J Korean Med Sci. | 36(10) | e65. doi:10.3346/jkms.2021.36.e65. | 2021 |
| Uemura K, Hamada H, Ando W, Takao M, Sugano N. | Minimum 10 years clinical results of an anatomical short stem with a proximal hydroxyapatite coating. | Mod Rheumatol. | 31(5) | 1066-1072. doi: 10.1080/14397595.2020.1868129. | 2021 |
| Tani T, Ando W, Hamada H, Takao M, Sugano N. | Hip subluxation and osteophyte morphology are related to coronal contracture of the hip. | J Orthop Res. | 39(8) | 1691-1699. doi:10.1002/jor.24891. | 2021 |

| | | | | | |
|--|---|-----------------|-------|--|------|
| Yin X, Kim K, Suetsugu H, Bang SY, Wen L, Koido M, Ha E, Liu L, Sakamoto Y, Jo S, Leng RX, Otomo N, Laurd, Kottyan LC, Weirauch MT, Parameswaran S, Eswar S, Salim H, Chen X, Yamamoto K, Harley JB, Ohmura K, Kim TH, Yang S, Yamamoto T, Kim BJ, Shen N, Ikegawa S, Lee HS, Zhang X, Terao C, Cui Y, Bae SC. Meta-analysis of 208370 East Asians identifies 11 susceptibility loci for systemic lupus erythematosus. | Japanese Research Committee on Idiopathic Osteonecrosis of the Femoral Head. Arthritis Rheumatism. 2020;62(11):2083-2093. | Ann Rheum Dis | 80(5) | 632-640. doi:10.1136/annrheumdis-2020-219209. | 2021 |
| Hamada H, Ando W, Takao M, Sugano N. | Gamma-glutamyltransferase: A useful marker of habitual drinking in cases of alcohol-associated osteonecrosis of the femoral head. Alcohol and Alcoholism. 2020;45(1):1-6. | Alcohol Alcohol | 56(2) | 175-180. doi:10.1093/alcalc/aga117. | 2021 |
| Ando W, Sakai T, Fukushima W, Kaneuji A, Ueshima K, Yamasaki T, Yamamoto T, Niishi T, Working group for ONFH guidelines, Sugano N. | Japanese Orthopaedic Association 2019 Guidelines for Osteonecrosis of the femoral head. | J Orthop Sci. | 26(1) | 46-68. doi:10.1016/j.jos.2020.06.013. | 2021 |
| Itsuji T, Tonomura H, Ishibashi H, Mikami Y, Nagae M, Takatori R, Tanida T, Matsuda KI, Tanaka M, Kubo T. | Hepatocyte growth factor regulates HIF-1 α -induced nucleus pulposus cell proliferation through MAPK-, PI3K/Akt-, and STAT3-mediated signaling. | J Orthop Res. | 39(6) | 1184-1191. doi:10.1002/jor.24679. | 2021 |
| Oba Y, Sonohata M, Kitajima M, Kawano S, Eto S, Mawatairi M | Conventional cementless total hip arthroplasty in patients with dwarfism with height less than 140cm and minimum 10-year follow-up: A clinical study. | J Orthop Sci. | 26(1) | 128-134. doi:10.1016/j.jos.2020.02.001. | 2021 |

| | | | | | |
|--|---|---------------------------|-------|---|------|
| Hashimoto A, Sonohata M, Hirata H, Kawano S, Eto S, Ueno M, Mawatari M | Periarticular analgesic injection containing a corticosteroid after total hip arthroplasty may prevent deep venous thrombosis: a retrospective comparative cohort study. | BMC Musculoskelet Disord. | 22(1) | 19. doi:10.1186/s12891-020-03879-x. | 2021 |
| Nakayama Y, Sonohata M, Kitajima M, Sugino H, Hashimoto A, Hirata H, Kii S, Nakashima T, Kawano S, Eto S, Ueno M, Motegi S, Mawatari M | Fragility fracture following total hip arthroplasty. | J Orthop Sci. | 27(2) | 4. doi:10.1016/j.jos.2020.12.016. | 2022 |
| Kobayashi T, Morimoto T, Yoshihara T, Sonohata M, Riviere C, Mawatari M | The relationship between pelvic incidence and anatomical acetabular anteversion in female Japanese patients with hip osteoarthritis: a retrospective iconographic study. | Surg Radiol Anat. | 43(7) | 1141-1147. doi:10.1007/s00276-021-02710-z. | 2021 |
| Hashimoto A, Miyamoto H, Kii S, Kobatake T, Shobuike T, Noda I, Sonohata M, Mawatari M: | Time-dependent efficacy of combination of silver-containing hydroxyapatite coating and vancomycin methicillin-resistant Staphylococcus aureus biofilm formation in vitro. | BMC Res Notes | 14(1) | 81. doi:10.1186/s13104-021-05499-7. | 2021 |
| Sonohata M, Nakashima T, Kitajima M, Kawano S, Eto S, Mawatari M: | Total hip arthroplasty using hydroxyapatite-coated cementless cup for rapidly destructive coxarthrosis: Minimum 10-year follow-up. | J Orthop Sci. | 26(2) | 225-229. doi:10.1016/j.jos.2020.03.009. | 2021 |
| Kobayashi T, Morimoto T, Sonohata M, Mawatari M: | Is dislocation following total hip arthroplasty caused while suffering from delirium? | Nagoya J Med Sci | 83(3) | 601-607. doi:10.18999/nagjms.83.3.601. | 2021 |
| Hashimoto A, Miyamoto H, Kii S, Kobatake T, Shobuike T, Noda I, Sonohata M, Mawatari M: | Time-dependent efficacy of combination of silver-containing hydroxyapatite coating and vancomycin methicillin-resistant Staphylococcus aureus biofilm formation in vitro. | BMC Res Notes | 14(1) | 81. doi:10.1186/s13104-021-05499-7. | 2021 |
| Hasegawa M, Morikawa M, Seaman M, Cheng VK, Sudo A. | Population-based prevalence of femoroacetabular impingement in Japan. | Mod Rheumatol | 31(4) | 899-903. doi:10.1080/14397595.2020.18166 | 2021 |

| | | | | | |
|---|--|--------------------------------|----------|--|------|
| Hasegawa M, Tone S, Naito Y, Wakabayashi H, Sudo A. | Minimum ten-year results in revision total hip arthroplasty using titanium fully porous long stem. | Int Orthop. | 45(7) | 1727-1733. doi:10.1007/s00264-021-05030-4. | 2021 |
| Cheng VK, Hasegawa M, Hattori T, Ito N, Linn E, Cheng K, Hughes-Austin J, Masuda K, Sudo A. | Prevalence of radiographic hip dysplasia in Japanese population-based study. | Mod Rheumatol. | In press | doi:10.1080/14397595.2021.1918884. | 2021 |
| Naito Y, Hasegawa M, Tone S, Wakabayashi H, Sudo A. | Thirteen-year results of cementless total hip arthroplasty with zirconia heads on highly cross-linked polyethylene. | Mod Rheumatol. | 31(5) | 1045-1049. doi:10.1080/14397595.2020.1847753. | 2021 |
| Hasegawa M, Tone S, Naito Y, Wakabayashi H, Sudo A. | Comparison of the accuracies of computed tomography-based navigation and image-free navigation for acetabular cup insertion in total hip arthroplasty in the lateral decubitus position. | Comput Assist Surg (Abingdon). | 26(1) | 69-76. doi:10.1080/24699322.2021.1949494. | 2021 |
| Naito Y, Hasegawa M, Tone S, Wakabayashi H, Sudo A. | The accuracy of acetabular cup placement in primary total hip arthroplasty using an image-free navigation system. | BMC Musculoskelet Disord. | 22(1) | 1016. doi:10.1186/s12891-021-04902-5. | 2021 |
| Tone S, Hasegawa M, Naito Y, Wakabayashi H, Sudo A. | Accuracy of image-free navigation in intraoperative leg length change from total hip arthroplasty using evaluations from 2D and 3D measurements. | BMC Musculoskelet Disord. | 22(1) | 1021. doi:10.1186/s12891-021-04906-1. | 2021 |
| Tanaka T, Moro T, Asai S, Hashikura K, Ishikura H, Tanaka S. | Comparison between cup implantations during total hip arthroplasty with or without a history of rotational acetabular osteotomy. | Arch Orthop Trauma Surg | In press | doi:10.1007/s00402-021-04253-9. | 2021 |
| Ishikura H, Nakamura M, Oka H, Yonezawa K, Hidaka R, Tanaka T, Kawano H, Tanaka S. | A new classification system for evaluating fatty infiltration of the gluteus minimus in hip osteoarthritis using plain computed tomography. | J Orthop Sci. | In press | doi:10.1016/j.jos.2021.04.005. | 2021 |
| Ishikura H, Nakamura M, Matsuda K, Tanaka T, Kawano H, Tanaka S. | Severe osteolysis and periprosthetic femoral fracture 45 years after acrylic hemiarthroplasty of the hip: a case report. | BMC Musculoskelet Disord. | 22(1) | 482. doi:10.1186/s12891-021-04380-9 | 2021 |

| | | | | | |
|--|--|---------------------------|----------|---|------|
| Tanaka T, Moro T, Ishikura H, Hashikura K, Kaneko T, Tanaka S. | Characteristics of three-dimensional acetabular morphology with excellent outcome after rotational acetabular osteotomy. | J Orthop Surg Res. | 16(1) | 192. doi:10.1186/s13018-021-12346-0. | 2021 |
| Mitsutake R, Tanino H, Ito H. | The effect of range of motion simulated with a patient-specific three-dimensional simulation analysis on dislocation after total hip arthroplasty. | Hip Int. 2021 Sep | In press | 11207000211044807. | 2021 |
| Tanino H, Nishida Y, Mitsutake R, Ito H. | Accuracy of a portable accelerometer-based navigation system for cup placement and intraoperative leg length measurement in total hip arthroplasty: a cross-sectional study. | BMC Musculoskelet Disord. | 22(1) | 299. doi:10.1186/s12891-021-04167-y. | 2021 |
| Okuzu Y, Goto K, Shimizu Y, Kawai T, Kuroda Y, Matsuda S. | Sacroiliac joint degeneration is common in patients with end-stage hip osteoarthritis secondary to unilateral developmental dysplasia of the hip: Factors associated with its severity and laterality. | J Orthop Sci. | (1) | 135-140. doi:10.1016/j.jos.2020.02.005. | 2021 |
| Murao M, Nankaku M, Hamada R, Kawano T, Goto K, Kuroda Y, Ikeguchi R, Matsuda S. | Hip Abductor Strength and Lower Limb Load on Nonoperating Predict Functional Mobility in Women Patients With Total Hip Arthroplasty. | Am J Phys Med Rehabil. | 100(1) | 72-76. doi:10.1097/PHM.0000000000001523. | 2021 |
| Mitsuzawa S, Kuroda Y, Okuzu Y, Kawai T, Goto K, Kaido T, Uemoto S, Matsuda S. | Corticosteroid-associated osteonecrosis of the femoral head after orthotopic liver transplantation and the outcomes of subsequent total hip arthroplasty. | J Orthop Sci. | In press | doi:10.1016/j.jos.2020.12.014. | 2021 |
| Okuzu Y, Goto K, Kuroda Y, Kawai T, Matsuda S. | Preoperative Factors Associated With Low Back Pain Improvement After Total Hip Arthroplasty in a Japanese Population. | J Arthroplasty. | In press | doi:10.1016/j.art.2021.08.025 | 2021 |
| Ishikawa T, Kawai T, Goda N, Goto K, Kuroda Y, Matsuda S. | Chronic Expanding Hematomas That Occurred 20 Years or More After Primary Total Hip Arthroplasty: A Report of 2 Cases. | JBJS Case Connect. | 11(1) | doi:10.2106/JBJS.CC.20.00619. | 2021 |

| | | | | | |
|--|--|----------------------------|-----------|---|------|
| Kuroda Y, Okuzu Y, Kawai T, Goto K, Matsuda S. | Difference in Therapeutic Strategies for Joint-Preserving Surgery for Non-Traumatic Osteonecrosis of the Femoral Head between the United States and Japan: A Review of the Literature. | Orthop Surg. | 13(3) | 742-748. doi:10.1111/os.12979. | 2021 |
| Kawai T, Shimizu T, Goto K, Kuroda Y, Okuzu Y, Fujibayashi S, Matsuda S. | Number of Levels of Spinal Fusion Associated with the Rate of Joint-Space Narrowing in the Hip. | J Bone Joint Surg Am | 103(11) | 953-960. doi:10.2106/JBJS.20.01578. | 2021 |
| Kuroda Y, Koumoto S, Okuzu Y, Kawai T, Goto K, Matsuda S. | Similarity in Sequential Bilateral Transient Osteoporosis of the Hip: A Report of 3 Cases. | JBJS Case Connect. | 11(2) | doi:10.2106/JBJS.CC.20.00722. | 2021 |
| Goto K, Aoyama T, Toguchida J, Kuroda Y, Kawai T, Okuzu Y, Matsuda S. | Ten-year results of mesenchymal stromal cell transplantation augmented with vascularised bone grafts for advanced osteonecrosis of the femoral head. | J Orthop. | 26 | 67-71 doi:10.1016/j.jor.2021.07.006. | 2021 |
| Kawano T, Nankaku M, Muraom M, Goto K, Kuroda Y, Kawai T, Ikeguchi R, Matsuda S. | Functional characteristics associated with hip abductor torque in severe hip osteoarthritis. | Musculoskeletal Sci Pract. | 55 | 102431. doi:10.1016/j.msksp.2021.102431. | 2021 |
| Takaoka Y, Goto K, Tamura J, Okuzu Y, Kawai T, Kuroda Y, Orita K, Matsuda S. | Radiolucent lines do not affect the longevity of highly cross-linked polyethylene cemented components in total hip arthroplasty. | Bone Joint J. | 103-B(10) | 1604-1610. doi:10.1302/0301-620X | 2021 |
| Kuroda Y, Nankaku M, Okuzu Y, Kawai T, Goto K, Matsuda S. | Percutaneous autologous iliac impaction bone graft for advanced femoral head osteonecrosis: a retrospective observational study of unsatisfactory short-term outcomes. | J Orthop Surg Res. | 16(1) | 141 doi:10.1186/s13018-021-02288-7. | 2021 |
| Tateuchi H, Akiyama H, Goto K, So K, Kuroda Y, Ichihashi N. | Strategies for increasing gait speed in patients with hip osteoarthritis: their clinical significance and effects on hip loading. | Arthritis Res Ther. | 23(1) | 129. doi:10.1186/s13075-021-02514-x. | 2021 |
| Tateda K, Nagoya S, Suzuki D, Kosukegawa I, Yamashita T. | Acetabular Morphology in Patients with Developmental Dysplasia of the Hip with High Dislocation. | Hip Pelvis | 33(1) | doi:10.5371/hp.2021.33.1.25. | 2021 |

| | | | | | |
|--|--|--------------------------------------|----------|--|------|
| Fukushima W, Hara M, Kitamura Y, Shibata M, Ugawa Y, Hirata K, Oka A, Miyamoto S, Kusunoki S, Kuwabara S, Hashimoto S, Sobue T. | A nationwide epidemiological survey of adolescent patients with diverse symptoms similar to those following human papillomavirus vaccination: background prevalence and incidence for considering vaccine safety in Japan. | J Epidemiol. | In press | doi:10.2188/jea.JE20210277. | 2021 |
| Murata K, Onoyama S, Yamamura K, Mizuno Y, Furuno K, Matsubara K, Hatae K, Masuda K, Nomura Y, Ohno T, Kinumaki A, Miura M, Sakai Y, Ohga S, Fukushima W, Kishimoto J, Nakamura Y, Hara T. | Kawasaki Disease and Vaccination: Prospective Case-Control and Case-Crossover Studies among Infants in Japan. | Vaccines (Basel). | 9(8) | 839. doi:10.3390/vaccines9080839. | 2021 |
| Kakiuchi M, Wada E, Harada T, Yamasaki R, Fukushima W. | Relief of Low Back Pain After Posterior Decompression for Lumbar Spinal Stenosis. | Spine (Phila Pa 1976). | 46(18) | E966-E975. doi:10.1097/BRS.00000000000004006. | 2021 |
| Yamada K, Aota Y, Saito T, Inaba Y: | Correlation between leg pain at rest and spinal nerve edema in symptomatic lumbar foraminal stenosis. | Orthop Traumatol Surg Res. | | 103119: doi:10.1016/j.otsr.2021.103119. | 2021 |
| Choe H, Kobayashi N, Oba M, Morita A, Abe K, Inaba Y: | Application of Ultrasonography During Sternocleidomastoid Muscle Release for Congenital Muscular Torticollis: A Case Report | J. JBJS Case Connect. | 11(4) | doi:10.2106/JBJS.CC.21.00309. | 2021 |
| Choe H, Kobayashi N, Kobayashi D, Watanabe S, Abe K, Tezuka T, Kawabata Y, Takeyama M, Inaba Y: | Postoperative excessive external femoral rotation in revision total hip arthroplasty is associated with muscle weakness in iliopsoas and gluteus medius and risk for hip dislocation. | J Orthop Surg Res. | 16(1) | 582. doi:10.1186/s13018-021-02744-4. | 2021 |
| Nejima S, Kumagai K, Fujimaki H, Yamada S, Sotozawa M, Matsubara J, Inaba Y | Increased contact area of flange and decreased wedge volume of osteotomy site by open wedge distal tibial tuberosity arc osteotomy compared to the conventional technique. | Knee Surg Sports Traumatol Arthrosc. | 29(10) | 3450-3457. doi:10.1007/s00167-020-06296-8. | 2021 |
| Otoshi A, Kumagai K, Yamada S, Nejima S, Fujisawa T, Miyatake K, Inaba Y: | Return to sports activity after opening wedge high tibial osteotomy in patients aged 70 years and older. | J Orthop Surg Res. | 16(1) | 576. doi:10.1186/s13018-021-02718-6. | 2021 |

| | | | | | |
|---|--|--|----------|---|------|
| Kobayashi N, Kamono E, Maeda K, Misumi T, Yukizawa Y, Inaba Y: | Effectiveness of diluted povidone-iodine lavage for preventing periprosthetic joint infection: an updated systematic review and meta-analysis. | J Orthop Surg Res. | 16(1) | 569. doi:10.1186/s13018-021-02703-z. | 2021 |
| Inoue T, Higashi T, Kobayashi N, Ide M, Harigane K, Mochida Y, Inaba Y | Associated with Aggravation of Cervical Spine Lesions in Patients with Rheumatoid Arthritis: A Retrospective Longitudinal Cohort Study. | A retrospective longitudinal cohort study. Spine | In press | doi:10.1097/BRS.0000000000000420. PMID: 34524270. | 2021 |
| Higashihira S, Inasaka R, Iwamura Y, Kohno M, Akiyama G, Inaba Y: | Patient-reported outcome measures after total knee arthroplasty using knee injury and osteoarthritis outcome score. | JRPMS. | 5(3) | 93-100. | 2021 |
| Fujimaki H, Nakazawa A, Hirano M, Takeuchi T, Kadowaki A, Kusayama Y, Ide M, Kanai K, Kim Y, Matsubara J, Kumagai K, Inaba Y. | Status of fracture risk assessment and osteoporosis treatment in Japanese patients with rheumatoid arthritis. | Mod Rheumatol. | 31(5) | 987-991. doi:10.1080/14397595.2020.1847752. | 2021 |
| Thiengwittayaporn S, Budhiparاما N, Tanavalee C, Tantavirut S, Sorial RM, Li C, Kim KI; | Asia-Pacific (AP) Region Venous Thromboembolism (VTE) Consensus Group (Choe H, Inaba Y, et al): Asia-Pacific venous thromboembolism consensus in knee and hip arthroplasty and hip fracture surgery: Part 3. Pharmacological venous thromboembolism prophylaxis. | Knee Surg Relat Res. | 33(1) | 24. doi:10.1186/s43019-021-00100-8. | 2021 |
| Kato S, Mihara H, Niimura T, Watanabe K, Kawai T, Choe H, Inaba Y: | Impact of focal apex angle on postoperative decompression status of the spinal cord and neurological recovery after cervical laminoplasty. | J Neurosurg Spine. | 35(4) | 410-418. doi:10.3171/2020.12.SPINE201831. | 2021 |
| Kumagai K, Fujimaki H, Yamada S, Nejima S, Matsubara J, Inaba Y: | Changes of synovial fluid biomarker levels after opening wedge high tibial osteotomy in patients with knee osteoarthritis. | Osteoarthritis Cartilage. | 29(7) | 1020-1028. doi:10.1016/j.joca.2021.03.013. | 2021 |
| Nejima S. Kumagai K. Kobayashi H. Fujimaki H. Yamada S. Sotozawa M. Hisatomi K. Inaba Y: | Medialization of the mechanical axis of the tibia is related to lateralization of the tibial tuberosity in knee osteoarthritis. | Knee. | 30 | 134-140. doi:10.1016/j.knee.2021.03.023. | 2021 |

| | | | | | |
|--|---|-----------------------------|-------|--|------|
| Amarase C, Tanavalee A, Larbpaiboonpong V, Lee MC, Crawford RW, Matsubara M, Zhou Y; | Asia-Pacific (AP) Region Venous Thromboembolism (VTE) Consensus Group (Choe H, Inaba Y, et al): Asia-Pacific venous thromboembolism consensus in knee and hip arthroplasty and hip fracture surgery: Part 2. Mechanical venous thromboembolism prophylaxis. | Knee Surg Rel at Res. | 33(1) | 20. doi:10.1186/s43019-021-00101-7. | 2021 |
| Ngarmukos S, Kim KI, Wongsak S, Chotanaphuti T, Inaba Y, Chen CF, Liu D; | Asia-Pacific Region Venous Thromboembolism (VTE) Consensus Group (Choe H, Inaba Y, et al): Asia-Pacific venous thromboembolism consensus in knee and hip arthroplasty and hip fracture surgery: Part 1. Diagnosis and risk factors. | Knee Surg Rel at Res. | 33(1) | 18. doi:10.1186/s43019-021-00099-y. | 2021 |
| Watanabe S, Choe H, Kobayashi N, Ike H, Kobayashi D, Inaba Y; | Prediction of pelvic mobility using whole-spinal and pelvic alignment in standing and sitting position in total hip arthroplasty patients. | J Orthop Surg (Hong Kong). | 29(2) | 23094990211019099. doi:10.1177/23094990211019099. | 2021 |
| Akamatsu T, Kumagai K, Yamada S, Nejima S, Sotozawa M, Ogino T, Inaba Y; | No differences in clinical outcomes and cartilage repair between opening wedge and closed wedge high tibial osteotomies at short-term follow-up: A retrospective case series analysis. | J Orthop Surg (Hong Kong). | 29(2) | 23094990211020366. doi:10.1177/23094990211020366. | 2021 |
| Watanabe S, Kobayashi N, Tomoyama A, Choe H, Yamazaki E, Inaba Y; | Clinical characteristics and risk factors for culture-negative periprosthetic joint. | J Orthop Surg Res. | 16(1) | 292. doi:10.1186/s13018-021-02450-1. | 2021 |
| Uchino Y, Higashi T, Kobayashi N, Inoue T, Mochida Y, Inaba Y; | Risk factors associated with cervical spine lesions in patients with rheumatoid arthritis: an observational study. | BMC Musculoskeletal Disord. | 22(1) | 408. doi:10.1186/s12891-021-04285-7. | 2021 |
| Takagawa S, Kobayashi N, Yukizawa Y, Oishi T, Tsuji M, Misumi T, Inaba Y; | Identifying factors predicting prolonged rehabilitation after simultaneous bilateral total knee arthroplasty: a retrospective observational study. | BMC Musculoskeletal Disord. | 22(1) | 368. doi:10.1186/s12891-021-04211-x. | 2021 |

| | | | | | |
|--|---|--------------------------------------|--------|--|------|
| Kobayashi N, Higashihira S, Kitayama H, Kamono E, Yukizawa Y, Oishi T, Takagawa S, Honda H, Choe H, Inaba Y. | Effect of decreasing the anterior pelvic tilt on range of motion in femoroacetabular impingement: A computer-simulation study. Orthop | J Sports Med. | 9(4) | 2325967121999464. doi:10.1177/2325967121999464. | 2021 |
| Morita A, Kobayashi N, Choe H, Tezuka T, Higashihira S, Inaba Y: | Preoperative factors predicting the severity of BMD loss around the implant after Total Hip Arthroplasty. | BMC Musculoskelet Disord. | 22(1) | 290. doi:10.1186/s12891-021-04161-4. | 2021 |
| Nejima S, Kumagai K, Kobayashi H, Yamada S, Akamatsu T, Ogino T, Sotozawa M, Inaba Y: | Coronal shaft bowing of the femur affects varus inclination of the surgical transepicondylar axis in varus knee osteoarthritis. | Knee Surg Sports Traumatol Arthrosc. | 29(3) | 814-819. doi:10.1007/s00167-020-06025-1. | 2021 |
| Sasaki T, Akamatsu Y, Kobayashi H, Mitsuhashi S, Nejima S, Kumagai K, Saito T, Inaba Y | Rotational Changes in the Distal Tibial Fragment Relative to the Proximal Tibial Fragment at the Osteotomy Site after Open-Wedge High-Tibial Osteotomy | Biomed Res Int. | 2021 | 6357109. doi:10.1155/2021/6357109. | 2021 |
| Nejima S, Kumagai K, Kobayashi H, Yamada S, Akamatsu T, Ogino T, Sotozawa M, Inaba Y | Varus inclination of the tibia is related to patellofemoral osteoarthritis in Japanese female patients with moderate knee osteoarthritis. | Knee Surg Sports Traumatol Arthrosc. | 29(2) | 652-658. doi:10.1007/s00167-020-06000-w. | 2021 |
| Yang F, Choe H, Kobayashi N, Tezuka T, Oba M, Miyamae Y, Morita A, Abe K, Inaba Y. | An automated real-time PCR assay for synovial fluid improves the preoperative etiological diagnosis of periprosthetic joint infection and septic arthritis. | J Orthop Res. | 39(2) | 348-355. doi:10.1002/jor.24959. | 2021 |
| Kumagai K, Fujimaki H, Yamada S, Nejima S, Matsubara J, Inaba Y. | Difference in the early postoperative change of the joint line convergence angle between opening wedge and closed wedge high tibial osteotomies. | J Orthop Surg Res. | 16(1) | 66. doi:10.1186/s13018-021-02214-x. | 2021 |
| Kumagai K, Yamada S, Akamatsu T, Nejima S, Ogino T, Sotozawa M, Inaba Y. | Intraoperatively accurate limb alignment after opening wedge high tibial osteotomy can be lost by large knee joint line convergence angle during surgery. | Arch Orthop Trauma Surg. | 141(1) | 23-28. doi:10.1007/s00402-020-03419-1. | 2021 |

| | | | | | |
|---|---|------------------------|----------------|---|------|
| Kamada S, Kise N, Kinoshita K, Shiota E, Yamamoto T. | Femoral Neck Fracture in a Hemodialysis Patient after Liver Transplantation: A Case Report. | Prog Rehabil Med. | 6 | 20210003. doi:10.2490/prm.20210003. | 2021 |
| Matsunaga T, Kamachi Y, Kinoshita K, Sakamoto T, Yamamoto T. | Magnetic Resonance Imaging Assessment of Abductor Muscles Shortly After Curved Periacetabular Osteotomy. | J Arthroplasty. | 36(2) | 429-433. doi:10.1016/j.arthro.2020.08.041. | 2021 |
| Doi N, Kinoshita K, Sakamoto T, Minokawa A, Setoguchi D, Yamamoto T. | Incidence and clinical outcome of lateral femoral cutaneous nerve injury after periacetabular osteotomy. | Bone Joint J. | 103-B(4) | 659-664. doi:10.1302/0301-620X.103B4.BJJ-2020-0990.R2. | 2021 |
| Kosaka H, Maeyama A, Nishio J, Nabeshima K, Yamamoto T. | Histopathologic evaluation of bone marrow lesions in early stage subchondral insufficiency fracture of the medial femoral condyle. | Int J Clin Exp Pathol. | 14(7) | 819-826. | 2021 |
| Suzuki M, Kinoshita K, Sakamoto T, Seo H, Kinoshita S, Yamamoto T. | Leg Length Change After Curved Periacetabular Osteotomy and Its Impact on the Clinical Outcomes. | J Arthroplasty. | 36(9) | 3089-3096. doi:10.1016/j.arthro.2021.04.019. | 2021 |
| Kinoshita S, Ishimatsu T, Suzuki M, Seo H, Sakamoto T, Kinoshita K, Yamamoto T. | Femoral nerve status during the anterolateral approach for total hip arthroplasty: Motor-evoked potential analysis and an influencing factor. | J Orthop Sci. | S0949-2658(21) | 00334-1. doi:10.1016/j.jos.2021.09.007. | 2021 |
| Kamachi Y, Kinoshita K, Sakamoto T, Matsunaga T, Yamamoto T. | Bone union status of all osteotomy sites one year after curved periacetabular osteotomy based on computed tomography. Orthop Traumatol | Surg Res. | 107(8) | 102955. doi:10.1016/j.otsr.2021.102955. | 2021 |
| Baba S, Motomura G, Ikemura S, Yamaguchi R, Hamai S, Fujii M, Kawano K, Nakashima | Risk factors for radiological changes after bipolar hemiarthroplasty for osteonecrosis of the femoral head. | Modern Rheumatology | 31(3) | 725-732. doi:10.1080/14397595.2020.1775959. | 2021 |
| Harada S, Hamai S, Shiimoto K, Hara D, Fujii M, Ikemura S, Motomura G, Nakashima Y. | Patient-reported outcomes after primary or revision total hip arthroplasty: A propensity score-matched Asian cohort study. | PLoS One. | 16(5) | e0252112. doi:10.1371/journal.pone.0252112. | 2021 |
| Fujii M, Nakashima Y, Kitamura K, Motomura G, Hamai S, Ikemura S, Noguchi Y. | Preoperative Rather Than Postoperative Intra-Articular Cartilage Degeneration Affects Long-Term Survivorship of Periacetabular Osteotomy. | Arthroscopy | 37(7) | 2112-2122. doi:10.1016/j.arthro.2021.01.060. | 2021 |

| | | | | | |
|--|--|--|----------|--|------|
| Kitamura K, Fujii M, Iwamoto M, Ikemura S, Hamai S, Motomura G, Nakashima Y. | Is Anterior Rotation of the Acetabulum Necessary to Normalize Joint Contact Pressure in Periacetabular Osteotomy? A Finite-element Analysis Study. | Clin Orthop Relat Res. | In Press | doi:10.1097/CORR.0000000000001893. | 2021 |
| Komiyama K, Hamai S, Motomura G, Ikemura S, Fujii M, Kawahara S, Nakashima Y. | Total hip arthroplasty after periacetabular osteotomy versus primary total hip arthroplasty: a propensity-matched cohort study. | Arch Orthop Trauma Surg | 141(8) | 1411-1417. doi:10.1007/s00402-021-03817-z. | 2021 |
| Kitamura K, Fujii M, Ikemura S, Hamai S, Motomura G, Nakashima Y. | Does Patient-specific Functional Pelvic Tilt Affect Joint Contact Pressure in Hip Dysplasia? A Finite-element Analysis Study. | Clinical Orthopaedics and Related Research | 479(8): | 1712-1724. doi:10.1097/CORR.0000000000001737. | 2021 |
| Fujii M, Kitamura K, Ikemura S, Hamai S, Motomura G, Nakashima Y. | Pneumatic femoral broaching decreases post-operative subsidence of a cementless taper-wedge stem. | International Orthopaedics | In press | doi:10.1007/s00264-021-05196-x. | 2021 |
| Kuwakado S, Kawaguchi K, Sakagawa A, Takahashi M, Odaguchi T, Shimamoto N, Motomura G, Hamai S, Nakashima Y. | Factors Affecting the Length of Convalescent Hospital Stay Following Total Hip and Knee Arthroplasty. | Progress in Rehabilitation Medicine | 6 | 20210033. doi:10.2490/prm.20210033. | 2021 |
| Harada S, Hamai S, Shiimoto K, Hara D, Motomura G, Ikemura S, Fujii M, Nakashima Y. | Wear analysis of the first-generation cross-linked polyethylene at minimum 10 years follow-up after THA: no significant effect of sports participation. | Journal of Artificial Organs | In press | doi:10.1007/s10047-021-01297-x. | 2021 |
| Shiimoto K, Hamai S, Ikebe S, Higaki H, Hara D, Gondo H, Komiyama K, Yoshimoto K, Harada S, Nakashima Y. | Computer simulation based on in vivo kinematics of a replaced hip during chair-rising for elucidating target cup and stem positioning with a safety range of hip rotation. | Clinical biomechanics (Bristol, Avon) | In press | doi:10.1016/j.clinbiomech.2021.105537. | 2021 |
| Ogawa T, Yoshii T, Okawa A, Fushimi K, Jinno T. | Association between hospital surgical volume and complications after total hip arthroplasty in femoral neck fracture: A propensity score-matched cohort study. | Injury. | 52(10) | 3002-3010. doi:10.1016/j.injury.2021.02.092. | 2021 |

| | | | | | |
|---|--|------------------------------|----------------|---|------|
| Ogawa T, Yoshii T, Okawa A, Fushimi K, Jinno T. | Association Between Cemented vs Cementless Hemiarthroplasty and Short-Term Change of In-Hospital Mortality in Elderly Patients with Femoral Neck Fracture: A Propensity-Score Matching Analysis in a Multicenter Database. | Clin Interv Agi | 16 | 1151-1159. doi:10.2147/CIA.S315090. | 2021 |
| Ogawa T, Schermann H, Kobayashi H, Fushimi K, Okawa A, Jinno T. | Age and clinical outcomes after hip fracture surgery: do octogenarian, nonagenarian and centenarian classifications matter? | Age Ageing. | 50(6) | 1952-1960. doi:10.1093/ageing/afab137. | 2021 |
| Takada R, Jinno T, Miyatake K, Watanabe N, Koga H, Yoshii T, Okawa A. | Difference in tapered wedge stem alignment between supine and lateral position in cementless total hip arthroplasty via modified Watson-Jones anterolateral approach. | Eur J Orthop Surg Traumatol. | In press | doi:10.1007/s00590-021-03001-4. | 2021 |
| Taniguchi N, Jinno T, Ohba T, Endo H, Wako M, Fujita K, Koyama K, Ichikawa J, Ando T, Ochiai S, Haro H. | Differences of two-year longitudinal changes of locomotive syndrome among patients treated with thoracolumbar interbody fusion, total hip arthroplasty, and total knee arthroplasty for degenerative diseases. | Modern Rheumatology | In press | doi:10.1093/mr/rwab033. | 2021 |
| Ogawa T, Tachibana T, Yamamoto N, Udagawa K, Kobayashi H, Fushimi K, Yoshii T, Okawa A, Jinno T. | Patient body mass index modifies the association between waiting time for hip fracture surgery and in-hospital mortality: A multicenter retrospective cohort study. | J Orthop Sci. | S0949-2658(21) | 00234-7. doi:10.1016/j.jos.2021.07.015 | 2021 |
| Koyano G, Jinno T, Koga D, Hoshino C, Okawa A. | Intra-articular Injections of Cross-linked Hyaluronic Acid in Japanese Patients with Symptomatic Osteoarthritis of the Hip | Prog Rehabil Med. | 6 | 20210038. doi:10.2490/prm.20210038. PMID: 34632157 | 2021 |
| Ikeda T, Sato R, Ninomiya K, Suzuki K, Hirakawa K, Jinno T. | Nutritional Factors Related to Muscle Weakness 1 Year after Total Hip Arthroplasty: A Case-Control Study. | Ann Nutr Metab | 1 | 7. doi:10.1159/000520551. | 2021 |

| | | | | | |
|---|--|-------------------------------------|---------|---|------|
| Hirata M, Oe K, Kaneuji A, Uozu R, Shintani K, Saito T. | Relationship between the Surface Roughness of Material and Bone Cement: An Increased "Polished" Stem May Result in the Excessive Taper-Slip. | Materials (Base | 14(13) | 3702. doi: 10.3390/ma 14133702. | 2021 |
| Kaneuji A, Takahashi E, Fukui M, Ichiseki T, Fukui K, Kawahara N. | Long-Term Outcomes for Cementless Anatomic Femoral Components, Compared by Area of Porous Coating, in Patients Younger Than 50 Years Treated for Hip Dysplasia. | J Arthroplasty. | 36(8) | 2864-2870. doi:10.1016/j.art h.2021.03.024. | 2021 |
| Kaneuji A, Hara T, Takahashi E, Fukui K, Ichiseki T, Kawahara N. | A Novel Minimally Invasive Spherical Periacetabular Osteotomy: Pelvic Ring Preservation and Patient-Specific Osteotomy by Preoperative 3-Dimensional Templating. | J Bone Joint Surg Am. | 103(18) | 1724-1733. doi:10.2106/BJJ S.20.00940. | 2021 |
| Kawaguchi M, Segawa A, Shintani K, Nakamura Y, Ishigaki Y, Yonezawa K, Sasamoto T, Kaneuji A, Kawahara N. | Bone formation at Ti-6Al-7Nb scaffolds consisting of 3D honeycomb frame and diamond-like carbon coating implanted into the femur of beagles. | J Biomed Mater Res B Appl Biomater. | 109(9) | 1283-1291. doi:10.1002/jbm. b.34789. | 2021 |
| Kitajima H, Kaneuji A, Soma D, Fukui M, Kawahara N | Bone ingrowth observed in a cup removed during revision surgery for early dislocation after primary THA: A case report. | Int J Surg Cas Rep | 83 | 105941. doi:10.1016/j.ijsc r.2021.105941. | 2021 |
| Hirata H, Ueda S, Ichiseki T, Shimasaki M, Ueda Y, Kaneuji A, Kawahara N. | Taurine inhibits glucocorticoid-induced bone mitochondrial injury, preventing osteonecrosis in rabbits and cultured osteocytes. | Int J Mol Sci. | 21(18) | E6892, doi:10.3390/ijms 21186892. | 2021 |
| Seki K, Ogasa H, Tokushige A, Imagama T, Seki T, Nakashima D, Sakai T. | Ultrasonography-detected synovitis of hand is associated with the presence of synovitis in the forefoot of patients with rheumatoid arthritis. | J Orthop Sci | 26(5) | 850-3. doi:10.1016/j.jos. 2020.09.002. | 2021 |
| Imagama T, Seki K, Seki T, Tokushige A, Matsuki Y, Yamazaki K, Nakashima D, Okazaki T, Hirata K, Yamamoto M, Tanaka H, Sakai T. | Synovial fluid presepsin as a novel biomarker for rapid differential diagnosis of native joint septic arthritis from crystal arthritis. | Int J Infect Dis | 102 | 472-477. doi:10.1016/j.ijid. 2020.10.030. | 2021 |

| | | | | | |
|--|---|------------------------|----------|--|------|
| Imagama T, Nakashima D, Seki K, Seki T, Matsuki Y, Yamazaki K, Sakai T. | Comparison of bacterial culture results of preoperative synovial fluid and intraoperative specimens in patients with joint infection. | J Infect Chemo | 27(4) | 562-567 doi:10.1016/j.jiac.2020.11.006. | 2021 |
| Yamazaki K, Imagama T, Matsuki Y, Seki T, Seki K, Sakai T. | Evaluation of femoral anteversion, hip rotation, and lateral patellar tilt after total hip arthroplasty using a changeable neck system. | J Artif Organs | 24 | 492-497. doi:10.1007/s10047-021-01269-1. | 2021 |
| Imagama T, Seki K, Seki T, Matsuki Y, Yamazaki K, Sakai T. | Low frequency of local findings in periprosthetic hip infection caused by low-virulent bacteria compared to periprosthetic knee infection. | Sci Rep | 11(1) | 11714. doi:10.1038/s41598-021-91139-w. | 2021 |
| Yukata K, Uchiyama Y, Nishida N, Ohgi J, Sakai T, Fujii H. | Greater trochanteric fracture with lesser trochanteric extension. A report of 2 cases and CT-based finite element analysis. | JBJS Case Connect | 11(3) | doi:10.2106/BJB.S.CC.20.00356. | 2021 |
| Nishida N, Izumiyama T, Asahi R, Jiang F, Ohgi J, Yamagata H, Imajo Y, Suzuki H, Funaba M, Sugimoto S, Fukushima F, Chen X, Sakai T. | Analysis of Individual Differences in Pelvic and spine alignment in seated posture and Impact on the Seatbelt Kinematics using human body model. | Plos One, | 16(7) | 0254120. doi:10.1371/journal.pone.0254120. | 2021 |
| Suzuki H, Imajo Y, Funaba M, Nishida N, Sakamoto T, Sakai T. | Current Concepts of Neural Stem/Progenitor Cell Therapy for Chronic Spinal Cord Injury. | Front Cell Neurosci. | In press | doi:10.3389/fncel.2021.794692. | 2021 |
| Mihara A, Yukata K, Seki T, Iwanaga R, Nishida N, Fujii K, Nagao Y, Sakai T. | Effects of sclerostin antibody on bone healing. | World J Orthop. | 12(9) | 651-659. doi:10.5312/wjo.v12.i9.651. | 2021 |
| Sakai T. | CORR Insights®: Adverse Local Tissue Reactions are Common in Asymptomatic Individuals After Hip Resurfacing Arthroplasty: Interim Report from a Prospective Longitudinal Study. | Clin Orthop Relat Res. | 479(12) | 2651-2652. doi:10.1097/CORR.0000000000001926. | 2021 |

| | | | | | |
|--|--|----------------------------|----------------|--|------|
| Matsuki Y, Imagama T, Tokuhige A, Yamazaki K, Sakai T. | Accuracy of cup placement using computed tomography-based navigation system in total hip arthroplasty through the direct anterior approach. | J Orthop Sci. | S0949-2658(21) | 00364-X. doi:10.1016/j.jos.2021.10.018. | 2021 |
| Tanaka K, Shimizu Y, Kamada H, Aikawa S, Mishima H, Kanamori A, Nishino T, Sakane M, Ochiai N, Yamazaki M. | Feasibility and Safety of a Novel Leg Exercise Apparatus for Venous Thromboembolism Prophylaxis after Total Joint Arthroplasty of the Lower Extremities-A Pilot Study. | Tomography. | 7(4) | 734-746. doi:10.3390/tomography7040061. | 2021 |
| Nishino T, Mishima H, Kawamura H, Yoshizawa T, Miyakawa S, Yamazaki M. | Ten-year results of 55 dysplasia hips of hip offset and leg length reconstruction in total hip arthroplasty with cementless tapered stems having a high offset option designed for dysplastic femur. | J Orthop Surg (Hong Kong). | 28(1) | 2309499020909499. doi:10.1177/2309499020909499. | 2021 |
| Totsuka S, Nishino T, Watanabe R, Yamazaki M, Mishima H. | New Evaluation Method for Bone Formation around a Fully Hydroxyapatite-Coated Stem Using Digital Tomosynthesis: A Retrospective Cross-Sectional Study. | Diagnostics (Basel). | 11(11) | 2094. doi:10.3390/diagnostics11112094. | 2021 |
| Takehashi H, Nishino T, Mishima H, Wada H, Yamazaki M, Hyodo K. | Stress distribution of cementless stems with unique flanges in a rectangular cross-section: thermoelastic stress imaging study. | J Rural Med. | 16(2) | 83-90. doi:10.2185/jrm.2020-058. | 2021 |
| Kawamura H, Watanabe Y, Nishino T, Mishima H. | Effects of lower limb and pelvic pin positions on leg length and offset measurement errors in experimental total hip arthroplasty. | J Orthop Surg Res. | 16(1) | 193. doi:10.1186/1186-13018-021-02347-z. | 2021 |
| Tomaru Y, Yoshioka T, Nanakamura J, Sugaya H, Hagiwara S, Nawata K, Ohtori S, Yamazaki M, Mishima H. | Concentrated autologous bone marrow aspirate transplantation versus conservative treatment for corticosteroid-associated osteonecrosis of the femoral head in systemic lupus erythematosus. | J Rural Med. | 16(1) | 1-7 doi:10.2185/jrm.2020-033. | 2021 |

| | | | | | |
|--|---|---------------------------|----------|--|------|
| Yamamuro Y, Kabata T, Kajino Y, Inoue D, Hasegawa K, Tsuchiya H. | Does intraoperative periprosthetic occult fracture of the acetabulum affect clinical outcomes after primary total hip arthroplasty? | Arch Orthop Trauma Surg. | In press | doi:10.1007/s00402-021-04226-y. | 2021 |
| Saiki Y, Ojima T, Kabata T, Hayashi S, Tsuchiya H. | Accuracy of different navigation systems for femoral and tibial implantation in total knee arthroplasty: a randomised comparative study. | Arch Orthop Trauma Surg. | 141(12) | 2267-2276. doi:10.1007/s00402-021-04205-3. | 2021 |
| Hasegawa K, Kabata T, Kajino Y, Inoue D, Sakamoto J, Tsuchiya H. | The influence of pelvic tilt on stress distribution in the acetabulum: finite element analysis. | BMC Musculoskelet Disord. | 22(1) | 13750. doi:10.1038/s41598-021-93348-9. | 2021 |
| Yoshitani J, Kabata T, Kajino Y, Inoue D, Ohmori T, Ueoka K, Yamamuro Y, Taninaka A, Tsuchiya H. | Correlation between lag screw route and the ideal insertion point of the intramedullary nail. | Sci Rep. | 11(1) | 13750. doi:10.1038/s41598-021-93348-9. | 2021 |
| Yamamuro Y, Kabata T, Goshima K, Kajino Y, Inoue D, Ohmori T, Ueno T, Takagi T, Yoshitani J, Ueoka K, Taninaka A, Tsuchiya H. | Change in leg length after open-wedge high tibial osteotomy can be predicted from the opening width: A three-dimensional analysis | Knee. | 30 | 185-194. doi:10.1016/j.knee.2021.04.010. | 2021 |
| Ueoka K, Kabata T, Tokoro M, Kajino Y, Inoue D, Takagi T, Ohmori T, Yoshitani J, Ueno T, Yamamuro Y, Taninaka A, Tsuchiya H. | Antibacterial Activity in Iodine-coated Implants Under Conditions of Iodine Loss: Study in a Rat Model Plus In Vitro Analysis. | Clin Orthop Relat Res. | 479(7) | 1613-1623. doi:10.1097/CORR.0000000000001753. | 2021 |
| Ohmori T, Kabata T, Kajino Y, Inoue D, Kato S, Tsuchiya H. | Contralateral Lower-Limb Functional Status Before Total Hip Arthroplasty: An Important Indicator for Postoperative Gait Speed. | J Bone Joint Surg Am. | 103(12) | 1093-1103. doi:10.2106/JBJS.20.00949. | 2021 |
| Ueoka K, Kabata T, Kajino Y, Inoue D, Ohmori T, Ueno T, Yoshitani J, Yamamuro Y, Taninaka A, Kato S, Yahata T, Tsuchiya H. | The prevalence and impact of sarcopenia in females undergoing total hip arthroplasty: A prospective study. | Mod Rheumatol. | 30 | 1-6. doi:10.1080/14397595.2021.1899603. | 2021 |
| Kato S, Demura S, Shinmura K, Yokogawa N, Kabata T, Matsubara H, Kajino Y, Igarashi K, Inoue D, Kurokawa Y, Okuno N, Tsuchiya H. | Association of low back pain with muscle weakness, decreased mobility function, and malnutrition in older women: A cross-sectional study. | PLoS One. | 16(1) | e0245879. doi:10.1371/journal.pone.0245879. | 2021 |

| | | | | | |
|--|---|---------------------------|----------|--|------|
| Inoue D, Kabata T, Kajino Y, Ohmori T, Ueoka K, Tsuchiya H. | Comparison of mid-term clinical results between cementless and cemented femoral stems in total hip arthroplasty with femoral shortening osteotomy for Crowe type IV hips. | Arch Orthop Trauma Surg. | 141(6) | 1064. doi:10.1007/s00402-020-03749-0. | 2021 |
| Saiki Y, Ojima T, Kabata T, Kubo N, Hayashi S, Tsuchiya H. | Gradual exacerbation of knee flexion angle after total knee arthroplasty in patients with diabetes mellitus. | Mod Rheumatol. | 31(6) | 1215-1220. doi:10.1080/14397595.2020.1868688. | 2021 |
| Ohmori T, Kabata T, Kato S, Takagi T, Kajino Y, Inoue D, Taga T, Yamamoto T, Kurokawa Y, Yoshitani J, Ueno T, Ueoka K, Yamamuro Y, Yahata T, Tsuchiya H. | The efficacy of total hip arthroplasty on locomotive syndrome and its related physical function in patients with hip osteoarthritis. | J Orthop Sci. | 26(3) | 389-395. doi:10.1016/j.jos.2020.04.011. | 2021 |
| Ueno T, Kabata T, Kajino Y, Inoue D, Ohmori T, Yoshitani J, Ueoka K, Yamamuro Y, Tsuchiya H. | Anterior pelvic plane tilt poorly estimates the sagittal body alignment due to internal rotation of innominate bone. | J Orthop Res. | 39(3) | 589. doi:10.1002/jor.24760. | 2021 |
| Osawa Y, Seki T, Takegami Y, Makida K, Ochiai S, Imagama S. | Collapse Progression or Cessation Affects the Natural History of Contralateral Osteonecrosis of the Femoral Head | Head.J Arthrop lasty. | 36(12) | 3839-3844. doi:10.1016/j.art h.2021.08.005. | 2021 |
| Ochiai S, Seki T, Osawa Y, Kawasaki M, Yamaguchi J, Ishiguro N. | Pelvic incidence affects postoperative dislocation rate in total hip arthroplasty patients with spinal fusion. | Hip Int. | In press | doi: 10.1177/11207000211054332. | 2021 |
| Asai H, Takegami Y, Seki T, Ishiguro N. | Pelvic Tilt Reduces the Accuracy of Acetabular Component Placement When Using a Portable Navigation System: An In Vitro Study. | Arthroplast Today. | 7 | 177-181. doi:10.1016/j.artd.2020.12.012. | 2021 |
| Kataoka Y, Shimizu T, Takeda R, Tadano S, Saito Y, Osuka S, Ishida T, Samukawa M, Irie T, Takahashi D, Iwasaki N, Tohyama H. | Effects of unweighting on gait kinematics during walking on a lower-body positive-pressure treadmill in patients with hip osteoarthritis. | BMC Musculoskelet Disord. | 22(1) | 46. doi:10.1186/s12891-020-03909-8. | 2021 |
| Yokota S, Sakamoto K, Shimizu Y, Asano T, Takahashi D, Kudo K, Iwasaki N, Shimizu T. | Evaluation of whole-body modalities for diagnosis of multifocal osteonecrosis-a pilot study. | Arthritis Res Ther. | 23(1) | 3. doi:10.1186/s13075-021-02473-3. | 2021 |

| | | | | | |
|---|---|--------------------------|----------|---|------|
| Nakamura Y, Shimizu T, Asano T, Shimodan S, Ishizu H, Takahashi D, Takahata M, Iwasaki N. | Short-term efficacy and safety of zoledronate acid or denosumab in Japanese patients with postmenopausal osteoporosis. | J Bone Miner Metab | 39(5) | 824-832. doi:10.1007/s00774-021-01221-6. | 2021 |
| Takahashi D, Noyama Y, Asano T, Shimizu T, Irie T, Terkawi MA, Iwasaki N. | Finite element analysis of double-plate fixation using reversed locking compression-distal femoral plates for Vancouver B1 periprosthetic femoral fractures. | BMC Musculoskelet Disord | 22(1) | 276. doi:10.1186/s12891-021-04152-5. | 2021 |
| Ebata T, Terkawi MA, Hamasaki M, Matsumae G, Onodera T, Aly MK, Yokota S, Alhasan H, Shimizu T, Takahashi D, Homan K, Kadoya K, Iwasaki N. | Flightless I is a catabolic factor of chondrocytes that promotes hypertrophy and cartilage degeneration in osteoarthritis. | iScience. | 24(6) | 102643. doi:10.1016/j.isci.2021.102643. | 2021 |
| Matsumae G, Shimizu T, Yuan T, Takahashi D, Ebata T, Alhasan H, Yokota S, Kadoya K, Terkawi MA, Iwasaki N. | Targeting thymidine phosphorylase as a potential therapy for bone loss associated periprosthetic osteolysis | Bioeng Transl Med | 6(3) | e10232. doi:10.1002/btm2.10232. | 2021 |
| Shimizu T, Arita K, Murota E, Hiratsuka S, Fujita R, Ishizu H, Asano T, Takahashi D, Takahata M, Iwasaki N. | Effects after starting or switching from bisphosphonate to romosozumab or denosumab in Japanese postmenopausal patients. | J Bone Miner Metab. | 39(5) | 868-875. doi:10.1007/s00774-021-01226-1. | 2021 |
| Ogawa T, Shimizu T, Asano T, Iwasaki N, *Takahashi D. | Radiological predictors associated with success of treatment for developmental dysplasia of hip using the Pavlik harness: A retrospective study. | J Orthop Sci. | In press | doi:10.1016/j.jos.2021.06.010. | 2021 |
| Yokota S, Matsumae G, Shimizu T, Hasegawa T, Ebata T, Takahashi D, Heguo C, Tian Y, Alhasan H, Takahata M, Kadoya K, Terkawi MA, Iwasaki N. | Cardiotrophin Like Cytokine Factor 1 (CLCF1) alleviates bone loss in osteoporosis mouse models by suppressing osteoclast differentiation through activating interferon signaling and repressing the nuclear factor- κ B signaling pathway. | Bone. | In press | doi:10.1016/j.bone.2021.116140. | 2021 |
| Tani T, Ando W, Hamada H, Takao M, Sugano N. | Hip subluxation and osteophyte morphology are related to coronal contracture of the hip. | J Orthop Res. | 39(8) | 1691-9. doi:10.1002/jor.24891. | 2021 |

| | | | | | |
|---|---|-------------------------|--------|---|------|
| Hevesi M, Jacob G, Shimomura K, Ando W, Nakamura N, Krych AJ. | Current hip cartilage regeneration/repair modalities: a scoping review of biology and surgery | Int Orthop. | 45(2) | 319-33. doi:10.1007/s00264-020-04789-2. | 2021 |
| Kobayashi M, Chijimatsu R, Hart DA, Hamamoto S, Jacob G, Yano F, Saito T, Shimomura K, Ando W, Chung UI, Tanaka S, Yoshikawa H, Nakamura N. | Evidence that TD-198946 enhances the chondrogenic potential of human synovium-derived stem cells through the NOTCH3 signaling pathway. | J Tissue Eng Regen Med. | 15(2) | 103-15. doi:10.1002/term.3149. | 2021 |
| Shimomura K, Hamada H, Hart DA, Ando W, Nishii T, Trauttnig S, Nehrer S, Nakamura N. | Histological Analysis of Cartilage Defects Repaired with an Autologous Human Stem Cell Construct 48 Weeks Postimplantation Reveals Structural Details Not Detected by T2-Mapping MRI. | Cartilage. | 29 | 694S-706S. doi:10.1177/1947603521989423. | 2021 |
| 伊藤重治, 高窪祐弥, 平山朋幸, 門馬亮介, 高橋健大, 大木弘治, 佐々木幹, 石井政次, 高木理彰 | Pinnacle カップを用いた metal-on-metal 人工股関節の中長期成績. | Hip Joint. | 50 | 723-724 | 2021 |
| 伊藤重治, 高窪祐弥, 門馬亮介, 大木弘治, 佐々木幹, 小林真司, 石井政次, 高木理彰 | Bacteroides 感染のため複数回の人工股関節再置換術を要した 1 例 | 日本人工関節学会誌 | 50 | 723-724 | 2021 |
| 伊藤重治, 高窪祐弥, 門馬亮介, 高木理彰 | ポリエチレン摩耗に伴う骨溶解と人工股関節再置換術. | 関節外科 | 40(2). | 186-192 | 2021 |
| 名越 智 | 整形外科手術 名人の know-how 弛みのないステムを再置換するにはステムの抜去と人工股関節再置換術の実際. | 整災外 | 64 | 1356-1359 | 2021 |
| 名越 智, 鈴木大輔, 岡崎俊一郎, 舘田健児, 小助川維摩, 清水淳也, 金泉新, 佐々木幹人, 山下敏彦 | 臼蓋骨欠損 に対するセメントレスインプラントを用いた再建術の工夫-同種骨がない場合- | 人工関節学会誌 | 50 | 873-874 | 2021 |
| 小助川維摩, 舘田健児, 金泉新, 山下敏彦, 名越 智, 佐々木幹人, 清水淳也 | 患者の術前状態が人工股関節全置換術の術後評価に及ぼす影響. | 人工関節学会誌 | 50 | 905-906 | 2021 |
| 堀田和志 舘田健児 小助川維摩 鈴木大輔 名越智 山下敏彦 | 重度な寛骨臼形成不全を有する歩行可能な脳性麻痺患者に対し,骨切り術を行った 1 例. | 整災外 | 64 | 373-376 | 2021 |
| 小林千益, 菅野伸彦 | 特発性大腿骨頭壊死症(ONFH)に対する人工物置換術の最新情報.特集『大腿骨頭壊死症の基礎と臨床 Up to date』 | 関節外科 | 40(12) | 62(1296)-66(1300). | 2021 |

| | | | | | |
|-------------------------------------|--|---------------|-------|---------|------|
| 池 裕之、稲葉 裕 | 牽引台を使用する抑臥位. | 整形外科看護. | 26 | 182-183 | 2021 |
| 川端佑介、竹山昌伸、崔 賢民、林田健太、稲葉 裕 | 1.画像診断, 骨・軟部腫瘍における PET/CT 検査の有用性. II.診断, 骨・軟部腫瘍のマネジメント (その1). | 別冊整形外科. | 79 | 29-33 | 2021 |
| 島崎 貴幸、持田 勇一、針金 健吾、長岡 亜紀子、小林 直実、稲葉 裕 | 関節リウマチの外反母趾変形に対する DLMO 法後の外反母趾再発に及ぼすリスク因子の解析. | 関節外科. | 48(1) | | 2021 |
| 稲澤 真、関屋 辰洋、稲葉 裕 | 保存治療で治癒し得た化膿性椎間関節炎の1例 | 関東整形災害外科学会雑誌. | 52(3) | 117-120 | 2021 |
| 大庭真俊、水野祐介、稲葉 裕 | 1 術前の患者全身評価. 第2章 THA:手術までの準備. BHA THA 人工股関節置換術. | 羊土社 | | 38-43 | 2021 |
| 大庭真俊、稲葉 裕 | 2 術前の準備. 第2章 THA:手術までの準備. BHA THA 人工股関節置換術. | 羊土社 | | 44-47 | 2021 |
| 大庭真俊、稲葉 裕 | 3 インフォームドコンセントの実際と注意点. 第2章 THA:手術までの準備. BHA THA 人工股関節置換術. | 羊土社 | | 48-52 | 2021 |
| 池 裕之、稲葉 裕 | 4 インプラントの分類～セメントレスシステム～. 第2章 THA:手術までの準備. BHA THA 人工股関節置換術. | 羊土社 | | 52-57 | 2021 |
| 池 裕之、稲葉 裕 | 5 側方進入. §1. BHA・THA に用いられる進入法. 第3章 手術手技の実際. BHA THA 人工股関節置換術パーフェクト | 羊土社 | | 101-112 | 2021 |
| 石田 崇、稲葉 裕 | 6 前方進入. §1. BHA・THA に用いられる進入法. 第3章 手術手技の実際. BHA THA 人工股関節置換術. | 羊土社 | | 113-121 | 2021 |
| 池 裕之 稲葉 裕 | 4 Zweymuller 型ステムを用いた手術. §4. THA:ステム. 第3章 手術手技の実際. BHA THA 人工股関節置換術. | 羊土社 | | 170-176 | 2021 |
| 崔 賢民、稲葉 裕 | 3 感染予防法. 第4章合併症対策・術後管理 より安全な手術を行うために. BHA THA 人工股関節置換術. | 羊土社 | | 224-228 | 2021 |

| | | | | | |
|-----------|--|-------------|--------|---------|------|
| 手塚太郎、稲葉 裕 | 8 周術期 VTE 予防の実際と VTE への対応. 第 4 章合併症対策・術後管理 より安全な手術を行うために. BHA THA 人工股関節置換術. | 羊土社 | | 255-260 | 2021 |
| 手塚太郎、稲葉 裕 | 術前患者教育から術後療法～クリニカルパスの活用～. 第 4 章合併症対策・術後管理 より安全な手術を行うために. BHA THA 人工股関節置換術. | 羊土社 | | 266-272 | 2021 |
| 手塚太郎、稲葉 裕 | 1 術後 VTE 発症のメカニズム, 疫学など. 第 1 章 VTE (予防法, 発生時の対応). スペシャリストがすすめる人工関節手術合併症対策. | 南江堂 | | 2-4 | 2021 |
| 手塚太郎、稲葉 裕 | 2 術前検査 (エコー, 血液検査など). 第 1 章 VTE (予防法, 発生時の対応). スペシャリストがすすめる人工関節手術合併症対策. | 南江堂 | | 5-7 | 2021 |
| 手塚太郎、稲葉 裕 | 3 術中・術後管理 (抗凝固薬投与, 弾性ストッキング, フットポンプの使用). 第 1 章 VTE (予防法, 発生時の対応). スペシャリストがすすめる人工関節手術合併症対策. | 南江堂 | | 8-11 | 2021 |
| 池 裕之、稲葉 裕 | 整形外科学レビュー 2021-2022. III章 下肢 7-2 変形性股関節症. | 総合医学社 | | 150-157 | 2021 |
| 稲葉 裕 | 今日の整形外科治療指針. 23 股関節の疾患. 大腿骨頭壊死症. | 医学書院 | | 737-739 | 2021 |
| 兼氏 歩 | 成人寛骨臼形成不全股に対する骨切り術の適応と問題点 | 日本整形外科学会雑誌 | 95 | 523-534 | 2021 |
| 兼氏 歩 | 人工股関節インプラントの成績向上を目指したバイオメカニクス研究 | 金沢大学十全医学会雑誌 | 130(2) | 63-66 | 2021 |

| | | | | | |
|--|--|--------------------|--------|-----------|------|
| 高橋詠二, 兼氏 歩 | 【大腿骨頸部骨折に対する人工骨頭置換術 エビデンスに基づく手術手技と機種選択の実際】手術手技 セメント人工骨頭手術手技、注意点および機種選択の実際 整形外科 | Surgical Technique | 11(3) | 318-325 | 2021 |
| 植田修右, 市堰 徹, 兼氏 歩 | ステロイド関連骨壊死に対する病態研究の最新情報 | 関節外科 | 40(12) | 1261-1265 | 2021 |
| 坂井孝司 | 特発性大腿骨頭壊死症の診断基準と鑑別診断. | 関節外科基礎と臨床 | 40 | 32-42 | 2021 |
| 加畑多文 | Crowe group III/IV に対する人工股関節全置換術術前計画と手術の実際. | 日本整形外科学会雑誌 | 95(9) | 622-627. | 2021 |
| 西村愛世, 加畑多文, 吉谷純哉, 楫野良知, 井上大輔, 大森隆昭, 上岡顕, 山室裕紀, 谷中惇, 土屋弘行, 徳海裕史 | Short femoral nail 理想的挿入位置の 3D テンプレートニングソフトによる検討. | 日本関節病学会誌 | 40(2) | 89-93 | 2021 |
| 黒川由貴, 加藤仁志, 加畑多文, 出村諭, 松原秀憲, 八幡徹太郎, 土屋弘行 | 【ロコモティブシンドロームの現況】病態・診断ロコモ度判定と障害関節 | 整形外科 | 72(6) | 543-546 | 2021 |
| 櫻井吾郎, 吉田信也, 麦井直樹, 加畑多文, 八幡徹太郎 | 重度の両股・両膝関節変形に対し、4 ヶ月連続で人工関節置換術を施行された関節リウマチの 1 症例. | 国立大学リハビリテーション | 42 | 18-12 | 2021 |
| 井上大輔, 加畑多文, 楫野良知, 山室裕紀, 谷中惇, 片岡大往, 齊木理友, 土屋弘行 | 【インプラント周囲感染の対処法-人工関節・脊椎・骨折-】患者因子からみたインプラント周囲感染予防. | 関節外科 | 40(4) | 10-15 | 2021 |
| 井上大輔, 加畑多文, 楫野良知, 尾島朋宏, 土屋弘行. | 【低侵襲手術における感染対策マニュアル】Surgical Site Infection(SSI)の予防 術前手指消毒と術中創洗浄のエビデンスと実際. | 整形外科最小侵襲手術ジャーナル | 98 | 39-45 | 2021 |
| 加畑多文, 楫野良知, 井上大輔, 大森隆昭, 吉谷純哉, 上野琢郎, 高木知治, 上岡顕, 山室裕紀, 谷中惇, 齊木理友, 土屋弘行 | 【人工股関節再置換術-最前線から難易度の高い症例まで-】寛骨臼の骨欠損に対する再建法 金属製補填材料による再建. | 関節外科 | 40 (2) | 198-204 | 2021 |
| 関泰輔 | 大腿骨転子間弯曲内反骨切り術の適応と臨床成績. | 関節外科 | 40(12) | 49-56 | 2021 |
| 高橋大介, 清水智弘, 宮崎拓自, 中村夢志郎, 岩崎倫政 | 大腿骨頭軟骨下脆弱性骨折の特徴と治療方針. | 関節外科 | 40(12) | 43-47 | 2021 |