

<i>Demographics</i>			
Variable	Control group	HS group	<i>p</i>-value
n	880	42	
Sex (%)			
Male	473 (53.8)	25 (59.5)	<i>p</i> =0.5276
Female	407 (46.3)	15 (40.5)	
Age, median (IQR)*, y	30 (22 – 44)	28 (23.0– 39.5)	<i>p</i> =0.6047
BMI, median (IQR)*	21.6 (19.9– 23.9)	23.1 (20.4 – 25.0)	<i>p</i>=0.0398
Smoker (%)			
Yes	68 (7.8)	5 (11.5)	<i>p</i> = 0.7350
No	812 (92.2)	37 (88.5)	

有病率: 42/922 = 4.7 % (95% CI 2.4 % - 6.2%)

IQR: Interquartile range

Table 1
 欧州の診断基準で計算

<i>Demographics</i>			
Variable	Control group	HS group	<i>p</i>-value
n	915	7	
Sex (%)			
Male	495 (54.1)	3 (42.9)	<i>p</i> =0.7092
Female	420 (45.9)	4 (57.1)	
Age, median (IQR)*, y	30 (24 – 44)	27 (23– 33)	<i>p</i> =0.5173
BMI, median (IQR)*	21.7 (19.9– 23.9)	21.5 (19.7 – 23.8)	<i>p</i> =0.9309
Smoker (%)			
Yes	72 (7.9)	1 (14.2)	<i>p</i> = 0.4397
No	843 (92.1)	6 (85.7)	

有病率: 7/922 = 0.76 % (95% CI 0.22 % - 1.41%)

IQR: Interquartile range

Table 2
日本の診断基準で計算