

採用文献

(英文 49)

1. Imamura Y, et al. Effect of multidisciplinary care of dialysis initiation for outpatients with chronic kidney disease. *Int Urol Nephrol*. 2021 Jul;53(7):1435-1444.
2. Bravo-Zúñiga J, et al. Effectiveness of adherence to a renal health program in a health network in Peru. *Rev Saude Publica*. 2020;54:80. 8787.2020054002109. Epub 2020 Aug 14.
3. Helou N, et al. A Multidisciplinary Approach for Improving Quality of Life and Self-Management in Diabetic Kidney Disease: A Crossover Study. *J Clin Med*. 2020;9(7):2160.
4. Chen JH, et al. Effect of nephrology referrals and multidisciplinary care programs on renal replacement and medical costs on patients with advanced chronic kidney disease: A retrospective cohort study. *Medicine (Baltimore)*. 2019;98(33):e16808.
5. Imamura Y, et al. Usefulness of multidisciplinary care to prevent worsening renal function in chronic kidney disease. *Clin Exp Nephrol*. 2019;23(4):484-492.
6. Nicoll R, et al. Models of care for chronic kidney disease: A systematic review. *Nephrology (Carlton)*. 2018;23(5):389-396.
7. Jiamjariyapon T, et al. Effectiveness of Integrated Care on Delaying Progression of stage 3-4 Chronic Kidney Disease in Rural Communities of Thailand (ESCORT study): a cluster randomized controlled trial. *BMC Nephrol*. 2017;18(1):83.
8. Fogelfeld L, et al. Combined diabetes-renal multifactorial intervention in patients with advanced diabetic nephropathy: Proof-of-concept. *J Diabetes Complications*. 2017;31(3):624-630.
9. Cooney D, et al. A pharmacist based intervention to improve the care of patients with CKD: a pragmatic, randomized, controlled trial. *BMC Nephrol*. 2015 Apr 16;16:56.
10. Howden EJ, et al. Exercise training in CKD: efficacy, adherence, and safety. *Am J Kidney Dis*. 2015 Apr;65(4):583-91.
11. Cueto-Manzano AM, et al. Comparison of primary health-care models in the management of chronic kidney disease. *Kidney Int Suppl (2011)*. 2013 May;3(2):210-214.
12. Chen YR, et al. Effectiveness of multidisciplinary care for chronic kidney disease in Taiwan: a 3-year prospective cohort study. *Nephrol Dial Transplant*. 2013 Mar;28(3):671-82.
13. Cho EJ, et al. Effect of multidisciplinary pre-dialysis education in advanced chronic kidney disease: Propensity score matched cohort analysis. *Nephrology (Carlton)*. 2012 Jul;17(5):472-9.
14. Barrett BJ, et al. A nurse-coordinated model of care versus usual care for stage 3/4 chronic kidney disease in the community: a randomized controlled trial. *Clin J Am Soc Nephrol*. 2011 Jun;6(6):1241-7.

15. Hopkins RB, et al, Barrett BJ, Goeree R. Cost-effectiveness analysis of a randomized trial comparing care models for chronic kidney disease. *Clin J Am Soc Nephrol*. 2011 Jun;6(6):1248-57.
16. Senior PA, et al. Delivery of multifactorial interventions by nurse and dietitian teams in a community setting to prevent diabetic complications: a quality-improvement report. *Am J Kidney Dis*. 2008 Mar;51(3):425-34.
17. Yamagata K, Makino H, Iseki K, et al. Effect of Behavior Modification on Outcome in Early-to Moderate-Stage Chronic Kidney Disease : A Cluster-Randomized Trial. *PLoS One*. 2016 ; 11 : e0151422.
18. Paes-Barreto JG, et al. *J Ren Nutr* 2013 ; 23 : 164–71.
19. de Waal D, et al. *J Ren Nutr* 2016 ; 26 : 1–9.
20. Bayliss EA, Bhardwaja B, Ross C, et al. Multidisciplinary team care may slow the rate of decline in renal function. *Clin J Am Soc Nephrol* 2011 ; 6 : 704-10.
21. Slinin Y, et al. *Am J Kidney Dis* 2011 ; 58 : 583–90.
22. Kelly JT, et al. *Clin J Am Soc Nephrol* 2017 ; 12 : 272–9.
23. Jing Z, et al. *Clin Nutr* 2016 ; 35 : 117–24.
24. Chiavaroli L, et al. *Eur J Clin Nutr* 2015 ; 69 : 761–8.
25. Qin X, et al. *Atherosclerosis* 2012 ; 222 : 307–13.
26. van Zuilen AD, et al. *Kidney Int* 2012 ; 82 : 710–7.
27. Peeters MJ, et al. *J Am Soc Nephrol* 2014 ; 25 : 390–8.
28. Zhang AH, Tam P, LeBlanc D, et al. Natural history of CKD stage 4 and 5 patients following referral to renal management clinic. *Int urol nephrol*. 2009 ; 41 : 977-82.
29. Lei CC, Lee PH, Hsu YC, et al. Educational intervention in CKD retards disease progression and reduces medical costs for patients with stage 5 CKD. *Ren Fail*. 2013 ; 35 : 9-16.
30. Wu IW, Wang SY, Hsu KH, et al. Multidisciplinary predialysis education decreases the incidence of dialysis and reduces mortality--a controlled cohort study based on the NKF/DOQI guidelines. *Nephrol Dial Transplant*. 2009 ; 24 : 3426-33.
31. Fenton A, Sayar Z, Dodds A, et al. Multidisciplinary care improves outcome of patients with stage 5 chronic kidney disease. *Nephron Clin Pract*. 2010 ; 115 : c283-8.
32. Cho EJ, Park HC, Yoon HB, et al. Effect of multidisciplinary pre-dialysis education in advanced chronic kidney disease : Propensity score matched cohort analysis. *Nephrology (Carlton)* . 2012 ; 17 : 472-9.
33. Strand H, Parker D. Effects of multidisciplinary models of care for adult pre-dialysis patients with chronic kidney disease: a systematic review. *Int j Evid - Based Healthc* 2012 ; 10 : 53-9.

34. Chang AR, Evans M, Yule C, Bohn L, Young A, Lewis M, Graboski E, Gerdy B, Ehmann W, Brady J, Lawrence L, Antunes N, Green J, Snyder S, Kirchner HL, Grams M, Perkins R. Using pharmacists to improve risk stratification and management of stage 3A chronic kidney disease: a feasibility study. *BMC Nephrol*. 2016 Nov 8;17(1):168.
35. Molnar AO, Jeyakumar N, McArthur E, Battistella M, Garg AX, Sood MM, Brimble KS. Potentially inappropriate prescribing in older adults with advanced chronic kidney disease. *PLoS One*. 2020 Aug 20;15(8):e0237868.
36. Lin E, et al. Cost-effectiveness of multidisciplinary care in mild to moderate chronic kidney disease in the United States: A modeling study. *PLOS medicine* 2018.
37. Verberne WR, et al. Value-based evaluation of dialysis versus conservative care in older patients with advanced chronic kidney disease: a cohort study. *BMC Nephrology* 2018.
38. Hansen RA, Chin H, Blalock S, Joy MS. Predialysis chronic kidney disease: evaluation of quality of life in clinic patients receiving comprehensive anemia care. *Research in Social and Administrative Pharmacy* 2009.
39. Chan KY, et al. Enhanced Psychosocial Support for Caregiver Burden for Patients With Chronic Kidney Failure Choosing Not to Be Treated by Dialysis or Transplantation: A Pilot Randomized Controlled Trial. *2016 Am J Kidney Dis* 2016.
40. Zala P ,et al. Experiences of Patients with Chronic Kidney Disease and Their Family Members in an Advanced Practice Nurse-Led Counseling Service. *Nephrology Nursing Journal* 2017
41. Nakanishi M, Mizuno T, Mizokami F, Koseki T, Takahashi K, Tsuboi N, Katz M, Lee JK, Yamada S. Impact of pharmacist intervention for blood pressure control in patients with chronic kidney disease: A meta-analysis of randomized clinical trials. *J Clin Pharm Ther*. 2021 Feb;46(1):114-120.
42. Imasawa T, Saito C, Kai H, Iseki K, Kazama JJ, Shibagaki Y, Sugiyama H, Nagata D, Narita I, Nishino T, Hasegawa H, Honda H, Maruyama S, Miyazaki M, Mukoyama M, Yasuda H, Wada T, Ishikawa Y, Tsunoda R, Nagai K, Okubo R, Kondo M, Hoshino J, Yamagata K. Long-term Effectiveness of a Primary Care Practice Facilitation Program for Chronic Kidney Disease Management: An Extended Follow-up of a Cluster-Randomized FROM-J Study. *Nephrol Dial Transplant*. 2022 Feb 23; gfac041. doi: 10.1093/ndt/gfac041. Online ahead of print.
43. Tsuchida-Nishiwaki M, Uchida HA, Takeuchi H, Nishiwaki N, Maeshima Y, Saito C, Sugiyama H, Wada J, Narita I, Watanabe T, Matsuo S, Makino H, Hishida A, Yamagata K. Association of blood pressure and renal outcome in patients with chronic kidney disease; a post hoc analysis of FROM-J study. *Sci Rep*. 2021 Jul 22;11(1):14990.
44. Okubo R, Kondo M, Hoshi SL, Okada M, Doi M, Takahashi H, Kai H, Saito C, Iseki K, Iseki C, Watanabe T, Narita I, Matsuo S, Makino H, Hishida A, Yamagata K. Cost-Effectiveness

of Behavior Modification Intervention for Patients With Chronic Kidney Disease in the FROM-J Study. *J Ren Nutr*. 2021 Sep;31(5):484-493.

45. Yoshida K, Shimizu S, Kita Y, Takagi WH, Shibagaki Y, Sakurada T. Impact of inpatient educational programs on mortality after the start of dialysis therapy. *Clin Exp Nephrol*. 2022 Aug;26(8):819-826. doi: 10.1007/s10157-022-02211-2. Epub 2022 Mar 25.
46. Machida S, Shibagaki Y, Sakurada T. An inpatient educational program for chronic kidney disease. *Clin Exp Nephrol*. 2019 Apr;23(4):493-500. doi: 10.1007/s10157-018-1660-5. Epub 2018 Oct 19.
47. Takagi WH, Osako K, Machida S, Koitabashi K, Shibagaki Y, Sakurada T. Inpatient educational program delays the need for dialysis in patients with chronic kidney disease stage G5. *Clin Exp Nephrol*. 2021 Feb;25(2):166-172. doi: 10.1007/s10157-020-01979-5. Epub 2020 Oct 11.
48. Imamura Y et al. Relationship between compliance with management target values and renal prognosis in multidisciplinary care for outpatients with chronic kidney disease. *CEN* 2022, in press.
49. Shi Y, et al. The effectiveness of multidisciplinary care models for patients with chronic kidney disease: a systematic review and meta-analysis. *Int Urol Nephrol* 2018;50:301-312.

(和文 6)

1. 野口亨秀. 保存期慢性腎不全患者の低たんぱく・高カロリー食事療法は透析導入後の生存率を改善する可能性. 岐阜県医師会医学雑誌 2019.
2. 菅原寿美麗、他. 保存期腎不全患者の腎教育入院の指導の受け止めに関する研究日本看護学会論文集：慢性期看護 2019.
3. 新井英之、他. 重篤な高 K 血症のため血液透析が必要であった保存期腎不全症例の検討. 長崎医学会雑誌 2018.
4. 中澤純、他. 当院における CKD 教育入院の腎機能変化速度 ($\Delta eGFR$) に対する影響の検討. 大津市民病院雑誌 2017.
5. 内藤毅郎、他. わが国における慢性腎臓病診療と地域連携の現状と地域差－かかりつけ医を対象とした全国アンケート調査結果（第一報）－ 日本腎臓学会誌 2013.
6. 上野里紗、他. 当院における保存期腎不全検査教育入院の効果. 日本腎臓学会誌 2013.

(他の参考文献)

1. National Kidney F (2002) K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis* 39(2 Suppl 1):S1–S266.
2. Coresh J, Selvin E, Stevens LA, Manzi J, Kusek JW, Eggers P, Van Lente F, Levey AS (2007) Prevalence of chronic kidney disease in the United States. *JAMA* 298(17):2038–2047. doi:10.1001/jama.298.17.2038.
3. Go AS, Chertow GM, Fan D, McCulloch CE, Hsu CY (2004) Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N Engl J Med* 351(13):1296–1305. doi:10.1056/NEJMoa041031.
4. Yamagata K, Ishida K, Sairenchi T, Takahashi H, Ohba S, Shiigai T, Narita M, Koyama A (2007) Risk factors for chronic kidney disease in a community-based population: a 10-year follow-up study. *Kidney Int* 71(2):159–166. doi:10.1038/sj.ki.5002017.
5. Hallan SI, Coresh J, Astor BC, Asberg A, Powe NR, Romundstad S, Hallan HA, Lydersen S, Holmen J (2006) International comparison of the relationship of chronic kidney disease prevalence and ESRD risk. *J Am Soc Nephrol* 17(8):2275 – 2284. doi:10.1681/ASN.2005121273.
6. Chen YR, Yang Y, Wang SC, Chiu PF, Chou WY, Lin CY, Chang JM, Chen TW, Ferng SH, Lin CL (2013) Effectiveness of multidisciplinary care for chronic kidney disease in Taiwan: a 3-year prospective cohort study. *Nephrol Dial Transplant* 28(3):671 – 682. doi:10.1093/ndt/gfs469.
7. Vassalotti JA, Centor R, Turner BJ, Greer RC, Choi M, Sequist TD, National Kidney Foundation Kidney Disease Outcomes Quality Initiative (2016) Practical approach to detection and management of chronic kidney disease for the primary care clinician. *Am J Med* 129(2):153–162, e157. doi:10.1016/j.amjmed.2015.08.025.
8. Levey AS, Stevens LA, Coresh J (2009) Conceptual model of CKD: applications and implications. *Am J Kidney Dis* 53(3 Suppl 3):S4–S16. doi:10.1053/j.ajkd.2008.07.048.
9. Levey AS, Atkins R, Coresh J, Cohen EP, Collins AJ, Eckardt KU, Nahas ME, Jaber BL, Jadoul M, Levin A, Powe NR, Rossert J, Wheeler DC, Lameire N, Eknoyan G (2007) Chronic kidney disease as a global public health problem: approaches and initiatives—a position statement from Kidney Disease Improving Global Outcomes. *Kidney Int* 72(3):247–259. doi:10.1038/sj.ki.5002343.
10. Wang SM, Hsiao LC, Ting IW, Yu TM, Liang CC, Kuo HL, Chang CT, Liu JH, Chou CY, Huang CC (2015) Multidisciplinary care in patients with chronic kidney disease: a systematic review and meta-analysis. *Eur J Intern Med* 26(8):640 – 645. doi:10.1016/j.ejim.2015.07.002.
11. Rettig RA, Norris K, Nissenson AR (2008) Chronic kidney disease in the United States: a public policy imperative. *Clin J Am Soc Nephrol* 3(6):1902 – 1910.

doi:10.2215/CJN.02330508.

12. Ducharme A, Doyon O, White M, Rouleau JL, Brophy JM (2005) Impact of care at a multidisciplinary congestive heart failure clinic: a randomized trial. *CMAJ* 173(1):40–45. doi:10.1503/cmaj.1041137.
13. Sidhom MA, Poulsen MG (2006) Multidisciplinary care in oncology: medicolegal implications of group decisions. *Lancet Oncol* 7(11):951 – 954. doi:10.1016/S1470-2045(06)70942-1.
14. Chen YR, Yang Y, Wang SC, Chou WY, Chiu PF, Lin CY, Tsai WC, Chang JM, Chen TW, Ferng SH, Lin CL (2014) Multidisciplinary care improves clinical outcome and reduces medical costs for pre-end-stage renal disease in Taiwan. *Nephrology (Carlton)* 19(11):699–707. doi:10.1111/nep.12316.
15. Wu IW, Wang SY, Hsu KH, Lee CC, Sun CY, Tsai CJ, Wu MS (2009) Multidisciplinary predialysis education decreases the incidence of dialysis and reduces mortality — a controlled cohort study based on the NKF/DOQI guidelines. *Nephrol Dial Transplant* 24(11):3426–3433. doi:10.1093/ndt/gfp259.
16. Curtis BM, Ravani P, Malberti F, Kennett F, Taylor PA, Djurdjev O, Levin A (2005) The short- and long-term impact of multi-disciplinary clinics in addition to standard nephrology care on patient outcomes. *Nephrol Dial Transplant* 20(1):147 – 154. doi:10.1093/ndt/gfh585.
17. Fenton A, Sayar Z, Dodds A, Dasgupta I (2010) Multidisciplinary care improves outcome of patients with stage 5 chronic kidney disease. *Nephron Clin Pract* 115(4):c283–c288. doi:10.1159/000313487.
18. Wei SY, Chang YY, Mau LW, Lin MY, Chiu HC, Tsai JC, Huang CJ, Chen HC, Hwang SJ (2010) Chronic kidney disease care program improves quality of pre-end-stage renal disease care and reduces medical costs. *Nephrology (Carlton)* 15(1):108 – 115. doi:10.1111/j.1440-1797.2009.01154.x.
19. Barrett BJ, Garg AX, Goeree R, Levin A, Molzahn A, Rigatto C, Singer J, Soltys G, Soroka S, Ayers D, Parfrey PS (2011) A nurse-coordinated model of care versus usual care for stage 3/4 chronic kidney disease in the community: a randomized controlled trial. *Clin J Am Soc Nephrol* 6(6):1241–1247. doi:10.2215/CJN.07160810.
20. Rognant N, Alamartine E, Aldigier JC, Combe C, Vendrely B, Deteix P, Cluzel P, Juillard L, Vrtovsnik F, Maurice C, Fave S, Laville M (2013) Impact of prior CKD management in a renal care network on early outcomes in incident dialysis patients: a prospective observational study. *BMC Nephrol* 14:41. doi:10.1186/1471-2369-14-41.
21. Chen PM, Lai TS, Chen PY, Lai CF, Yang SY, Wu V, Chiang CK, Kao TW, Huang JW, Chiang WC, Lin SL, Hung KY, Chen YM, Chu TS, Wu MS, Wu KD, Tsai TJ (2015) Multidisciplinary care program for advanced chronic kidney disease: reduces renal

- replacement and medical costs. *Am J Med* 128(1):68 – 76. doi:10.1016/j.amjmed.2014.07.042.
22. Peeters MJ, van Zuilen AD, van den Brand JA, Bots ML, van Buren M, Ten Dam MA, Kaasjager KA, Ligtenberg G, Sijpkens YW, Sluiter HE, van de Ven PJ, Vervoort G, Vleming LJ, Blankestijn PJ, Wetzels JF (2014) Nurse practitioner care improves renal outcome in patients with CKD. *J Am Soc Nephrol* 25(2):390–398. doi:10.1681/ASN.201212122222.
 23. Chan JC, So WY, Yeung CY, Ko GT, Lau IT, Tsang MW, Lau KP, Siu SC, Li JK, Yeung VT, Leung WY, Tong PC, SURE Study Group (2009) Effects of structured versus usual care on renal endpoint in type 2 diabetes: the SURE study: a randomized multicenter translational study. *Diabetes Care* 32(6):977–982. doi:10.2337/dc08-1908.
 24. Jessup RL (2007) Interdisciplinary versus multidisciplinary care teams: do we understand the difference? *Aust Health Rev* 31(3):330–331.
 25. Ethier J, Mendelssohn DC, Elder SJ, Hasegawa T, Akizawa T, Akiba T, Canaud BJ, Pisoni RL (2008) Vascular access use and outcomes: an international perspective from the Dialysis Outcomes and Practice Patterns Study. *Nephrol Dial Transplant* 23(10):3219–3226. doi:10.1093/ndt/gfn261.
 26. Xue JL, Dahl D, Ebben JP, Collins AJ (2003) The association of initial hemodialysis access type with mortality outcomes in elderly Medicare ESRD patients. *Am J Kidney Dis* 42(5):1013–1019.
 27. Liem YS, Weimar W (2009) Early living-donor kidney transplantation: a review of the associated survival benefit. *Transplantation* 87(3):317–318. doi:10.1097/TP.0b013e3181952710.
 28. Cho EJ, Park HC, Yoon HB, Ju KD, Kim H, Oh YK, Yang J, Hwang YH, Ahn C, Oh KH (2012) Effect of multidisciplinary pre-dialysis education in advanced chronic kidney disease: Propensity score matched cohort analysis. *Nephrology (Carlton)* 17(5):472–479. doi:10.1111/j.1440-1797.2012.01598.x.
 29. Goovaerts T, Jadoul M, Goffin E (2005) Influence of a pre-dialysis education programme (PDEP) on the mode of renal replacement therapy. *Nephrol Dial Transplant* 20(9):1842–1847. doi:10.1093/ndt/gfh905.
 30. Andrassy KM (2013) Comments on ‘KDIGO 2012 clinical practice guideline for the evaluation and management of chronic kidney disease’. *Kidney Int* 84(3):622–623. doi:10.1038/ki.2013.243.
 31. Yu YJ, Wu IW, Huang CY, Hsu KH, Lee CC, Sun CY, Hsu HJ, Wu MS (2014) Multidisciplinary predialysis education reduced the inpatient and total medical costs of the first 6 months of dialysis in incident hemodialysis patients. *PLoS ONE* 9(11):e112820. doi:10.1371/journal.pone.0112820.
 32. Goldstein M, Yassa T, Dacouris N, McFarlane P (2004) Multidisciplinary predialysis care

- and morbidity and mortality of patients on dialysis. *Am J Kidney Dis* 44(4):706–714.
33. Hemmelgarn BR, Manns BJ, Zhang J, Tonelli M, Klarenbach S, Walsh M, Culleton BF (2007) Association between multidisciplinary care and survival for elderly patients with chronic kidney disease. *J Am Soc Nephrol* 18(3):993–999. doi:10.1681/ASN.2006080860.
 34. Yeoh HH, Tiquia HS, Abcar AC, Rasgon SA, Idroos ML, Daneshvari SF (2003) Impact of predialysis care on clinical outcomes. *Hemodial Int* 7(4):338–341. doi:10.1046/j.1492-7535.2003.00059.x.
 35. Harris LE, Luft FC, Rudy DW, Kesterson JG, Tierney WM (1998) Effects of multidisciplinary case management in patients with chronic renal insufficiency. *Am J Med* 105(6):464–471.
 36. Devins GM, Mendelssohn DC, Barre PE, Binik YM (2003) Predialysis psychoeducational intervention and coping styles influence time to dialysis in chronic kidney disease. *Am J Kidney Dis* 42(4):693–703.
 37. Levin A, Lewis M, Mortiboy P, Faber S, Hare I, Porter EC, Mendelssohn DC (1997) Multidisciplinary predialysis programs: quantification and limitations of their impact on patient outcomes in two Canadian settings. *Am J Kidney Dis* 29(4):533–540.
 38. Bayliss EA, Bhardwaja B, Ross C, Beck A, Lanese DM (2011) Multidisciplinary team care may slow the rate of decline in renal function. *Clin J Am Soc Nephrol* 6(4):704–710. doi:10.2215/CJN.06610810.
 39. Xiao-yu Z, Xin-hong L, Feng Z (2013) The effects of predialysis CKD management on hemodialysis patients. *Chin Nurs Manag* 13(7):75–78. doi:10.3969/j.issn.1672-1756.2013.07.026.