

発表論文

1. Kawashima M, Hisamatsu T, Harada A, Kadota A, Kondo K, Okami Y, Hayakawa T, Kita Y, Okayama A, Ueshima H, Okamura T, Miura K; NIPPON DATA90 Research Group. Relationship Between Hemoglobin Concentration and Cardiovascular Disease Mortality in a 25-Year Follow-up Study of a Japanese General Population - NIPPON DATA90. *Circ J.* 2024 Apr 25;88(5):742-750. doi: 10.1253/circj.CJ-23-0725.
2. Kubo K, Hirata A, Kadota A, Harada A, Nakamura Y, Hayakawa T, Takashima N, Fujiyoshi A, Okami Y, Kita Y, Okayama A, Miura K, Ueshima H, Okamura T; NIPPON DATA90 Research Group. Risk Factors for Heart Failure and Coronary Artery Disease Mortality Based on the National Vital Statistics During a 25-Year Follow-up in Japan - NIPPON DATA90. *Circ J.* 2024 Aug 23;88(9):1478-1487. doi: 10.1253/circj.CJ-23-0847.
3. Hirata T, Hirata A, Torii S, Takashima N, Kadota Aya, Choudhury SR, Okayama A, Miura K, Okamura T. NIPPON DATA90 Study Group. Association between alcohol intake and death from cardiovascular diseases and its subtypes stratified by dyslipidemia in Japanese men: 20-years follow-up of NIPPON DATA90. *Environ Health Prev Med.* 2024;29:61. doi: 10.1265/ehpm.24-00164.
4. Hisamatsu T, Kogure M, Tabara Y, Hozawa A, Sakima A, Tsuchihashi T, Yoshita K, Hayabuchi H, Node K, Takemi Y, Ohkubo T, Miura K; Japanese Society of Hypertension Working Group on Urine Sodium-to-Potassium Ratio. Practical use and target value of urine sodium-to-potassium ratio in assessment of hypertension risk for Japanese: Consensus Statement by the Japanese Society of Hypertension Working Group on Urine Sodium-to-Potassium Ratio. *Hypertens Res.* 2024 Dec;47(12):3288-3302. doi: 10.1038/s41440-024-01861-x.
5. Tsukinoki R, Murakami Y, Hayakawa T, Kadota A, Harada A, Kita Y, Okayama A, Miura K, Okamura T, Ueshima H. Comprehensive assessment of the impact of blood pressure, body mass index, smoking, and diabetes on healthy life expectancy in Japan: NIPPON DATA90. *J Epidemiol.* 2025 Jan 11. doi: 10.2188/jea.JE20240298. Online ahead of print.